

Ymir

Contents

Topology and bornology	5
1. Introduction	5
2. Nets	5
3. Paracompact spaces	7
4. Bornology	7
Bibliography	9

Topology and bornology

1. Introduction

In the whole project, a neighbourhood in a topology space is taken in Bourbaki's sense. In particular, a neighbourhood is not necessarily open.

We follow Bourbaki's convention about compact space. A compact space is always Hausdorff.

On the other hand, we do not require locally compact spaces and paracompact spaces be Hausdorff.

References to this chapter include [\[Ber93\]](#).

2. Nets

Let X be a set, $Y \subseteq X$ be a subset. Consider a collection τ of subsets of X , we write

$$\tau|_Y := \{V \in \tau : V \subseteq Y\}.$$

Definition 2.1. Let X be a topology space and τ be a collection of subsets of X . We say τ is

- (1) *dense* if for any $V \in \tau$ and any $x \in V$, there is a fundamental system of neighbourhoods of x in V consisting of sets from $\tau|_V$;
- (2) a *quasi-net* on X if for each $x \in X$, there exist $n \in \mathbb{Z}_{>0}$, $V_1, \dots, V_n \in \tau$ such that $x \in V_1 \cap \dots \cap V_n$ and that $V_1 \cup \dots \cup V_n$ is a neighbourhood of x in X ;
- (3) a *net* on X if it is a quasi-net and if for any $U, V \in \tau$, $\tau|_{U \cap V}$ is a quasi-net on $U \cap V$;
- (4) *locally finite* if for any $x \in X$, there is a neighbourhood U of x in X such that $\{V \in \tau : V \cap U \neq \emptyset\}$ is finite.

We observe that if τ is a net, $\tau|_{U \cap V}$ is in fact a net.

Lemma 2.2. Let X be a topological space and τ be a quasi-net on X .

- (1) A subset $U \subseteq X$ is open if and only if for each $V \in \tau$, $U \cap V$ is open in V .
- (2) Suppose that τ consists of compact sets. Then X is Hausdorff if and only if for any $U, V \in \tau$, $U \cap V$ is compact.

We remind the readers that a compact space is Hausdorff by our convention.

PROOF. (1) The direct implication is trivial. Suppose that $U \cap V$ is open in V for all $V \in \tau$. We want to show that U is open. Take $x \in U$, we can find $n \in \mathbb{Z}_{>0}$, $V_1, \dots, V_n \in \tau$ all containing x such that $V_1 \cup \dots \cup V_n$ is a neighbourhood of x in X . By our hypothesis, we can find open sets W_1, \dots, W_n in W such that $W \cap V_i = U \cap V_i$

for $i = 1, \dots, n$. Then $W = W_1 \cap \dots \cap W_n$ is an open neighbourhood of x in X . But then

$$U \cap (V_1 \cup \dots \cup V_n) \supseteq W \cap (V_1 \cup \dots \cup V_n),$$

the latter is a neighbourhood of x hence so is the former. It follows that U is open.

(2) The direct implication is trivial. Consider the quasi-net $\tau \times \tau := \{U \times V : U, V \in \tau\}$ on $X \times X$. By (1), it suffices to verify that the intersection of the diagonal with $U \times V$ is closed in $U \times V$ for any $U, V \in \tau$. But this intersection is homeomorphic to $U \cap V$, which is compact by our assumption and hence closed as U, V are both Hausdorff. \square

Lemma 2.3. Let X be a Hausdorff space. Assume that X admits a quasi-net τ consisting of compact sets. Then X is locally compact.

PROOF. Take $x \in X$. By assumption, we can find $n \in \mathbb{N}$ and $V_1, \dots, V_n \in \tau$ all containing x such that $V_1 \cup \dots \cup V_n$ is a neighbourhood of x . This neighbourhood is clearly compact. \square

Lemma 2.4. Let X be a Hausdorff space and τ be a collection of compact subsets of X . Then the following are equivalent:

- (1) τ is a quasi-net;
- (2) For each $x \in X$, there are $n \in \mathbb{N}$ and $V_1, \dots, V_n \in \tau$ such that $V_1 \cup \dots \cup V_n$ is a neighbourhood of x in X .

PROOF. (1) \implies (2): This is trivial.

(2) \implies (1): Given $x \in X$, take V_1, \dots, V_n as in (2). We may assume that $x \in V_1, \dots, V_m$ and $x \notin V_{m+1}, \dots, V_n$ for some $1 \leq m \leq n$. Then $V_1 \cup \dots \cup V_m$ is a neighbourhood of x in X : if U is an open neighbourhood of x in X contained in $V_1 \cup \dots \cup V_n$, then $U \setminus (V_{m+1} \cup \dots \cup V_n)$ is an open neighbourhood of x in X contained in $V_1 \cup \dots \cup V_m$. \square

Lemma 2.5. Let X be a topological space and τ be a net on X consisting of compact sets. Then

- (1) for any pair $U, V \in \tau$, the intersection $U \cap V$ is locally closed in U and in V ;
- (2) If $n \in \mathbb{Z}_{>0}$, $V, V_1, \dots, V_n \in \tau$ are such that

$$V \subseteq V_1 \cup \dots \cup V_n,$$

then there are $m \in \mathbb{Z}_{>0}$ and $U_1, \dots, U_m \in \tau$ such that

$$V = U_1 \cup \dots \cup U_m$$

and each U_j is contained in some V_i .

PROOF. (1) It suffices to show that $U \cap V$ is locally compact in the induced topology. This follows from [Lemma 2.3](#).

(2) For each $x \in V$ and each $i = 1, \dots, n$ such that $x \in V_i$, we take a neighbourhood of x in $V \cap V_i$ of the form $W_i V_{i1} \cup \dots \cup V_{im_i}$ for some $m_i \in \mathbb{Z}_{>0}$ and $V_{ij} \in \tau$ for $j = 1, \dots, m_i$. Then the union of all W_i 's is a neighbourhood of x of the form $U_1 \cup \dots \cup U_m$, where U_j belongs to τ and is contained in some V_i . Using the compactness of V , we conclude. \square

3. Paracompact spaces

Definition 3.1. A topological space X is *paracompact* if any open covering of X admits a locally finite refinement.

A paracompact space is not necessarily Hausdorff according to our definition.

Proposition 3.2. Let X be a locally compact topological space.

- (1) Assume that each connected component of X is σ -compact, then X is paracompact.
- (2) If X is paracompact and Hausdorff, then each connected component of X is σ -compact.

If the conditions in (2) are satisfied, for any basis of neighbourhoods \mathcal{B} of X , every open covering \mathcal{U} of X can be refined into a locally finite covering \mathcal{V} consisting of elements in \mathcal{B} .

We do not assume that the elements in \mathcal{B} be open. The covering \mathcal{V} is not necessarily open.

Proposition 3.3. Let X be a paracompact space and $Y \subseteq X$ be a closed subspace. Then Y is paracompact.

Proposition 3.4. Let X be a locally compact Hausdorff space and $Y \subseteq X$ be a subspace, then the following are equivalent:

- (1) Y is locally compact and Hausdorff;
- (2) Y is a locally closed subspace of X .

4. Bornology

Definition 4.1. Let X be a set. A *bornology* on X is a collection \mathcal{B} of subsets of X such that

- (1) For any $x \in X$, there is $B \in \mathcal{B}$ such that $x \in B$;
- (2) For any $B \in \mathcal{B}$ and any subset $A \subseteq B$, $A \in \mathcal{B}$;
- (3) \mathcal{B} is stable under finite union.

The pair (X, \mathcal{B}) is called a *bornological set*. The elements of \mathcal{B} are called the *bounded subsets* of (X, \mathcal{B}) . When \mathcal{B} is obvious from the context, we omit it from the notations.

A morphism between bornological sets (X, \mathcal{B}_X) and (Y, \mathcal{B}_Y) is a map of sets $f : X \rightarrow Y$ such that for any $A \in \mathcal{B}_X$, $f(A) \in \mathcal{B}_Y$. Such a map is called a *bounded map*.

Definition 4.2. Let (X, \mathcal{B}) be a bornological set. A *basis* for \mathcal{B} is a subset $\mathcal{A} \subseteq \mathcal{B}$ such that for any $B \in \mathcal{B}$, there are $A_1, \dots, A_n \in \mathcal{A}$ such that $B \subseteq A_1 \cup \dots \cup A_n$.

Bibliography

- [Ber93] V. G. Berkovich. Étale cohomology for non-Archimedean analytic spaces.
Publications Mathématiques de l'Institut des Hautes Études Scientifiques
78.1 (1993), pp. 5–161.