

Ymir

Contents

Berkovich analytic spaces	5
1. Introduction	5
2. The category of Berkovich analytic spaces	5
3. Analytic domains	16
4. Berkovich site	19
5. Closed immersions	23
6. Separated morphisms	25
7. Analytic germs	26
8. Reduction	28
9. Some results whose proofs I do not understand	36
Bibliography	41

Berkovich analytic spaces

1. Introduction

The main references of this chapter: [Ber93], [Ber12], [Tem04], [Tem00], [Duc18].

2. The category of Berkovich analytic spaces

Let $(k, |\bullet|)$ be a complete non-Archimedean valued field and H be a subgroup of $\mathbb{R}_{>0}$ such that $|k^\times| \cdot H \neq \{1\}$.

Definition 2.1. Let X be a locally Hausdorff space and τ be a net of compact subsets. A k_H -affinoid atlas \mathcal{A} on X with the net τ is a map which assigns

- (1) to each $V \in \tau$, a k_H -affinoid algebra A_V and a homeomorphism $\varphi_V : \mathrm{Sp} A_V \rightarrow V$;
- (2) to each $U, V \in \tau$, $U \subseteq V$, a morphism of k_H -affinoid algebras $\alpha_{V/U} : A_V \rightarrow A_U$ representing a k_H -affinoid domain $\mathrm{Sp} A_U$ in $\mathrm{Sp} A_V$ such that the following diagram commutes

$$\begin{array}{ccc} \mathrm{Sp} A_U & \xrightarrow{\mathrm{Sp} \alpha_{V/U}} & \mathrm{Sp} A_V \\ \downarrow \varphi_U & & \downarrow \varphi_V \\ U & \longrightarrow & V \end{array} .$$

The triple (X, \mathcal{A}, τ) as above is called a k_H -analytic space.

A *morphism* between atlases \mathcal{A} and \mathcal{A}' on X with the net τ is an assignment that with each $V \in \tau$, one associates a morphism of k_H -affinoid algebras $\beta_V : A_V \rightarrow A'_V$ such that

- (1) for each $V \in \tau$, the following diagram is commutative:

$$\begin{array}{ccc} \mathrm{Sp} A'_V & \xrightarrow{\mathrm{Sp} \beta_V} & \mathrm{Sp} A_V \\ \downarrow \varphi'_V & \swarrow \varphi_V & \\ V & & \end{array} ;$$

- (2) for each $U, V \in \tau$, $U \subseteq V$, the following diagram is commutative:

$$\begin{array}{ccc} A_V & \xrightarrow{\alpha_{V/U}} & A_U \\ \downarrow \beta_V & & \downarrow \beta_U \\ A'_V & \xrightarrow{\alpha'_{V/U}} & A'_U \end{array}$$

Here we have denoted the data associated with \mathcal{A}' with a prime. In this way, the atlases on X with the net τ form a category.

We remind the readers that by our convention a compact space is Hausdorff.

By Condition (2), if $W \subseteq U \subseteq V$ are three sets in τ , then $\alpha_{V/U} \circ \alpha_{U/W} = \alpha_{V/W}$.

Remark 2.2. As a convention, we will denote the atlas by capital letters in caligraphic font and the affinoid algebras by the same letter in roman font. We will usually omit the maps φ_U 's by identifying $\mathrm{Sp} A_U$ with U . We will say U is a k_H -affinoid domain in V .

Remark 2.3. Our definition is a special case of the original definitions in [Ber93]. This seems to be the most important case though.

Lemma 2.4. Let (X, \mathcal{A}, τ) be a k_H -analytic space, $U \in \tau$ and W is a k_H -affinoid domain in U . Then for any $V \in \tau$ containing W , W is a k_H -affinoid domain in V .

PROOF. As $\tau|_{U \cap V}$ is a net and W is compact, we can find $U_1, \dots, U_n \in \tau|_{U \cap V}$ with $W \subseteq U_1 \cup \dots \cup U_n$. As W, U_i are k_H -affinoid domains in U , $W_i = W \cap U_i$ is a k_H -affinoid domain in U_i for all $i = 1, \dots, n$ by Corollary 13.12 in Affinoid algebras. It follows from Corollary 10.7 and Corollary 13.12 in Affinoid algebras that W_i and $W_i \cap W_j$ are both k_H -affinoid domains in V for $i, j = 1, \dots, n$. So W is a compact k_H -analytic domain in V .

By Proposition 13.25 in Affinoid algebras,

$$A_W := \ker \left(\prod_{i=1}^n A_{W_i} \rightarrow \prod_{i,j=1}^n A_{W_i \cap W_j} \right)$$

is k_H -affinoid and $\mathrm{Sp} A_W \rightarrow \mathrm{Sp} A$ induces a homeomorphism $\mathrm{Sp} A_W \rightarrow W$ by Proposition 10.6 in Affinoid algebras. By Proposition 13.25 in Affinoid algebras again, W is affinoid in V . \square

Definition 2.5. Let (X, \mathcal{A}, τ) be a k_H -analytic space. We define $\bar{\tau}$ as the set of all $W \subseteq X$ such that there is $U \in \tau$ containing W and W is k_H -affinoid in U .

Lemma 2.6. Let (X, \mathcal{A}, τ) be a k_H -analytic space. Then $\bar{\tau}$ is a net on X and there is a k_H -affinoid atlas $\bar{\mathcal{A}}$ on X with the net $\bar{\tau}$ extending \mathcal{A} . Moreover, the k_H -affinoid atlas $\bar{\mathcal{A}}$ on X with the net $\bar{\tau}$ extending \mathcal{A} is unique up to a canonical isomorphism.

PROOF. **Step 1.** We first show that $\bar{\tau}$ is a net. Let $U, V \in \bar{\tau}$ and $x \in U \cap V$. Take $U', V' \in \tau$ containing U and V respectively. Take $n \in \mathbb{Z}_{>0}$ and $W_1, \dots, W_n \in \tau$ such that

- (1) $x \in W_1 \cap \dots \cap W_n$;
- (2) $W_1 \cup \dots \cup W_n$ is a neighbourhood of x in $U' \cap V'$.

This is possible because $\tau|_{U' \cap V'}$ is a quasi-net by assumption.

By Lemma 2.4, U (resp. V) and W_1, \dots, W_n are k_H -affinoid domains in U' (resp. V').

By Corollary 13.12 in Affinoid algebras, $U_i := U \cap W_i$ (resp. $V_i := V \cap W_i$) is a k_H -affinoid domain in W_i for $i = 1, \dots, n$. By Corollary 13.12 in Affinoid algebras again, $U_i \cap V_i$ is a k_H -affinoid domain in W_i for $i = 1, \dots, n$. So $U_i \cap V_i \in \bar{\tau}|_{U \cap V}$ for $i = 1, \dots, n$. But

$$\bigcup_{i=1}^n U_i \cap V_i = (U \cap V) \cap \bigcup_{i=1}^n W_i,$$

so $\bigcup_{i=1}^n U_i \cap V_i$ is a neighbourhood of x in $U \cap V$ and $x \in \bigcap_{i=1}^n U_i \cap V_i$. It follows that $\bar{\tau}$ is a net.

Step 2. We extend the k_H -affinoid atlas \mathcal{A} .

For each $V \in \bar{\tau}$, we fix a $V' \in \tau$ containing V .

By [Lemma 2.4](#), V is a k_H -affinoid domain in V' . Let $A_{V'} \rightarrow A_V$ be the morphism of k_H -affinoid algebras representing the k_H -affinoid domain V in $\mathrm{Sp} A_{V'}$. We define the homeomorphism $\varphi_V : \mathrm{Sp} A_V \rightarrow V$ as the morphism induced by $\mathrm{Sp} A_V \rightarrow \mathrm{Sp} A$.

For $U, V \in \bar{\tau}$ with $U \subseteq V$, we want to define $\alpha_{V/U} : A_V \rightarrow A_U$. We handle two cases. When $V \in \tau$, as $\tau|_{U' \cap V}$ is a quasi-net, we can find $n \in \mathbb{Z}_{>0}$ and $U_1, \dots, U_n \in \tau|_{U' \cap V}$ such that

$$U = \bigcup_{i=1}^n U_i.$$

By [Lemma 2.4](#), U_1, \dots, U_n are k_H -affinoid domains in U' and in V . By [Theorem 13.19](#) in [Affinoid algebras](#),

$$A_U \xrightarrow{\sim} \ker \left(\prod_{i=1}^n A_{U_i} \rightarrow \prod_{i,j=1}^n A_{U_i \cap U_j} \right).$$

So the morphism $\alpha_{V/U_i} : A_V \rightarrow A_{U_i}$ and $\alpha_{V/U_i \cap U_j} : A_V \rightarrow A_{U_i \cap U_j}$ for $i = 1, \dots, n$ and $j = 1, \dots, n$ induces a morphism $\alpha_{V/U} : A_V \rightarrow A_U$. Observe that $\alpha_{V/U}$ represents the k_H -affinoid domain U in V , so it is independent of the choice of U_1, \dots, U_n .

More generally, when $V \in \bar{\tau}$, we have constructed a morphism $\alpha_{V'/U} : A_{V'} \rightarrow A_U$ representing the k_H -affinoid domain U in V' , it follows that U is a k_H -affinoid domain in V , and we therefore get the desired morphism $\alpha_{V/U} : A_V \rightarrow A_U$.

It is easy to verify that the constructions gives a k_H -affinoid atlas with the net $\bar{\tau}$ extending \mathcal{A} . The uniqueness of the extension is immediate. \square

Definition 2.7. Let (X, \mathcal{A}, τ) and $(X', \mathcal{A}', \tau')$ be k_H -analytic spaces. A *strong morphism* $\varphi : (X, \mathcal{A}, \tau) \rightarrow (X', \mathcal{A}', \tau')$ is a pair consisting of

- (1) a continuous map $\varphi : X \rightarrow X'$ such that for each $V \in \tau$, there is $V' \in \tau'$ with $\varphi(V) \subseteq V'$;
- (2) for each $V \in \tau$, $V' \in \tau'$ with $\varphi(V) \subseteq V'$, a morphism of k_H -affinoid spectra $\varphi_{V/V'} : V \rightarrow V'$

such that for each $V, W \in \tau$, $V', W' \in \tau'$ satisfying $V \subseteq W$, $W' \subseteq V'$, $\varphi(V) \subseteq V'$ and $\varphi(W) \subseteq W'$, the following diagram commutes:

$$\begin{array}{ccc} V & \xrightarrow{\varphi_{V/V'}} & V' \\ \downarrow & & \downarrow \\ W & \xrightarrow{\varphi_{W/W'}} & W' \end{array}.$$

Recall our convention [Remark 2.2](#), the morphism $\varphi_{V/V'}$ means a morphism $A'_{V'} \rightarrow A_V$ of k_H -affinoid algebras making the following diagram commutative

$$\begin{array}{ccc} \mathrm{Sp} A_V & \longrightarrow & \mathrm{Sp} A'_{V'} \\ \downarrow \varphi_V & & \downarrow \varphi'_{V'} \\ V & \xrightarrow{\varphi} & V' \end{array}.$$

We will continue our identifications as in [Remark 2.2](#) to simplify our notations.

Proposition 2.8. Let (X, \mathcal{A}, τ) and $(X', \mathcal{A}', \tau')$ be k_H -analytic spaces. Let $\varphi : (X, \mathcal{A}, \tau) \rightarrow (X', \mathcal{A}', \tau')$ be a strong morphism. Then φ extends uniquely to a strong morphism $\varphi : (X, \bar{\mathcal{A}}, \bar{\tau}) \rightarrow (X', \bar{\mathcal{A}}', \bar{\tau}')$.

PROOF. Let $U \in \bar{\tau}$, $U' \in \bar{\tau}'$ with $\varphi(U) \subseteq U'$. Take $V \in \tau$ and $V' \in \tau'$ containing U and U' respectively. By [Lemma 2.4](#), U (resp. V) is a k_H -affinoid domain in V (resp. V'). Take $W \in \tau'$ with $\varphi(V) \subseteq W'$. Then in particular, $\varphi(U) \subseteq W'$. As $\tau'|_{V' \cap W'}$ is a quasi-net and $\varphi(U)$ is compact, we can find $n \in \mathbb{Z}_{>0}$ and $W_1, \dots, W_n \in \tau'|_{V' \cap W'}$ such that

$$\varphi(U) \subseteq W_1 \cup \dots \cup W_n.$$

Now W_i is a k_H -affinoid domain in W' by [Lemma 2.4](#), so $V_i := \varphi_{V/W'}^{-1}(W_i)$ is an affinoid domain in V by [Corollary 13.12](#) in [Affinoid algebras](#), and we have an induced morphism $V_i \rightarrow W_i$ for $i = 1, \dots, n$. This morphism in turn induces a morphism of k_H -affinoid spectra

$$U_i := U \cap V_i \rightarrow U'_i := U' \cap W_i \rightarrow U'$$

for $i = 1, \dots, n$. These morphisms are compatible on their intersections by construction. So by [Theorem 13.19](#) in [Affinoid algebras](#), they glue together to a morphism of k_H -affinoid spectra $\bar{\varphi}_{U/U'} : U \rightarrow U'$. It is easy to see that this construction defines a strong morphism.

As for the uniqueness, it suffices to show that the morphism $U_i \rightarrow U'_i$ is uniquely determined for $i = 1, \dots, n$. In other words, we need to show that the dotted arrow that makes the following diagram commutes is unique:

$$\begin{array}{ccc} U_i & \cdots \cdots \cdots & U'_i \\ \downarrow & & \downarrow \\ V & \xrightarrow{\varphi_{V/W'}} & W' \end{array}$$

for $i = 1, \dots, n$. It suffices to apply the universal property of the k_H -affinoid domain $U'_i \rightarrow W'$. \square

Definition 2.9. Let (X, \mathcal{A}, τ) , $(X', \mathcal{A}', \tau')$, $(X'', \mathcal{A}'', \tau'')$ be k_H -analytic spaces. Let

$$\varphi : (X, \mathcal{A}, \tau) \rightarrow (X', \mathcal{A}', \tau'), \quad \psi : (X', \mathcal{A}', \tau') \rightarrow (X'', \mathcal{A}'', \tau'')$$

be strong morphisms. We will define their *composition* $\chi = \psi \circ \varphi$ as follows. The underlying map of topological spaces is just the composition of the underlying maps of topological spaces corresponding to ψ and φ .

Let $\bar{\varphi}$ and $\bar{\psi}$ be the extensions of φ and ψ to $\bar{\tau}$ and $\bar{\tau}'$ as in [Proposition 2.8](#).

Given $V \in \tau$ and $V'' \in \tau''$ with $\chi(V) \subseteq V''$, we need to define a morphism of k_H -affinoid spectra $\chi_{V/V''} : V \rightarrow V''$. Take $V' \in \tau'$ and $U'' \in \tau''$ such that $\varphi(V) \subseteq V'$ and $\psi(V') \subseteq U''$. Since $\chi(V) \subseteq U'' \cap V''$ and V is compact, we can take $n \in \mathbb{Z}_{>0}$ and $V''_1, \dots, V''_n \in \tau''|_{U'' \cap V''}$ with $\chi(V) \subseteq V''_1 \cup \dots \cup V''_n$. Then $V'_i := \psi_{V'/U''}^{-1}(V''_i)$ and $V_i := \varphi_{V/V'}^{-1}(V'_i)$ are k_H -affinoid domains in V' and V respectively for $i = 1, \dots, n$ and $V = V_1 \cup \dots \cup V_n$. The morphisms $\bar{\varphi}$ and $\bar{\psi}$ then induce a morphism $V_i \rightarrow V''_i \rightarrow V''$ of k_H -affinoid spectra. These morphisms are clearly compatible on the intersections and hence induce a morphism $V \rightarrow V''$ of k_H -affinoid spectra by [Theorem 13.19](#) in [Affinoid algebras](#).

It is easy to verify that $\psi \circ \varphi$ is a strong morphism.

In this way, we get a category $k_H\text{-}\widetilde{\mathcal{A}n}$ of k_H -analytic spaces.

Definition 2.10. Let (X, \mathcal{A}, τ) and $(X', \mathcal{A}', \tau')$ be k_H -analytic spaces. A strong morphism $\varphi : (X, \mathcal{A}, \tau) \rightarrow (X', \mathcal{A}', \tau')$ is said to be a *quasi-isomorphism* if

- (1) φ is a homeomorphism between X and X' ;
- (2) for any pair $V \in \tau$ and $V' \in \tau'$ with $\varphi(V) \subseteq V'$, $\mathrm{Sp} \varphi_{V/V'}$ identifies V with an affinoid domain in V' .

Lemma 2.11. Let (X, \mathcal{A}, τ) and $(X', \mathcal{A}', \tau')$ be k_H -analytic spaces and $\varphi : (X, \mathcal{A}, \tau) \rightarrow (X', \mathcal{A}', \tau')$ be a strong morphism. Then for any $V \in \tau$ and $V' \in \tau'$, the intersection $V \cap \varphi^{-1}(V')$ is a compact k_H -analytic domain in V .

PROOF. Take $U' \in \tau'$ with $\varphi(V) \subseteq U'$. As $\tau|_{U' \cap V'}$ is a quasi-net, we can find $n \in \mathbb{Z}_{>0}$ and $U'_1, \dots, U'_n \in \tau|_{U' \cap V'}$ with $\varphi(V) \subseteq U'_1 \cup \dots \cup U'_n$ and

$$V \cap \varphi^{-1}(V') = \bigcup_{i=1}^n \varphi_{V/U}^{-1}(U'_i).$$

□

Lemma 2.12. The system of quasi-isomorphisms in $k_H\text{-}\widetilde{\mathcal{A}n}$ is a right multiplicative system.

For the notion of right multiplicative system, we refer to [Stacks, Tag 04VC].

PROOF. We verify the three axioms as in [Stacks, Tag 04VC].

RMS1. The identity is clear a quasi-isomorphism. It remains to verify that the composition of quasi-isomorphisms is still a quasi-isomorphism.

We take φ, ψ as in Definition 2.9. We will use the same notations as in Definition 2.9. We need to show that $V \rightarrow V''$ identifies V with a k_H -affinoid domain in V'' . From the construction, we know that φ identifies V_i with a k_H -affinoid domain in V'_i and ψ identifies V'_i with a k_H -affinoid domain in V''_i for $i = 1, \dots, n$. In particular, $\chi(V)$ is a compact k_H -analytic domain in V'' . It follows from Proposition 13.25 in Affinoid algebras that $\chi(V)$ is a k_H -affinoid domain in V'' .

RMS2. If $\varphi : (X, \mathcal{A}, \tau) \rightarrow (X', \mathcal{A}', \tau')$ and $f : (\widetilde{X}', \widetilde{\mathcal{A}}', \widetilde{\tau}') \rightarrow (X', \mathcal{A}', \tau')$ are given strong morphisms of k_H -analytic spaces and g is a quasi-isomorphism, then there are k_H -analytic space $(\widetilde{X}, \widetilde{\mathcal{A}}, \widetilde{\tau})$ and strong morphisms $\tilde{\varphi} : (\widetilde{X}, \widetilde{\mathcal{A}}, \widetilde{\tau}) \rightarrow (\widetilde{X}', \widetilde{\mathcal{A}}', \widetilde{\tau}')$ and $f : (\widetilde{X}, \widetilde{\mathcal{A}}, \widetilde{\tau}) \rightarrow (X, \mathcal{A}, \tau)$ such that f is a quasi-isomorphism and the following diagram commutes:

$$\begin{array}{ccc} (\widetilde{X}, \widetilde{\mathcal{A}}, \widetilde{\tau}) & \xrightarrow{\tilde{\varphi}} & (\widetilde{X}', \widetilde{\mathcal{A}}', \widetilde{\tau}') \\ \downarrow f & & \downarrow g \\ (X, \mathcal{A}, \tau) & \xrightarrow{\varphi} & (X', \mathcal{A}', \tau') \end{array}.$$

We may assume that $\widetilde{X}' = X'$. Then $\widetilde{\tau}' \subseteq \tau'$. We let $\widetilde{X} = X$. Let $\tilde{\tau}$ be the family of all $V \in \tau$ for which there is $\widetilde{V}' \in \widetilde{\tau}'$ with $\varphi(V) \subseteq \widetilde{V}'$. By Lemma 2.11, $\tilde{\tau}$ is a net on \widetilde{X} . The k_H -atlas $\widetilde{\mathcal{A}}$ defines a k_H -affinoid atlas $\widetilde{\mathcal{A}}$ with the net $\tilde{\tau}$. The strong morphism $\tilde{\varphi}$ induces $\tilde{\varphi}$. The morphism f is the canonical quasi-isomorphism. It is immediate that these constructions satisfy the desired conditions.

RMS3. If $\varphi, \psi : (X, \mathcal{A}, \tau) \rightarrow (X', \mathcal{A}', \tau')$ are strong morphisms of k_H -analytic spaces and there is a quasi-isomorphism $g : (X', \mathcal{A}', \tau') \rightarrow (\widetilde{X}', \widetilde{\mathcal{A}}', \widetilde{\tau}')$ of k_H -analytic

spaces such that $g \circ \varphi = g \circ \psi$, then there is a quasi-isomorphism $f : (\tilde{X}, \tilde{\mathcal{A}}, \tilde{\tau}) \rightarrow (X, \mathcal{A}, \tau)$ with $\varphi \circ f = \psi \circ f$.

We will in fact show that $\varphi = \psi$. It is clear that they coincide as maps of topological spaces. Let $V \in \tau$, $V' \in \tau'$ such that $\varphi(V) \subseteq V'$. Take $\tilde{V}' \in \tilde{\tau}'$ with $g(V') \subseteq \tilde{V}'$. Then we have two morphisms of k -affinoid spectra $\varphi_{V/V'}, \psi_{V/V'} : V \rightarrow V'$ such that their compositions with $g_{V'/\tilde{V}'}$ coincide. As V' is an affinoid domain in \tilde{V}' , it follows that $\varphi_{V/V'} = \psi_{V/V'}$ by the universal property. \square

Definition 2.13. The category $k_H\text{-}\mathcal{A}\mathfrak{n}$ is the right category of fractions of $k_H\text{-}\widetilde{\mathcal{A}\mathfrak{n}}$ with respect to the system of quasi-isomorphisms. A morphism in $k_H\text{-}\mathcal{A}\mathfrak{n}$ is called a *morphism* between k_H -analytic spaces.

We refer to [Stacks, Tag 04VB] for the definition of right category of fractions. For later references, we explicitly write down the morphisms in $k_H\text{-}\mathcal{A}\mathfrak{n}$.

Lemma 2.14. Let $\varphi : (X, \mathcal{A}, \tau) \rightarrow (X', \mathcal{A}', \tau')$ be a morphism of k_H -analytic spaces. We define a partial order on the set of nets on X : $\tau_1 \preceq \tau_0$ if $\tau_1 \subseteq \overline{\tau_0}$. Then the set of nets is a directed set and

$$\mathrm{Hom}_{k_H\text{-}\mathcal{A}\mathfrak{n}}((X, \mathcal{A}, \tau), (X', \mathcal{A}', \tau')) = \lim_{\substack{\longrightarrow \\ \sigma \preceq \tau}} \mathrm{Hom}_{k_H\text{-}\widetilde{\mathcal{A}\mathfrak{n}}}((X, \mathcal{A}_\sigma, \sigma), (X', \mathcal{A}', \tau'))$$

in the category of sets, where \mathcal{A}_σ is induced by $\overline{\mathcal{A}}$. The transition maps are all injective.

PROOF. This follows immediately from the definition. \square

Definition 2.15. Let (X, \mathcal{A}, τ) be a k_H -analytic space. We say a subset $W \subseteq X$ is τ -special if it is compact and there exist $n \in \mathbb{Z}_{>0}$ and a covering $W = W_1 \cup \dots \cup W_n$ with $W_i \in \tau$, $W_i \cap W_j \in \tau$ for all $i, j = 1, \dots, n$ and the natural map

$$A_{W_i} \hat{\otimes}_k A_{W_j} \rightarrow A_{W_i \cap W_j}$$

is an admissible epimorphism.

The covering W_1, \dots, W_n is called a τ -special covering of W .

Under our convention, the assumption means that $W_i \cap W_j \rightarrow W_i \times W_j$ is a closed immersion of k_H -affinoid spectra.

Example 2.16. Let (X, \mathcal{A}, τ) be a k_H -analytic space. Suppose that $V \in \tau$ and W is a compact k_H -analytic domain in V . Let $n \in \mathbb{Z}_{>0}$ and $W = W_1 \cup \dots \cup W_n$ with $W_i \in \tau$, $W_i \cap W_j \in \tau$ for all $i, j = 1, \dots, n$. Then $\{W_i\}_i$ is a τ -special covering of W . This follows from Corollary 13.14 in Affinoid algebras.

Lemma 2.17. Let (X, \mathcal{A}, τ) be a k_H -analytic space and W be a τ -special subset of X . If $U, V \in \tau|_W$, then $U \cap V \in \tau$ and the natural map

$$A_U \hat{\otimes}_k A_V \rightarrow A_{U \cap V}$$

is an admissible epimorphism.

PROOF. Let $n \in \mathbb{Z}_{>0}$ and W_1, \dots, W_n be a τ -special covering of W . As $U \cap W_i$ and $V \cap W_i$ are compact for $i = 1, \dots, n$, we can find $m_i \in \mathbb{Z}_{>0}$ (resp. $k_i \in \mathbb{Z}_{>0}$) and finite coverings $U_{i1}, \dots, U_{im_i} \in \tau$ of $U \cap W_i$ (resp. $V_{i1}, \dots, V_{ik_i} \in \tau$ of $V \cap W_i$).

Observe that $U_{ik} \cap V_{jl}$ is a k_H -affinoid domain in $U \cap V$, hence $U_{ik} \cap V_{jl} \in \tau$ for any $i, j = 1, \dots, n$, $k = 1, \dots, m_i$ and $l = 1, \dots, k_l$. By Proposition 12.3 in Affinoid

algebras, $U_{ik} \cap V_{jl} \rightarrow U_{ik} \times V_{jl}$ is a closed immersion since $W_i \cap W_j \rightarrow W_i \times W_j$ is by our assumption.

Consider the finite covering

$$\mathcal{U} := \{U_{ik} \times V_{jl} : i, j = 1, \dots, n; k = 1, \dots, m_i; l = 1, \dots, k_l\}$$

of $U \times V$. For each tuple (i, j, k, l) , $A_{U_{ik} \cap V_{jl}}$ is a finite $A_{U_{ik} \times V_{jl}}$ -algebra. By [Theorem 14.1](#) in [Affinoid algebras](#), we can construct a finite $A_{U \times V}$ -algebra $A_{U \cap V}$ inducing all of these $A_{U_{ik} \cap V_{jl}}$'s. By [Proposition 8.1](#) in [Affinoid algebras](#), $A_{U \cap V}$ is k_H -affinoid.

As \mathcal{U} is a finite k_H -affinoid covering of $U \times V$, $\{A_{U_{ik} \cap V_{jl}}\}_{i,k,j,l}$ is a finite k_H -affinoid covering of $U \cap V$ by [Corollary 13.12](#) in [Affinoid algebras](#). In particular, we have a natural homeomorphism

$$\mathrm{Sp} A_{U \cap V} \xrightarrow{\sim} U \cap V.$$

Observe that $A_U \hat{\otimes}_k A_V \rightarrow A_{U \cap V}$ is surjective. We endow $A_{U \cap V}$ with the structure of finite $A_U \hat{\otimes}_k A_V$ -Banach algebras by [Proposition 9.10](#) in [Affinoid algebras](#). Then $A_U \hat{\otimes}_k A_V \rightarrow A_{U \cap V}$ is an admissible epimorphism by [Proposition 9.7](#) in [Affinoid algebras](#).

On the other hand $U \cap V$ is a compact k_H -analytic domain in U , so by [Proposition 13.25](#) in [Affinoid algebras](#), $U \cap V$ is a k_H -affinoid in U . In particular, $U \cap V \in \bar{\tau}$. \square

Lemma 2.18. Let (X, \mathcal{A}, τ) be a k_H -analytic space and $W \subseteq X$ be a τ -special set. Then for any finite covering $\{W_i\}_{i \in I}$ of W with $W_i \in \tau$ for $i \in I$, the Banach k -algebra

$$A_W := \ker \left(\prod_{i \in I} A_{W_i} \rightarrow A_{W_i \cap W_j} \right)$$

does not depend on the choice of $\{W_i\}_{i \in I}$ up to canonical isomorphisms.

Moreover, we have a canonical map $W \rightarrow \mathrm{Sp} A_W$, which does not depend on the choice of the covering modulo the canonical isomorphism between A_W .

PROOF. It follows from [Lemma 2.17](#) that the covering $\{W_i\}_{i \in I}$ is τ -special. It suffices to apply the same argument of [Lemma 13.22](#) in [Affinoid algebras](#). \square

Definition 2.19. Let (X, \mathcal{A}, τ) be a k_H -analytic space. Let $\hat{\tau}$ denote the collection of $\bar{\tau}$ -special subsets $W \subseteq X$ such that

- (1) A_W is k -affinoid;
- (2) the natural map $W \rightarrow \mathrm{Sp} A_W$ is bijective;
- (3) there is a $\bar{\tau}$ -special covering $\{W_i\}_{i \in I}$ of W such that W_i is a k -affinoid domain in W for $i \in I$.

The sets from $\hat{\tau}$ are called *k_H -affinoid domains in (X, \mathcal{A}, τ)* .

Observe that W is k_H -affinoid and W_i is a k_H -affinoid domain in W by [Corollary 13.20](#) in [Affinoid algebras](#). Condition (3) holds for any $\bar{\tau}$ -special covering.

Proposition 2.20. Let (X, \mathcal{A}, τ) be a k_H -analytic space. Then $\hat{\tau}$ is a net. For any net σ on X contained in $\bar{\tau}$, we have $\hat{\sigma} = \hat{\tau}$.

Moreover, $\hat{\hat{\tau}} = \hat{\tau}$.

PROOF. Let $U, V \in \hat{\tau}$. Take $\bar{\tau}$ -special coverings $\{U_i\}_{i \in I}$, $\{V_j\}_{j \in J}$ of U and V respectively. In order to show that $\hat{\tau}|_{U \cap V}$ is a quasi-net, it suffices to show that $\hat{\tau}|_{U_i \cap V_j}$ is for any $i \in I$ and $j \in J$. This follows simply from the fact that $\bar{\tau}|_{U_i \cap V_j}$ is a quasi-net. Similarly, as $\hat{\tau}$ is a quasi-net as $\bar{\tau}$ is. So $\hat{\tau}$ is a net.

Let σ be a net on X contained in $\bar{\tau}$. By [Lemma 2.17](#), it suffices to verify that for any $V \in \bar{\tau}$, there are $n \in \mathbb{Z}_{>0}$ and $U_1, \dots, U_n \in \bar{\sigma}$ with $V = U_1 \cup \dots \cup U_n$. As σ is a net on X , we can find $m \in \mathbb{Z}_{>0}$, $W_1, \dots, W_m \in \sigma$ such that

$$V \subseteq W_1 \cup \dots \cup W_m.$$

As $V, W_j \in \bar{\tau}$ for $j = 1, \dots, m$, by [Lemma 2.5](#) in [Topology and bornology](#), we can find $U_1, \dots, U_n \in \bar{\tau}$ such that $V = U_1 \cup \dots \cup U_n$ and each U_i is contained in some W_j . As $W_j \in \sigma$ for $j = 1, \dots, m$, it follows that $U_i \in \bar{\sigma}$ for $i = 1, \dots, n$.

By [Lemma 2.17](#),

$$\bar{\hat{\tau}} = \hat{\tau}.$$

Let $V \in \hat{\hat{V}}$. Let $\{V_i\}_{i \in I}$ be a $\hat{\tau}$ -special covering of V . For each $i \in I$, take a $\bar{\tau}$ -special covering $\{V_{ij}\}_{j \in J_i}$ of V_i . Then $\{V_{ij}\}_{i,j}$ is a $\bar{\tau}$ -special covering of V . It follows that $V \in \hat{\tau}$. \square

Proposition 2.21. Let (X, \mathcal{A}, τ) be a k_H -analytic space. There is a k_H -analytic atlas $\hat{\mathcal{A}}$ on X with the net $\hat{\tau}$ extending \mathcal{A} . Moreover, $\hat{\mathcal{A}}$ is unique up to a canonical isomorphism.

PROOF. For each $V \in \hat{\tau}$, Fix a $\bar{\tau}$ -special covering $\{V_i\}_{i \in I_V}$.

We define A_V using this covering as in [Lemma 2.18](#). By definition, the canonical map $V \rightarrow \mathrm{Sp} A_V$ is a homeomorphism.

Next take $U, V \in \hat{\tau}$ with $U \subseteq V$. We want to identify U with a k_H -affinoid domain in V . First assume that $U \in \tau$, then $U \cap V_i$ is a k_H -affinoid domain in V_i for $i \in I_V$ by [Lemma 2.17](#). Hence, U is a k_H -affinoid domain in V . If we only know $U \in \hat{\tau}$, we know that U_i is a k_H -affinoid domain in V for any $i \in I_U$. It follows that U is a k_H -affinoid domain in V by [Proposition 13.25](#) in [Affinoid algebras](#).

The uniqueness is immediate. \square

Definition 2.22. Let (X, \mathcal{A}, τ) be a k_H -analytic space. A $\hat{\tau}$ -special set is called a k_H -special domain in X .

Observe that a k_H -special domain inherits a structure of k_H -analytic space from (X, \mathcal{A}, τ) .

Proposition 2.23. Let $\varphi : (X, \mathcal{A}, \tau) \rightarrow (X', \mathcal{A}', \tau')$ be a morphism of k_H -analytic spaces. Then for any k_H -affinoid domains $V \subseteq X$ and $V' \subseteq X'$, the intersection $V \cap \varphi^{-1}(V')$ is a k_H -special domain in X .

PROOF. By [Proposition 2.20](#), we may assume that φ is a strong morphism. In this case, it suffices to apply [Lemma 2.11](#). \square

Lemma 2.24. Let (X, \mathcal{A}, τ) and $(X', \mathcal{A}', \tau')$ be k_H -analytic spaces. Let $\varphi : (X, \mathcal{A}, \tau) \rightarrow (X', \mathcal{A}', \tau')$ be a strong morphism. Then φ extends uniquely to a strong morphism $\varphi : (X, \hat{\mathcal{A}}, \hat{\tau}) \rightarrow (X', \hat{\mathcal{A}}', \hat{\tau}')$.

PROOF. Let $V \in \hat{\tau}$ and $V' \in \hat{\tau}'$ with $\varphi(V) \subseteq V'$. We want to define $\varphi_{V/V'} : V \rightarrow V'$ of k_H -affinoid spectra. By [Proposition 2.8](#), we may extend φ uniquely to $\bar{\tau}$. Take a $\bar{\tau}$ -special covering of V , we may reduce to the case where $V \in \bar{\tau}$. Take

$W' \in \tau'$ such that $\varphi(V) \subseteq W'$. As $\tau|_{W' \cap V'}$ is a quasi-net, we can find $n \in \mathbb{Z}_{>0}$ and $W_1, \dots, W_n \in \tau'|_{V' \cap W}$ such that $\varphi(V) \subseteq W_1 \cup \dots \cup W_n$. Considering the inverse images of W_i 's and $W_i \cap W_j$'s using [Lemma 2.17](#), we are reduced to the case where $V' \in \bar{\tau}'$. This is already handled in [Proposition 2.8](#). The uniqueness of the extension is clear. \square

Proposition 2.25. Let (X, \mathcal{A}, τ) , $(X', \mathcal{A}', \tau')$ be k_H -analytic spaces.

- (1) There is a canonical bijection between

$$\text{Hom}_{k_H\text{-An}}((X, \mathcal{A}, \tau), (X', \mathcal{A}', \tau'))$$

and the set of pairs consisting of

- (a) a continuous map $\varphi : X \rightarrow X'$ such that for all $x \in X$, there exist $n \in \mathbb{Z}_{>0}$, neighbourhoods $V_1 \cup \dots \cup V_n$ of x and $V'_1 \cup \dots \cup V'_n$ of $\varphi(x)$ with $x \in V_1 \cap \dots \cap V_n$ and $\varphi(V_i) \subseteq V'_i$ for $i = 1, \dots, n$, where $V_i \subseteq X$ and $V'_i \subseteq X'$ are k_H -affinoid domains;
- (b) for each pair of k_H -affinoid domains $V \subseteq X$, $V' \subseteq X'$ with $\varphi(V) \subseteq V'$, a morphism of k_H -affinoid spectra $\varphi_{V/V'} : V \rightarrow V'$ such that if $V, W \subseteq X$ and $V', W' \subseteq X'$ are k_H -affinoid domains with $\varphi(V) \subseteq V'$, $\varphi(W) \subseteq W'$, the diagram below commutes

$$\begin{array}{ccc} V & \xrightarrow{\varphi_{V/V'}} & V' \\ \downarrow & & \downarrow \\ W & \xrightarrow{\varphi_{W/W'}} & W' \end{array}.$$

- (2) Under the bijection in (1), an isomorphism corresponds to the pair where φ is a hoemomorphism such that $\varphi(\hat{\tau}) = \hat{\tau}'$ and for any $V \in \hat{\tau}$, $\varphi_{V/\varphi(V)}$ is an isomorphism of k_H -affinoid spectra.

PROOF. (2) follows immediately from (1). So it suffices to prove (1).

We construct the forward map. Let $\varphi : (X, \mathcal{A}, \tau) \rightarrow (X', \mathcal{A}', \tau')$ be a morphism. Take a subnet σ of $\bar{\tau}$ such that φ is represented by a strong morphism

$$\varphi : (X, \mathcal{A}_\sigma, \sigma) \rightarrow (X', \mathcal{A}', \tau').$$

By [Lemma 2.24](#), this extends to a strong morphism

$$\varphi : (X, \widehat{\mathcal{A}}_\sigma, \hat{\sigma}) \rightarrow (X', \widehat{\mathcal{A}}', \hat{\tau}').$$

We get an injective map from the first set into the second set.

Conversely, we need to show that any given map from the second map comes from the first set. It suffices to show that

$$\sigma := \left\{ V \in \hat{\tau} : \varphi(V) \subseteq V' \text{ for some } V' \in \hat{\tau}' \right\}$$

is a net. Take $x \in X$ and neighbourhoods $V_1 \cup \dots \cup V_n$ of x and $V'_1 \cup \dots \cup V'_n$ of $\varphi(x)$ as in the statement of (1). Then $V_i \in \sigma$, so we conclude. \square

In practice, we do not distinguish a k_H -analytic space from the isomorphic k_H -analytic spaces. In particular, we will write (X, \mathcal{A}, τ) as X and always endow it with the structure $(X, \hat{\mathcal{A}}, \hat{\tau})$ of k_H -analytic space. If necessarily, we will write $|X|$ for the underlying topological space.

Corollary 2.26. The natural functor $k_H\text{-Aff} \rightarrow k_H\text{-An}$ is fully faithful.

PROOF. Let $X = \mathrm{Sp} A$ be a k_H -affinoid spectrum. We endow it with the net $\tau = \{X\}$. The k_H -atlas with the net τ assigns $X \in \tau$ with A . It is easily verified that this is a functor. By [Proposition 2.25](#), the functor is fully faithful. \square

Definition 2.27. A k_H -affinoid space is an object of $k_H\text{-An}$ lying in the essential image of the functor $k_H\text{-Aff} \rightarrow k_H\text{-An}$.

The category of k_H -affinoid spaces is denoted by $k_H\text{-Aff}$.

The notation for the category of k_H -affinoid spaces is the same as the notation for the category of k_H -affinoid spectra, as the two categories are canonically equivalent.

Definition 2.28. A k_H -analytic space X is *good* if any point $x \in X$ admits a k_H -affinoid neighbourhood.

Example 2.29. Fix $n \in \mathbb{N}$. Let \mathbb{A}_k^n denote the set of all semi-valuations on $k[T_1, \dots, T_n]$ whose restriction to k coincides with the given valuation on k . We provide \mathbb{A}_k^n with the weakest topology such that for any $f \in k[T_1, \dots, T_n]$, the map $|\bullet| \mapsto |f|$ is continuous.

Observe that as a topological space,

$$(2.1) \quad \mathbb{A}_k^n \xrightarrow{\sim} \varinjlim_{r \in \mathbb{R}_{>0}^n} \mathrm{Sp} k\{r^{-1}T\}.$$

As a set, this is clear: if $|\bullet| \in \mathbb{A}_k^n$, we take $r = (|T_1|, \dots, |T_n|)$, then $|\bullet| \leq \|\bullet\|_r$, so $|\bullet| \in \mathrm{Sp} k\{r^{-1}T\}$. As

$$\bigcap_{r \in \mathbb{R}_{>0}^n} k\{r^{-1}T\} = k[T_1, \dots, T_n],$$

so the topology on the right-hand side of (2.1) is the weakest topology making $|\bullet| \mapsto |f|$ continuous for any $f \in k[T_1, \dots, T_n]$. It follows immediately that (2.1) is an identification of topological spaces.

It is clear that \mathbb{A}_k^n has a structure of good k_H -analytic space.

Proposition 2.30. Let X be a k_H -analytic space, $x \in X$ and U be a neighbourhood of x in X . Then there is a neighbourhood V of x in X contained in U such that V is open connected locally compact paracompact and Hausdorff. Moreover, we can guarantee that $\bar{V} \subseteq U$ and V is a countable union of k_H -affinoid domains.

PROOF. Take $n \in \mathbb{Z}_{>0}$ and k_H -affinoid spaces V_1, \dots, V_n containing x and $V_1 \cup \dots \cup V_n$ is a neighbourhood of x in X . If we have proved the proposition for V_i in place of X and $U \cap V_i$ in place of U for $i = 1, \dots, n$, namely, if we have found open connected locally compact paracompact and Hausdorff sets W_i containing x and contained in $U \cap V_i$ whose closure in V_i is contained in $U \cap V_i$, then we can take $V = W_1 \cup \dots \cup W_n$.

So we may assume that X is a k_H -affinoid space, say $X = \mathrm{Sp} A$. Choose a k_H -rational neighbourhood

$$W = \mathrm{Sp} A\{r^{-1} \frac{f}{g}\}$$

of x in U , where $n \in \mathbb{N}$, $f = (f_1, \dots, f_n) \in A^n$, $r \in \sqrt{|k^\times| \cdot H}^n$, $g \in A$ and f_1, \dots, f_n, g generate the unit ideal in A . This is possible by [Corollary 10.9](#) and [Proposition 10.13](#) in [Affinoid algebras](#). Take $\delta > 0$ so that $x \in \mathrm{Sp} A\{((1 - \delta)r)^{-1} \frac{f}{g}\}$.

Choose a strictly increasing sequence $\epsilon_i \in (0, 1) \cap \sqrt{|k^\times| \cdot H}$ converging to $1 - \delta/2$ for $i \in \mathbb{Z}_{>0}$. Let

$$W_i = \operatorname{Sp} A \left\{ (\epsilon_i r)^{-1} \frac{f}{g} \right\}$$

for $i \in \mathbb{Z}_{>0}$. Then W_i lies in the interior of W_{i+1} for $i \in \mathbb{Z}_{>0}$. Choose a connected component V_i of W_i so that $V_1 \subseteq V_2 \subseteq \dots$ and $x \in V := \bigcup_{i=1}^{\infty} V_i$. If $x \in V_i$ for some $i \in \mathbb{Z}_{>0}$, then x lies in the topological interior of V_{i+1} . Hence, x lies in the interior of V . By construction, V is open connected paracompact locally compact and Hausdorff. Moreover, $\bar{V} \subseteq U$ by our construction. \square

Proposition 2.31. Let $\{X_i\}_{i \in I}$ be a family of k_H -analytic spaces. Suppose that for $i, j \in I$, we are given a k_H -analytic domain $X_{ij} \subseteq X_i$ and an isomorphism $\nu_{ij} : X_{ij} \rightarrow X_{ji}$ satisfying the *cocycle condition*: $X_{ii} = X_i$, $\nu_{ij}(X_{ij} \cap X_{il}) = X_{ji} \cap X_{jl}$ and $\nu_{il} = \nu_{jl} \circ \nu_{ij}$ on $X_{ij} \cap X_{il}$ for $i, j, l \in I$.

Assume that either of the following conditions holds:

- (1) X_{ij} is open in X_i for all $i, j \in I$;
- (2) for any $i \in I$, all X_{ij} 's are closed in X_i and the number of $j \in I$ with $X_{ij} \neq \emptyset$ is finite.

Then there is a k_H -analytic space X and morphisms $\mu_i : X_i \rightarrow X$ for $i \in I$ such that

- (1) μ_i is an isomorphism of X_i with a k_H -analytic domain in X ;
- (2) $X = \bigcup_{i \in I} \mu_i(X_i)$;
- (3) $\mu_i(X_{ij}) = \mu_i(X_i) \cap \mu_j(X_j)$ for $i, j \in I$;
- (4) $\mu_i = \mu_j \circ \nu_{ij}$ on X_{ij} for $i, j \in I$.

The space X is unique up to a canonical isomorphism. Moreover, under Condition (1), $\mu_i(X_i)$ is open in X for $i \in I$; under Condition (2), $\mu_i(X_i)$ is closed in X for $i \in I$.

Under both conditions, if all X_i 's are Hausdorff (resp. paracompact), then so is X .

We will call X the *gluing* of the X_i 's along the X_{ij} 's.

PROOF. By [Proposition 3.12](#), the uniqueness of X is clear.

Let

$$\tilde{X} = \coprod_{i \in I} X_i$$

in $k_H\text{-An}$. Observe that

$$|\tilde{X}| = \coprod_{i \in I} |X_i|$$

in the category $\mathcal{T}\text{op}$. The system ν_{ij} 's defines an equivalence relation R on $|\tilde{X}|$. Let $|X| = |\tilde{X}|/R$ and $\mu_i : |X_i| \rightarrow |X|$ be the induced map for $i \in I$.

Under Condition (1), $\mu_i(|X_i|)$ is open in $|X|$ for $i \in I$. Under Condition (2), $\mu_i(|X_i|)$ is closed in $|X|$ for $i \in I$.

Under both conditions, the map μ_i induces a homeomorphism $|X_i| \rightarrow \mu_i(|X_i|)$ for $i \in I$. If all $|X_i|$'s are Hausdorff (resp. paracompact), so is $|X|$.

All these claims follow from well-known results in general topology.

We will endow $|X|$ with a structure of k_H -analytic space. Let τ be the set of $V \subseteq |X|$ for which there is $i \in I$ such that $V \subseteq \mu_i(X_i)$ and $\mu_i^{-1}(V)$ is a k_H -affinoid domain in X_i . Then τ is a net on X . There is an obvious k -affinoid atlas on X with the net τ . All properties in the proposition are satisfied by $X = (|X|, \mathcal{A}, \tau)$. \square

Definition 2.32. Let X be a k_H -analytic space and $x \in X$, take a k_H -affinoid domain $\mathrm{Sp} A$ in X containing x , we define the *completed residue field* $\mathcal{H}(x)$ of x in X as the completed residue field of x in $\mathrm{Sp} A$.

By [Corollary 13.16](#) in [Affinoid algebras](#), $\mathcal{H}(x)$ does not depend on the choice of $\mathrm{Sp} A$ up to an isomorphism of complete valuation fields over k .

3. Analytic domains

Let $(k, |\bullet|)$ be a complete non-Archimedean valued field and H be a subgroup of $\mathbb{R}_{>0}$ such that $|k^\times| \cdot H \neq \{1\}$.

Definition 3.1. Let X be a k_H -analytic space. A subset $Y \subseteq X$ is called a *k_H -analytic domain* if for any $y \in Y$, there exist $n \in \mathbb{Z}_{>0}$, k_H -affinoid domains V_1, \dots, V_n contained in Y such that

- (1) $y \in V_1 \cap \dots \cap V_n$;
- (2) $V_1 \cup \dots \cup V_n$ is a neighbourhood of y in Y .

Observe that the net of k_H -affinoid domains in X that are contained in Y form a net on Y . In particular, Y inherits a k_H -analytic space structure from X , and we have a canonical morphism $Y \rightarrow X$ in $k_H\text{-An}$.

Lemma 3.2. Let X be a k_H -analytic space, Y be a k_H -analytic domain in X and $x \in Y$. Then the completed residue field of x in X is the same as the completed residue field of x in Y modulo isomorphisms of completed valuation fields over k .

PROOF. This follows immediately from [Corollary 13.16](#) in [Affinoid algebras](#). \square

Proposition 3.3. Let X be a k -analytic space and $x \in X$. Let $\lambda \in \widetilde{\mathcal{H}(x)}$ be a non-zero homogeneous element. Then we can find a k -affinoid domain $\mathrm{Sp} B$ of x in X and an invertible function $f \in B$ such that

$$\lambda = \widetilde{f(x)}.$$

If X is good, we may assume that $\mathrm{Sp} B$ is a k -affinoid neighbourhood of x in X .

PROOF. We may assume that X is k -affinoid, say $X = \mathrm{Sp} A$. Let $\chi_x : A \rightarrow \mathcal{H}(x)$ be the character corresponding to x . Let $|\bullet|_x$ be the bounded semi-valuation on A corresponding to x . As $\mathrm{Frac} A / \ker |\bullet|_x$ is dense in $\mathcal{H}(x)$ by definition, we can find $g, h \in A$ such that $g(x) \neq 0$, $h(x) \neq 0$ and

$$\lambda = \widetilde{g(x)} / \widetilde{h(x)}.$$

Let Y be the open k -analytic domain in X defined by $g(x) \neq 0$ and $h(x) \neq 0$. We take a k -affinoid domain $\mathrm{Sp} B$ of X containing x such that $\mathrm{Sp} B \subseteq Y$. If X is good, we may assume that $\mathrm{Sp} B$ is a neighbourhood of x in X . Then the images of g and h in B are invertible by [Corollary 6.11](#) in [Banach rings](#). Now $f = g/h \in B$ satisfies our assumptions. \square

Example 3.4. Let X be a k_H -analytic space. Then any open subset U of X is a k_H -analytic domain.

In fact, for $x \in U$, take V_1, \dots, V_n as in [Definition 3.1](#). By [Proposition 10.13](#) in [Affinoid algebras](#), up to replacing V_i 's by k_H -Laurent domains in them, we may guarantee that $V_i \subseteq U$ for all $i = 1, \dots, n$.

Proposition 3.5. Let X, X' be k_H -analytic spaces and $\varphi : X' \rightarrow X$ a morphism of k_H -analytic spaces.

- (1) Let Y, Z be k_H -analytic domains in X , then so is $Y \cap Z$.
- (2) Let Y be a k_H -analytic domain in X , then $\varphi^{-1}(Y)$ is a k_H -analytic domain in X' .

PROOF. (1) Let $x \in Y \cap Z$. Take k_H -affinoid domains V_1, \dots, V_n contained in Y and k_H -affinoid domains W_1, \dots, W_m contained in Z such that

$$x \in V_1 \cap \dots \cap V_n, \quad x \in W_1 \cap \dots \cap W_m$$

and $V_1 \cup \dots \cup V_n$ is a neighbourhood of x in Y , $W_1 \cup \dots \cup W_m$ is a neighbourhood of x in Z . For each $i = 1, \dots, n$ and $j = 1, \dots, m$, $\hat{\tau}|_{V_i \cap W_j}$ is a quasi-net, so we can find a neighbourhood of x in $V_i \cap W_j$ of the form $U_1^{ij} \cup \dots \cup U_{m_{ij}}^{ij}$ with $U_1^{ij}, \dots, U_{m_{ij}}^{ij}$ being k_H -affinoid domains in X containing x . Then each element in the collection $\{U_k^{ij}\}$ contains x and the union is a neighbourhood of x in $Y \cap Z$.

(2) Let $x' \in \varphi^{-1}(Y)$ and $x = \varphi(x')$. By [Proposition 2.25](#), we can find $n \in \mathbb{Z}_{>0}$, k_H -affinoid domains V'_1, \dots, V'_n on X' and k_H -affinoid domains V_1, \dots, V_n on X such that

$$x' \in V'_1 \cap \dots \cap V'_n, \quad x \in V_1 \cap \dots \cap V_n, \\ \varphi(V'_i) \subseteq V_i \text{ for } i = 1, \dots, n,$$

and $V'_1 \cup \dots \cup V'_n$ (resp. $V_1 \cup \dots \cup V_n$) is a neighbourhood of x' (resp. x) in X' (resp. X). Take k_H -affinoid domains W_1, \dots, W_m in X contained in Y , each containing x such that $W_1 \cup \dots \cup W_m$ is a neighbourhood of x in Y .

Then for each $i = 1, \dots, n$, $j = 1, \dots, m$, we can find k_H -affinoid domains W_{ij}^k for $k = 1, \dots, r_{ij}$ contained in $W_j \cap V_i$ and containing x such that $\cup_k W_{ij}^k$ is a neighbourhood of x in $W_j \cap V_i$. Thus, $\cup_{j,k} W_{ij}^k$ is a neighbourhood of x in $V_i \cap Y$. Then $U_{ij}^k := \varphi^{-1}(V_i^k) \cap V'_i$ is a k_H -affinoid domain in V'_i by [Corollary 13.12](#) in [Affinoid algebras](#). Moreover, $\cup_{j,k} U_{ij}^k$ is a neighbourhood of x' in $V'_i \cap Y'$. So $\cup_{i,j,k} U_{ij}^k$ is a neighbourhood of x' in Y' . \square

Proposition 3.6. Let X be a k_H -analytic space and Y be a k_H -analytic domain in X . Then for any k_H -analytic space Z and any morphism $\varphi : Z \rightarrow X$ whose image is contained in Y , there is a unique morphism $\psi : Z \rightarrow Y$ such that the following diagram commutes:

$$\begin{array}{ccc} Z & & \\ \downarrow \psi & \searrow \varphi & \\ Y & \longrightarrow & X \end{array}.$$

PROOF. The uniqueness of ψ is obvious. We only need to prove the existence. This is an immediate consequence of [Proposition 2.25](#) and [Proposition 3.5](#).

To be more precise, assume that φ is given by a data as in [Proposition 2.25](#), we only have to show that each k_H -affinoid domain V in X , $V \cap Y$ is a k_H -affinoid domain in Y . This follows from [Proposition 3.5](#). \square

Corollary 3.7. Let $\varphi : X' \rightarrow X$ be a morphism of k_H -analytic spaces and Y be a k_H -analytic domain in X . Then $X' \times_Y X$ in the category $k_H\text{-An}$ exists and $\varphi^{-1}(Y)$ represents $X' \times_Y X$.

PROOF. This follows from [Proposition 3.6](#) and [Proposition 3.5](#). \square

Corollary 3.8. Let $\mathrm{Sp} B$ be a k_H -affinoid space, then we have a functorial isomorphism

$$\mathrm{Hom}_{k_H\text{-}\mathcal{A}\mathrm{n}}(\mathrm{Sp} B, \mathbb{A}_k^1) \xrightarrow{\sim} B.$$

PROOF. As $\mathrm{Sp} B$ is compact as a topological space, its image in \mathbb{A}_k^1 is contained in $\mathrm{Sp} k\{r^{-1}T\}$ for some $r > 0$. By [Proposition 3.6](#), we have natural bijections

$$\mathrm{Hom}_{k_H\text{-}\mathcal{A}\mathrm{n}}(\mathrm{Sp} B, \mathbb{A}_k^1) \xrightarrow{\sim} \varinjlim_{r>0} \mathrm{Hom}_{k_H\text{-}\mathcal{A}\mathrm{n}}(\mathrm{Sp} B, \mathrm{Sp} k\{r^{-1}T\}) \xrightarrow{\sim} \varinjlim_{r>0} \mathrm{Hom}_{k\text{-}\mathcal{A}\mathrm{ff}\mathcal{A}\mathrm{lg}}(k\{r^{-1}T\}, B).$$

By [Corollary 6.5](#) in [Affinoid algebras](#), the right-hand side is identified with B . \square

Proposition 3.9. Let X be a k_H -analytic space, Y be a k_H -analytic domain in X . For a subset $Z \subseteq Y$, the following are equivalent:

- (1) Z be a k_H -analytic domain in X ;
- (2) Z is a k_H -analytic domain in Y .

PROOF. (1) \implies (2): Let $z \in Z$, we can find $n \in \mathbb{Z}_{>0}$ and k_H -affinoid domains V_1, \dots, V_n in X containing z and contained in Z such that $V_1 \cup \dots \cup V_n$ is a neighbourhood of z in Z . But observe that V_1, \dots, V_n are k_H -affinoid domains in Y as well, so we conclude.

(2) \implies (1): This follows from the same argument. It suffices to observe that a k_H -affinoid domain in Y is necessarily k_H -affinoid in X , as can be seen from [Definition 2.19](#). \square

Definition 3.10. Let X, Y be k_H -analytic spaces and $\varphi : Y \rightarrow X$ be a morphism. We say φ is an *open immersion* if $\varphi(Y)$ is open in X and φ induces an isomorphism between Y and $\varphi(Y)$ as k_H -analytic spaces.

By [Example 3.4](#), $\varphi(Y)$ is a k_H -analytic domain in X and by [Proposition 3.6](#), we have a morphism of k_H -analytic spaces $Y \rightarrow \varphi(Y)$.

Proposition 3.11. Let X be a k_H -analytic space and Y be a k_H -analytic domain in X . Assume that Y is a k_H -affinoid space, then Y is a k_H -affinoid domain in X .

PROOF. As Y is a k_H -affinoid space, we know that $|Y|$ is compact. Take finitely many k_H -affinoid domains V_1, \dots, V_n in X such that

$$Y = V_1 \cup \dots \cup V_n.$$

Then V_1, \dots, V_n are k_H -affinoid domains in Y : let $\mathrm{Sp} D \rightarrow Y$ be a morphism of k_H -affinoid spectra, whose image lies in V_i for some $i = 1, \dots, n$. Consider the following commutative diagram

$$\begin{array}{ccccc} & & \mathrm{Sp} D & & \\ & \swarrow \text{dotted} & \downarrow & \searrow & \\ V_i & \longrightarrow & Y & \longrightarrow & X \end{array}$$

By [Proposition 3.6](#), there is a unique dotted morphism making the outer triangle commutative, hence making the whole diagram commutative. We have therefore shown that V_i is a k_H -affinoid domain in Y .

So the covering $\{V_1, \dots, V_n\}$ of Y satisfies the assumptions in [Definition 2.19](#) and Y is k_H -affinoid. \square

Proposition 3.12. Let X be a k_H -analytic space and $\{Y_i\}_{i \in I}$ be a family of k_H -analytic domains in X which forms a quasi-net on X . Then for any k_H -analytic space X' , the following sequence is exact

$$\mathrm{Hom}_{k_H\text{-An}}(X, X') \rightarrow \prod_{i \in I} \mathrm{Hom}_{k_H\text{-An}}(Y_i, X') \rightrightarrows \prod_{i, j \in I} \mathrm{Hom}_{k_H\text{-An}}(Y_i \cap Y_j, X').$$

PROOF. Let $\{\varphi_i : Y_i \rightarrow X'\}_{i \in I}$ be a family of morphisms such that φ_i, φ_j coincides on $Y_i \cap Y_j$ for $i, j \in I$. We need to glue the φ_i 's into a single morphism $\varphi : X \rightarrow X'$. Clearly, the underlying maps glue together to a continuous map $\varphi : X \rightarrow X'$ by [Lemma 2.2 in Topology and bornology](#).

Let τ be the collection of k_H -affinoid domains V in X such that there is $i \in I$ and a k_H -affinoid domain $V' \subseteq X'$ with $V \subseteq Y_i$ and $\varphi_i(V) \subseteq V'$. Then τ is a net on X , and we have a morphism $X \rightarrow X'$. \square

4. Berkovich site

Let $(k, |\bullet|)$ be a complete non-Archimedean valued field and H be a subgroup of $\mathbb{R}_{>0}$ such that $|k^\times| \cdot H \neq \{1\}$.

Lemma 4.1. Let X be a k_H -analytic space. Consider the category \mathcal{C} of k_H -analytic domains in X , where the morphisms are inclusions of k_H -analytic domains. For each $Y \in \mathcal{C}$, consider the set of coverings $\mathrm{Cov}(Y)$ consisting of all $\{Y_i \rightarrow Y\}_{i \in I}$ such that Y_i is a k_H -analytic domain in Y and $\{Y_i\}_{i \in I}$ is a quasi-net on Y . The class of coverings $\{\mathrm{Cov}(Y)\}_Y$ defines a Grothendieck pretopology.

PROOF. It suffices to verify the axioms in [\[Stacks, Tag 03NH\]](#).

(1) An isomorphism $Y' \rightarrow Y$ in \mathcal{C} is in $\mathrm{Cov}(Y)$.

This is trivial as an isomorphism in \mathcal{C} is necessarily identity.

(2) If $\{Y_i \rightarrow Y\}_{i \in I}$ and $\{Y_{ij} \rightarrow Y_i\}_{j \in J_i}$ for all $i \in I$ are in $\mathrm{Cov}(Y)$ and $\mathrm{Cov}(Y_i)$ respectively, then $\{Y_{ij} \rightarrow Y\}_{i,j}$ is in $\mathrm{Cov}(Y)$.

By [Proposition 3.9](#), Y_{ij} is a k_H -analytic domain in Y for any $i \in I, j \in J_i$. It suffices to show that $\{Y_{ij}\}_{i,j}$ is a quasi-net on Y . Let $y \in Y$, we can find finitely many elements among $\{Y_i\}_{i \in I}$, say Y_1, \dots, Y_n so that $y \in Y_i$ for each $i = 1, \dots, n$ and $Y_1 \cup \dots \cup Y_n$ is a neighbourhood of y in Y . Similarly, for each $i = 1, \dots, n$, we can find finitely many Y_{i1}, \dots, Y_{ij_i} among $\{Y_{ij}\}_{j \in J_i}$ so that y is contained in each of them and $Y_{i1} \cup \dots \cup Y_{ij_i}$ is a neighbourhood of y in Y_i . Then each element in $\{Y_{ij}\}_{i=1, \dots, n; j=1, \dots, j_i}$ contains y and the union is a neighbourhood of y in Y .

(3) If $\{Y_i \rightarrow Y\}_{i \in I}$ lies in $\mathrm{Cov}(Y)$ and $Z \rightarrow Y$ is a k_H -analytic domain in Y , then the fiber products $Y_i \times_Y Z$ exist and $\{Y_i \times_Y Z \rightarrow Z\}_{i \in I}$ lies in $\mathrm{Cov}(Z)$.

By [Corollary 3.7](#), $Y_i \times_Y Z$ exists and is represented by the inverse image of Z in Y_i , which is a k_H -analytic domain in Y_i by [Proposition 3.5](#). It is clear that $\{Y_i \times_Y Z\}_{i \in I}$ is a quasi-net on Z . \square

Definition 4.2. Let X be a k_H -analytic space. We will write the site constructed in [Lemma 4.1](#) as X and call it the *Berkovich site* of X . The corresponding Grothendieck topology is called the *Berkovich Grothendieck topology*. The topos $\mathrm{Sh}(X)$ associated with X is called the *Berkovich topos* of X .

Observe that the Berkovich Grothendieck topology is subcanonical by [Proposition 3.12](#).

Definition 4.3. Let X be a k_H -analytic space. We define a sheaf of rings \mathcal{O}_X on X as follows: let Y be a k_H -analtic domain in X , we set

$$\mathcal{O}_X(Y) = \text{Hom}_{k_H\text{-An}}(X, \mathbb{A}_k^1).$$

By [Corollary 3.8](#) and [Proposition 3.12](#), \mathcal{O}_X defines a sheaf of rings. We call \mathcal{O}_X the structure sheaf of X . The corresponding ringed site (X, \mathcal{O}_X) is called the *Berkovich ringed site*. The induced ringed topos $(\text{Sh}(X), \mathcal{O}_X)$ is called the *Berkovich ringed topos*.

Given any morphism $f : Y \rightarrow X$ of k_H -analytic spaces, we have an induced morphism of the corresponding ringed sites, still denoted by φ .

Definition 4.4. Let X be a k_H -analytic space. An \mathcal{O}_X -module \mathcal{M} is *coherent* if there is an admissible covering $\{Y_i\}_{i \in I}$ of X such that $\mathcal{M}|_{Y_i}$ is isomorphic to the cokernel of a homomorphism of finite free \mathcal{O}_{Y_i} -modules.

Example 4.5. Let A be a k_H -affinoid algebra and M be a finite A -module. Then M induces a coherent sheaf of $\mathcal{O}_{\text{Sp } A}$ -modules \tilde{M} as follows:

$$\tilde{M}(V) = M \otimes_A A_V.$$

Conversely, we can reformulate Kiehl's theorem.

Theorem 4.6. Let A be a k_H -affinoid algebra and \mathcal{M} be a coherent sheaf of $\mathcal{O}_{\text{Sp } A}$ -modules. Set $M = H^0(X, \mathcal{M})$, then M is a finite A -module and we have a canonical isomorphism

$$\tilde{M} \xrightarrow{\sim} \mathcal{M}.$$

The left-hand side is defined in [Example 4.5](#).

PROOF. This is just a reformulation of [Theorem 14.1](#) in [Affinoid algebras](#). \square

Corollary 4.7. Let $\varphi : \text{Sp } B \rightarrow \text{Sp } A$ be a morphism of k_H -affinoid spaces. Then the following are equivalent:

- (1) $\varphi_* \mathcal{O}_{\text{Sp } B}$ is a coherent $\mathcal{O}_{\text{Sp } A}$ -module;
- (2) B is a finite Banach A -module.

PROOF. Observe that for any k_H -affinoid domain $\text{Sp } C$ in $\text{Sp } A$,

$$\varphi_* \mathcal{O}_{\text{Sp } B}(\text{Sp } C) = \mathcal{O}_{\text{Sp } B}(\varphi^{-1}(\text{Sp } C)) = \mathcal{O}_{\text{Sp } B}(\text{Sp } C \hat{\otimes}_A B) = C \hat{\otimes}_A B \xrightarrow{\sim} C \otimes_A B.$$

Here we applied [Corollary 13.12](#) in ?? and [Proposition 9.6](#) in ?? . So $\varphi_* \mathcal{O}_{\text{Sp } B} \cong \tilde{B}$.

From this (2) trivially implies (1).

Conversely, assume (1), let $B = H^0(\text{Sp } A, \varphi_* \mathcal{O}_{\text{Sp } B})$. By [Theorem 4.6](#), B is a finite A -module. Let B' denote the ring B endowed with the finite Banach A -algebra structure as in [Proposition 9.10](#) in [Affinoid algebras](#). We need to show that the identity map $B' \rightarrow B$ is admissible. Observe that the identity map is bounded by [Proposition 9.4](#) in ?? . By [Proposition 10.5](#) in ?? , it suffices to show that the induced map $\text{Sp } B \rightarrow \text{Sp } B'$ is surjective. Let $\varphi' : \text{Sp } B' \rightarrow \text{Sp } A$ be the natural morphism of k_H -affinoid spaces. Then

$$\varphi_*(\mathcal{O}_{\text{Sp } B}) \xrightarrow{\sim} \varphi'_*(\mathcal{O}_{\text{Sp } B'}).$$

It follows that $\varphi^{-1}(x) = \varphi'^{-1}(x)$ for any $x \in \text{Sp } A$. We conclude. \square

Corollary 4.8. Let $\varphi : \text{Sp } B \rightarrow \text{Sp } A$ be a morphism of k_H -affinoid spaces. Then the following are equivalent:

- (1) $\varphi_* \mathcal{O}_{\mathrm{Sp} B}$ is a coherent $\mathcal{O}_{\mathrm{Sp} A}$ -module and $\mathcal{O}_{\mathrm{Sp} A} \rightarrow \varphi_* \mathcal{O}_{\mathrm{Sp} B}$ is surjective;
- (2) $A \rightarrow B$ is an admissible epimorphism.

PROOF. Assume (2). By [Corollary 4.7](#), $\varphi_* \mathcal{O}_{\mathrm{Sp} B}$ is a coherent $\mathcal{O}_{\mathrm{Sp} A}$ -module. To see that $\mathcal{O}_{\mathrm{Sp} A} \rightarrow \varphi_* \mathcal{O}_{\mathrm{Sp} B}$ is surjective, it suffices to show that for each k_H -affinoid space $\mathrm{Sp} C$ in $\mathrm{Sp} A$,

$$C \rightarrow C \otimes_A B$$

is surjective. This follows from the assumption.

Assume (1). We know that B is a finite Banach A -module. In particular, $A \rightarrow B$ is admissible by [Proposition 9.7](#) in [Affinoid algebras](#). As $\mathcal{O}_{\mathrm{Sp} A} \rightarrow \varphi_* \mathcal{O}_{\mathrm{Sp} B}$ is surjective, by [Theorem 4.6](#), $A \rightarrow B$ is surjective. [Include details](#) \square

Definition 4.9. Let $\mathrm{Sp} A$ be a k_H -affinoid space and $\mathcal{M} = \tilde{M}$ is a coherent sheaf of \mathcal{O}_X -modules on X , where M is a finite A -module. The *support* $\mathrm{Supp} M$ of \mathcal{M} is the closed subset $\mathrm{Sp} A / \mathrm{Ann}_A(M)$ of $\mathrm{Sp} A$.

Let X be a k_H -analytic space and \mathcal{M} be a coherent sheaf of \mathcal{O}_X -modules. Then the *support* $\mathrm{Supp} \mathcal{M}$ of \mathcal{M} is a subset of X such that a point $x \in X$ lies in $\mathrm{Supp} \mathcal{M}$ if and only if for some k_H -affinoid domain V in X containing x , $x \in \mathrm{Supp} \mathcal{M}|_V$.

Here $\mathrm{Ann}_A(M)$ is the annihilator of M in A .

Lemma 4.10. Let X be a k_H -analytic space and \mathcal{M} be a coherent sheaf of \mathcal{O}_X -modules. Take $x \in \mathrm{Supp} \mathcal{M}|_V$ and a k_H -affinoid domain V in X containing x . Then $x \in \mathrm{Supp} \mathcal{M}|_V$.

PROOF. By assumption, there is a k_H -affinoid domain U in X containing x such that $x \in \mathrm{Supp} \mathcal{M}|_U$.

Let $W \subseteq U \cap V$ be a k_H -affinoid domain in X containing x . We claim that $x \in \mathrm{Supp} \mathcal{M}|_W$. Let $M = H^0(U, \mathcal{M})$, then $M \otimes_{A_U} A_W = H^0(W, \mathcal{M})$. By [\[Stacks, Tag 07T8\]](#) and [Theorem 13.18](#) in [Affinoid algebras](#),

$$\mathrm{Ann}_{A_U}(M) \otimes_{A_U} A_W = \mathrm{Ann}_{A_W}(M \otimes_{A_U} A_W)$$

and $\mathrm{Supp}(\mathcal{M}|_W) = \mathrm{Supp}(\mathcal{M}|_U) \cap W$. The claim follows. We may assume that $U \subseteq V$. In this case, the same argument shows that $x \in \mathrm{Supp} \mathcal{M}|_V$. \square

Proposition 4.11. Let X be a Hausdorff k_H -analytic space. Then the following are equivalent:

- (1) X is paracompact;
- (2) X admits a locally finite covering by k_H -affinoid domains.

Note that the covering in (2) is necessarily a G-covering.

PROOF. Assume (1). Then (2) follows from [Proposition 3.2](#) in [Topology and bornology](#). We take \mathcal{B} to be the collection of finite unions of k_H -affinoid domains that contain an open subset of X .

Assume (2). Let $\{X_i\}_{i \in I}$ be a locally finite covering of X by k_H -affinoid domains. Define an equivalence relation on I generated by $i \sim j$ if $X_i \cap X_j \neq \emptyset$. We say X_i and X_j are elementarily linked in this case. Fix $C \in I / \sim$ and $i \in C$. For any $n \in \mathbb{Z}_{>0}$, C_n denotes the union of X_j where j and i are linked through a chain of elementary links of length at most n . As the covering is locally finite, we see that C_n is compact. So

$$X_C = \bigcup_{i=1}^{\infty} C_i$$

is σ -compact. The space X is clearly the coproduct of X_C 's, hence paracompact by [Proposition 3.2](#) in [Topology and bornology](#). \square

Proposition 4.12. The category $k_H\text{-An}$ admits finite limits.

PROOF. By general abstract nonsense, it suffices to show that $k_H\text{-An}$ admits finite fiber products.

Let $\varphi : Y \rightarrow X$ and $f : X' \rightarrow X$ be morphisms of k_H -affinoid spaces. We want to construct $Y \times_X X'$.

Step 1. We assume that X, Y, X' are all paracompact and Hausdorff.

By [Proposition 4.11](#), we can find a locally finite G-covering $\{X_i\}_{i \in I}$ of X consisting of k_H -affinoid domains in X . By [Proposition 4.11](#) again, we can find a locally finite G-covering $\{Y_{ij}\}_j$ $\varphi^{-1}(X_i)$ consisting of k_H -affinoid domains in Y and a locally finite G-covering $\{X'_{il}\}_l$ consisting of k_H -affinoid domains in X' for each $i \in I$.

We can glue $Y_{ij} \times_{X_i} X'_{il}$'s by [Proposition 2.31](#) to get a k_H -analytic space Y' . By [Proposition 3.12](#), Y' represents the fiber product $Y \times_X X'$.

Step 2. Assume only that X is a paracompact and Hausdorff.

Take open paracompact Hausdorff coverings $\{Y_i\}_{i \in I}$ of Y and $\{X'_j\}_{j \in J}$ of X' . The existence of these coverings follows from [Proposition 2.30](#). Similar to Step 1, we glue the $Y_i \times_X X'_j$'s along the open subsets $(Y_i \cap Y_k) \times_X (X'_j \cap X'_l)$'s by [Proposition 2.31](#), we get a locally Hausdorff k_H -analytic space Y' . Then by [Proposition 3.12](#) again, Y' represents the fiber product $Y \times_X X'$.

Step 3. We handle the general case.

Take a covering $\{X_i\}_{i \in I}$ by open paracompact Hausdorff subsets. Let Y' be the gluing of $\varphi^{-1}(X_i) \times_{X_i} f^{-1}(X_i)$'s along $\varphi^{-1}(X_i \cap X_j) \times_{X_i \cap X_j} f^{-1}(X_i \cap X_j)$'s by [Proposition 2.31](#). Then by [Proposition 3.12](#) again, Y' represents the fiber product $Y \times_X X'$. \square

Remark 4.13. The original proof in [\[Ber93\]](#) does not make any sense to me. Please contact me if you understand the details of Berkovich's argument.

In a similar vein, we prove

Proposition 4.14. If K/k is an analytic field extension, then there is a natural functor of base extension $k_H\text{-An} \rightarrow K_H\text{-An}$ extending the functor $k_H\text{-Aff} \rightarrow K_H\text{-Aff}$ defined by $\text{Sp } A \mapsto \text{Sp } A \hat{\otimes}_k K$.

We will denote the image of a k_H -analytic space X by X_K .

PROOF. Fix a k_H -analytic space X , we want to construct functorially a K_H -analytic space X_K .

Step 1. We assume that X is paracompact and Hausdorff.

By [Proposition 4.11](#), we can find a locally finite G-covering $\{X_i\}_{i \in I}$ of X consisting of k_H -affinoid domains in X . We can glue $X_{i,K}$'s by [Proposition 2.31](#) to get X_K .

Step 2. In general, let $\{Y_i\}_{i \in I}$ be an open covering of X by paracompact Hausdorff subsets. We glue $Y_{i,K}$'s by [Proposition 2.31](#) to get X_K .

These constructions are clearly functorial and defines a functor $k_H\text{-An} \rightarrow K_H\text{-An}$. \square

5. Closed immersions

Let $(k, |\bullet|)$ be a complete non-Archimedean valued field and H be a subgroup of $\mathbb{R}_{>0}$ such that $|k^\times| \cdot H \neq \{1\}$.

Lemma 5.1. Let $\varphi : Y \rightarrow X$ be a morphism of k_H -analytic spaces. Then the following are equivalent:

- (1) for any $x \in X$, there are $n \in \mathbb{Z}_{>0}$ and k_H -affinoid domains V_1, \dots, V_n in X containing x such that $V_1 \cup \dots \cup V_n$ is a neighbourhood of x in X and the restriction $\varphi^{-1}(V_i) \rightarrow V_i$ is a closed immersion for any $i = 1, \dots, n$;
- (2) for any k_H -affinoid domain V in X , $\varphi^{-1}(V) \rightarrow V$ is a closed immersion.

Recall that closed immersions between k_H -affinoid spaces are defined in [Definition 12.1](#) in [Affinoid algebras](#).

The statement in [\[Ber93, Lemma 1.3.7\]](#) is not correct.

PROOF. Only (1) \implies (2) is non-trivial. Assume (1). Let τ be the collections of $V \subseteq X$ satisfying (2). Then we claim that τ is a net.

Observe that τ is a quasi-net by our assumption. To see that it is a net, take $U, V \in \tau$ and $x \in U \cap V$, then we can find $n \in \mathbb{Z}_{>0}$ and k_H -affinoid domains W_1, \dots, W_n in $U \cap V$ containing x such that $W_1 \cup \dots \cup W_n$ is a neighbourhood of x in $U \cap V$. In order to show that $\tau|_{U \cap V}$ is a quasi-net, it suffices to show that $\varphi^{-1}(W_i) \rightarrow W_i$ is a closed immersion for $i = 1, \dots, n$. This follows from [Proposition 12.3](#) in [Affinoid algebras](#).

Let V be a k_H -affinoid domain in X . By (1) and the compactness of V , we can find $n \in \mathbb{Z}_{>0}$ and $V_1, \dots, V_n \in \tau$ such that $V \subseteq V_1 \cup \dots \cup V_n$. By [Lemma 2.5](#) in [Topology and bornology](#), we can find $m \in \mathbb{Z}_{>0}$ and $U_1, \dots, U_m \in \tau$ such that

$$V = U_1 \cup \dots \cup U_m$$

and each U_j is contained in some V_i , where $j = 1, \dots, m$ and $i = 1, \dots, n$. By [Proposition 12.3](#) in [Affinoid algebras](#) again, $U_j \in \tau$ for each $j = 1, \dots, m$. It suffices to apply [Corollary 4.8](#) to conclude that $V \in \tau$. \square

Definition 5.2. Let $\varphi : Y \rightarrow X$ be a morphism of k_H -analytic spaces. We say φ is a *closed immersion* if the equivalent conditions in [Lemma 5.1](#) are satisfied.

Observe that this definition extends [Definition 12.1](#) in [Affinoid algebras](#).

Proposition 5.3. Let $\varphi : Y \rightarrow X$, $\psi : Z \rightarrow X$ be a morphism of k_H -analytic spaces. Assume that $\varphi : Y \rightarrow X$ is a closed immersion. Consider the Cartesian diagram

$$\begin{array}{ccc} Z \times_X Y & \longrightarrow & Y \\ \downarrow & \square & \downarrow \varphi \\ Z & \xrightarrow{\psi} & X \end{array}$$

Then $Z \times_X Y \rightarrow Z$ is a closed immersion.

PROOF. Taking a G -covering of Z , we may assume that Z is compact. We could cover the images of Z in X by finitely many k_H -affinoid domains V_1, \dots, V_n in X , considering their preimages in Z , we could reduce to the case where the image of Z in X is contained in a k_H -affinoid domain. We could then assume that X is a k_H -affinoid space and hence so is Y . By taking a G -covering of Z again, we may assume that Z is affinoid. It suffices to apply [Proposition 12.3](#) in [Affinoid algebras](#). \square

Proposition 5.4. Let $\varphi : Y \rightarrow X$ be a morphism of k_H -analytic spaces. Then the following are equivalent:

- (1) φ is a closed immersion;
- (2) for any G-covering $\{X_i\}_{i \in I}$ of X , the restriction of φ to $\varphi^{-1}(X_i) \rightarrow X_i$ is a closed immersion for all $i \in I$;
- (3) for some G-covering $\{X_i\}_{i \in I}$ of X , the restriction of φ to $\varphi^{-1}(X_i) \rightarrow X_i$ is a closed immersion for all $i \in I$.

In other words, being a closed immersion is a G-local property on the target.

PROOF. Assume (1). Let $\{X_i\}_{i \in I}$ be a G-covering of X . Then the restriction of φ to $\varphi^{-1}(X_i) \rightarrow X_i$ is a closed immersion for all $i \in I$ by [Proposition 5.3](#). So (2) holds.

(2) trivially implies (3).

Assume (3). Using the fact that (1) implies (2) as we already proved, we may refine the G-covering $\{X_i\}_{i \in I}$ and assume that each X_i is k_H -affinoid. It follows from [Lemma 5.1](#) that φ is a closed immersion, so (1) holds. \square

Corollary 5.5. Let $H' \supseteq H$ is a subgroup of $\mathbb{R}_{>0}$. Let $\varphi : Y \rightarrow X$ be a morphism of k_H -analytic spaces. Then the following are equivalent:

- (1) φ is a closed immersion;
- (2) φ is a closed immersion when view as a morphism of $k_{H'}$ -affinoid spaces.

PROOF. By [Proposition 5.4](#), we may assume that X is k_H -affinoid. In this case, Y is also k_H -affinoid and the result is clear. \square

Corollary 5.6. Let $\varphi : Y \rightarrow X$ be a morphism of k_H -analytic spaces and K/k be an analytic field extension.

- (1) If φ is a closed immersion, so is φ_K ;
- (2) If $K = k_r$ for some k -free polyray r and φ_K is a closed immersion, then so is φ .

PROOF. By [Proposition 5.4](#), we may assume that X is a k_H -affinoid space in both cases. Then so is Y . Now (1) is obvious and (2) follows from [Proposition 3.11](#) in [Affinoid algebras](#). \square

Proposition 5.7. Let $\varphi : X \rightarrow Y$, $\psi : Y \rightarrow Z$ be closed immersions of k_H -affinoid spaces. Then $\psi \circ \varphi : X \rightarrow Z$ is also a closed immersion.

PROOF. By [Proposition 5.4](#), we may assume that Z is k_H -affinoid, so Y and X are also k_H -affinoid. In this case, the result is clear, as the composition of admissible epimorphisms is clearly admissible epimorphic. \square

Proposition 5.8. Let $\varphi : Y \rightarrow X$ be a morphism of k_H -analytic spaces. Then the following are equivalent:

- (1) φ is a closed immersion;
- (2) $\varphi_* \mathcal{O}_Y$ is a coherent \mathcal{O}_X -module and $\mathcal{O}_X \rightarrow \varphi_* \mathcal{O}_Y$ is surjective.

PROOF. As both properties are G-local on X , we may assume that X is a k_H -affinoid space and hence so is Y . This result then follows from [Corollary 4.8](#). \square

6. Separated morphisms

Let $(k, |\bullet|)$ be a complete non-Archimedean valued field and H be a subgroup of $\mathbb{R}_{>0}$ such that $|k^\times| \cdot H \neq \{1\}$.

Definition 6.1. Let $\varphi : X \rightarrow Y$ be a morphism of k_H -analytic spaces. The *diagonal morphism* of φ is the morphism $\Delta_\varphi = \Delta_{X/Y} : X \rightarrow X \times_Y X$ defined as follows: let $\{Y_i\}_{i \in I}$ be a G-covering of Y by k_H -affinoid domains and $\{X_{ij}\}_{j \in J_i}$ be a G-covering of $\varphi^{-1}(Y_i)$ by k_H -affinoid domains in X . Then we have a diagonal morphism $\Delta_{X_{ij}/Y_i} : X_{ij} \rightarrow X_{ij} \times_{Y_i} X_{ij}$ defined by the codiagonal morphism of k_H -affinoid algebras. The induced morphisms $X_{ij} \rightarrow X \times_Y X$ can be glued together by [Proposition 3.12](#) to get Δ_φ . By [Proposition 3.12](#) does not depend on the choices of the G-coverings.

Definition 6.2. A morphism $\varphi : X \rightarrow Y$ of k_H -analytic spaces is *separated* if $\Delta_{X/Y} : X \rightarrow X \times_Y X$ is a closed immersion.

Example 6.3. A morphism between k_H -affinoid spaces is always separated. This follows from [Example 12.2](#) in [Affinoid algebras](#) by base change.

Proposition 6.4. Let $\varphi : Y \rightarrow X$, $\psi : Z \rightarrow X$ be a morphism of k_H -analytic spaces. Assume that $\varphi : Y \rightarrow X$ is separated. Consider the Cartesian diagram

$$\begin{array}{ccc} Z \times_X Y & \longrightarrow & Y \\ \downarrow & \square & \downarrow \varphi \\ Z & \xrightarrow{\psi} & X \end{array}$$

Then $Z \times_X Y \rightarrow Z$ is separated.

PROOF. By general abstract nonsense, we have a Cartesian diagram

$$\begin{array}{ccc} Z \times_X Y & \xrightarrow{\Delta_{Z \times_X Y/Z}} & (Z \times_X Y) \times_Z (Z \times_X Y) = Z \times_X (Y \times_X Y) \\ \downarrow & \square & \downarrow \\ Y & \xrightarrow{\Delta_{Y/X}} & Y \times_X Y \end{array}$$

So the assertion follows from [Proposition 5.3](#). \square

Proposition 6.5. Let $\varphi : Y \rightarrow X$ be a morphism of k_H -analytic spaces. Then the following are equivalent:

- (1) φ is separated;
- (2) for any G-covering $\{X_i\}_{i \in I}$ of X , the restriction of φ to $\varphi^{-1}(X_i) \rightarrow X_i$ is separated for all $i \in I$;
- (3) for some G-covering $\{X_i\}_{i \in I}$ of X , the restriction of φ to $\varphi^{-1}(X_i) \rightarrow X_i$ is separated for all $i \in I$.

PROOF. (1) \implies (2) by [Proposition 6.4](#).

(2) \implies (3) is trivial.

Assume (3). Let $Y_i = \varphi^{-1}(X_i)$. Then $Y_i \times_{X_i} Y_i$ is a G-covering of $Y \times_X Y$, and we have a Cartesian diagram

$$\begin{array}{ccc} Y_i & \xrightarrow{\Delta_{Y_i/X_i}} & Y_i \times_{X_i} Y_i \\ \downarrow & \square & \downarrow \\ Y & \xrightarrow{\Delta_{Y/X}} & Y \times_X Y \end{array}$$

for $i \in I$. So the assertion follows from [Proposition 5.4](#). \square

Proposition 6.6. Let $\varphi : X \rightarrow Y$, $\psi : Y \rightarrow Z$ be separated morphisms of k_H -affinoid spaces. Then $\psi \circ \varphi : X \rightarrow Z$ is also separated.

PROOF. We have a Cartesian diagram

$$\begin{array}{ccc} X \times_Y X & \xrightarrow{\psi} & X \times_Z X \\ \downarrow & \square & \downarrow \\ Y & \xrightarrow{\Delta_{Y/Z}} & Y \times_Z Y \end{array}.$$

By [Proposition 6.4](#), $\psi : X \times_Y X \rightarrow X \times_Z X$ is a closed immersion. On the other hand, $\Delta_{X/Z} : X \rightarrow X \times_Z X$ factorizes as $\psi \circ \Delta_{X/Y}$. It follows from [Proposition 5.7](#) that $\Delta_{X/Z}$ is a closed immersion. \square

Proposition 6.7. Let $H' \supseteq H$ is a subgroup of $\mathbb{R}_{>0}$. Let $\varphi : Y \rightarrow X$ be a morphism of k_H -analytic spaces. Then the following are equivalent:

- (1) φ is separated;
- (2) φ is separated when view as a morphism of $k_{H'}$ -affinoid spaces.

PROOF. This follows immediately from [Corollary 5.5](#). \square

Proposition 6.8. Let $\varphi : Y \rightarrow X$ be a morphism of k_H -analytic spaces and K/k be an analytic field extension.

- (1) If φ is separated, so is φ_K ;
- (2) If $K = k_r$ for some k -free polyray r and φ_K is separated, then so is φ .

We will prove later on that the assumption in (2) is unnecessary.

PROOF. This follows immediately from [Corollary 5.6](#). \square

7. Analytic germs

Let $(k, |\bullet|)$ be a complete non-Archimedean valued field and H be a subgroup of $\mathbb{R}_{>0}$ such that $|k^\times| \cdot H \neq \{1\}$.

Definition 7.1. A *punctured k_H -analytic space* (X, x) is a k_H -analytic space X together with a point $x \in X$.

A morphism between punctured k_H -analytic spaces (X, x) and (Y, y) is a morphism $\varphi : X \rightarrow Y$ of k_H -analytic spaces sending x to y .

The category of punctured k_H -analytic spaces is denoted by $k_H\text{-}\mathcal{An}_*$.

Definition 7.2. A morphism of punctured k_H -analytic spaces $(X, x) \rightarrow (Y, y)$ is said to be *separated* (resp. a *closed immersion*) if the underlying morphism of k_H -analytic spaces is separated (resp. a closed immersion).

Definition 7.3. The category $k_H\text{-}\mathcal{Ger}$ is the category of right fractions of $k_H\text{-}\mathcal{An}_*$ with respect to the system of morphisms

$$\varphi : (X, x) \rightarrow (Y, y)$$

that induces an isomorphism of X with an open neighbourhood of y in Y in $k\text{-}\mathcal{Ger}$.

When we view (X, x) as an object in $k_H\text{-}\mathcal{Ger}$, we write it as X_x . An object in $k_H\text{-}\mathcal{Ger}$ is called a k_H -analytic germ.

Be careful, we require φ to induce an isomorphism in $k\text{-Ger}$ instead of $k_H\text{-Ger}$, although eventually, we will show that these notions coincide.

By definition,

$$\text{Hom}_{k_H\text{-Ger}}(X_x, Y_y) = \varinjlim_U \text{Hom}_{k_H\text{-An}^*}((U, x), (Y, y)),$$

where U runs over all open neighbourhoods of x in X .

Definition 7.4. A k_H -analytic germ X_x is *good* if x admits an affinoid neighbourhood in X .

Note that this condition does not depend on the representative (X, x) . To see this, let $U \subseteq x$ be an open subset containing x . We need to show that if x admits a k_H -affinoid neighbourhood in X , then it admits one in U . This follows from [Proposition 10.13](#) in [Affinoid algebras](#).

Definition 7.5. A morphism of k_H -analytic germs $\varphi : X_x \rightarrow Y_y$ is said to be *separated* (resp. *boundaryless*, *a closed immersion*) if it is induced by a separated morphism (resp. boundaryless, resp. a closed immersion) of punctured k_H -analytic spaces $(U, x) \rightarrow (Y, y)$, where U is an open neighbourhood of x in X .

Definition 7.6. Let X_x be a k_H -analytic germ. A k_H -analytic domain in X_x is a k_H -analytic germ V_x , where V is a k_H -analytic domain in X containing x .

We say a finite family of k_H -analytic germs $\{V_{ix}\}_{i \in I}$ *covers* X_x if there is a representative (X, x) of X_x such that V_{ix} can be represented by a k_H -analytic domain $V_i \subseteq X$ for $i \in I$ and

$$X = \bigcup_{i \in I} V_i.$$

Definition 7.7. Let $\phi : Y_y \rightarrow X_x$ be a morphism of k_H -analytic germs and V_x be a k_H -analytic domain in X_x . Represent ϕ by a morphism $\phi : (Y, y) \rightarrow (X, x)$ and represent V_x by a k_H -analytic domain V in X . Then the k_H -analytic domain $\phi^{-1}(V)$ in Y determines a k_H -analytic germ $\phi^{-1}(V)_y$, which does not depend on the choices we made. This k_H -analytic germ is denoted by $\phi^{-1}(V_x)$.

Recall that $\phi^{-1}(V)$ is a k_H -analytic domain in Y by [Proposition 3.5](#).

Definition 7.8. Let X be a good k_H -analytic space and $x \in X$, we define

$$\mathcal{O}_{X,x} := \varinjlim_V A_V,$$

where V runs over all k_H -affinoid neighbourhoods of x in X . [Include the definition of affinoid neighbourhoods](#)

Observe that k_H -affinoid neighbourhoods of x in X are cofinal in the directed set of k -affinoid neighbourhoods of x in X . This follows from [Proposition 10.13](#) in [Affinoid algebras](#). So we may let V runs over all k -affinoid neighbourhoods of x in X as well.

Example 7.9. Let X_x be a k_H -analytic germ. Take a k_H -affinoid domain $\text{Sp } A$ of X containing x . Given $r \in \sqrt{|k^\times|} \cdot \bar{H}^n$ and $f \in A^n$, we write

$$X_x\{r^{-1}f\} := (\text{Sp } A\{r^{-1}f\})_x.$$

Then $X_x\{r^{-1}f\}$ is a k_H -analytic germ. Observe that $X_x\{r^{-1}f\}$ is independent of the choice of $\text{Sp } A$. This construction depends only on the classes of f in $\mathcal{O}_{X,x}^n$.

Given $\bar{f} \in \mathcal{O}_{X,x}^n$, we define $X_x\{r^{-1}\bar{f}\} = X_x\{r^{-1}f\}$ for any $f \in A^n$ lifting \bar{f} as above.

8. Reduction

Let $(k, |\bullet|)$ be a complete non-Archimedean valued field and H be a subgroup of $\mathbb{R}_{>0}$ such that $|k^\times| \cdot H \neq \{1\}$.

In this section, when we do not specify the grading of a graded object, we mean it is $\mathbb{R}_{>0}$ -graded. In particular \tilde{k} means $\tilde{k}^{\mathbb{R}_{>0}}$.

Definition 8.1. Let $X = \mathrm{Sp} A$ be a k -affinoid space and $x \in X$, we define the *reduction* $\widetilde{(X, x)}$ of X at x as follows: let $\chi_x : A \rightarrow \mathcal{H}(x)$ be the character corresponding to x , we define

$$\widetilde{(X, x)} := \mathbf{P}_{\widetilde{\mathcal{H}(x)/\tilde{k}}} \{ \widetilde{\chi_x(\tilde{A})} \} \subseteq \mathbf{P}_{\widetilde{\mathcal{H}(x)/\tilde{k}}}.$$

Observe that $\widetilde{(X, x)}$ is an affine open subset of $\mathbf{P}_{\widetilde{\mathcal{H}(x)/\tilde{k}}}$. This follows from [Corollary 11.11](#) in [Affinoid algebras](#).

Lemma 8.2. Let $X = \mathrm{Sp} A$ be a k -affinoid space and $x \in X$. Let $U = \mathrm{Sp} B$ be a k -affinoid space. Let $\iota : U \rightarrow X$ be an isomorphism of U with an open neighbourhood of x . We still write $\iota^{-1}(x) \in U$ as x . Then the natural morphism

$$\widetilde{(U, x)} = \widetilde{(X, x)}.$$

PROOF. We first recall that $\mathcal{H}(x)$ does not depend on if we view x as in $\mathrm{Sp} A$ or in $\mathrm{Sp} B$ by [Corollary 13.16](#) in [Affinoid algebras](#).

Observe that the morphism $\chi_x : B \rightarrow \mathcal{H}(x)$ is boundaryless with respect to A by [Proposition 15.9](#) in [Affinoid algebras](#). By [Proposition 15.2](#) in [Affinoid algebras](#), $\widetilde{\chi_x(\tilde{B})}$ is finite over $\widetilde{\chi_x(\tilde{A})}$. By [Lemma 4.5](#) in [Commutative algebras](#), we have

$$\mathbf{P}_{\widetilde{\mathcal{H}(x)/\tilde{k}}} \{ \widetilde{\chi_x(\tilde{A})} \} = \mathbf{P}_{\widetilde{\mathcal{H}(x)/\tilde{k}}} \{ \widetilde{\chi_x(\tilde{B})} \}.$$

□

Definition 8.3. Let X_x be a good k -analytic germs. Take an affinoid neighbourhood U of x in X , then we define

$$\widetilde{X_x} := \widetilde{(U, x)} \subseteq \mathbf{P}_{\widetilde{\mathcal{H}(x)/\tilde{k}}}.$$

By [Lemma 8.2](#), $\widetilde{X_x}$ depends only on X_x .

The construction is clearly functorial in X_x .

Lemma 8.4. Let X_x be a good k -analytic germ and Y_x be a k -analytic domain in X_x . Then Y_x can be covered by finitely many k -analytic domains in X_x of the form

$$X_x \{ r^{-1} f \},$$

where $n \in \mathbb{N}$, $f = (f_1, \dots, f_n) \in \mathcal{O}_{X,x}^\times$ is a tuple of invertible elements and $r_i = |f_i(x)|$.

PROOF. We may assume that X is k -affinoid, say $X = \mathrm{Sp} A$. By [Corollary 12.8](#) in [Affinoid algebras](#), Y can be covered by finitely many k -rational domains in X , say of the form $\mathrm{Sp} A\{r^{-1}g/h\}$, where $m \in \mathbb{N}$, $r = (r_1, \dots, r_m) \in \mathbb{R}_{>0}^m$, $g = (g_1, \dots, g_m) \in A^m$, $h \in A$ and g_1, \dots, g_m, h generates the unit ideal. We may assume that $Y = \mathrm{Sp} A\{r^{-1}g/h\}$.

By shrinking X , we may assume that h is invertible. Set $f_i = g_i/h$, then

$$Y = \operatorname{Sp} A\{r_1^{-1}f_1, \dots, r_m^{-1}f_m\}.$$

By further shrinking X , it suffices to consider those i with $|f_i(x)| = r_i$. \square

Lemma 8.5. Let X_x be a good k -analytic germ. Given $n \in \mathbb{N}$ and $f = (f_1, \dots, f_n) \in \mathcal{O}_{X,x}^\times$, then

$$\widetilde{X_x\{r^{-1}f\}} = \widetilde{X_x}\{\widetilde{\chi_x}(f_1), \dots, \widetilde{\chi_x}(f_n)\},$$

where $r = (r_1, \dots, r_n)$ and $r_i = |f_i(x)|$ for $i = 1, \dots, n$.

PROOF. We may assume that X is k -affinoid, say $X = \operatorname{Sp} A$. By induction on n , we may assume that $n = 1$. Consider the admissible epimorphism

$$\phi : A\{r^{-1}T\} \rightarrow A\{r^{-1}f\}$$

sending T to f . By [Theorem 11.10](#) in [Affinoid algebras](#),

$$\tilde{\phi} : \tilde{A}[r^{-1}T] \rightarrow \widetilde{A\{r^{-1}f\}}$$

is finite. Let $\chi_x : A\{r^{-1}f\} \rightarrow \mathcal{H}(x)$ be the character defined by x .

Then $\widetilde{\chi_x(A\{r^{-1}f\})}$ is finite over $\widetilde{\chi_x(\tilde{A})[f]}$. So the assertion follows from [Lemma 4.5](#) in [Commutative algebras](#). \square

Lemma 8.6. Let X_x be a good k -analytic germs and Y_x be a good k -analytic domain in X_x . Then we can find $n \in \mathbb{N}$, $f_1, \dots, f_n \in \mathcal{O}_{X,x}^\times$ such that

$$\widetilde{Y_x} = \widetilde{X_x}\{\widetilde{\chi_x}(f_1), \dots, \widetilde{\chi_x}(f_n)\}.$$

In particular, we can identify $\widetilde{Y_x}$ with an open subset of $\widetilde{X_x}$.

PROOF. The same argument as in [Lemma 8.4](#) that we can assume that $X = \operatorname{Sp} A$ and $Y = \operatorname{Sp} A\{r^{-1}f\}$ for some $n \in \mathbb{N}$, $r = (r_1, \dots, r_n) \in \mathbb{R}_{>0}^n$, $f = (f_1, \dots, f_n) \in A^n$ with $r_i = |f_i(x)|$ for $i = 1, \dots, n$. So the assertion follows from [Lemma 8.5](#). \square

Lemma 8.7. Let X_x be a good k -analytic germ, $n \in \mathbb{Z}_{>0}$ and Y_{1x}, \dots, Y_{nx} be a covering of X_x by good k -analytic domains. Then

$$\widetilde{X_x} = \bigcup_{i=1}^n \widetilde{Y_{ix}}.$$

PROOF. Observe that we are free to replace $\{Y_{ix}\}_i$ by its refinements by coverings by good k -analytic domains. We may assume that X is k -affinoid, say $X = \operatorname{Sp} A$. Then by [Lemma 13.3](#) in [Affinoid algebras](#), we may assume that the covering is k -rational and is generated by $r_1^{-1}f_1, \dots, r_n^{-1}f_n$. Up to shrinking X , we may guarantee that $|f_i(x)| = r_i$ for $i = 1, \dots, n$. In this case, the assertion follows from [Lemma 8.5](#). \square

Lemma 8.8. Let $\phi : Y_y \rightarrow X_x$ be a morphism of good k -analytic germs. Let X'_x be a good k -analytic domain in X_x and set $Y'_y = \phi^{-1}(X'_x)$, then

$$\widetilde{Y'_y} = \tilde{\phi}^{-1}(\widetilde{X'_x}).$$

PROOF. By [Lemma 8.4](#), we may find $m \in \mathbb{Z}_{>0}$, $n_1, \dots, n_m \in \mathbb{N}$, $g_{i1}, \dots, g_{in_i} \in \mathcal{O}_{X,x}^\times$ for $i = 1, \dots, m$ such that X'_x is covered by $X\{r_{i1}^{-1}g_{i1}, \dots, r_{in_i}^{-1}g_{in_i}\}$ for $i = 1, \dots, m$, where $r_{ij} = |g_{ij}(x)|$ for $i = 1, \dots, m$, $j = 1, \dots, n_i$.

Then Y'_x is covered by $Y\{r_{i1}^{-1}g'_{i1}, \dots, r_{in_i}^{-1}g'_{in_i}\}$ for $i = 1, \dots, m$, where g'_{ij} is the image of g_{ij} in $\mathcal{O}_{Y,y}^\times$ for $i = 1, \dots, m$, $j = 1, \dots, n_i$.

By [Lemma 8.5](#), we have

$$\widetilde{X}'_x = \bigcup_{i=1}^m \widetilde{X}_x \{ \widetilde{\chi}_x(\widetilde{g_{i1}}), \dots, \widetilde{\chi}_x(\widetilde{g_{in_i}}) \}$$

and

$$\widetilde{Y}'_y = \bigcup_{i=1}^m \widetilde{Y}_y \{ \widetilde{\chi}_y(\widetilde{g'_{i1}}), \dots, \widetilde{\chi}_y(\widetilde{g'_{in_i}}) \}.$$

Our assertion is now clear. \square

Definition 8.9. Let X_x be a k -analytic germ. By [Lemma 8.6](#), the reduction defines a functor from the category of good k -analytic germs in X_x (with inclusions as the morphisms) to the category of open affine subsets of $\mathbf{P}_{\widetilde{\mathcal{H}(x)/k}}$. We define

$$\widetilde{X}_x := \varinjlim_{Y_x} \widetilde{Y}_x,$$

where Y_x runs over the filtered category of good k -analytic germs in X_x and the colimit is taken in the category $\mathcal{T}_{\widetilde{\mathcal{H}(x)/k}}$.

The object \widetilde{X}_x is called its *reduction* of X_x .

Theorem 8.10. Let X_x be a k -analytic germ. Then the reduction functor

$$k\text{-Ger} \rightarrow \mathcal{T}_{\widetilde{\mathcal{H}(x)/k}}$$

establishes a bijection between the k -analytic domains and non-empty quasi-compact open subsets of \widetilde{X}_x .

This bijection commutes with finite unions and finite intersections.

In [\[Tem04\]](#), the author forgot the non-emptiness assumption.

PROOF. The last assertion is obvious by construction.

Step 1. We prove the theorem under the additional assumption that X_x is good.

Step 1.1. Let $l, m \in \mathbb{N}$ and $f = (f_1, \dots, f_l) \in \mathcal{O}_{X,x}^l$, $g = (g_1, \dots, g_m) \in \mathcal{O}_{X,x}^m$. Assume that

$$\widetilde{X}_x\{\widetilde{f}\} \subseteq \widetilde{X}_x\{\widetilde{g}\},$$

then we prove that

$$X_x\{r^{-1}f\} \subseteq X_x\{s^{-1}g\},$$

where $r = (r_1, \dots, r_l)$, $s = (s_1, \dots, s_m)$,

$$r_i = |f_i(x)|, \quad s_j = |g_j(x)|$$

for $i = 1, \dots, l$, $j = 1, \dots, m$.

We may assume that X is k -affinoid, say $X = \text{Sp } A$ and $f_1, \dots, f_l, g_1, \dots, g_m \in A$. Let $\chi_x : A \rightarrow \mathcal{H}(x)$ be the character of x . Let

$$B = \widetilde{\chi}_x(\widetilde{A}) \subseteq \widetilde{\mathcal{H}(x)}.$$

By definition,

$$\widetilde{X}_x = \mathbf{P}_{\widetilde{\mathcal{H}}(x)}\{B\}.$$

By [Lemma 8.5](#), we have

$$\begin{aligned} X_x\{\widetilde{r^{-1}f}\} &= \widetilde{X}_x\{B[\tilde{f}]\}, \\ X_x\{\widetilde{r^{-1}f, s^{-1}g}\} &= \widetilde{X}_x\{B[\tilde{f}, \tilde{g}]\}. \end{aligned}$$

The right-hand sides are equal by our assumption, so by [Lemma 4.5](#) in [Commutative algebras](#), $B[\tilde{f}, \tilde{g}]$ is finite over $B[\tilde{f}]$. We take monic polynomials of \tilde{g}_j over $B[\tilde{f}]$:

$$T^{n_j} + \tilde{a}_{j,1}T^{n_j-1} + \cdots + \tilde{a}_{j,n_j} \in B[\tilde{f}][T]$$

with $\tilde{a}_{j,1}, \dots, \tilde{a}_{j,n_j}$ homogeneous of degree $|g_j(x)|^1, \dots, |g_j(x)|^{n_j}$ respectively. This is possible by [Proposition 2.18](#) in [Commutative algebras](#). We lift $\tilde{a}_{j,k}$ to $a_{j,k} \in A\{r^{-1}f\}$ with $\rho(a_{j,k}) = \rho(g_j)^k$ for $j = 1, \dots, m$, $k = 1, \dots, n_j$. It follows that

$$|(g_j^{n_j} + a_{j,1}g_j^{n_j-1} + \cdots + a_{j,n_j})(x)| < |g_j(x)|^n$$

for $j = 1, \dots, m$. Up to shrinking X , we may assume that this inequality holds everywhere on $X\{r^{-1}f\}$.

By then $|g_j(y)| \leq |g_j(x)|$ for any $y \in X\{r^{-1}f\}$. Our assertion follows.

Step 1.2. Suppose that Y_x is a k -analytic domain in X_x with $\tilde{Y}_x = \widetilde{X}_x$, then $Y_x = X_x$.

We may assume that X is k -affoid, say $X = \mathrm{Sp} A$.

By [Lemma 8.4](#), we can write Y_x as a finite union of $V_{i,x} := X_x\{r_i^{-1}f_i\}$ for $i = 1, \dots, m$, where $n_i \in \mathbb{N}$, $f_i = (f_{i1}, \dots, f_{in_i}) \in \mathcal{O}_{X,x}^{\times, n_i}$ and $r_i = (r_{i1}, \dots, r_{in_i})$ with $r_{ij} = |f_{ij}(x)|$.

By [Lemma 8.7](#), $\widetilde{V}_{i,x}$ for $i = 1, \dots, m$ covers \widetilde{X}_x .

By [Lemma 4.8](#) in [Commutative algebras](#), we can refine this covering to a Laurent covering

$$\mathcal{U} := \left\{ \tilde{U}_j = \widetilde{X}_x\{\tilde{g}_1^{j_1}, \dots, \tilde{g}_l^{j_l}\} \right\}_{j=(j_1, \dots, j_l) \in \{\pm 1\}^l},$$

where $l \in \mathbb{N}$ and $\tilde{g}_1, \dots, \tilde{g}_l$ are homogeneous elements in $\widetilde{\mathcal{H}}(x)$. Lift $\tilde{g}_1, \dots, \tilde{g}_l$ to $g_1, \dots, g_l \in A$. We consider the k -Laurent covering of X generated by

$$\rho(\tilde{g}_1)^{-1}g_1, \dots, \rho(\tilde{g}_l)^{-1}g_l.$$

The reduction of this covering is clearly \mathcal{U} . By Step 1.1, the germs of \mathcal{U} at x is a refinement of $\{V_{1,x}, \dots, V_{m,x}\}$, so the latter is a covering of X_x , namely $X_x \xrightarrow{\sim} Y_x$.

Step 1.3. We prove that each quasi-compact open subset \tilde{Y}_x of \widetilde{X}_x is the reduction of some k -analytic domain Y_x in X_x .

We can write

$$\tilde{Y}_x = \bigcup_{i=1}^m \widetilde{X}_x\{\widetilde{f_{i1}}, \dots, \widetilde{f_{in_i}}\},$$

where $m \in \mathbb{Z}_{>0}$, $n_1, \dots, n_m \in \mathbb{N}$, $\widetilde{f_{ij}} \in \widetilde{\mathcal{H}}(x)$ are homogeneous elements for $i = 1, \dots, m$, $j = 1, \dots, n_i$. We lift $\widetilde{f_{ij}}$ to $f_{ij} \in \mathcal{O}_{X,x}$, it suffices to take

$$\bigcup_{i=1}^m X_x\{r_{i1}^{-1}f_{i1}, \dots, r_{in_i}^{-1}f_{in_i}\},$$

where $r_{ij} = |f_{ij}(x)|$ for $i = 1, \dots, m$, $j = 1, \dots, n_i$.

Step 1.4. Suppose that Y_x, Z_x are k -analytic domains in X_x with $\widetilde{Y}_x = \widetilde{Z}_x$. Then we prove that $Y_x = Z_x$.

Take $p, q \in \mathbb{N}$, good k -analytic domains Y_x^1, \dots, Y_x^p in Y_x and good k -analytic domains Z_x^1, \dots, Z_x^q in Z_x such that

$$\widetilde{Y}_x = \bigcup_{i=1}^p \widetilde{Y}_x^i = \bigcup_{i=1}^q \widetilde{Z}_x^i.$$

Therefore, for any $i = 1, \dots, p$, $\{\widetilde{Y}_x^i \cap \widetilde{Z}_x^j\}_{j=1, \dots, q}$ is a covering of \widetilde{Y}_x^i . By Step 1.2, $\{Y_x^i \cap Z_x^j\}_{j=1, \dots, q}$ is a covering of Y_x^i for $i = 1, \dots, p$. So $Y_x \subseteq Z_x$. By symmetry $Y_x = Z_x$.

We have finished the proof when X_x is good.

Step 2. We handle the general case.

Step 2.1. We prove that each quasi-compact open subset \widetilde{Y}_x of \widetilde{X}_x is the reduction of some k -analytic domain Y_x in X_x .

Take $p \in \mathbb{N}$, good k -analytic domains X_x^1, \dots, X_x^p in X_x such that

$$\widetilde{X}_x = \bigcup_{i=1}^p \widetilde{X}_x^i.$$

By Step 1, $\widetilde{Y}_x \cap \widetilde{X}_x^i$ can be lifted to a k -analytic domain Y_x^i in X_x^i for $i = 1, \dots, p$. The union of Y_x^i 's for $i = 1, \dots, p$ is a lifting of \widetilde{Y}_x .

Step 2.2. Suppose that Y_x, Z_x are k -analytic domains in X_x with $\widetilde{Y}_x = \widetilde{Z}_x$. Then we prove that $Y_x = Z_x$.

For each $i = 1, \dots, p$, we have

$$\widetilde{Y_x \cap X_x^i} = \widetilde{Y_x} \cap \widetilde{X_x^i} = \widetilde{Z_x} \cap \widetilde{X_x^i} = \widetilde{Z_x \cap X_x^i}.$$

By Step 1, $Y_x \cap X_x^i = Z_x \cap X_x^i$ coincides for $i = 1, \dots, p$, so $Y_x = Z_x$. \square

Corollary 8.11. Let $\varphi : Y_y \rightarrow X_x$ be a morphism of k -analytic germs, then the following are equivalent:

- (1) φ is a closed immersion;
- (2) $\tilde{\varphi} : \widetilde{Y}_y \rightarrow \widetilde{X}_x$ is an isomorphism and φ is represented by a G-locally closed immersion.

Include the notion of G-locally closed immersion somewhere

PROOF. (1) \implies (2): This is obvious.

(2) \implies (1): After shrinking X and Y , we can take a k -analytic domain X' in X , a neighbourhood Y' of y in Y such that $\varphi(Y') \subseteq X'$ and the restriction $Y' \rightarrow X'$ is a closed immersion. It suffices to show that φ is boundaryless at y . In other words, we need to show that $X'_x = X_x$. By [Theorem 8.10](#), this is equivalent to $\widetilde{X}'_x = \widetilde{X}_x$. By (1) \implies (2) direction of this corollary, $\widetilde{X}'_x = \widetilde{Y}'_y$. But clearly $\widetilde{Y}_y = \widetilde{Y}'_y$. So our assertion follows. \square

Lemma 8.12. Let $Y_y \rightarrow X_x, Z_z \rightarrow X_x$ be morphisms of k -analytic germs. Let $T = Y \times_X Z$. Take a point $t \in T$ whose image in Y is y and whose image in Z is z . Then the natural map

$$\widetilde{T}_t \cong \left(\widetilde{Y}_y \times_{\mathbf{P}_{\widetilde{\mathcal{H}}(y)/\tilde{k}} \widetilde{\mathcal{H}}(t)/\tilde{k}} \right) \times_{\left(\widetilde{X}_x \times_{\mathbf{P}_{\widetilde{\mathcal{H}}(x)/\tilde{k}} \widetilde{\mathcal{H}}(t)/\tilde{k}} \right)} \left(\widetilde{Z}_z \times_{\mathbf{P}_{\widetilde{\mathcal{H}}(z)/\tilde{k}} \widetilde{\mathcal{H}}(t)/\tilde{k}} \right)$$

is a homeomorphism.

Existence of t needs to be proved somewhere following Ducros

PROOF. As both sides commute with colimits, we may assume that X, Y, Z are all k -affinoid, say $X = \mathrm{Sp} A$, $Y = \mathrm{Sp} B$ and $Z = \mathrm{Sp} C$.

Let $\chi_x : A \rightarrow \mathcal{H}(x)$ (resp. $\chi_y : B \rightarrow \mathcal{H}(y)$, resp. $\chi_z : C \rightarrow \mathcal{H}(z)$) be the character corresponding to x (resp. y , resp. z). Let \tilde{A}_0 (resp. \tilde{B}_0 , resp. \tilde{C}_0) be the image of $\widetilde{\chi_x(\tilde{A})}$ (resp. $\widetilde{\chi_y(\tilde{B})}$, resp. $\widetilde{\chi_z(\tilde{C})}$) in $\widetilde{\mathcal{H}(t)}$. The character corresponding to t is given by

$$\chi_t : B \hat{\otimes}_A C \rightarrow \mathcal{H}(t).$$

So

$$\tilde{T}_t = \mathbf{P} \widetilde{\mathcal{H}(t)/\bar{k}} \{ \mathrm{Im} \chi_t \}.$$

As $\tilde{B} \otimes_{\tilde{A}} \tilde{C} \rightarrow \widetilde{B \hat{\otimes}_A C}$ is finite by [Lemma 11.13](#) in [Affinoid algebras](#), by [Lemma 4.5](#) in [Commutative algebras](#), we have

$$\tilde{T}_t = \mathbf{P} \widetilde{\mathcal{H}(t)/\bar{k}} \{ \tilde{D}_0 \},$$

where \tilde{D}_0 is the image of the natural map $\tilde{B} \otimes_{\tilde{A}} \tilde{C} \rightarrow \widetilde{\mathcal{H}(t)}$,

We are supposed to prove that

$$\mathbf{P} \widetilde{\mathcal{H}(t)/\bar{k}} \{ \tilde{D}_0 \} \cong \mathbf{P} \widetilde{\mathcal{H}(t)/\bar{k}} \{ \tilde{B}_0 \} \times_{\mathbf{P} \widetilde{\mathcal{H}(t)/\bar{k}} \{ \tilde{A}_0 \}} \mathbf{P} \widetilde{\mathcal{H}(t)/\bar{k}} \{ \tilde{C}_0 \}.$$

Equivalently,

$$\mathbf{P} \widetilde{\mathcal{H}(t)/\bar{k}} \{ \tilde{D}_0 \} \cong \mathbf{P} \widetilde{\mathcal{H}(t)/\bar{k}} \{ \tilde{B}_0, \tilde{C}_0 \}.$$

This is obvious as \tilde{D}_0 is generated by \tilde{B}_0 and \tilde{C}_0 . \square

Corollary 8.13. Let $Y_y \rightarrow X_x$ be a morphism of k -analytic germs and V_x be a k -analytic domain in X_x . Let $W_y = Y_y \times_{X_x} V_x$. Then \widetilde{W}_y is the preimage of \widetilde{V}_y in \widetilde{Y}_y .

PROOF. This follows immediately from [Corollary 8.11](#). \square

Corollary 8.14. Let $\varphi : Y_y \rightarrow X_x$ be a morphism of k -analytic germs, then the following are equivalent:

- (1) φ is separated;
- (2) $\tilde{\varphi} : \widetilde{Y}_y \rightarrow \widetilde{X}_x$ is separated.

PROOF. Observe that the diagonal morphism $\Delta_{Y/X} : Y \rightarrow Y \times_X Y$ is a G-locally closed immersion, so by [Corollary 8.11](#), φ is separated if and only if

$$\widetilde{\Delta_{Y/X}} : \widetilde{Y}_y \rightarrow (\widetilde{Y \times_X Y})_{(y,y)}$$

is an isomorphism.

By [Lemma 8.12](#), the natural map

$$\widetilde{Z}_z \rightarrow \widetilde{Y}_y \times_{X'} \widetilde{Y}_y$$

is a homeomorphism, where

$$X' = \widetilde{X}_x \times_{\mathbf{P} \widetilde{\mathcal{H}(x)/\bar{k}}} \mathbf{P} \widetilde{\mathcal{H}(y)/\bar{k}}.$$

Thus, $\widetilde{\Delta_{Y/X}}$ is an isomorphism if and only if $\widetilde{Y_y} \rightarrow X'$ is injective, namely, $\widetilde{Y_y} \rightarrow \widetilde{X_x}$ is separated. \square

Lemma 8.15. Let X_x be a *good* k -analytic germ. Then the following are equivalent:

- (1) x admits a k -affinoid neighbourhood V in X which admits a k_H -analytic structure;
- (2) $\widetilde{X_x}$ is an H -strict affine open subset of $\mathbf{P}_{\widetilde{\mathcal{H}(x)}/\bar{k}}$.

If Temkin's argument of [Lemma 9.1](#) is OK, we can remove the good assumption!

PROOF. Assume (1). Let $V = \mathrm{Sp} A$ be as in (1). Let $\chi_x : A \rightarrow \mathcal{H}(x)$ be the character defined by x . It follows from [Theorem 8.4](#) in [Affinoid algebras](#) that $\widetilde{\chi_x(A)}$ is contained in $\widetilde{\mathcal{H}(x)}^{\sqrt{|k \times| \cdot H}}$. It follows that $\widetilde{X_x}$ is H -strict by [Corollary 5.4](#) in [Commutative algebras](#).

Assume (2). Take $n \in \mathbb{N}$ and non-zero homogeneous elements $f_1, \dots, f_n \in \widetilde{\mathcal{H}(x)}$ with degree $r_1, \dots, r_n \in H$ such that

$$\widetilde{X_x} = \mathbf{P}_{\widetilde{\mathcal{H}(x)}/\bar{k}}\{f_1, \dots, f_n\}.$$

By assumption, X_x is good. * So we can find a k -affinoid neighbourhood V of x in X . Up to shrinking $V = \mathrm{Sp} B$, we make find invertible elements h_1, \dots, h_n in B such that $\widetilde{h_i(x)} = f_i$ for $i = 1, \dots, n$.

Let $h : V \rightarrow \mathbb{A}_k^n$ be the morphism induced by h_1, \dots, h_n . **Include the functor of points of An** Let $t = h(x)$ and W be the affinoid domain in \mathbb{A}_k^n defined by $|T_i| \leq r_i$ for $i = 1, \dots, n$. We have a commutative diagram

$$\begin{array}{ccc} (V, x) & \xrightarrow{h} & (\mathbb{A}_k^n, t) \\ & \searrow & \uparrow \\ & & (W, t) \end{array}.$$

Also observe that the morphism of k -analytic germs $X_x \rightarrow (\mathbb{A}_k^n)_t$ factorizes through W_t , as can be seen on the level of reduction. So up to shrinking V , we can find a k -affinoid neighbourhood W' of t in \mathbb{A}_k^n such that $h(V) \subseteq W \cap W'$. We may assume that W' is a k_H -analytic domain. As $\widetilde{V_x} = \widetilde{X_x}$ is the preimage of $(W \cap W')_t = \widetilde{W_t}$, the morphism $V \rightarrow W \cap W'$ is boundaryless at x . As $W \cap W'$ is k_H -analytic, it follows from [Proposition 15.5](#) in [Affinoid algebras](#) that x admits a k_H -affinoid neighbourhood in X . \square

Corollary 8.16. Let X_x be a k -analytic germ. The following are equivalent:

- (1) the germ X_x is k_H -analytic;
- (2) the reduction $\widetilde{X_x}$ is H -strict.

We say X_x is k_H -analytic in the sense that it lies in the essential image of $k_H\text{-Ger} \rightarrow k\text{-Ger}$.

PROOF. (1) \implies (2) follows immediately from [Lemma 8.15](#).

* Assume that [Lemma 9.1](#) is correct, by [Theorem 9.2](#), the germ X_x is automatically good.

Assume (2). Let $\{U_i\}_{i \in I}$ be an H -strict atlas of \widetilde{X}_x . For each $i \in I$, we can find a k -analytic domain $X_{i,x}$ in X_x such that $U_i = \widetilde{X_{i,x}}$ by [Theorem 8.10](#). By [Theorem 8.10](#),

$$\widetilde{X_{i,x}} \cap \widetilde{X_{j,x}} = U_i \cap U_j$$

for all $i, j \in I$. So we may assume that \widetilde{X}_x is an H -strict quasi-compact open subset of $\mathbf{P}^{\widetilde{\mathcal{H}(x)}/\bar{k}}$.

Cover \widetilde{X}_x by finitely many H -strict affine open subset V_1, \dots, V_m . By [Theorem 8.10](#), we can lift V_i to a k_H -analytic germ $W_{i,x}$ in X_x for $i = 1, \dots, m$. Moreover $W_{i,x} \cap W_{j,x}$ is k_H -analytic for any $j = 1, \dots, m$. It follows that X_x is k_H -analytic. \square

Definition 8.17. Let X_x be a k_H -analytic germ. We define

$$\widetilde{X}_x^H := \left(\widetilde{X}_x \right)^H.$$

This makes sense by [Corollary 8.16](#).

Proposition 8.18. Let Y_y, X_x be k_H -analytic germs. Then for any morphism $Y_y \rightarrow X_x$ in $k\text{-}\mathcal{G}\text{er}$, there is a unique continuous map $\widetilde{Y}_y^H \rightarrow \widetilde{X}_x^H$ making the diagram commutative:

$$\begin{array}{ccccc}
 & \widetilde{Y}_y & \xrightarrow{\quad} & \widetilde{X}_x & \\
 & \swarrow & & \swarrow & \\
 \widetilde{Y}_y^H & \xrightarrow{\quad} & \widetilde{X}_x^H & & \\
 \downarrow & \cdots & \downarrow & & \downarrow \\
 & \mathbf{P}^{\widetilde{\mathcal{H}(y)}/\bar{k}} & \xrightarrow{\quad} & \mathbf{P}^{\widetilde{\mathcal{H}(x)}/\bar{k}} & \\
 \downarrow & \swarrow & \downarrow & \swarrow & \\
 \mathbf{P}^{\widetilde{\mathcal{H}(y)}^H/\bar{k}^H} & \xrightarrow{\quad} & \mathbf{P}^{\widetilde{\mathcal{H}(x)}^H/\bar{k}^H} & &
 \end{array}$$

PROOF. This follows immediately from [Proposition 6.5](#) in [Commutative algebras](#). \square

Proposition 8.19. Let X_x be a k -analytic germ. If X_x lies in the essential image of $k_H\text{-}\mathcal{G}\text{er} \rightarrow k\text{-}\mathcal{G}\text{er}$, then the k_H -analytic germ whose image in $k\text{-}\mathcal{G}\text{er}$ is isomorphic to X_x is unique up to a canonical isomorphism in $k_H\text{-}\mathcal{G}\text{er}$.

PROOF. Let $\tau = \{V_{i,x}\}_{i \in I}$ be a k_H -affinoid atlas defining the k_H -analytic structure on X_x . Let V_x be a k -analytic domain in X_x that admits a k_H -analytic structure. By [Corollary 8.16](#), \widetilde{V}_x and $\widetilde{V_{i,x}}$ are all H -strict. But

$$\widetilde{V_x} \cap \widetilde{V_{i,x}} = \widetilde{V_x} \cap \widetilde{V_{i,x}},$$

so $V_x \cap V_{i,x}$ admits a k_H -analytic structure. But $V_{i,x}$ is separated so the k_H -analytic structure is unique. Therefore, $V_x \cap V_{i,x}$ is k_H -analytic with respect to τ for any $i \in I$. So V_x is H -strict with respect to τ . \square

Corollary 8.20. Let X be a k -analytic space. Assume that X admits a k_H -analytic structure, then the k_H -analytic structure is unique up to a unique isomorphism.

PROOF. This is a consequence of [Proposition 8.19](#). \square

Theorem 8.21. Let $H' \supseteq H$ be a subgroup of $\mathbb{R}_{>0}$. The natural functor

$$k_H\text{-An} \rightarrow k_{H'}\text{-An}$$

is fully faithful.

PROOF. We may assume that $H' = \mathbb{R}_{>0}$. It suffices to prove the result at the level of germs, namely, it suffices to show that

$$k_H\text{-Ger} \rightarrow k\text{-Ger}$$

is fully faithful.

Let X_x, Y_y be k_H -analytic germs and suppose that we are given a morphism $Y_y \rightarrow X_x$ in $k\text{-Ger}$. Let V_x be a k_H -analytic domain in X_x . We need to show that $W_y := Y_y \times_{X_x} V_x$ is a k_H -analytic domain in Y_y . We draw the Cartesian diagram:

$$\begin{array}{ccc} W_y & \longrightarrow & V_x \\ \downarrow & \square & \downarrow \\ Y_y & \longrightarrow & X_x \end{array}$$

By [Corollary 8.16](#), $\widehat{X}_x, \widehat{Y}_y$ are H -strict and by [Proposition 8.18](#), the preimage of \widehat{V}_x in \widehat{Y}_x is open quasi-compact and H -strict. But this preimage is just \widehat{W}_y by [Corollary 8.13](#), so we conclude that W_y is k_H -analytic by [Corollary 8.16](#). \square

Corollary 8.22. Let X be a k -analytic space. Then there is at most one k_H -analytic space X' up to isomorphisms in $k_H\text{-An}$ whose image under

$$k_H\text{-An} \rightarrow k\text{-An}$$

is isomorphic to X .

In particular, we can and will view k_H -analytic spaces as k -analytic spaces that admit (necessarily unique) structures of k_H -analytic spaces.

PROOF. This follows immediately from [Theorem 8.21](#). \square

9. Some results whose proofs I do not understand

We introduce a lemma allowing one to tell when the gluing of two affinoid spaces in a suitable position is good.

Lemma 9.1. [This is probably problematic, we will avoid using this result!](#) Let X be a separated compact k -analytic space and $x \in X$. Assume that

- (1) $\widetilde{X}_x \subseteq \mathbf{P}_{\widehat{\mathcal{H}(x)}/\widehat{k}}$ is an affine subset;
- (2) X is the union of two k -affinoid domains $Y = \text{Sp } B$ and $Z = \text{Sp } C$ both containing x ;
- (3) there is a non-zero homogeneous element $\lambda \in \widehat{\mathcal{H}(x)}$ such that

$$\widetilde{Y}_x = \widetilde{X}_x\{\lambda\}, \quad \widetilde{Z}_x = \widetilde{X}_x\{\lambda^{-1}\}.$$

Then X_x is good.

[This proof does not make much sense to me. I am just reproducing the arguments of Temkin. Need some reflection!](#)

PROOF. We observe that we are free to shrink X to k -analytic domains of the following form: $Y' \cup Z'$, where Y' and Z' are k -affinoid neighbourhoods of x in Y and Z respectively. We will express this procedure as *shrinking* X .

Step 1. We show that after shrinking X , we may assume that $Y \cap Z = \mathrm{Sp} A$, where

$$A = B\{tf^{-1}\} = C\{t^{-1}g\}$$

for some $f \in B$ and $g \in C$ and $\rho_A(f - g) < t := \rho(\lambda)$ and $\lambda = \widetilde{f(x)}$.

By [Proposition 3.3](#), up to shrinking X , we can find $f \in B$, $g \in C$ both invertible such that

$$\lambda = \widetilde{f(x)} = \widetilde{g(x)}.$$

It therefore follows that

$$|(f - g)(x)| < t.$$

After shrinking X , we can make sure that

$$\sup_{y \in Y \cap Z} |(f - g)(y)| < t.$$

After shrinking X , we can guarantee that

$$\sup_{y \in Y} |f(y)| \leq t, \quad \inf_{z \in Z} |g(z)| \geq t.$$

Why? This is not an open condition on Y or Z !!!!

In particular,

$$Y \cap Z \subseteq Y\{tf^{-1}\} \cap Z\{t^{-1}g\}.$$

This relies on the unjustified claim

By [Lemma 8.5](#), we have

$$Y\{\widetilde{tf^{-1}}\}_x = Z\{\widetilde{t^{-1}g}\}_x = (\widetilde{Y \cap Z})_x.$$

Applying [Theorem 8.10](#), we can find k -affinoid neighbourhoods $Y' = \mathrm{Sp} B'$ and $Z' = \mathrm{Sp} C'$ of x in Y and Z respectively such that

$$Y' \cap Z = Y'\{tf^{-1}\}, \quad Y \cap Z' = Z'\{t^{-1}g\}.$$

As $Y' \cap Z'$ is a k -affinoid neighbourhood of x in $Y \cap Z$, we can find a k -Laurent neighbourhood W of x in $Y \cap Z$ which is contained in $Y' \cap Z'$ and which is of the form

$$(Y \cap Z) \{s^{-1}u, s'v^{-1}\},$$

where $n, m \in \mathbb{N}$, $s = (s_1, \dots, s_n) \in \mathbb{R}_{>0}^n$, $s' = (s'_1, \dots, s'_m) \in \mathbb{R}_{>0}^m$ and $u = (u_1, \dots, u_n) \in A^n$, $v = (v_1, \dots, v_m) \in A^m$. This follows from [Proposition 10.13](#) in [Affinoid algebras](#).

As $Y' \cap Z = Y'\{tf^{-1}\}$ is a k -Weierstrass domain in Y' , by [Proposition 10.5](#) in [Affinoid algebras](#), we can find $u'_i, v'_j \in B'$ sufficiently close to u_i, v_j over $Y' \cap Z$ for $i = 1, \dots, n$, $j = 1, \dots, m$ so that $Y'' := Y'\{s^{-1}u', s'v'^{-1}\}$ is a neighbourhood of x in Y' . Similarly, we can find $Z'' = Z'\{s^{-1}u'', s'v''^{-1}\}$ for suitable perturbations of u and v . Moreover,

$$W = Y'' \cap (Y' \cap Z) = Z'' \cap (Y \cap Z').$$

It follows that $W = Y''\{tf^{-1}\} = Z''\{t^{-1}g\} = Y'' \cap Z''$. Replacing Y and Z by Y'' and Z'' and X by $Y'' \cup Z''$, we reduce to the situation stated in this step.

Step 2. We show that after shrinking X , we may guarantee that there are admissible epimorphisms

$$(9.1) \quad \begin{aligned} k\{r^{-1}T, t^{-1}S_1, pS_2\} &\rightarrow B, & T_i &\mapsto f_i \text{ for } i = 1, \dots, n, S_1 \mapsto f, S_2 \mapsto f^{-1}, \\ k\{r^{-1}T, q^{-1}S_1, tS_2\} &\rightarrow C, & T_i &\mapsto g_i \text{ for } i = 1, \dots, n, S_1 \mapsto g, S_2 \mapsto g^{-1}, \end{aligned}$$

where $n \in \mathbb{N}$, $p < t < q$, $r = (r_1, \dots, r_n) \in \mathbb{R}_{>0}^n$, $f_1, \dots, f_n \in B$, $g_1, \dots, g_n \in C$ and

$$\|f_i - g_i\| < r_i \text{ for } i = 1, \dots, n; \quad \|f - g\| < t,$$

where the norm $\|\bullet\|$ on A is the quotient norm A induced by

$$k\{r^{-1}T, t^{-1}S_1, tS_2\} \rightarrow B, \quad T_i \mapsto f_i \text{ for } i = 1, \dots, n, S_1 \mapsto f, S_2 \mapsto f^{-1}.$$

In order to guarantee that r in both morphisms are the same, we need an argument as in Step 1, which is problematic! This is unfortunately essential to Step 3. I cannot make sense of this proof anymore!

As \widetilde{X}_x is affine, we can write it as

$$\mathbf{P}_{\widetilde{\mathcal{H}(x)}/\bar{k}}\{\alpha_1, \dots, \alpha_m\}$$

for some non-zero homogeneous elements $\alpha_i \in \widetilde{\mathcal{H}(x)}$ for $i = 1, \dots, m$.

By [Proposition 3.3](#), after shrinking X , we may assume that $\alpha_i = \widetilde{f_i(x)}$ for some invertible $f_i \in B$ for $i = 1, \dots, m$.

Let $r_i = |\rho(f_i)|$ for $i = 1, \dots, m$. Set

$$D = k\{r^{-1}T, t^{-1}S_1, pS_2\}.$$

Let $\phi : D \rightarrow B$ be the first morphism in the beginning of this step and $\chi_x : B \rightarrow \mathcal{H}(x)$ be the character corresponding to x . Then $\varphi := \chi_x \circ \phi$ satisfies

$$\tilde{\varphi}(\tilde{D}) = \tilde{k}[\alpha_1, \dots, \alpha_m, \lambda].$$

On the other hand, $\widetilde{Y}_x = \mathbf{P}_{\widetilde{\mathcal{H}(x)}/\bar{k}}\{\alpha_1, \dots, \alpha_m, \lambda\}$ by [Lemma 8.5](#). It follows from [Lemma 4.5](#) in [Commutative algebras](#) and [Proposition 15.2](#) in [Affinoid algebras](#) that χ_x is inner with respect to D . It follows from [Proposition 15.2](#) in [Affinoid algebras](#) that ϕ can be extended to a continuous epimorphism

$$k\{r^{-1}T, t^{-1}S_1, pS_2, U_1, \dots, U_a\} \rightarrow B, \quad U_i \mapsto u_i \text{ for } i = 1, \dots, a$$

with $|u_i(x)| < 1$ for $i = 1, \dots, a$.

As $Y \cap Z$ is a k -Weierstrass domain in Z , we can find $g_1, \dots, g_m, g' \in C$ close enough to f_1, \dots, f_m, f on $Y \cap Z$. Up to shrinking Z , we may also guarantee that they are close on Z . In particular, we can replace g by g' but guaranteeing that $Y \cap Z = Z\{t^{-1}g\}$ still holds.

Similarly, one constructs

$$k\{r^{-1}T, q^{-1}S_1, tS_2, V_1, \dots, V_b\} \rightarrow C, \quad V_i \mapsto v_i \text{ for } i = 1, \dots, b$$

with $|v_i(x)| < 1$ for $i = 1, \dots, b$. By perturbation, we can find $u'_1, \dots, u'_a \in C$, v'_1, \dots, v'_b close to u_1, \dots, u_a and v_1, \dots, v_b on $Y \cap Z$ such that

$$Y' = Y\{v'_1, \dots, v'_b\}, \quad Z' = Z\{u'_1, \dots, u'_a\}$$

are neighbourhoods of x in Y and Z respectively. Up to replacing Y and Z by Y' and Z' , we conclude this step.

Step 3. We show that X is k -affinoid after the reduction in Step 2.

We first assume that the two maps in (9.1) are both isomorphisms.

Let B_+ denote the subspace of B consisting of elements of the form:

$$\sum_{\alpha \in \mathbb{N}^n} \sum_{j=0}^{\infty} \lambda_{\alpha,j} f^{\alpha} f^j.$$

Let C_- denote the subspace of C consisting of elements of the form:

$$\sum_{\alpha \in \mathbb{N}^n} \sum_{j=-\infty}^0 \lambda_{\alpha,j} g^{\alpha} g^j.$$

We observe that each element $A = B_+ + C_-$. Take $a \in A$ and expand

$$a = \sum_{\alpha \in \mathbb{N}^n} \sum_{j=-\infty}^{\infty} \lambda_{\nu,j} f^{\alpha} f^j.$$

Then

$$a = \sum_{\alpha \in \mathbb{N}^n} \sum_{j=-\infty}^0 \lambda_{\nu,j} f^{\alpha} f^j + \sum_{\alpha \in \mathbb{N}^n} \sum_{j=1}^{\infty} \lambda_{\nu,j} f^{\alpha} f^j$$

is the desired decomposition.

In particular, for $i = 1, \dots, n$, we can write

$$f_i - g_i = b_i + c_i, \quad f - g = b + c$$

with $b, b_1, \dots, b_n \in B_+$, $c, c_1, \dots, c_n \in C_-$. Then $h_i := f_i - b_i$ and $h := f - b$ are contained in $D := B \cap C$. It follows that

$$\begin{aligned} k\{r^{-1}T, t^{-1}S, pS^{-1}\} &\rightarrow B, & T_i &\mapsto h_i \text{ for } i = 1, \dots, n, S \mapsto h, \\ k\{r^{-1}T, q^{-1}S, tS\} &\rightarrow C, & T_i &\mapsto h_i \text{ for } i = 1, \dots, n, S \mapsto h, \end{aligned}$$

are isomorphisms. It follows that $D \cong k\{r^{-1}T, pS^{-1}, q^{-1}S\}$ and $X \cong \text{Sp } D$.

Now we handle the general case. Let

$$A' = k\{r^{-1}T, t^{-1}S, tS^{-1}\}$$

and consider the epimorphism

$$A' \rightarrow A, \quad T_i \mapsto f_i \text{ for } i = 1, \dots, n; S \mapsto f.$$

We can find preimages $G_1, \dots, G_n, G \in A'$ of g_1, \dots, g_n, g such that the norms of $T_i - G_i$ and $S - G$ are small enough. Then the map

$$k\{r^{-1}T, t^{-1}S, tS^{-1}\} \rightarrow A', \quad T_i \mapsto G_i, S \mapsto G$$

is an isomorphism. Let

$$\begin{aligned} B' &= k\{r^{-1}T, t^{-1}S, pS^{-1}\}, \\ C' &= k\{r^{-1}G, q^{-1}G, tG^{-1}\}. \end{aligned}$$

Set $Y' = \text{Sp } B'$ and $Z' = \text{Sp } C'$. By construction, we have canonical isomorphisms

$$Y'\{S^{-1}\} \cong Z'\{G\} \cong \text{Sp } A'.$$

By the previous case, X' obtained by gluing Y' and Z' along $\text{Sp } A'$ is k -affinoid. The homomorphisms $B' \rightarrow A$ and $C' \rightarrow A$ factorize through B and C and gives closed immersions $Y \rightarrow Y'$, $Z \rightarrow Z'$. The latter gives rise to a closed immersion $Y \cap Z \rightarrow Y' \cap Z'$. Hence, we get a closed immersion $X \rightarrow X'$. So X is k -affinoid. \square

Theorem 9.2. This is probably problematic, we will avoid using this result! Let X_x be a k -analytic germ. Then the following are equivalent:

- (1) X_x is good;
- (2) \widetilde{X}_x is an affine open subset of $\mathbf{P}_{\widetilde{\mathcal{H}(x)}/\tilde{k}}$.

PROOF. (1) \implies (2). This follows from the definition.

(2) \implies (1). In proving this direction, we are free to replace X by its k -analytic domain that contains x in the interior. We will express this procedure simply as *shrinking* X .

By [Corollary 8.14](#), we may assume that X is a compact separated k -analytic space. Let $\{Y_i\}_{i \in I}$ be a finite k -affinoid covering of X . After shrinking X , we may assume that $x \in Y_i$ for each $i \in I$.

By [Theorem 8.10](#), $\{\widetilde{Y_{i,x}}\}_{i \in I}$ is a finite affine covering of \widetilde{X}_x .

By [Lemma 4.8](#) in [Commutative algebras](#), we can find a Laurent covering

$$\mathcal{V} = \left\{ \widetilde{X}_x \{f_1^{\pm 1}, \dots, f_n^{\pm 1}\} \right\}$$

of \widetilde{X}_x refining $\{\widetilde{Y_{i,x}}\}_{i \in I}$, where $n \in \mathbb{N}$ and f_1, \dots, f_n are homogeneous elements in $\widetilde{\mathcal{H}(x)}$. By [Theorem 8.10](#), we can lift each element in \mathcal{V} to a k -analytic domain $V_{i,x}$ in X_x . After shrinking X , we may assume that $X = V_1 \cup \dots \cup V_n$.

By induction on n , we reduce easily to the case $n = 1$. Now it suffices to apply [Lemma 9.1](#). \square

Bibliography

- [Ber12] V. G. Berkovich. Spectral theory and analytic geometry over non-Archimedean fields. 33. American Mathematical Soc., 2012.
- [Ber93] V. G. Berkovich. Étale cohomology for non-Archimedean analytic spaces. *Publications Mathématiques de l'Institut des Hautes Études Scientifiques* 78.1 (1993), pp. 5–161.
- [Duc18] A. Ducros. Families of Berkovich spaces. 400. 2018, pp. vii+262.
- [Stacks] T. Stacks Project Authors. Stacks Project. <http://stacks.math.columbia.edu>. 2020.
- [Tem00] M. Temkin. On local properties of non-Archimedean analytic spaces. *Math. Ann.* 318.3 (2000), pp. 585–607. URL: <https://doi.org/10.1007/s002080000123>.
- [Tem04] M. Temkin. On local properties of non-Archimedean analytic spaces. II. *Israel J. Math.* 140 (2004), pp. 1–27. URL: <https://doi.org/10.1007/BF02786625>.