

Mingchen Xia

# Singularities in global pluripotential theory

– Lectures at Zhejiang University –

Latest update: May 16, 2024. The latest version can be found [here](#).



# Preface

This book is an expanded version of my lecture notes at the Institute for Advanced Study in Mathematics (IASM) at Zhejiang university. My initial goal was to write a self-contained reference for the participants of the lectures. But I soon realized that many results have never been rigorously proved in any literature. When trying to fix these loose ends, the length of the notes becomes uncontrollable, eventually leading to the current book.

In this book, I would like to present my point of view towards the *global* pluripotential theories. There are three different but interrelated theories which deserve this name. They are

- (1) the pluripotential theory on compact Kähler manifolds,
- (2) the pluripotential theory on the Berkovich analytification of projective varieties, and
- (3) the toric pluripotential theory on toric varieties.

We will begin by explaining the picture in the first case. Let us fix a compact Kähler manifold  $X$ . The central objects are the *quasi-plurisubharmonic functions* on  $X$ .

We are mostly interested in the *singularities* of such functions, that is, the places where a quasi-plurisubharmonic function  $\varphi$  tends to  $-\infty$  and how it tends to  $-\infty$ .

Singularities occur naturally in mathematics. In geometric applications,  $X$  should be regarded as the compactified moduli space of certain geometric objects. A Zariski open subset  $U \subseteq X$  would parametrize smooth objects. The natural metric on the associated polarizing line bundle is usually smooth only on  $U$ , not on  $X$ . In case we have suitable positivities, the classical Grauert–Riemann extension theorem ([Theorem B.2.2](#)) allows us to extend the metric outside  $U$ , but at the cost of introducing singularities.

The classification of singularities is a huge project. Locally near the singularities we know that quasi-plurisubharmonic functions present very complicated behaviours. There are many local invariants associated with the singularities. The most notable ones are the Lelong numbers and the multiplier ideal sheaves. These invariants only reflect the rough behaviour of a quasi-plurisubharmonic function. As an example,

a quasi-plurisubharmonic function with log-log singularities have the same local invariants as a bounded one.

The situation changes drastically in the global setting, namely on compact manifolds. In the global setting, there are three different ways to classify quasi-plurisubharmonic functions according to their singularities:

- (1) The singularity type characterizing the singularities up to a bounded term.
- (2) The  $P$ -singularity type associated with global masses.
- (3) The  $I$ -singularity type associated with all non-Archimedean data.

The classification becomes rougher and rougher as we go downward. In the first case, we say two quasi-plurisubharmonic functions have the same singularity type if their difference lies in  $L^\infty$ . The corresponding equivalence class gives us essentially the finest information of the singularities we can expect. The other two relations are more delicate, we will study them in detail in [Chapter 6](#).

A natural idea to study the singularities would consist of the following steps:

- (1) Classify the  $I$ -singularity types.
- (2) Classify the  $P$ -singularity types within a given  $I$ -singularity class.
- (3) Classify the singularity types within a given  $P$ -equivalence class.

The Step 3 is well-studied in the literature in the last decade under the name of pluripotential theory with prescribed singularities. There are numerous excellent results in this direction. In some sense, this step is already well-understood.

We will give a complete answer to Step 1 in [Chapter 7](#), where we show that  $I$ -singularity types can be described very explicitly.

It remains to consider Step 2. This is not an easy task. It is easy to construct examples where a given  $I$ -equivalence class consists of a huge amount of  $P$ -equivalence classes.

On the other hand, by contrast, in the toric pluripotential theory and non-Archimedean pluripotential theory, Step 2 is essentially trivial: An  $I$ -equivalence class consists of a single  $P$ -equivalence class. In the toric situation, an  $I$  or  $P$ -equivalence class is simply a sub-convex body of the Newton body, while in the non-Archimedean situation, an  $I$  or  $P$ -equivalence class is a homogeneous plurisubharmonic metric.

This apparent anomaly and numerous examples show that in the pluripotential theory on compact Kähler manifolds, certain singularities are pathological. Within each  $I$ -equivalence, we could pick up a canonical  $P$ -equivalence class, the quasi-plurisubharmonic functions in which are said to be  $I$ -good. We will study the theory of  $I$ -good singularities in [Chapter 7](#). As we will see later on, almost all (if not all) singularities occurring naturally are  $I$ -good.

My personal impression is that we are in a situation quite similar to the familiar one in real analysis. There are many non-measurable functions, but in real life, unless you construct a pathological function by force, you only encounter measurable functions. Similarly, although there exist many non- $I$ -good singularities, you would never encounter them in reality!

Having established this general principle, we could content ourselves in the framework of  $I$ -good singularities. Then Step 2 is essentially solved, and we have a pretty good understanding of the classification of singularities.

Of course, this classification is a bit abstract. To put it into use, we will introduce two general techniques allowing us to make induction on  $\dim X$ . For a prime divisor  $Y$  in general position, we have the so-called analytic Bertini theorem relation quasi-plurisubharmonic functions on  $X$  and on  $Y$ . For a non-generic  $Y$ , we have the technique of trace operators. These techniques will be explained in [Chapter 8](#).

In the toric situation, these constructions and methods are quite straightforward and are likely known to experts before I entered this field, see [Chapter 5](#) for the toric pluripotential theory on ample line bundles.

The corresponding toric pluripotential theory on big line bundles has never been written down in the literature. We will develop the theory of partial Okounkov bodies in [Chapter 10](#) and the general toric pluripotential theory will be developed as an application in [Chapter 12](#).

Finally, we give applications to non-Archimedean pluripotential theory in [Chapter 13](#) based on the theory of test curves developed in [Chapter 9](#). We also prove the convergence of the partial Bergman kernels in [Chapter 14](#).

The readers are only supposed to be familiar with the basic pluripotential theory. The excellent book [\[GZ17\]](#) is more than enough.

*Minghen Xia*  
in Hangzhou, March 2024



## Acknowledgements

I would like to express my gratitude to Bing Wang and Song Sun for their gracious invitations to China and for providing me with the opportunity to deliver a series of lectures.

Furthermore, I am indebted to the dedicated researchers and secretaries of the University of Science and Technology of China (USTC) and the Institute for Advanced Study in Mathematics (IASM) for cultivating an exceptional research environment. Their commitment to excellence has allowed me to immerse myself fully in the field of mathematics during my time in China.

I am also immensely thankful to the participants of the course, including Song Sun, Mingyang Li, Xin Fu, Jiyuan Han, Junsheng Zhang, Yifan Chen, Yueqing Feng, Minghao Miao, and Federico Giust. Their active engagement and insightful discussions have greatly enriched my lectures and enhanced my understanding of the subject matter.

Special appreciation goes to Yi Yao and Kewei Zhang for their invaluable contributions to discussions on toric geometry, which ultimately inspired the theory developed in [Chapter 12](#).

Most results in this book are developed in collaboration with Tamás Darvas and Kewei Zhang, whose insights are always crucial in the development of the theories. I would like to thank them for the collaborations over years.

A substantial part of the current book was essentially contained in my PhD thesis. I would like to thank my advisor Robert Berman for his guidance and my colleagues in Göteborg and Paris for constant discussions, especially Bo Berndtsson, David Witt Nyström, Sébastien Boucksom and Elizabeth Wulcan.

This work would not have been possible without the unwavering support and encouragement of all those mentioned above. Thank you for your generosity, guidance, and camaraderie throughout this endeavor.

Enfin, je tiens à exprimer ma gratitude à Sébastien Boucksom et Madame Natalia Hristic de la Sorbonne Université, qui m'ont aidé à contacter le ministère de l'intérieur en France. Sans leur intervention, je serais encore coincé en France, échoué par l'efficacité extraordinaire du gouvernement français, en particulier de la préfecture de Créteil et ce livre n'aurait jamais vu le jour.





# Contents

## Part I Preliminaries

<b>1</b>	<b>Plurisubharmonic functions</b>	3
1.1	The definition of plurisubharmonic functions	3
1.1.1	The 1-dimensional case	3
1.1.2	The higher dimensional case	4
1.1.3	The manifold case	6
1.2	Properties of plurisubharmonic functions	7
1.3	Plurifine topology	11
1.3.1	Plurifine topology on domains	11
1.3.2	Plurifine topology on manifolds	14
1.4	Lelong numbers and multiplier ideal sheaves	15
1.5	Quasi-plurisubharmonic functions	19
1.6	Analytic singularities	20
1.7	The space of currents	23
1.8	Plurisubharmonic metrics on line bundles	24
<b>2</b>	<b>Non-pluripolar products</b>	27
2.1	Bedford–Taylor theory	27
2.2	The non-pluripolar products	28
2.3	Properties of non-pluripolar products	30
<b>3</b>	<b>The envelope operators</b>	35
3.1	The $P$ -envelope	35
3.1.1	Rooftop operator and the definition of the $P$ -envelope	35
3.1.2	Properties of the $P$ -envelope	39
3.1.3	Relative full mass classes	42
3.2	The $\mathcal{I}$ -envelope	45
3.2.1	$\mathcal{I}$ -equivalence	45
3.2.2	The definition of the $\mathcal{I}$ -envelope	47
3.2.3	Properties of the $\mathcal{I}$ -envelope	49

<b>4</b>	<b>Geodesic rays in the space of potentials</b>	53
4.1	Subgeodesics	53
4.2	Geodesics in the space of potentials	55
<b>5</b>	<b>Toric pluripotential theory on ample line bundles</b>	63
5.1	Toric setup	63
5.2	Toric plurisubharmonic functions	64
5.3	Toric pluripotential theory	67

## Part II The theory of $\mathcal{I}$ -good singularities

<b>6</b>	<b>Comparison of singularities</b>	75
6.1	The $P$ and $\mathcal{I}$ -partial orders	75
6.1.1	The definitions of the partial orders	75
6.1.2	Properties of the partial orders	79
6.2	The $d_S$ -pseudometric	83
6.2.1	The definition of the $d_S$ -pseudometric	83
6.2.2	Convergence theorems	89
6.2.3	Continuity of invariants	97
<b>7</b>	<b><math>\mathcal{I}</math>-good singularities</b>	99
7.1	The notion of $\mathcal{I}$ -good singularities	99
7.2	Properties of $\mathcal{I}$ -good singularities	102
7.3	The volume of Hermitian big line bundles	105
<b>8</b>	<b>The trace operator</b>	109
8.1	The definition of the trace operator	109
8.2	Properties of the trace operator	111
8.3	Restricted volumes	115
8.4	Analytic Bertini theorems	120
<b>9</b>	<b>Test curves</b>	127
9.1	The notion of test curves	127
9.2	Ross–Witt Nyström correspondence	130
9.3	$\mathcal{I}$ -model test curves	136
9.4	Operations on test curves	137
<b>10</b>	<b>The theory of Okounkov bodies</b>	147
10.1	Flags and valuations	147
10.1.1	The algebraic setting	147
10.1.2	The transcendental setting	148
10.2	Algebraic partial Okounkov bodies	152
10.2.1	The spaces of sections	152
10.2.2	Algebraic Okounkov bodies	153
10.2.3	Construction of partial Okounkov bodies	155
10.2.4	Basic properties of partial Okounkov bodies	157

10.2.5	The Hausdorff convergence property of partial Okounkov bodies	160
10.2.6	Recover Lelong numbers from partial Okounkov bodies	164
10.3	Transcendental partial Okounkov bodies	165
10.3.1	The traditional approach to the Okounkov body problem	165
10.3.2	Definitions of partial Okounkov bodies	166
10.3.3	The valuative characterization	171
10.4	Okounkov test curves	176
<b>11</b>	<b>The theory of b-divisors</b>	<b>185</b>
11.1	The intersection theory of b-divisors	185
11.2	The singularity b-divisors	187
11.3	Okounkov bodies of b-divisors	190
<b>Part III Applications</b>		
<b>12</b>	<b>Toric pluripotential theory on big line bundles</b>	<b>197</b>
12.1	Toric setup	197
12.2	Toric partial Okounkov bodies	198
12.2.1	Newton bodies	198
12.2.2	Partial Okounkov bodies	198
12.3	The pluripotential theory	202
<b>13</b>	<b>Non-Archimedean pluripotential theory</b>	<b>207</b>
13.1	The definition of non-Archimedean metrics	207
13.2	Operations on non-Archimedean metrics	210
13.3	Duistermaat–Heckman measures	216
<b>14</b>	<b>Partial Bergman kernels</b>	<b>219</b>
14.1	Partial envelopes	219
14.2	Quantization of partial equilibrium measures	228
14.2.1	Bernstein–Markov measures	228
14.2.2	Partial Bergman kernels	229
<b>Comments</b>		<b>237</b>
<b>A</b>	<b>Convex functions and convex bodies</b>	<b>243</b>
A.1	The notion of convex functions	243
A.2	Legendre transform	246
A.3	Classes of convex functions	248
A.4	Monge–Ampère measures	250
A.5	Separation lemmata	251

<b>B</b>	<b>Pluripotential theory on unibranch spaces</b>	253
B.1	Complex spaces	253
B.2	Plurisubharmonic functions	254
B.3	Extensions of the results in the smooth setting	257
<b>C</b>	<b>Almost semigroups</b>	259
C.1	Convex bodies	259
C.2	The Okounkov bodies of almost semigroups	261
C.2.1	Generalities on semigroups	261
C.2.2	Okounkov bodies of semigroups	263
C.2.3	Okounkov bodies of almost semigroups	265
	<b>Index</b>	269
	<b>References</b>	273

## Conventions

In the whole book we adopt the following conventions:

- A complex space is always assumed to be *reduced*, *paracompact* and *Hausdorff*.
- A *modification* of a complex space  $X$  is proper bimeromorphic morphism  $\pi: Y \rightarrow X$  that is obtained from a finite composition of blow-ups with smooth centers.
- A *subnet* of a net refers to a cofinal subnet.
- A *domain* in  $\mathbb{C}^n$  refers to a connected open subset.
- A *complex manifold* is assumed to be paracompact.
- A *submanifold* of a complex manifold means a complex submanifold.
- A *neighbourhood* is not necessarily open.
- The set  $\mathbb{N}$  of natural numbers includes 0.

We will use the following notations throughout the book:

- If  $I$  is a non-empty set, then  $\text{Fin}(I)$  denote the net of finite non-empty subsets of  $I$ , ordered by inclusion.
- $\text{dd}^c$  means  $(2\pi)^{-1}i\partial\bar{\partial}$ .



# **Part I**

## **Preliminaries**

In the first two chapters [Chapter 1](#) and [Chapter 2](#) of this part, we recall a few preliminaries about the notion of plurisubharmonic functions and the non-pluripolar products of plurisubharmonic functions.

All materials in these chapters are standard and are well-documented in other textbooks, so we will be rather sketchy. The readers are encouraged to consult the excellent textbook [\[GZ17\]](#).

In [Chapter 3](#), we develop the techniques of envelope operators. All results in this section are known and are written in various articles.

In [Chapter 4](#), we develop the theory of geodesics in the space of quasi-plurisubharmonic functions. Most results in this chapter are known to different degrees, but not in the fully general form as we present. Most proofs are similar to the known proofs in the literature, but the presence of singularities requires a very careful treatment.

In [Chapter 5](#), we recall the basic results about the toric pluripotential theory on ample line bundles, which will be generalized to big line bundles in [Chapter 12](#).

Experienced readers may safely skip the whole part.



# Chapter 1

## Plurisubharmonic functions

chap:psh

In this chapter, we recall the notion of plurisubharmonic functions and a few basic properties of these functions. The main purpose is to fix the notations for later chapters, so we refer to the literature for most proofs.

We give some details about the plurifine topology in [Section 1.3](#), since the related proofs are scattered in a number of articles.

In the literature related to multiplier ideal sheaves and Lelong numbers, there are several different conventions about their normalizations. The readers could find more about the conventions that we adopt in the whole book in [Section 1.4](#).

### 1.1 The definition of plurisubharmonic functions

sec:pshdef

In this section, we recall the notion of plurisubharmonic functions. We will also take care of the 0-dimensional case, which makes a number of induction arguments easier to carry out. None of our references treats the 0-dimensional case, but the readers can easily verify that the results in this section hold in this exceptional case.

#### 1.1.1 The 1-dimensional case

Let  $\Omega$  be a domain (a connected open subset) in  $\mathbb{C}$ .

def:subhar1

**Definition 1.1.1** A *subharmonic function* on  $\Omega$  is a function  $\varphi: \Omega \rightarrow [-\infty, \infty)$  satisfying the following three conditions:

- (1)  $\varphi \not\equiv -\infty$ ;
- (2)  $\varphi$  is upper semi-continuous;
- (3)  $\varphi$  satisfies the *sub-mean value inequality*: For any  $a \in \Omega$  and  $r > 0$  such that  $B_1(a, r) \Subset \Omega$ , we have

$$\varphi(a) \leq \frac{1}{2\pi} \int_0^{2\pi} \varphi(a + re^{i\theta}) d\theta.$$

We will denote the set of subharmonic functions on  $\Omega$  as  $\text{SH}(\Omega)$ .

Here  $B_1(a, r)$  denotes the open ball with center  $a$  and radius  $r$ . See (1.1).

In fact, for each  $a \in \Omega$ , in (3), it suffices to require the sub-mean value inequality for all small enough  $r > 0$ .

Intuitively, at a specific point  $a \in \Omega$ , the Condition (2) gives a lower bound of the value of  $\varphi(a)$  using the nearby values of  $\varphi$ , while the Condition (3) gives an upper bound. This intuition leads to the following rigidity theorem:

thm:sh\_rigid

**Theorem 1.1.1** *Let  $\varphi: \Omega \rightarrow [-\infty, \infty)$  be a measurable function. Then the following are equivalent:*

- (1)  $\varphi$  is locally integrable and  $\Delta\varphi \geq 0$ .
- (2)  $\varphi$  coincides almost everywhere with a subharmonic function  $\psi$  on  $\Omega$ .

Moreover, the subharmonic function  $\psi$  is unique.

Here in condition 1,  $\Delta\varphi$  is the Laplacian in the sense of currents. This is a special case of [Theorem 1.1.2](#) below.

This theorem gives a very useful way to construct subharmonic functions.

### 1.1.2 The higher dimensional case

We will fix  $n \in \mathbb{N}$  and a domain  $\Omega$  (a connected open subset) in  $\mathbb{C}^n$ .

def:psh

**Definition 1.1.2** When  $n \geq 1$ , a *plurisubharmonic function* on  $\Omega$  is a function  $\varphi: \Omega \rightarrow [-\infty, \infty)$  satisfying the following three conditions:

- (1)  $\varphi \not\equiv -\infty$ ;
- (2)  $\varphi$  is upper semi-continuous;
- (3) for any complex line  $L \subseteq \mathbb{C}^n$  and any connected component  $U$  of  $L \cap \Omega$ , the restriction  $\varphi|_U$  is either subharmonic or constantly  $-\infty$ .

When  $n = 0$ , the only domain  $\Omega$  is the singleton. A *plurisubharmonic function* on  $\Omega$  is a real-valued function on  $\Omega$ .

The set of plurisubharmonic functions on  $\Omega$  is denoted by  $\text{PSH}(\Omega)$ .

A plurisubharmonic function is also called a psh function for short.

*Example 1.1.1* When  $n = 0$ , we have a canonical bijection  $\text{PSH}(\Omega) \cong \mathbb{R}$ .

*Example 1.1.2* When  $n = 1$ , we have  $\text{PSH}(\Omega) = \text{SH}(\Omega)$ .

Similar to [Theorem 1.1.1](#), we have a rigidity theorem for plurisubharmonic functions as well.

thm:psh\_rigid

**Theorem 1.1.2** *Let  $\varphi: \Omega \rightarrow [-\infty, \infty)$  be a measurable function. Then the following are equivalent:*

- (1)  $\varphi$  is locally integrable and  $dd^c \varphi \geq 0$ ;
- (2)  $\varphi$  coincides almost everywhere with a plurisubharmonic function  $\psi$  on  $\Omega$ .

Moreover, the plurisubharmonic function  $\psi$  is unique.

Here the operator  $dd^c$  is normalized so that

$$dd^c = \frac{i}{2\pi} \partial \bar{\partial}.$$

For the proof, we refer to [GZ17, Proposition 1.43].

Plurisubharmonic functions have nice functorialities:

prop:func\_domain

**Proposition 1.1.1** *Let  $n' \in \mathbb{N}$  and  $\Omega' \subseteq \mathbb{C}^{n'}$  be a domain. Given any holomorphic map  $f: \Omega \rightarrow \Omega'$  and any  $\varphi \in \text{PSH}(\Omega')$  exactly one of the following cases occurs:*

- (1)  $f^* \varphi \equiv -\infty$ ;
- (2)  $f^* \varphi \in \text{PSH}(\Omega)$ .

We refer to [GZ17, Proposition 1.44] for the proof<sup>1</sup>.

For each  $n \in \mathbb{N}$ ,  $a \in \mathbb{C}^n$  and  $r > 0$ , we write

$$B_n(a, r) = \{z \in \mathbb{C}^n : |z - a| < r\}. \quad (1.1)$$

{eq:Bnar}

prop:ballpshconvex

**Proposition 1.1.2** *Let  $\varphi \in \text{PSH}(B_n(a, r_0))$  for some  $r_0 > 0$ . Then the function*

$$(-\infty, \log r_0) \rightarrow \mathbb{R}, \quad \log r \mapsto \sup_{B_n(a, r)} \varphi$$

*is convex and increasing.*

See [Bou17, Corollary 2.4].

prop:subhimpilyconv

**Proposition 1.1.3** *Let  $a < b$  be two real numbers. Let  $f: (a, b) \rightarrow [-\infty, \infty)$  be a function. Define*

$$g: \{z \in \mathbb{C} : e^{-b} < |z| < e^{-a}\} \rightarrow [-\infty, \infty), \quad z \mapsto f(-\log |z|).$$

*Suppose that  $g$  is harmonic, then  $f$  is convex. In particular,  $f$  takes real values only.*

See [HK76, Theorem 2.12] for a more general result.

<sup>1</sup> We remind the readers that the statement of [GZ17, Proposition 1.44] is flawed.

### 1.1.3 The manifold case

Let  $X$  be a complex manifold. In the whole book, complex manifolds are assumed to be paracompact, namely all connected components have countable bases.

def:pshmfd

**Definition 1.1.3** A *plurisubharmonic function* on  $X$  is a function  $\varphi: X \rightarrow [-\infty, \infty)$  such that for any  $x \in X$ , there exists an open neighbourhood  $U$  of  $x$  in  $X$ , an integer  $n \in \mathbb{N}$ , a domain  $\Omega \subseteq \mathbb{C}^n$  and a biholomorphic map  $F: \Omega \rightarrow U$  such that  $F^*(\varphi|_U) \in \text{PSH}(\Omega)$ .

The set of plurisubharmonic functions on  $X$  is denoted by  $\text{PSH}(X)$ .

*Example 1.1.3* When  $X$  is a domain in  $\mathbb{C}^n$ , the notions of plurisubharmonic functions in [Definition 1.1.3](#) and in [Definition 1.1.2](#) coincide.

*Example 1.1.4* Write  $\{X_i\}_{i \in I}$  for the set of connected components of  $X$ . Then we have a natural bijection

$$\text{PSH}(X) \cong \prod_{i \in I} \text{PSH}(X_i).$$

Here the product is in the category of sets. In particular, if  $X = \emptyset$ , then  $\text{PSH}(X) = \emptyset$ .

This example allows us to reduce to the case of connected manifolds when studying general plurisubharmonic functions.

prop:pullbackpsh

**Proposition 1.1.4** Let  $Y$  be another complex manifold and  $f: Y \rightarrow X$  be a holomorphic map. Then for any  $\varphi \in \text{PSH}(X)$ , exactly one of the following cases occurs:

- (1)  $f^*\varphi$  is identically  $-\infty$  on some connected component of  $Y$ ;
- (2)  $f^*\varphi \in \text{PSH}(Y)$ .

This proposition follows easily from [Proposition 1.1.1](#). We leave the details to the readers.

[Theorem 1.1.2](#) implies immediately the general form of the rigidity theorem:

thm:psh\_rigid\_gen

**Theorem 1.1.3** Let  $\varphi: X \rightarrow [-\infty, \infty)$  be a measurable function. Then the following are equivalent:

- (1)  $\varphi$  is locally integrable and  $\text{dd}^c \varphi \geq 0$ ;
- (2)  $\varphi$  coincides almost everywhere with a plurisubharmonic function  $\psi$  on  $X$ .

Moreover, the plurisubharmonic function  $\psi$  is unique.

def:pluripolarsets

**Definition 1.1.4** A subset  $E \subseteq X$  is *pluripolar* if for any  $x \in X$ , there is an open neighbourhood  $U$  of  $x$  in  $X$  and a function  $\psi \in \text{PSH}(U)$  such that

$$\psi|_{E \cap U} \equiv -\infty.$$

A subset  $E \subseteq X$  is *non-pluripolar* if  $E$  is not pluripolar.

A subset  $F \subseteq X$  is *co-pluripolar* if  $X \setminus F$  is pluripolar.

thm:Josefson

**Theorem 1.1.4 (Josefson's theorem)** *Let  $E \subseteq \mathbb{C}^n$  be a pluripolar set. Then there is  $\varphi \in \text{PSH}(\mathbb{C}^n)$  such that  $\varphi|_E \equiv -\infty$ .*

See [GZ17, Corollary 4.41] for the proof of a more general result.

There is also a global version of Josefson's theorem:

thm:gloJosefson

**Theorem 1.1.5** *Assume that  $X$  is a compact complex manifold and  $E \subseteq X$  is a pluripolar set. Then there is a quasi-plurisubharmonic function  $\varphi$  on  $X$  with  $\varphi|_E \equiv -\infty$ .*

For a proof, see [Vu19].

prop:pluripolarunion

**Proposition 1.1.5** *Let  $(E_i)_{i \in \mathbb{Z}_{>0}}$  be a sequence of pluripolar sets in  $X$ . Then*

$$E := \bigcup_{i=1}^{\infty} E_i$$

*is pluripolar.*

**Proof** The problem is local, so we may assume that  $X \subseteq \mathbb{C}^n$  is a domain. In this case, by Theorem 1.1.4 for each  $i \in \mathbb{Z}_{>0}$  we can choose  $\psi_i \in \text{PSH}(\mathbb{C}^n)$  such that

$$\psi_i|_{E_i} \equiv -\infty, \quad \psi_i \leq 0$$

for all  $i > 0$ . After shrinking  $X$ , we may guarantee that  $\psi_i|_X \in L^1(X)$  for all  $i > 0$ . After rescaling, we may also assume that  $\|\psi_i\|_{L^1(X)} \leq 1$  for all  $i > 0$ .

We then define

$$\psi = \sum_{i=1}^{\infty} 2^{-i} \psi_i|_X.$$

Then  $\psi \in \text{PSH}(X)$  according to Proposition 1.2.1 below and  $\psi|_E = -\infty$ .  $\square$

## 1.2 Properties of plurisubharmonic functions

In this section, we explore the basic properties of plurisubharmonic functions.

Let  $X$  be a complex manifold.

prop:pshfunction\_closedseq

**Proposition 1.2.1**

- (1) *Assume that  $(\varphi_i)_{i \in I}$  is a non-empty family in  $\text{PSH}(X)$  that is locally uniformly bounded from above. Then  $\sup^*_i \varphi_i \in \text{PSH}(X)$ .*
- (2) *Assume that  $(\varphi_i)_{i \in I}$  is a decreasing net in  $\text{PSH}(X)$  such that  $\lim_{i \in I} \varphi_i$  is not identically  $-\infty$  on each connected component of  $X$ , then  $\lim_{i \in I} \varphi_i \in \text{PSH}(X)$ .*

Here  $\sup^*$  denotes the upper semicontinuous regularization of the supremum. When  $I$  is a finite family, observe that

$$\sup_{i \in I}^* \varphi_i = \sup_{i \in I} \varphi_i.$$

When  $I = \{1, \dots, m\}$ , we write

$$\varphi_1 \vee \dots \vee \varphi_m := \sup_{i \in I} \varphi_i.$$

We refer to [GZ17, Proposition 1.28, Proposition 1.40]<sup>2</sup>.

prop:Choquet

**Proposition 1.2.2 (Choquet's lemma)** *Assume that  $X$  has countably many connected components. Assume that  $(\varphi_i)_{i \in I}$  is a non-empty family in  $\text{PSH}(X)$  that is locally uniformly bounded from above. There exists a countable subset  $J \subseteq I$  such that*

$$\sup_{i \in I}^* \varphi_i = \sup_{j \in J}^* \varphi_j.$$

**Proof** We may assume that  $X$  is connected. Since by our convention, the complex manifold  $X$  is paracompact, it can be covered by countably many open balls, so we can easily reduce to the case where  $X$  is an open ball. In this case, the result is proved in [GZ17, Lemma 4.31].  $\square$

prop:supsupstardiff

**Proposition 1.2.3** *Let  $(\varphi_i)_{i \in I}$  be a non-empty family in  $\text{PSH}(X)$  that is locally uniformly bounded from above. Then the set*

$$\left\{ x \in X : \sup_{i \in I} \varphi_i < \sup_{i \in I}^* \varphi_i \right\}$$

*is pluripolar.*

See [GZ17, Corollary 4.28].

prop:pshlocIp

**Proposition 1.2.4** *Let  $\varphi \in \text{PSH}(X)$ , then for any  $p \geq 1$ ,  $\varphi \in L_{\text{loc}}^p(X)$ .*

See [GZ17, Theorem 1.46, Theorem 1.48].

prop:pshfuncdetdense

**Proposition 1.2.5** *Suppose that  $\varphi, \psi \in \text{PSH}(X)$ . Assume that there is a dense subset  $E \subseteq X$  such that  $\varphi|_E \leq \psi|_E$ , then  $\varphi \leq \psi$ .*

**Proof** The problem is local, so we may assume that  $X$  is a domain in  $\mathbb{C}^n$ .

We may assume that  $\varphi|_E = \psi|_E$  after replacing  $\varphi$  by  $\varphi \vee \psi$ . Then we need to show that  $\varphi = \psi$ .

It follows from [GZ17, Theorem 4.20] that this holds outside a pluripolar set  $Y \subseteq X$ . In particular,  $\varphi = \psi$  almost everywhere. It follows from the uniqueness statement in Theorem 1.1.3 that  $\varphi = \psi$ .  $\square$

thm:GRemmert

**Theorem 1.2.1 (Grauert–Remmert)** *Let  $Z$  be an analytic subset in  $X$  and  $\varphi \in \text{PSH}(X \setminus Z)$ . Then the function  $\varphi$  admits an extension to  $\text{PSH}(X)$  in the following two cases:*

<sup>2</sup> In [GZ17, Proposition 1.28], the second part is only stated for sequences, the net version is obvious using the sub-mean value inequality.

- (1) *The set  $Z$  has codimension at least 2 everywhere.*
- (2) *The set  $Z$  has codimension at least 1 everywhere and is locally bounded from above on an open neighbourhood of  $Z$ .*

*In both cases, the extension is unique.*

**Proof** The extension is unique thanks to [Proposition 1.2.5](#).

(2) Thanks to the uniqueness of the extension, the problem is local, so we may assume that  $X$  is a domain in  $\mathbb{C}^n$  with  $n > 0$  and there is a non-zero holomorphic function  $f$  vanishing identically on  $Z$ . For each  $\epsilon > 0$ , we claim that the function  $\varphi_\epsilon$  defined by

$$\varphi_\epsilon(x) := \begin{cases} \varphi(x) + \epsilon \log |f(x)|^2, & x \in X \setminus Z; \\ -\infty, & x \in Z \end{cases}$$

is plurisubharmonic on  $X$ . By [Definition 1.1.2](#), it suffices to verify the case  $n = 1$ . In this case, we may assume that  $Z = \{0\}$ . It is clear that  $\varphi_\epsilon \in \text{SH}(X \setminus Z)$ . It suffices to verify the sub-mean value inequality at 0, which is immediate.

Next observe that the sequence  $\varphi_\epsilon$  is increasing as  $\epsilon \searrow 0$  and  $\varphi_\epsilon$  is locally uniformly bounded from above. It follows from [Proposition 1.2.1](#) that  $\tilde{\varphi} := \sup_{\epsilon > 0}^* \varphi_\epsilon \in \text{PSH}(X)$ . Moreover,  $\tilde{\varphi}$  clearly extends  $\varphi$ .

(1) It suffices to verify that  $\varphi$  is locally bounded from above near each point of  $Z$ . The problem is local, so we may assume that  $X$  is a domain in  $\mathbb{C}^n$  with  $n \geq 2$ .

Assume that our assertion fails. Take  $z \in Z$  so that there exists a sequence  $(x_j)_j$  in  $X \setminus Z$  such that

$$\lim_{j \rightarrow \infty} \varphi(x_j) = \infty.$$

Since  $Z$  has codimension at least 2, we could take a complex line  $L$  passing through  $z$  and intersects  $Z$  only on a discrete set. After shrinking  $X$ , we may assume that

$$L \cap Z = \{z\}.$$

Take an open ball  $B_n(z, r) \Subset X$ . After adding a constant to  $\varphi$ , we may guarantee that  $\varphi < 0$  on  $L \cap \partial B_n(z, r)$ . Since  $\varphi$  is upper semi-continuous, we could find an open neighbourhood  $U$  of  $L \cap \partial B_n(z, r)$  such that

$$\varphi|_U < 0.$$

For each  $j \geq 1$ , take a complex line  $L_j$  passing through  $x_j$  such that  $L_j \rightarrow L$  as  $j \rightarrow \infty$ . Here the convergence is in the obvious sense. Then for large enough  $j$ , we know have

$$L_j \cap \partial B_n(z, r) \subseteq U.$$

It follows from the sub-mean value inequality that  $\varphi(x_j) < 0$  for large enough  $j$ , which is a contradiction.  $\square$

lma:invariantpshfunfinite

**Lemma 1.2.1** *Let  $\varphi \in \text{PSH}((\Delta^*)^n)$  be an  $(S^1)^n$ -invariant plurisubharmonic function. Then  $\varphi$  is finite everywhere.*

**Proof** It clearly suffices to handle the case  $n = 1$ . In this case, by [HK76](#), Theorem 2.12], the map

$$\log r \mapsto \int_0^1 \varphi(r \exp(2\pi i \theta)) d\theta = \varphi(r)$$

is a convex function of  $\log r$ . So the set  $\{r \in (0, 1) : \varphi(r) = -\infty\}$  is convex. But  $\varphi$  is almost everywhere finite by [Proposition 1.2.4](#). Since  $\varphi$  is  $S^1$ -invariant,  $\varphi|_{(0,1)}$  is almost everywhere finite. It follows from the convexity that it is everywhere finite.  $\square$

cor:L1limipp

**Corollary 1.2.1** *Let  $(\varphi_j)_{j \in \mathbb{Z}_{>0}}$  be a sequence in  $\text{PSH}(X)$  such that  $\varphi_j \xrightarrow{L^1_{\text{loc}}} \varphi \in \text{PSH}(X)$ . Then the set*

$$\left\{ x \in X : \varphi(x) \neq \overline{\lim}_{j \rightarrow \infty} \varphi_j(x) \right\}$$

*is pluripolar.*

**Proof** We first observe that  $(\varphi_j)_j$  is locally uniformly bounded from above. This follows from [GZ17](#), Exercise 1.20].

For each  $j \geq 1$ , let

$$\psi_j = \sup_{k \geq j}^* \varphi_k.$$

Then  $\psi_j \in \text{PSH}(X)$  by [Proposition 1.2.1](#). Moreover,  $(\psi_j)_j$  is a decreasing sequence and  $\psi_j \geq \varphi_j$  for all  $j$ . So by [Proposition 1.2.1](#) again,  $\psi := \inf_j \psi_j \in \text{PSH}(X)$ . On the other hand, by [Proposition 1.2.3](#), there is a pluripolar set  $Z \subseteq X$  such that for any  $x \in X \setminus Z$ , we have  $\psi(x) = \overline{\lim}_j \varphi_j(x)$ . Since  $\varphi_j \xrightarrow{L^1_{\text{loc}}} \varphi$ , we can find a set  $Y \subseteq X$  with zero Lebesgue measure such that  $\varphi_j(x) \rightarrow \varphi(x)$  for all  $x \in X \setminus Y$ .

In particular, for any  $x \in X \setminus (Y \cup Z)$ , we have

$$\psi(x) = \varphi(x).$$

But thanks to [Proposition 1.2.5](#), the equality holds everywhere. Therefore, for all  $x \in X \setminus Z$ ,

$$\varphi(x) = \overline{\lim}_{j \rightarrow \infty} \varphi_j(x).$$

prop:Kis

**Proposition 1.2.6 (Kiselman's principle)** *Let  $\Omega \subseteq \mathbb{C}^m \times \mathbb{C}^n$  be a pseudoconvex domain. Assume that for each  $z \in \mathbb{C}^m$ , the set*

$$\Omega_z := \{w \in \mathbb{C}^n : (z, w) \in \Omega\}$$

*has the form  $E + i\mathbb{R}^n$ , where  $E \subseteq \mathbb{R}^n$  is a subset. Let  $\varphi \in \text{PSH}(\Omega)$ , assume that  $\varphi$  is independent of the imaginary part of the variable in  $\mathbb{C}^n$ . Let  $\Omega'$  be the projection of  $\Omega$  to  $\mathbb{C}^m$ . Define  $\psi : \Omega' \rightarrow [-\infty, \infty)$  as follows:*

$$\psi(z) = \inf_{w \in \Omega_z} \varphi(z, w).$$



Then either  $\psi \equiv -\infty$  or  $\psi \in \text{PSH}(\Omega')$ .

See [DemBook](#) [DemT2b, Theorem 7.5].

## 1.3 Plurifine topology

sec:plurifine

### 1.3.1 Plurifine topology on domains

Let  $\Omega \subseteq \mathbb{C}^n$  ( $n \in \mathbb{N}$ ) be a domain.

def:pftopologydomain

**Definition 1.3.1** The *plurifine topology* on  $\Omega$  is the weakest topology making all  $\mathbb{R}$ -valued plurisubharmonic functions on  $\Omega$  continuous.

We want to distinguish the Euclidean topology from the plurifine topology. In the whole book, topological notions without adjectives refer to those with respect to the Euclidean topology. We include the symbol  $\mathcal{F}$  in order to denote those with respect to the plurifine topology. For example, we will say  $\mathcal{F}$ -open subset,  $\mathcal{F}$ -neighbourhood,  $\mathcal{F}$ -closure, etc. The  $\mathcal{F}$ -closure of a set  $E \subseteq \Omega$  will be denoted by  $\bar{E}^{\mathcal{F}}$ . We remind the readers that in the whole book, we follow Bourbaki's convention, a neighbourhood is not necessarily open. Similarly, an  $\mathcal{F}$ -neighbourhood is not necessarily  $\mathcal{F}$ -open.

A priori, we should include  $\Omega$  into the notations as well, but as we will see shortly in [Corollary 1.3.1](#), this is usually unnecessary.

prop:pf\_finer

**Proposition 1.3.1** The plurifine topology is finer than the Euclidean topology.

**Proof** It suffices to show that the unit ball  $\{z \in \mathbb{C}^n : |z| < 1\}$  is  $\mathcal{F}$ -open. This follows from the observation that this set can be written as

$$\{\psi < 0\} \text{ with } \psi(z) := (\log |z|) \vee (-1).$$

**Definition 1.3.2** A subset  $E \subseteq \Omega$  is *thin* at  $x \in \Omega$  if one of the following conditions holds:

- (1)  $x \notin \bar{E}$ ;
- (2)  $x \in \bar{E}$  and there is an open neighbourhood  $U \subseteq \Omega$  of  $x$  and  $\varphi \in \text{PSH}(U)$  such that

$$\overline{\lim_{y \rightarrow x, y \in E}} \varphi(y) < \varphi(x).$$

We say  $E$  is *thin* if it is thin at all  $x \in \Omega$ .

In the second case, the function  $\varphi$  can be very much improved.

prop:BTthin

**Proposition 1.3.2 (Bedford–Taylor)** Consider a set  $E \subseteq \Omega$  and  $x \in \bar{E}$ . Assume that  $E$  is thin at  $x$ , then there is  $\varphi \in \text{PSH}(\mathbb{C}^n)$  satisfying the following properties:

- (1)  $\varphi$  is locally bounded outside a neighbourhood of  $x$ ;

- (2)  $\varphi(x) > -\infty$ ;
- (3)  $\lim_{y \rightarrow x, y \in E} \varphi(y) = -\infty$ .

**Proof** By definition, there is an open neighbourhood  $U \subseteq \Omega$  of  $x$  and  $\psi \in \text{PSH}(U)$  such that

$$\overline{\lim}_{y \rightarrow x, y \in E} \psi(y) < \psi(x).$$

Without loss of generality, we may assume that  $x = 0$ ,  $U$  is the unit ball in  $\mathbb{C}^n$ ,  $\psi < 0$  and  $\psi|_{U \cap E} < -1$ , while  $\psi(0) = -\eta > -1$ .

As  $\psi$  is upper semicontinuous, we may choose  $\delta_j > 0$  for all large enough  $j \in \mathbb{Z}_{>0}$  such that  $\psi(y) < -\eta + 2^{-j-1}$  when  $y \in \mathbb{C}^n$  satisfies  $|y| < \delta_j$ . Now we let

$$\varphi_j(z) := \begin{cases} \left( \frac{2^{-j-1}}{\log |\delta_j|} \log |z| \right) \vee (\psi(z) + 2^{-j}), & \text{if } |z| < \delta_j, \\ \frac{2^{-j-1}}{\log |\delta_j|} \log |z|, & \text{if } |z| \geq \delta_j. \end{cases}$$

Then  $\varphi_j \in \text{PSH}(\mathbb{C}^n)$  and  $\varphi_j(0) = 2^{-j}$ . It suffices to take  $\varphi = \sum_j \varphi_j$ .

thm:Cartan

**Theorem 1.3.1 (Cartan)** Consider  $x \in \Omega$  and a set  $E \subseteq \Omega$ . Assume that  $x \in E$ . Then the following are equivalent:

- (1)  $E$  is an  $\mathcal{F}$ -neighbourhood of  $x$ ;
- (2)  $\Omega \setminus E$  is thin at  $x$ .

**Proof** (2)  $\implies$  (1). We may assume that  $x \in \overline{\Omega \setminus E}$ . Otherwise, our assertion follows from [Proposition 1.3.1](#).

By [Proposition 1.3.2](#), there is  $\varphi \in \text{PSH}(\mathbb{C}^n)$  and an open neighbourhood  $U \subseteq \Omega$  of  $x$  such that

$$\varphi(x) > \sup_{y \in U \cap (\Omega \setminus E)} \varphi(y) =: \lambda.$$

Let  $F = \{y \in \Omega : \varphi(y) > \lambda\}$ . Then  $x \in F$  and  $F$  is  $\mathcal{F}$ -open. Moreover,  $U \cap F \subseteq E$ . By [Proposition 1.3.1](#), we conclude (1).

(1)  $\implies$  (2). We may always replace  $E$  by smaller  $\mathcal{F}$ -neighbourhoods of  $x$ . In particular, we may assume that  $E$  has the following form

$$\{y \in U : \varphi_1(y) > \lambda_1, \dots, \varphi_m(y) > \lambda_m\},$$

where  $U \subseteq \Omega$  is an open neighbourhood of  $x$ ,  $\varphi_1, \dots, \varphi_m$  are  $\mathbb{R}$ -valued psh functions on  $\Omega$  and  $\lambda_1, \dots, \lambda_m \in \mathbb{R}$ . Since a finite union of thin sets is still thin, we may assume that  $m = 1$ . In this case,  $\Omega \setminus E$  is clearly thin at  $x$ .  $\square$

thm:pf\_basis

**Theorem 1.3.2** A basis of the plurifine topology on  $\Omega$  is given by sets of the following form:

$$\{x \in U : \varphi(x) > 0\}, \tag{1.2}$$

{eq:basis\_fine}

where  $U \subseteq \Omega$  is an open subset and  $\varphi \in \text{PSH}(U)$ .

**Proof** We first show that sets of the form (1.2) are  $\mathcal{F}$ -open. By Theorem 1.3.1, it suffices to show its complement in  $\Omega$  is thin at  $x$ , which is obvious.

Now consider  $x \in \Omega$  and an  $\mathcal{F}$ -open neighbourhood  $V \subseteq \Omega$  of  $x$ . We want to find a set of the form (1.2) contained in  $V$  and containing  $x$ .

Write  $E = \Omega \setminus V$ . In case  $x \in \text{Int } V$ , there is nothing to prove. So we may assume that  $x \in \bar{E}$ . By Theorem 1.3.1,  $E$  is thin at  $x$ . By definition, there is an open neighbourhood  $U \subseteq \Omega$  of  $x$  and  $\varphi \in \text{PSH}(U)$  such that

$$\overline{\lim}_{y \rightarrow x, y \in E \cap U} \varphi(y) < \varphi(x).$$

We may assume that  $\varphi|_{E \cap U} \leq 0 < \varphi(x)$ , Then the set  $\{y \in U : \varphi(y) > 0\}$  suffices for our purpose.  $\square$

cor:pf\_compatible

**Corollary 1.3.1** *Let  $\Omega_1 \subseteq \Omega_2 \subseteq \mathbb{C}^n$  be two non-empty open subsets. Then the plurifine topology on  $\Omega_1$  is the same as the subspace topology induced from the plurifine topology on  $\Omega_2$ .*

**Corollary 1.3.2** *Let  $L$  be an affine subspace of  $\mathbb{C}^n$ , then the plurifine topology on  $L$  is the same as the subspace topology induced from the plurifine topology on  $\mathbb{C}^n$ .*

**Proof** We may assume that  $L = \mathbb{C}^k \times \{0\}$  for some  $k \leq n$ . We write the coordinate  $z$  on  $\mathbb{C}^n$  as  $(z', z'')$  with  $z' \in \mathbb{C}^k$  and  $z'' \in \mathbb{C}^{n-k}$ .

Consider an  $\mathcal{F}$ -open set  $U \subseteq \mathbb{C}^n$  and  $x = (x', 0) \in U \cap L$ . We want to show that  $U \cap L$  (identified with a subset of  $\mathbb{C}^k$ ) is an  $\mathcal{F}$ -neighbourhood of  $x'$  in  $L$ . By Theorem 1.3.2, we may assume that there are open subsets  $U' \subseteq \mathbb{C}^k$  containing  $x'$  and  $U'' \subseteq \mathbb{C}^{n-k}$  containing 0 together with a psh function  $\psi$  on  $U' \times U''$  such that

$$x \in \{(z', z'') \in U' \times U'' : \psi(z', z'') > 0\} \subseteq \Omega.$$

It follows that

$$x' \in \{z' \in U' : \psi(z', 0) > 0\} \subseteq U \cap L.$$

Conversely, if  $U \subseteq \mathbb{C}^k$  is an  $\mathcal{F}$ -open subset, we claim that  $U \times \mathbb{C}^{n-k}$  is  $\mathcal{F}$ -open in  $\mathbb{C}^n$ . In fact, suppose that  $(x', x'') \in U \times \mathbb{C}^{n-k}$ . By Theorem 1.3.1, we can find an open neighbourhood  $V \subseteq \mathbb{C}^k$  of  $x'$  and a psh function  $\varphi$  on  $V$  such that

$$x' \in \{y \in V : \varphi(y) > 0\} \subseteq U.$$

We define  $\psi(z', z'') := \varphi(z')$ . Then

$$(x', x'') \in \{y \in U \times \mathbb{C}^{n-k} : \psi(y) > 0\} \subseteq U \times \mathbb{C}^{n-k}.$$

cor:compactnhformbase

**Corollary 1.3.3** *Let  $\Omega \subseteq \mathbb{C}^n$  be an  $\mathcal{F}$ -open subset and  $x \in \Omega$ . Then  $x$  has a compact  $\mathcal{F}$ -neighbourhood contained in  $\Omega$ .*

**Proof** By Theorem 1.3.2, we may assume that there is a locally compact open set  $U \subseteq \mathbb{C}^n$  and a psh function  $\varphi$  on  $U$  such that  $\Omega = \{y \in U : \varphi(y) > 0\}$ .

Take a compact neighbourhood  $K$  of  $x$  in  $U$ . Now  $\{y \in K : \varphi(y) \geq \varphi(x)/2\}$  is a compact  $\mathcal{F}$ -neighbourhood of  $x$  contained in  $\Omega$ .  $\square$

cor:holomappfcont

**Corollary 1.3.4** *Let  $\Omega \in \mathbb{C}^n$ ,  $\Omega' \subseteq \mathbb{C}^{n'}$  be two domains and  $F: \Omega' \rightarrow \Omega$  be a surjective holomorphic map. Then  $F$  is  $\mathcal{F}$ -continuous.*

**Proof** It suffices to show that the inverse image  $F^{-1}(U)$  of each  $\mathcal{F}$ -open subset  $U \subseteq \Omega$  is  $\mathcal{F}$ -open. By [Theorem 1.3.2](#), after possibly shrinking  $\Omega$  and  $\Omega'$ , we may assume that  $U$  has the form  $\{x \in \Omega : \psi(x) > 0\}$ , where  $\psi \in \text{PSH}(\Omega)$ . Since  $F^*\psi \in \text{PSH}(\Omega')$  by [Proposition 1.1.4](#), we find that

$$F^{-1}(U) = \{y \in \Omega' : F^*\psi(y) > 0\}$$

is  $\mathcal{F}$ -open.  $\square$

### 1.3.2 Plurifine topology on manifolds

Let  $X$  be a complex manifold.

def:pftopologygeneral

**Definition 1.3.3** The *plurifine topology* on  $X$  is the topology with a basis consisting of sets of the form  $F^{-1}(V)$ , where  $U \subseteq X$  is an open subset and  $F: U \rightarrow \Omega$  is a biholomorphic morphism with  $\Omega \subseteq \mathbb{C}^n$  is a domain for some  $n \in \mathbb{N}$  and  $V \subseteq \Omega$  is  $\mathcal{F}$ -open.

It follows from [Corollary 1.3.4](#) that the plurifine topologies on domains defined in [Definition 1.3.3](#) and in [Definition 1.3.1](#) coincide.

We refer to [Definition 1.5.1](#) for the notion of quasi-plurisubharmonic functions.

prop:pshfunFcont

**Proposition 1.3.3** *Let  $\varphi \in \text{QPSH}(X)$ , then  $\varphi|_{\{\varphi \neq -\infty\}}$  is  $\mathcal{F}$ -continuous.*

**Proof** The problem is local, so we may assume that  $X \subseteq \mathbb{C}^n$  is a domain and  $\varphi = \psi + g$ , where  $\psi \in \text{PSH}(X)$  and  $g \in C^\infty(X)$  and  $|g| \leq C$  for some  $C > 0$ . Take an open interval  $(a, b) \subseteq \mathbb{R}$ , it suffices to show that

$$U := \{x \in X : a < \varphi(x) < b\} = \{x \in X : a - g(x) < \psi(x) < b - g(x)\}$$

is  $\mathcal{F}$ -open. Take  $x \in U$ , we can find an open neighbourhood  $V$  of  $x$  in  $U$  such that

$$\sup_{y \in V} (a - g(y)) < \psi(x) < \inf_{y \in V} (b - g(y)).$$

Therefore,

$$\left\{ z \in V : \sup_{y \in V} (a - g(y)) < \psi(z) < \inf_{y \in V} (b - g(y)) \right\}$$

is an  $\mathcal{F}$ -open neighbourhood of  $z$  in  $U$ . We conclude that  $U$  is  $\mathcal{F}$ -open.  $\square$

ma:pshfunfinitelocuspfdense

**Lemma 1.3.1** *Let  $Z \subseteq X$  be a pluripolar subset. Then*

$$\overline{X \setminus Z}^{\mathcal{F}} = X.$$

**Proof** The problem is local, so we may assume that  $X$  is a domain in  $\mathbb{C}^n$  and  $Z = \{\varphi = -\infty\}$  for some  $\varphi \in \text{PSH}(X)$ . We need to show that  $\{\varphi > -\infty\}$  is  $\mathcal{F}$ -dense.

Let  $x \in X$  be a point with  $\varphi(x) = -\infty$  and  $U \subseteq X$  be an  $\mathcal{F}$ -open neighbourhood of  $x$  in  $X$ . We need to show that  $U \cap \{\varphi > -\infty\} \neq \emptyset$ .

Thanks to **Theorem 1.3.2**, after shrinking  $U$ , we may assume that there is  $\psi \in \text{PSH}(X)$  such that  $U = \{\psi > 0\}$ . Observe that  $U$  is not a pluripolar set: otherwise,  $\psi \leq 0$  almost everywhere hence everywhere by **Proposition 1.2.5**. So  $\varphi|_U \not\equiv -\infty$ . We conclude.  $\square$

r:diffsupinfindeppluripolar

**Corollary 1.3.5** *Let  $\varphi, \psi \in \text{QPSH}(X)$ . Set*

$$W = \{x \in X : \min\{\varphi(x), \psi(x)\} = -\infty\}$$

*Then for any pluripolar set  $Z \subseteq X$ , we have*

$$\sup_{X \setminus W} (\varphi - \psi) = \sup_{X \setminus W \cup Z} (\varphi - \psi), \quad \inf_{X \setminus W} (\varphi - \psi) = \inf_{X \setminus W \cup Z} (\varphi - \psi).$$

**Proof** This is an immediate consequence of **Lemma 1.3.1** and **Proposition 1.3.3**.  $\square$

## 1.4 Lelong numbers and multiplier ideal sheaves

sec:Lelongmis

Let  $X$  be a complex manifold.

**Definition 1.4.1** Let  $\varphi \in \text{PSH}(X)$  and  $x \in X$ . The *Lelong number*  $\nu(\varphi, x)$  of  $\varphi$  at  $x$  is defined as follows: take an open neighbourhood  $U$  of  $x$  in  $X$  and a biholomorphic map  $F: U \rightarrow \Omega$ , where  $\Omega$  is a domain in  $\mathbb{C}^n$ . Then we define

$$\nu(\varphi, x) := \sup \left\{ \gamma \in \mathbb{R}_{\geq 0} : \varphi|_U(F^{-1}(y)) \leq \gamma \log |y - F(x)|^2 + O(1) \text{ as } y \rightarrow F(x) \right\}. \quad (1.3)$$

{eq:nuvarphix}

Observe that  $\nu(\varphi, x)$  does not depend on the choices of  $U$  and  $F$ . Furthermore, it follows from **Proposition 1.4.1** below that the supremum in (1.3) is a maximum.

**Remark 1.4.1** Our definition of the Lelong number is not standard. It differs from the standard definition by a factor of 2.

prop:Lelongreform

**Proposition 1.4.1** *Let  $\varphi \in \text{PSH}(B_n(0, 1))$ . Then*

$$\nu(\varphi, 0) = \lim_{r \rightarrow 0+} \frac{\sup_{B_n(0, r)} \varphi}{\log r^2} \in [0, \infty). \quad (1.4)$$

{eq:Lelongnewdef}

**Proof** It follows from [Proposition 1.1.2](#) that the limit in (1.4) exists and is finite. We shall denote the limit by  $v'(\varphi, 0)$  for the time being.

We first observe that by [Proposition 1.1.2](#),

$$\varphi(x) \leq v'(\varphi, 0) \log |x|^2 + \sup_{B_n(0,1)} \varphi \quad (1.5)$$

{eq:varphixlocalupperbd}

when  $x \in B_n(0, 1)$ . In particular,  $v(\varphi, x) \geq v'(\varphi, 0)$ .

In order to argue the reverse inequality, we may assume that  $v(\varphi, x) > 0$ .

Next observe that by (1.3), for each small enough  $\epsilon > 0$ , we can find  $r_0 \in (0, 1)$  and  $C > 0$  so that for all  $x \in B_n(0, r_0)$ , we have

$$\varphi(x) \leq (v(\varphi, 0) - \epsilon) \log |x|^2 + C.$$

It follows that  $v'(\varphi, 0) \geq v(\varphi, 0) - \epsilon$ . Letting  $\epsilon \rightarrow 0+$ , we conclude.  $\square$

We recall Siu's semicontinuity theorem.

thm:Siusemi

**Theorem 1.4.1** *Let  $\varphi \in \text{PSH}(X)$ , then the map  $X \ni x \mapsto v(\varphi, x)$  is upper semicontinuous with respect to the Zariski topology.*

[Dem12](#)

For an elegant proof we refer to [\[Dem12a, Theorem 2.10\]](#).

prop:Lelongmax

**Proposition 1.4.2** *Let  $\varphi, \psi \in \text{PSH}(X)$ ,  $\lambda \in \mathbb{R}_{>0}$  and  $x \in X$ , then*

$$\begin{aligned} v(\varphi \vee \psi, x) &= \min\{v(\varphi, x), v(\psi, x)\}, \\ v(\varphi + \psi, x) &= v(\varphi, x) + v(\psi, x), \\ v(\lambda\varphi, x) &= \lambda v(\varphi, x). \end{aligned}$$

**Proof** All properties are local, so we may assume that  $X = B_n(0, 1)$  for some  $n \in \mathbb{N}$ . All properties follow directly from [Proposition 1.4.1](#).  $\square$

cor:supslelong

**Corollary 1.4.1** *Let  $(\varphi_i)_{i \in I}$  be a non-empty family in  $\text{PSH}(X)$  uniformly bounded from above and  $x \in X$ , then*

$$v\left(\sup_{i \in I}^* \varphi_i, x\right) = \inf_{i \in I} v(\varphi_i, x).$$

**Proof** We observe that the  $\leq$  inequality. It remains to argue the reverse inequality.

It follows from [Proposition 1.2.2](#) that we may assume that  $I$  is countable. When  $I$  is finite, this is already proved in [Proposition 1.4.2](#). Otherwise, we may further assume that  $I = \mathbb{Z}_{>0}$ . Thanks to [Proposition 1.4.2](#), we may further assume that  $(\varphi_i)_{i \in \mathbb{Z}_{>0}}$  is an increasing sequence. Furthermore, since the problem is local, we may assume that  $X = B_n(0, 1)$  for some  $n \in \mathbb{N}$ . In this case, by (1.5), we have

$$\varphi_i(x) \leq v(\varphi_i, 0) \log |x|^2 + C$$

for all  $x \in B_n(0, 1)$  and all  $i \geq 1$  and  $C$  is a constant independent of  $i$ . In particular, thanks to [Proposition 1.2.3](#), for almost all  $x \in B_n(0, 1)$ , we have

$$\varphi(x) \leq \lim_{i \rightarrow \infty} v(\varphi_i, 0) \log |x|^2 + C.$$

Thanks of [Proposition 1.2.5](#), the same holds for all  $x$  and hence

$$v(\sup_{i \in \mathbb{Z}_{>0}}^* \varphi_i, x) \geq \lim_{i \rightarrow \infty} v(\varphi_i, x).$$

We conclude. □

**Definition 1.4.2** Let  $F \subseteq X$  be a non-empty analytic subset. Then we define the *generic Lelong number* of  $\varphi$  along  $F$  as

$$v(\varphi, F) := \min_{x \in F} v(\varphi, x).$$

Note that the minimum is obtained by [Theorem 1.4.1](#).

**Definition 1.4.3** Let  $\varphi \in \text{PSH}(X)$ . Let  $E$  be a prime divisor over  $X$  (see [Definition B.1.1](#)). Take a proper bimeromorphic morphism  $\pi: Y \rightarrow X$  from a complex manifold  $Y$  such that  $E$  is a prime divisor on  $Y$ , then we define the *generic Lelong number* of  $\varphi$  along  $E$  as

$$v(\varphi, E) := v(\pi^* \varphi, E).$$

It follows from [Theorem 1.4.1](#) that  $v(\varphi, E)$  does not depend on the choice of  $\pi$ .

**Definition 1.4.4** Let  $\varphi \in \text{PSH}(X)$ , the *multiplier ideal sheaf*  $\mathcal{I}(\varphi)$  of  $\varphi$  is by definition the ideal sheaf given by

$$\Gamma(U, \mathcal{I}(\varphi)) = \{f \in \mathcal{O}_X(U) : |f|^2 \exp(-\varphi) \in L_{\text{loc}}^1(U)\}$$

for any open subset  $U \subseteq X$ .

*Remark 1.4.2* This definition is different from a few standard references, where instead of  $\exp(-\varphi)$ , they use  $\exp(-2\varphi)$ . The conventions adopted in the current book is the most convenient one as far as the author knows. It simplifies a number of formulae.

**Proposition 1.4.3 (Nadel)** Let  $\varphi \in \text{PSH}(X)$ . Then  $\mathcal{I}(\varphi)$  is coherent.

See [Dem12](#), [Proposition 5.7](#). □

thm:multisubadd

**Theorem 1.4.2** Let  $\varphi, \psi \in \text{PSH}(X)$ , then

$$\mathcal{I}(\varphi + \psi) \subseteq \mathcal{I}(\varphi) \cdot \mathcal{I}(\psi).$$

See [Dem12](#), [Theorem 14.2](#).

The two invariants are related by the following simple result:

prop:Lelongnumfrommis

**Proposition 1.4.4** Let  $\varphi \in \text{PSH}(X)$  and  $E$  be a prime divisor over  $X$ . Then

$$v(\varphi, E) = \lim_{k \rightarrow \infty} \frac{1}{k} \text{ord}_E \mathcal{I}(k\varphi).$$

See [DX21, Proposition 2.14]. We remind the readers that this particular form of the formula is compatible with our conventions of  $\nu$  and  $I$ .

Also observe the following simple lemma:

lma:blowupLelong

**Lemma 1.4.1** *Let  $x \in X$  and  $\varphi \in \text{PSH}(X)$ . Let  $\pi: Y \rightarrow X$  be the blow-up of  $X$  at  $x$  with exceptional divisor  $E$ . Then*

$$\nu(\varphi, x) = \nu(\varphi, E),$$

See [Bou02a, Corollaire 1.1.8].

Conversely, the information of the generic Lelong numbers determines the multiplier ideal sheaves:

thm:valuativemulti

**Theorem 1.4.3** *Let  $\varphi \in \text{PSH}(X)$ . Let  $x \in X$  and  $f \in \mathcal{O}_{X,x}$ . Then the following are equivalent:*

- (1)  $f \in \mathcal{I}(\varphi)_x$ ;
- (2) *there exists  $\epsilon > 0$  such that for any prime divisor  $E$  over  $X$  such that  $x$  is contained in the center of  $E$  on  $X$ , we have*

$$\text{ord}_E(f) \geq (1 + \epsilon)\nu(\varphi, E) - \frac{1}{2}A_X(E).$$

Here  $A_X$  denotes the log discrepancy. We refer to [Bou17, Corollary 10.18] for the proof and the precise definition of  $A_X$ .

thm:stongopen

**Theorem 1.4.4 (Guan–Zhou)** *Let  $\varphi, \psi_j \in \text{PSH}(X)$  ( $j \in \mathbb{Z}_{>0}$ ) such that  $\psi_j$  is an increasing sequence converging to  $\varphi$  almost everywhere. Then for any  $x \in X$ , the germs satisfy*

$$\mathcal{I}(\psi_j)_x = \mathcal{I}(\varphi)_x$$

when  $j$  is large enough.

See [GZ15, Hiep14] for the proof.

prop:pull-backmis

**Proposition 1.4.5** *Let  $\pi: Y \rightarrow X$  be a smooth morphism between complex manifolds. Assume that  $\varphi \in \text{PSH}(X)$ , then*

$$\mathcal{I}(\pi^* \varphi) = \pi^* \mathcal{I}(\varphi).$$

**Proof** It follows from [SHC6, Théorème 3.10] that locally  $\pi$  can be written as the composition of an étale morphism and a projection. It suffices to handle the two cases separately.

Recall that in the complex analytic setting, an étale morphism is locally biholomorphic, so there is nothing to prove in this case.

Next, assume that  $Y = X \times U$ , where  $U \subseteq \mathbb{C}^n$  is a domain and  $\pi$  is the natural projection. It follows from Fubini's theorem that

$$\mathcal{I}(\pi^* \varphi) \subseteq \pi^* \mathcal{I}(\varphi).$$



The reverse inequality is proved in [Dem12, Proposition 14.3]<sup>3</sup>.  $\square$

def:restidealsheaf

**Definition 1.4.5** Given a coherent ideal sheaf  $\mathcal{I}$  on  $X$ , the *restriction*  $\text{Res}_Y \mathcal{I}$  is the inverse image ideal sheaf given by

$$\text{Res}_Y \mathcal{I} := \mathcal{I} / (\mathcal{I} \cap \mathcal{I}_Y), \quad (1.6)$$

{eq:RestI}

where  $\mathcal{I}_Y$  is the ideal sheaf defining  $Y$ .

In the literature, it is common to denote this sheaf by the misleading notation  $\mathcal{I}|_Y$ .

There is a natural morphism

$$i_Y^* \mathcal{I} = \mathcal{I} / (\mathcal{I} \cdot \mathcal{I}_Y) \rightarrow \text{Res}_Y \mathcal{I}, \quad (1.7)$$

{eq:pullbacktoinverseimage}

where  $i_Y: Y \rightarrow X$  is the inclusion.

thm:OT

**Theorem 1.4.5 (Ohsawa–Takegoshi)** Let  $Y$  be a submanifold of  $X$  and  $\varphi \in \text{PSH}(X)$ . Assume that  $\varphi|_Y \not\equiv -\infty$ , then

$$\mathcal{I}(\varphi|_Y) \subseteq \text{Res}_Y \mathcal{I}(\varphi).$$

See [Dem12a, Theorem 14.1].

## 1.5 Quasi-plurisubharmonic functions

In practice, it is important to consider a variant of plurisubharmonic functions. We will fix a complex manifold  $X$  together with a closed real smooth  $(1, 1)$ -form  $\theta$  on  $X$ .

def:qpsh

**Definition 1.5.1** A  $\theta$ -*plurisubharmonic function* on  $X$  is a function  $\varphi: X \rightarrow [-\infty, \infty)$  such that for each  $x \in X$  and each open neighbourhood  $U$  of  $x$  in  $X$  satisfying the condition that  $\theta = \text{dd}^c g$  for some smooth function  $g$  on  $U$ , we have  $g + \varphi|_U \in \text{PSH}(U)$ . The set of  $\theta$ -psh functions on  $X$  is denoted by  $\text{PSH}(X, \theta)$ .

A *quasi-plurisubharmonic function* on  $X$  is a function  $\varphi: X \rightarrow [-\infty, \infty)$  such that there exists a smooth closed real  $(1, 1)$ -form  $\theta'$  on  $X$  such that  $\varphi \in \text{PSH}(X, \theta')$ . The set of quasi-plurisubharmonic functions on  $X$  is denoted by  $\text{QPSH}(X)$ .

There is a natural non-strict partial order on  $\text{QPSH}(X)$  defined as follows:

def:parorder

**Definition 1.5.2** Assume that  $X$  is compact. Given  $\varphi, \psi \in \text{QPSH}(X)$ , we say that  $\varphi$  is *more singular* than  $\psi$  and write  $\varphi \leq \psi$  if there is  $C \in \mathbb{R}$  such that  $\varphi \leq \psi + C$ . We also say  $\psi$  is *less singular* than  $\varphi$  and write  $\psi \leq \varphi$ .

In case  $\varphi \leq \psi$  and  $\psi \leq \varphi$ , we say  $\varphi$  and  $\psi$  has the same singularity types. We write  $\varphi \sim \psi$  in this case.

<sup>3</sup> In [Dem12a, Proposition 14.3], Demailly used the highly non-standard notation  $f^* \mathcal{I}(\varphi)$  to denote the image of  $f^* \mathcal{I}(\varphi) \rightarrow \mathcal{O}_X$ .

*Remark 1.5.1* The proceeding results concerning plurisubharmonic functions can be extended *mutatis mutandis* to quasi-plurisubharmonic functions. We will apply these extensions without further explanations.

prop:L1compa

**Proposition 1.5.1** Assume that  $X$  is compact. Let  $\theta$  be a closed real smooth  $(1, 1)$ -form on  $X$ . Then for any  $a, b \in \mathbb{R}$ ,  $a \leq b$ , the set

$$\left\{ \varphi \in \text{PSH}(X, \theta) : \sup_X \varphi \in [a, b] \right\}$$

is compact with respect to the  $L^1$ -topology. Moreover,  $\varphi \mapsto \sup_X \varphi$  is  $L^1$ -continuous for  $\varphi \in \text{PSH}(X, \theta)$ .

This is an immediate consequence of [GZ17, Proposition 8.5, Exercise 1.20].

prop:Lelongnumberupperbound

**Proposition 1.5.2** Assume that  $X$  is compact. Let  $\theta$  be a closed real smooth  $(1, 1)$ -form on  $X$  and  $E$  be a prime divisor over  $E$ . Then

$$\sup \{v(\varphi, E) : \varphi \in \text{PSH}(X, \theta)\} < \infty.$$

*Proof* It follows from the proof of Corollary 1.4.1 that  $v(\bullet, E)$  is upper semi-continuous with respect to the  $L^1$ -topology on  $\text{PSH}(X, \theta)$ . Thus, the desired upper bound follows from Proposition 1.5.1.  $\square$

prop:PSHpullbij

**Proposition 1.5.3** Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism from a compact Kähler manifold  $Y$ . Let  $\theta$  be a closed real smooth  $(1, 1)$ -form on  $X$ . Then the pull-back gives a bijection

$$\pi^*: \text{PSH}(X, \theta) \xrightarrow{\sim} \text{PSH}(Y, \pi^*\theta).$$

This follows from a more general result Theorem B.1.1.

## 1.6 Analytic singularities

def:neatanasing

**Definition 1.6.1** We say  $\varphi \in \text{QPSH}(X)$  has *analytic singularities* if for each  $x \in X$ , we can find an open neighbourhood  $U$  of  $x$  such that  $\varphi|_U$  has the form:

$$c \log(|f_1|^2 + \cdots + |f_N|^2) + R, \quad (1.8)$$

{eq:anasinglocal}

where  $f_1, \dots, f_N$  are holomorphic functions on  $U$ ,  $c \in \mathbb{Q}_{>0}$  and  $R$  is a bounded function on  $U$ .

When  $R$  can be taken to be smooth, we say  $\varphi$  has *neat analytic singularities*.

Suppose that there is a coherent ideal  $\mathcal{I} \subseteq \mathcal{O}_X$  on  $X$  such that we can choose  $U$  so that the  $f_1, \dots, f_N$  can be chosen as the generators of  $\Gamma(U, \mathcal{I})$  and  $c$  is independent of the choice of  $U$ , we say  $\varphi$  has analytic singularities of *type*  $(c, \mathcal{I})$ .

Each potential with analytic singularities has a type. The type is not uniquely determined. We refer to [Bou02a] and [Bou02b] for the details.

prop:analysingclosed

**Proposition 1.6.1** *Let  $\varphi, \psi \in \text{QPSH}(X)$  be potentials with analytic singularities, then so are  $\lambda\varphi$  ( $\lambda \in \mathbb{Q}_{>0}$ ),  $\varphi + \psi$  and  $\varphi \vee \psi$ .*

**Proof** The  $\lambda\varphi$  assertion is trivial. The  $\vee$  assertion is proved in [Dem15, Proposition 4.1.8]. The addition assertion is easy and is left to the readers.  $\square$

**Definition 1.6.2** Let  $D$  be an effective  $\mathbb{Q}$ -divisor on  $X$ . We say  $\varphi \in \text{QPSH}(X)$  has *log singularities* (along  $D$ ) on  $X$  if for each  $x \in X$ , there is an open neighbourhood  $U$  of  $x$  such that

- (1)  $D|_U$  has finitely many irreducible components and can be written as

$$D|_U = \sum_{i=1}^N a_i D_i$$

with  $D_i$  being prime divisors on  $D$ ,  $a_i \in \mathbb{Q}_{>0}$  and there is a holomorphic function  $s_i$  on  $U$  defining  $D_i$ , and

- (2) we have

$$\varphi|_U = a_i \sum_i \log |s_i|^2 + R, \quad (1.9)$$

{eq:logsingreminder}

where  $R$  is a bounded function on  $U$ .

By Proposition 1.6.1,  $\varphi$  has analytic singularities.

lma:logsingrem

**Lemma 1.6.1** *Suppose that  $\theta$  is a closed smooth real  $(1, 1)$ -form on  $X$ , a compact Kähler manifold and  $\varphi \in \text{PSH}(X, \theta)$ . Suppose that  $\varphi$  has log singularities along an effective  $\mathbb{Q}$ -divisor  $D$  on  $X$ . Then the cohomology class  $[\theta] - [D]$  is nef.*

*Moreover, if in addition  $\theta_\varphi$  is a Kähler current, then the cohomology class  $[\theta] - [D]$  is ample.*

**Proof** The first assertion follows immediately from the fact that  $R$  in (1.9) has bounded coefficients.

The second assertion follows immediately from the first.  $\square$

The following proposition follows immediate from the definitions:

**Proposition 1.6.2** *Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism from a complex manifold  $Y$ . Suppose that  $\varphi \in \text{QPSH}(X)$  has analytic singularities (resp. has log singularities along an effective  $\mathbb{Q}$ -divisor  $D$ ). Then  $\pi^*\varphi$  has analytic singularities (resp. has log singularities along  $\pi^*D$ ).*

thm:resolvelogsing

**Theorem 1.6.1** *Assume that  $X$  is compact. Suppose that  $\varphi \in \text{QPSH}(X)$  has analytic singularities. Then there is a modification  $\pi: Y \rightarrow X$  such that  $\pi^*\varphi$  has log singularities.*

For a proof, we refer to the arguments on [MM07, Page 104].

def:quasiequising

**Definition 1.6.3** Let  $X$  be a compact Kähler manifold and  $\theta$  be a closed real smooth  $(1, 1)$ -form on  $X$ . Consider  $\varphi \in \text{PSH}(X, \theta)$ . A sequence  $(\varphi_j)_{j \in \mathbb{Z}_{>0}}$  in  $\text{QPSH}(X)$  is *quasi-equisingular approximation* of  $\varphi$  if

- (1)  $\varphi_j$  has analytic singularities for each  $j$ ;
- (2)  $\varphi_j$  is decreasing with limit  $\varphi$ ;
- (3) there is a decreasing sequence  $\epsilon_j \geq 0$  with limit 0 and a Kähler form  $\omega$  on  $X$  such that  $\varphi_j \in \text{PSH}(X, \theta + \epsilon_j \omega)$ ;
- (4) for each  $\lambda' > \lambda > 0$ , there is  $j > 0$  such that

$$I(\lambda' \varphi_j) \subseteq I(\lambda \varphi).$$

We also say  $\theta_{\varphi_j}$  is a quasi-equisingular approximation of  $\theta_\varphi$ .

def:analy-sing

**Definition 1.6.4** Let  $I \subseteq \mathcal{O}_X$  be an analytic coherent ideal sheaf and  $c \in \mathbb{Q}_{>0}$ . A function  $\varphi \in \text{QPSH}(X)$  is said to have *gentle analytic singularities* (of type  $(c, I)$ ) if

- (1)  $\varphi$  has analytic singularities of type  $(c, I)$ ;
- (2)  $e^{\varphi/c} : X \rightarrow \mathbb{R}_{\geq 0}$  is a smooth function;
- (3) there is a proper bimeromorphic morphism  $\pi : \tilde{X} \rightarrow X$  from a Kähler manifold  $\tilde{X}$  and an effective  $\mathbb{Z}$ -divisor  $D$  on  $\tilde{X}$  such that one can write  $\pi^* \varphi$  locally as

$$\pi^* \varphi = c \log |g|^2 + h,$$

where  $g$  is a local equation of the divisor  $D$  and  $h$  is smooth.

thm:qequi

**Theorem 1.6.2** Let  $X$  be a compact Kähler manifold and  $\theta$  be a closed real smooth  $(1, 1)$ -form on  $X$ . Then any  $\varphi \in \text{PSH}(X, \theta)$  admits a quasi-equisingular approximation  $(\varphi_j)_{j \in \mathbb{Z}_{>0}}$ .

Moreover, we can guarantee that  $\varphi_j$  has gentle analytic singularities of type  $(2^{-j}, I(2^j \varphi))$ .

We refer to [DPS01] for the proof.

Quasi-equisingular approximations are essentially unique in the following sense:

prop:compqequi

**Proposition 1.6.3** Let  $X$  be a compact Kähler manifold and  $\theta$  be a closed real smooth  $(1, 1)$ -form on  $X$ . Consider  $\varphi \in \text{PSH}(X, \theta)$ . Let  $(\varphi_j)_j$  and  $(\psi_j)_j$  be two quasi-equisingular approximations of  $\varphi$ . Then for any  $\epsilon > 0$  and any  $j > 0$ , we can find  $k_0 > 0$  such that for any  $k \geq k_0$ , we have

$$\psi_k \leq (1 - \epsilon) \varphi_j.$$

See [Dem15, Corollary 4.1.7].

def:Iinfy

**Definition 1.6.5** Assume that  $X$  is compact. Let  $\varphi \in \text{QPSH}(X)$  be a potential with analytic singularities. Then we define  $\mathcal{I}_\infty(\varphi)$  as the ideal sheaf consisting of germs  $f$  of holomorphic functions such that  $|f|^2 \exp(-\varphi)$  is locally bounded.

**Lemma 1.6.2** Assume that  $X$  is compact. Let  $\varphi \in \text{QPSH}(X)$  be a potential with analytic singularities. The sheaf  $\mathcal{I}_\infty(\varphi)$  is a coherent sheaf.

**Proof** By [Theorem 1.6.1](#), we may find a modification  $\pi: Y \rightarrow X$  such that  $\pi^*\varphi$  has log singularities. Observe that

$$\mathcal{I}_\infty(\varphi) = \pi_* \mathcal{I}(\pi^*\varphi),$$

so we may replace  $X$  and  $\varphi$  by  $Y$  and  $\pi^*\varphi$  and assume that  $\varphi$  has log singularities along an effective  $\mathbb{Q}$ -divisor  $D$ . We decompose  $D$  into its irreducible components:

$$D = \sum_{i=1}^N a_i D_i.$$

In this case, observe that

$$\mathcal{I}_\infty(\varphi) = \mathcal{O}\left(-\sum_{i=1}^N ([a_i] D_i)\right)$$

is clearly coherent. □

lma:IandIinf

**Lemma 1.6.3** *Assume that  $X$  is compact. Let  $\varphi \in \text{QPSH}(X)$  be a potential with analytic singularities. Then for any  $\epsilon > 0$ , we can find  $k_0 > 0$  such that for each  $k \geq k_0$ , we have*

$$\mathcal{I}(k(1+\epsilon)\varphi) \subseteq \mathcal{I}_\infty(k\varphi).$$

See [Dem15](#), Proposition 4.1.6].

thm:CT-thm-refined'

**Theorem 1.6.3** *Let  $X$  be a connected compact Kähler manifold and  $Y \subseteq X$  be a connected submanifold. Take a Kähler form  $\omega$  on  $X$  and  $\varphi \in \text{PSH}(Y, \omega|_Y)$  such that  $\omega|_Y + \text{dd}^c \varphi$  is a Kähler current and that  $e^\varphi$  is a Hölder continuous function on  $Y$ . Then there exists  $\tilde{\varphi} \in \text{PSH}(X, \omega)$  satisfying*

- (1)  $\tilde{\varphi}|_Y = \varphi$ ;
- (2)  $\omega_{\tilde{\varphi}}$  is a Kähler current.

*In addition, if  $\varphi$  has analytic singularities, then so does  $\tilde{\varphi}$ .*

See [DRWNXZ](#), Theorem 6.1].

## 1.7 The space of currents

Let  $X$  be a connected compact Kähler manifold of dimension  $n$  and  $\alpha \in H^{1,1}(X, \mathbb{R})$ .

**Definition 1.7.1** Let  $Y$  be a complex manifold and  $m \in \mathbb{N}$ . We say an  $(m, m)$ -current  $T$  on  $Y$  is *positive* if either  $m > n$  or for any smooth  $(1, 0)$ -forms  $\beta_1, \dots, \beta_{n-m}$  on  $X$ , the measure

$$T \wedge i\beta_1 \wedge \overline{\beta_1} \wedge \dots \wedge i\beta_{n-m} \wedge \overline{\beta_{n-m}}$$

is positive.

**Definition 1.7.2** We say  $\alpha$  is *pseudo-effective* if there is a closed positive  $(1, 1)$ -current in  $\alpha$ .

We say  $\alpha$  is *big* if there is a closed positive  $(1, 1)$ -current  $T$  in  $\alpha$  dominating a Kähler form. Such currents are called *Kähler currents*.

def:spaceofcurrents

**Definition 1.7.3** We introduce the following notations:

- (1)  $\mathcal{Z}_+(X)$  denotes the space of closed positive  $(1, 1)$ -currents on  $X$ ;
- (2) given a pseudo-effective  $(1, 1)$ -class  $\alpha$  on  $X$ , we write  $\mathcal{Z}_+(X, \alpha)$  for the set of  $T \in \mathcal{Z}_+(X)$  such that  $[T] = \alpha$ ;

Given  $T, T' \in \mathcal{Z}_+(X)$ , we write  $T \leq T'$  and say  $T$  is *more singular* than  $T'$  if when we write  $T = \theta + \text{dd}^c \varphi$ ,  $T' = \theta' + \text{dd}^c \varphi'$ , we have  $\varphi \leq \varphi'$ . We write  $T \sim T'$  if  $T \leq T'$  and  $T' \leq T$ . In this case, we say  $T$  and  $T'$  have the same *singularity type*.

rmk:qpshtocurrents

**Remark 1.7.1** Observe that

$$\mathcal{Z}_+(X)/\sim \cong \text{QPSH}(X)/\sim$$

canonically. The correspondence sends the class of a closed positive current  $\theta_\varphi = \theta + \text{dd}^c \varphi$  to the class of  $\varphi$ .

We will adopt the following convention: whenever we have a notion for quasi-plurisubharmonic functions which depends only on the singularity type, we use the same notation and the same definition for closed positive  $(1, 1)$ -currents.

def:polarlocus

**Definition 1.7.4** Given  $T \in \mathcal{Z}_+(X)$ . We represent  $T$  as  $\theta + \text{dd}^c \varphi$  for some closed smooth real  $(1, 1)$ -form  $\theta$  on  $X$  and  $\varphi \in \text{PSH}(X, \theta)$ , then the *polar locus* of  $T$  is defined as the set  $\{\varphi = -\infty\}$ .

It is clear that the polar locus of  $T$  is independent of the choices of  $\theta$  and  $\varphi$ .

lma:Siudec

**Lemma 1.7.1 (Siu's decomposition)** *Let  $E$  be a prime divisor on  $X$ . Then for any closed positive  $(1, 1)$ -current  $T$  on  $X$ , the difference  $T - \nu(T, E)[E]$  is a closed positive  $(1, 1)$ -current.*

Here  $[E]$  is the current of integration associated with  $E$ . See [GH14, Page 386, Example 1] for the definition of  $[E]$ . See [Dem12a, Lemma 2.17] for the proof.

## 1.8 Plurisubharmonic metrics on line bundles

A natural source of quasi-plurisubharmonic functions is the metrics on line bundles.

Let  $X$  be a connected compact Kähler manifold and  $L$  be a holomorphic line bundle on  $X$ . Usually, we do not distinguish  $L$  from the associated invertible sheaf  $\mathcal{O}_X(L)$ .

**Definition 1.8.1** Let  $V$  be a 1-dimensional complex linear space. A *Hermitian form*  $h$  on  $V$  is a map  $h: V \times V \rightarrow \mathbb{C}$  such that

- (1)  $h$  is  $\mathbb{C}$ -linear in the second variable and conjugate linear in the first, and  
 (2)

$$|v|_h^2 := h(v, v) \in \mathbb{R}_{>0}$$

for each  $v \in V \setminus \{0\}$ .

We usually identify  $h$  with the quadratic form  $V \rightarrow \mathbb{R}$  sending  $v$  to  $|v|_h^2$ .

The *singular Hermitian form* on  $V$  is the map  $V \rightarrow \{0, \infty\}$  sending 0 to 0 and other elements to  $\infty$ .

We write  $|v|_h = \sqrt{|v|_h^2}$ .

def:Hermmetric

**Definition 1.8.2** A *Hermitian metric*  $h$  on  $L$  is a family of Hermitian forms  $(h_x)_{x \in X}$ , such that

- (1) for each  $x \in X$ ,  $h_x$  is a Hermitian form on  $L_x$ , and  
 (2) for each local section  $s$  of  $\mathcal{O}_X(L)$ , the map  $x \mapsto |s(x)|_{h_x}$  is smooth.

The pair  $(L, h)$  is called a *Hermitian line bundle*. We shall write  $\text{dd}^c h = c_1(L, h)$  for the first Chern form of  $h$ , normalized so that

$$[c_1(L, h)] = c_1(L).$$

The map  $x \mapsto |s(x)|_{h_x}$  will be denoted by  $|s|$ .

prop:LelongPoincare

**Proposition 1.8.1 (Lelong–Poincaré)** Let  $s \in H^0(X, L)$  be non-zero and  $h$  be a Hermitian metric on  $L$ . Then

$$c_1(L, h) + \text{dd}^c \log |s|_h^2 = [Z(s)],$$

where  $Z(s)$  is the prime divisor defined by  $s$  and  $[\bullet]$  denote the associated current of integration.

See [Dem12](#), (3.11).  
 See [Dem12a](#), (3.11).

**Definition 1.8.3** A *plurisubharmonic metric*  $h$  on  $L$  is a family  $(h_x)_x$  such that

- (1) for each  $x \in X$ ,  $h_x$  is either a Hermitian form on  $L_x$  or the singular Hermitian form, and  
 (2) there is a Hermitian metric  $h_0$  on  $L$  and  $\varphi \in \text{PSH}(X, c_1(L, h_0))$  such that for each  $x \in X$  and each  $v \in L_x$ , we have

$$|v|_{h_x}^2 = \begin{cases} 0, & \text{if } v = 0; \\ |v|_{h_{0,x}}^2 e^{-\varphi(x)}, & \text{if } v \neq 0. \end{cases} \quad (1.10) \quad \text{\texttt{\{eq:htwist\}}}$$

The (first) *Chern current* of  $h$  is by definition

$$\text{dd}^c h = c_1(L, h) := c_1(L, h_0) + \text{dd}^c \varphi.$$

We shall write the plurisubharmonic metric defined by (1.10) as  $h \exp(-\varphi)$ . As the readers can easily verify, our conventions guarantee that  $c_1(L, h)$  does not depend on the choice of  $h_0$ .

*Remark 1.8.1* In the literature, some people prefer the convention that in (1.10), neither side has the square.

We shall need the following Ohsawa–Takegoshi type extension theorem.

thm: OT\_ext

**Theorem 1.8.1** *Assume that  $L$  is big and  $T$  is a holomorphic line bundle on  $X$ . Fix a Hermitian metric  $r$  on  $T$ . Take a Kähler form  $\omega$  on  $X$ . Let  $Y \subseteq X$  be a connected submanifold of dimension  $m$ . Suppose that  $\varphi \in \text{PSH}(X, \theta - \delta\omega)$  for some  $\delta > 0$  and  $\varphi|_Y \not\equiv -\infty$ . Then there exists  $k_0(\delta, r) > 0$  such that for all  $k \geq k_0$  and  $s \in H^0(Y, T \otimes L|_Y^k \otimes \mathcal{I}(k\varphi|_Y))$ , there exists an extension  $\tilde{s} \in H^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi))$  such that*

$$\int_X (h^k \otimes r)(\tilde{s}, \tilde{s}) e^{-k\varphi} \omega^n \leq C \int_Y (h^k \otimes r)(s, s) e^{-k\varphi|_Y} \omega|_Y^m,$$

where  $C > 0$  is an absolute constant, independent of the data  $(\varphi, s, k)$ .

This is a special case of [His12](#), Theorem 1.4].

prop: Bergman\_approx

**Proposition 1.8.2** *Let  $(L, h)$  be a Hermitian line bundle on  $X$  and set  $\theta = c_1(L, h)$ . Let  $(T, h_T)$  be a Hermitian line bundle on  $X$ . Assume that  $\varphi \in \text{PSH}(X, \theta)$  is a potential with analytic singularities such that  $\theta_\varphi$  is a Kähler current. Fix a Kähler form  $\omega$  on  $X$ . For each  $k \geq 1$ , we let*

$$\varphi_k := \frac{1}{k} \log \sup_{\substack{s \in H^0(X, L^k \otimes T) \\ \int_X h^k \otimes h_T(s, s) e^{-k\varphi} \omega^n \leq 1}} h^k \otimes h_T(s, s). \quad (1.11)$$

{eq: Bergman\_seq\_def}

Then for any  $k \geq 0$ ,

$$\varphi \leq \varphi_k \leq \alpha_k \varphi,$$

where  $\alpha_k \in (0, 1)$  is an increasing sequence with limit 1.

Note that when  $k$  is large enough,  $\varphi_k \in \text{PSH}(X, \theta)$ . We refer to [DX21](#), Remark 2.9] for the proof.



## Chapter 2

### Non-pluripolar products

chap:npp

Let  $X$  be a complex manifold and  $\varphi_1, \dots, \varphi_p \in \text{PSH}(X)$  ( $p \in \mathbb{N}$ ). When the functions  $\varphi_1, \dots, \varphi_p$  are all smooth, there is an obvious definition of a current

$$\text{dd}^c \varphi_1 \wedge \dots \wedge \text{dd}^c \varphi_p \quad (2.1)$$

{eq:mixedMAtype}

by the usual differential calculus. It is of interest to extend this construction to the case where the  $\varphi_i$ 's have worse regularities.

There are a number of different approaches to this problem. In this book, we will choose the so-called non-pluripolar theory due to Bedford–Taylor, Guedj–Zeriahi and Boucksom–Eyssidieux–Guedj–Zeriahi. The reason is that the non-pluripolar theory is the only known theory satisfying the following two features: it is defined for all psh singularities (at least in the global setting) and it satisfies a monotonicity theorem.

We will recall the Bedford–Taylor theory in [Section 2.1](#) and the non-pluripolar theory in [Section 2.2](#).

Some key properties of the non-pluripolar products are recalled in [Section 2.3](#).

#### 2.1 Bedford–Taylor theory

sec:BTtheory

Let  $X$  be a complex manifold and  $\varphi_1, \dots, \varphi_p \in \text{PSH}(X)$  ( $p \in \mathbb{N}$ ) be locally bounded plurisubharmonic functions on  $X$ <sup>1</sup>. In this case, there is a canonical definition of the Monge–Ampère type product (2.1).

**Definition 2.1.1** We define the closed positive  $(p, p)$ -current (2.1) on  $X$  as follows: we make an induction on  $p \geq 0$ . When  $p = 0$ , we define (2.1) as the  $(0, 0)$ -current  $[X]$ . When  $p > 0$ , we let

$$\text{dd}^c \varphi_1 \wedge \dots \wedge \text{dd}^c \varphi_p := \text{dd}^c (\varphi_1 \text{dd}^c \varphi_2 \wedge \dots \wedge \text{dd}^c \varphi_p).$$

<sup>1</sup> In the literature, some people use  $\text{PSH}(X) \cap L_{\text{loc}}^\infty(X)$  to denote such functions, which is an abuse of notation. It is legitimate thanks to the rigidity [Theorem 1.1.3](#).

We call (2.1) the *Bedford–Taylor product*.

**Proposition 2.1.1** *The product  $\mathrm{dd}^c \varphi_1 \wedge \cdots \wedge \mathrm{dd}^c \varphi_p$  is a closed positive  $(p, p)$ -current on  $X$ . Moreover, the product is symmetric in the  $\varphi_i$ 's.*

See [GZ17, Proposition 3.3, Corollary 3.12].

The Bedford–Taylor theory has many satisfactory properties.

thm:contMA

**Theorem 2.1.1** *Let  $(\varphi_i^j)_j$  be decreasing sequences (resp. increasing sequences) of locally bounded psh functions on  $X$  converging (resp. converging a.e.) to locally bounded psh function  $\varphi_i$ , where  $i = 1, \dots, p$ . Then*

$$\varphi_0^j \mathrm{dd}^c \varphi_1^j \wedge \cdots \wedge \mathrm{dd}^c \varphi_p^j \rightarrow \varphi_0 \mathrm{dd}^c \varphi_1 \wedge \cdots \wedge \mathrm{dd}^c \varphi_p$$

as  $j \rightarrow \infty$ . In particular, if  $\varphi_0^j$  is the constant sequence 1, we have

$$\mathrm{dd}^c \varphi_1^j \wedge \cdots \wedge \mathrm{dd}^c \varphi_p^j \rightarrow \mathrm{dd}^c \varphi_1 \wedge \cdots \wedge \mathrm{dd}^c \varphi_p.$$

Here the notation  $\rightarrow$  denotes the weak-\* convergence of currents.

We refer to [GZ17, Theorem 3.18, Theorem 3.23] for the proofs.

## 2.2 The non-pluripolar products

sec:npp

The proof of all results in this section can be found in [BEGZ10].

Let  $X$  be a connected complex manifold of dimension  $n$ .

**Definition 2.2.1** Let  $\varphi_1, \dots, \varphi_p \in \mathrm{PSH}(X)$ . We set

$$O_k := \bigcap_{j=1}^p \{\varphi_j > -k\}, \quad k \in \mathbb{Z}_{>0}.$$

We say that  $\mathrm{dd}^c \varphi_1 \wedge \cdots \wedge \mathrm{dd}^c \varphi_p$  is *well-defined* if for each open subset  $U \subseteq X$  admitting a Kähler form  $\omega$  on  $U$ , for each compact subset  $K \subseteq U$ , we have

$$\sup_{k \geq 0} \int_{K \cap O_k} \left( \bigwedge_{j=1}^p \mathrm{dd}^c (\varphi_j \vee (-k)) \right) \wedge \omega^{n-p} < \infty. \quad (2.2) \quad \{\text{eq:welldefinepluri}\}$$

In this case, we define the *non-pluripolar product*  $\mathrm{dd}^c \varphi_1 \wedge \cdots \wedge \mathrm{dd}^c \varphi_p$  by

$$\mathbb{1}_{O_k} \mathrm{dd}^c \varphi_1 \wedge \cdots \wedge \mathrm{dd}^c \varphi_p = \mathbb{1}_{O_k} \bigwedge_{j=1}^p \mathrm{dd}^c (\varphi_j \vee (-k)) \quad (2.3) \quad \{\text{eq:npp}\}$$

on  $\bigcup_{k \geq 0} O_k$  and make a zero-extension to  $X$ .

prop:npp1

**Proposition 2.2.1** *Let  $\varphi_1, \dots, \varphi_p \in \text{PSH}(X)$ .*

- (1) *The product  $\text{dd}^c \varphi_1 \wedge \dots \wedge \text{dd}^c \varphi_p$  is local with respect to the plurifine topology in the following sense: Let  $O \subseteq X$  be a plurifine open subset and  $\psi_1, \dots, \psi_p \in \text{PSH}(X)$ . Assume that*

$$\varphi_j|_O = \psi_j|_O, \quad j = 1, \dots, p,$$

*and that*

$$\bigwedge_{j=1}^p \text{dd}^c \varphi_j \text{ and } \bigwedge_{j=1}^p \text{dd}^c \psi_j$$

*are both well-defined, then*

$$\left. \bigwedge_{j=1}^p \text{dd}^c \varphi_j \right|_O = \left. \bigwedge_{j=1}^p \text{dd}^c \psi_j \right|_O. \quad (2.4) \quad \{\text{eq:ppp1}\}$$

*If furthermore  $O$  is open in the usual topology, then the product*

$$\bigwedge_{j=1}^p \text{dd}^c \varphi_j|_O$$

*on  $O$  is well-defined and*

$$\left. \bigwedge_{j=1}^p \text{dd}^c \varphi_j \right|_O = \bigwedge_{j=1}^p \text{dd}^c \varphi_j|_O. \quad (2.5) \quad \{\text{eq:ppp2}\}$$

*Let  $\mathcal{U}$  be an open covering of  $X$ . Then  $\text{dd}^c \varphi_1 \wedge \dots \wedge \text{dd}^c \varphi_p$  is well-defined if and only if each of the following product is well-defined*

$$\bigwedge_{j=1}^p \text{dd}^c \varphi_j|_U, \quad U \in \mathcal{U}.$$

- (2) *The current  $\text{dd}^c \varphi_1 \wedge \dots \wedge \text{dd}^c \varphi_p$  and the fact that it is well-defined depend only on the currents  $\text{dd}^c \varphi_j$ , not on specific  $\varphi_j$ .*  
 (3) *When  $\varphi_1, \dots, \varphi_p \in L_{\text{loc}}^\infty(X)$ , the product  $\text{dd}^c \varphi_1 \wedge \dots \wedge \text{dd}^c \varphi_p$  is well-defined and is equal to the Bedford–Taylor product.*  
 (4) *Assume that  $\text{dd}^c \varphi_1 \wedge \dots \wedge \text{dd}^c \varphi_p$  is well-defined, then  $\text{dd}^c \varphi_1 \wedge \dots \wedge \text{dd}^c \varphi_p$  puts not mass on pluripolar sets.*  
 (5) *Assume that  $\text{dd}^c \varphi_1 \wedge \dots \wedge \text{dd}^c \varphi_p$  is well-defined, then  $\bigwedge_{j=1}^p \text{dd}^c \varphi_j$  is a closed positive  $(p, p)$ -current on  $X$ .*  
 (6) *The product is multilinear: Let  $\psi_1 \in \text{PSH}(X)$ , then*

$$\text{dd}^c(\varphi_1 + \psi_1) \wedge \bigwedge_{j=2}^p \text{dd}^c \varphi_j = \text{dd}^c \varphi_1 \wedge \bigwedge_{j=2}^p \text{dd}^c \varphi_j + \text{dd}^c \psi_1 \wedge \bigwedge_{j=2}^p \text{dd}^c \varphi_j \quad (2.6) \quad \{\text{eq:ppp6}\}$$

in the sense that left-hand side is well-defined if and only if both terms on right-hand side are well-defined, and the equality holds in that case.

**Definition 2.2.2** Let  $T_1, \dots, T_p$  be closed positive  $(1, 1)$ -currents on  $X$ . We say that  $T_1 \wedge \dots \wedge T_p$  is *well-defined* if there exists an open covering  $\mathcal{U}$  of  $X$ , such that on each  $U \in \mathcal{U}$ , we can find  $\varphi_j^U \in \text{PSH}(U)$  ( $j = 1, \dots, p$ ) such that

$$\text{dd}^c \varphi_j^U = T_j, \quad j = 1, \dots, p$$

and such that  $\text{dd}^c \varphi_1^U \wedge \dots \wedge \text{dd}^c \varphi_p^U$  is well-defined. In this case, we define the *non-pluripolar product*  $T_1 \wedge \dots \wedge T_p$  as the closed positive  $(p, p)$ -current on  $X$  defined by

$$(T_1 \wedge \dots \wedge T_p)|_U = \text{dd}^c \varphi_1^U \wedge \dots \wedge \text{dd}^c \varphi_p^U, \quad U \in \mathcal{U}. \quad (2.7)$$

{eq:ppp5}

**Proposition 2.2.1** can be formulated in terms of currents without any difficulty.

prop:nppwelldef

**Proposition 2.2.2** Let  $X$  be a compact Kähler manifold and  $T_1, \dots, T_p$  are closed positive  $(1, 1)$ -currents on  $X$ . Then  $T_1 \wedge \dots \wedge T_p$  is well-defined.

This proposition explains why we usually work in the setting of compact Kähler manifolds.

### 2.3 Properties of non-pluripolar products

sec:nppprop

Let  $X$  be a connected compact Kähler manifold of dimension  $n$  and  $\theta, \theta_1, \dots, \theta_n$  be closed real smooth  $(1, 1)$ -forms on  $X$ .

We write

$$\text{PSH}(X, \theta)_{>0} = \left\{ \varphi \in \text{PSH}(X, \theta) : \int_X \theta_\varphi^n > 0 \right\}. \quad (2.8)$$

{eq:PSHpos}

The non-pluripolar product  $\theta_\varphi^n$  is well-defined thanks to **Proposition 2.2.2**.

**Remark 2.3.1** Suppose that  $X$  is a connected complex manifold of dimension 0, namely,  $X$  is a single point. In this case, by definition, the non-pluripolar product  $\theta_\varphi^n$  is given by the current of integration at the unique point. So  $\text{PSH}(X, \theta)_{>0} = \text{PSH}(X, \theta) \cong \mathbb{R}$  in this case and  $\int_X \theta_\varphi^n = 1$  for all  $\varphi \in \text{PSH}(X, \theta)$ .

prop:nppmassinv

**Proposition 2.3.1** Let  $\pi : Y \rightarrow X$  be a proper bimeromorphic morphism from a Kähler manifold  $Y$  and  $\varphi_i \in \text{PSH}(X, \theta_i)$  for  $i = 1, \dots, n$ . Then

$$\int_Y \pi^* \theta_{1, \pi^* \varphi_1} \wedge \dots \wedge \pi^* \theta_{n, \pi^* \varphi_n} = \int_X \theta_{1, \varphi_1} \wedge \dots \wedge \theta_{n, \varphi_n}.$$

**Proof** This follows immediately from **Proposition 2.2.1** (1) and (4).  $\square$

We shall write

$$V_\theta = \sup \{ \varphi \in \text{PSH}(X, \theta) : \varphi \leq 0 \}. \quad (2.9)$$

{eq:Vtheta}

It follows from [Proposition 1.2.1](#) that  $V_\theta \in \text{PSH}(X, \theta)$  if  $\text{PSH}(X, \theta) \neq \emptyset$ .

thm:semicon

**Theorem 2.3.1 (Semicontinuity theorem)** *Let  $\varphi_j, \varphi_j^k \in \text{PSH}(X, \theta_j)$  ( $k \in \mathbb{Z}_{>0}$ ,  $j = 1, \dots, n$ ). Let  $\chi \geq 0$  be a bounded function such that there are  $\eta_1, \eta_2 \in \text{QPSH}(X)$  with  $\eta_1 + \chi = \eta_2$ .*

*Assume that for any  $j = 1, \dots, n$  and  $i = 1, \dots, m$ , as  $k \rightarrow \infty$ , either  $\varphi_j^k$  decreases to  $\varphi_j \in \text{PSH}(X, \theta)$  or increases to  $\varphi_j \in \text{PSH}(X, \theta)$  almost everywhere. Then for any open set  $U \subseteq X$ , we have*

$$\lim_{k \rightarrow \infty} \int_U \chi \theta_{1, \varphi_1^k} \wedge \dots \wedge \theta_{n, \varphi_n^k} \geq \int_U \chi \theta_{1, \varphi_1} \wedge \dots \wedge \theta_{n, \varphi_n}. \quad (2.10)$$

{eq:semicon1}

See [\[DDNL18mono, DDNL18b, Theorem 2.3\]](#).

thm:mono

**Theorem 2.3.2 (Monotonicity theorem)** *Let  $\varphi_j, \psi_j \in \text{PSH}(X, \theta_j)$  for  $j = 1, \dots, n$ . Assume that  $\varphi_j \geq \psi_j$  for every  $j$ , then*

$$\int_X \theta_{1, \varphi_1} \wedge \dots \wedge \theta_{n, \varphi_n} \geq \int_X \theta_{1, \psi_1} \wedge \dots \wedge \theta_{n, \psi_n}.$$

See [\[DDNL18mono, DDNL18b, Theorem 1.1\]](#).

As a corollary, we obtain that

cor:incseqnppcont

**Corollary 2.3.1** *Fix a directed set  $I$ . For each  $j = 1, \dots, n$ , take an increasing net  $(\varphi_j^i)_{i \in I}$  in  $\text{PSH}(X, \theta_j)$ , uniformly bounded from above. Set*

$$\varphi_j := \sup_{i \in I}^* \varphi_j^i.$$

Then

$$\lim_{i \in I} \int_X \theta_{1, \varphi_1^i} \wedge \dots \wedge \theta_{n, \varphi_n^i} = \int_X \theta_{1, \varphi_1} \wedge \dots \wedge \theta_{n, \varphi_n}. \quad (2.11)$$

{eq:increseqnppcont}

**Proof** We may assume that  $I$  is infinite as there is nothing to prove otherwise. Thanks to [Theorem 2.3.2](#), we already know the  $\leq$  inequality in (2.11). We prove the reverse inequality. When  $I \cong \mathbb{Z}_{>0}$  as directed sets, the reverse inequality follows from [Theorem 2.3.1](#). In general, by Choquet's lemma [Proposition 1.2.2](#), we can find a countable infinite subset  $R \subseteq I$  such that

$$\sup_{r \in R}^* \varphi_j^r = \sup_{i \in I}^* \varphi_j^i$$

for all  $j = 1, \dots, n$ . We fix a bijection  $R \cong \mathbb{Z}_{>0}$ . For any  $j = 1, \dots, n$ , we will then denote elements  $\varphi_j^r$  ( $r \in R$ ) by  $\varphi_j^1, \varphi_j^2, \dots$ . We shall write

$$\psi_j^a = \varphi_j^1 \vee \dots \vee \varphi_j^a$$

for each  $a \in \mathbb{Z}_{>0}$ .

It follows from the fact that  $I$  is a directed set and [Theorem 2.3.2](#) that

$$\lim_{i \in I} \int_X \theta_{1, \varphi_1^i} \wedge \cdots \wedge \theta_{n, \varphi_n^i} \geq \lim_{a \rightarrow \infty} \int_X \theta_{1, \psi_1^a} \wedge \cdots \wedge \theta_{n, \psi_n^a}.$$

From the special case mentioned above, we know that the right-hand side is exactly the right-hand side of [\(2.11\)](#), so we conclude.  $\square$

lma:pathoenvelope

**Lemma 2.3.1** *Let  $\varphi, \psi \in \text{PSH}(X, \theta)$ ,  $\varphi \leq \psi$  and  $\int_X \theta_\varphi^n > 0$ . Then for any*

$$a \in \left(1, \left(\frac{\int_X \theta_\psi^n}{\int_X \theta_\psi^n - \int_X \theta_\varphi^n}\right)^{1/n}\right), \quad (2.12) \quad \{\text{eq:arangetemp}\}$$

*there is  $\eta \in \text{PSH}(X, \theta)_{>0}$  such that*

$$a^{-1}\eta + (1 - a^{-1})\psi \leq \varphi.$$

The fraction in [\(2.12\)](#) is understood as  $\infty$  if  $\int_X \theta_\psi^n = \int_X \theta_\varphi^n$ . In particular, thanks to [Theorem 2.3.2](#), the interval [\(2.12\)](#) is non-empty.

We write

$$P_\theta(a\varphi + (1 - a)\psi) = \sup^* \{ \eta \in \text{PSH}(X, \theta) : a^{-1}\eta + (1 - a^{-1})\psi \leq \varphi \} \in \text{PSH}(X, \theta). \quad (2.13) \quad \{\text{eq:perversePtheta}\}$$

*Remark 2.3.2* The notation  $P_\theta(a\varphi + (1 - a)\psi)$  might lead to some potential confusions. But the author cannot come up with a better notation.

Observe that

$$a^{-1}P_\theta(a\varphi + (1 - a)\psi) + (1 - a^{-1})\psi \leq \varphi. \quad (2.14)$$

In fact, this equation holds outside a pluripolar set by [Proposition 1.2.3](#), hence it holds everywhere by [Proposition 1.2.5](#).

**Proof** Without loss of generality, we may assume that  $\varphi \leq \psi \leq 0$ .

We refer to [\[DDNL21b, Lemma 4.3\]](#) for the proof of the existence of  $\eta \in \text{PSH}(X, \theta)$  satisfying the given inequality. Next we argue that  $P_\theta(a\varphi + (1 - a)\psi) \in \text{PSH}(X, \theta)_{>0}$ . Choose

$$a' \in \left(a, \left(\frac{\int_X \theta_\psi^n}{\int_X \theta_\psi^n - \int_X \theta_\varphi^n}\right)^{1/n}\right).$$

It follows from [\(2.13\)](#) that

$$P_\theta(a\varphi + (1 - a)\psi) \geq \frac{a}{a'}P_\theta(a'\varphi + (1 - a')\psi) + \frac{a' - a}{a'}\varphi. \quad (2.15) \quad \{\text{eq:Pthetalowerbdttemp1}\}$$

Therefore, by [Theorem 2.3.2](#), we have

$$\int_X \theta_{P_\theta(a\varphi+(1-a)\psi)}^n \geq \frac{(a'-a)^n}{a'^n} \int_X \theta_\varphi^n > 0.$$

**Corollary 2.3.2** *Let  $\varphi, \psi \in \text{PSH}(X, \theta)_{>0}$ ,  $\varphi \leq \psi$ . Assume that  $\int_X \theta_\varphi^n = \int_X \theta_\psi^n$ . Then for any  $\epsilon \in (0, 1)$ , there is  $\eta \in \text{PSH}(X, \theta)$  such that*

- (1)  $\int_X \theta_\eta^n = \int_X \theta_\varphi^n$ ;
- (2)

$$\epsilon\eta + (1 - \epsilon^{-1})\psi \leq \varphi.$$

**Proof** Thanks to (2.15) and Theorem 2.3.2, for each  $a' > \epsilon^{-1}$ , we have

$$\int_X \theta_\eta^n > \left( \frac{(a' - \epsilon^{-1})}{a'} \right)^n \int_X \theta_\varphi^n,$$

where

$$\eta = P_\theta(\epsilon^{-1}\varphi + (1 - \epsilon^{-1})\psi).$$

Letting  $a' \rightarrow \infty$ , we conclude that

$$\int_X \theta_\eta^n \geq \int_X \theta_\varphi^n.$$

On the other hand, since  $\eta \leq \psi$ , we find that

$$\int_X \theta_\eta^n \leq \int_X \theta_\psi^n = \int_X \theta_\varphi^n.$$

Hence,

$$\int_X \theta_\eta^n = \int_X \theta_\varphi^n.$$

**Lemma 2.3.2** *For any  $\varphi \in \text{PSH}(X, \theta)_{>0}$ , there is  $\psi \in \text{PSH}(X, \theta)$  such that*

- (1)  $\theta_\psi$  is a Kähler current, and
- (2)  $\psi \leq \varphi$ .

*In particular, there is an increasing sequence  $(\varphi_i)_i$  in  $\text{PSH}(X, \theta)$  converging almost everywhere to  $\varphi$  such that  $\theta_{\varphi_i}$  is a Kähler current for all  $i \geq 1$ .*

**Proof** Using Lemma 2.3.1, we can find  $\epsilon > 0$  and  $\gamma \in \text{PSH}(X, \theta)$  such that

$$\frac{\epsilon}{1+\epsilon}V_\theta + \frac{1}{1+\epsilon}\gamma \leq \varphi.$$

We observe that the cohomology class  $[\theta]$  is big as a consequence of [BEGZ10, Proposition 1.22]. Therefore, we can take  $\eta \in \text{PSH}(X, \theta)$  such that  $\theta_\eta$  is a Kähler current and  $\eta \leq 0$ . Then we may take

$$\psi = \frac{\epsilon}{1+\epsilon}\eta + \frac{1}{1+\epsilon}\gamma.$$

cor:pathoenvlopeeqmass

lma:kahcurrentposmass

BEGZ10

[BEGZ10,

For the latter claim, it suffices to take

$$\varphi_i = (1 - (i + 1)^{-1})\varphi + (i + 1)^{-1}\psi.$$

lma:existsecposmass

**Lemma 2.3.3** *Let  $L$  be a holomorphic line bundle on  $X$  with  $\theta \in c_1(L)$ . Assume that  $\varphi \in \text{PSH}(X, \theta)_{>0}$ , then there exists  $k_0 > 0$  such that for each  $k \geq k_0$ , we have*

$$H^0(X, L^k \otimes \mathcal{I}(k\varphi)) \neq 0.$$

**Proof** By [Lemma 2.3.2](#), we may further assume that  $\theta_\varphi$  is a Kähler current. In this case, the result follows from [\[Dem12a, Theorem 13.21\]](#).  $\square$

thm:logconc

**Theorem 2.3.3** *Let  $\varphi_0, \varphi_1 \in \text{PSH}(X, \theta)$ . Then the map*

$$[0, 1] \ni t \mapsto \log \int_X \theta_{t\varphi_1 + (1-t)\varphi_0}^n$$

*is concave.*

See [\[DDNL19log\]](#) for the proof.

**Remark 2.3.3** Here and in the sequel, when we write expressions like  $t\varphi + (1-t)\psi$  for  $\varphi, \psi \in \text{QPSH}(X)$ , we will follow the convention that when  $t = 0$ , the value is  $\psi$  and when  $t = 1$ , the value is  $\varphi$ .



## Chapter 3

### The envelope operators

chap:enve

In this chapter, we study two envelope operators lying at the heart of the whole theory. The first envelope, called the  $P$ -envelope, is defined using the non-pluripolar masses, while the second, called the  $\mathcal{I}$ -envelope, is defined using the multiplier ideal sheaves. The corresponding theories are developed in [Section 3.1](#) and [Section 3.2](#) respectively.

Later on in [Chapter 6](#), we will develop corresponding  $P$  and  $\mathcal{I}$ -partial orders associated with these envelopes, allowing us to compare the singularities.

#### 3.1 The $P$ -envelope

sec:Penv

In this section,  $X$  will denote a connected compact Kähler manifold of dimension  $n$ .

##### 3.1.1 Rooftop operator and the definition of the $P$ -envelope

We will fix a smooth closed real  $(1, 1)$ -form  $\theta$  on  $X$ .

def:rooftop

**Definition 3.1.1** Given  $\varphi, \psi \in \text{PSH}(X, \theta)$ , we define their *rooftop operator* as follows:

$$\varphi \wedge \psi = \sup \{ \eta \in \text{PSH}(X, \theta) : \eta \leq \varphi, \eta \leq \psi \}.$$

When we want to be more specific, we could also write  $\varphi \wedge_{\theta} \psi$ . Suppose that  $\varphi \wedge \psi$  is not identically  $-\infty$ , then we have  $\varphi \wedge \psi \in \text{PSH}(X, \theta)$  by [Proposition 1.2.1](#).

lma:rooftopMA

**Lemma 3.1.1** Let  $\varphi, \psi \in \text{PSH}(X, \theta)$ . Assume that  $\varphi \wedge \psi \in \text{PSH}(X, \theta)$ . Then

$$\theta_{\varphi \wedge \psi}^n \leq \mathbb{1}_{\{\varphi \wedge \psi = \varphi\}} \theta_{\varphi}^n + \mathbb{1}_{\{\varphi \wedge \psi = \psi\}} \theta_{\psi}^n.$$

See [[DDNL18mono](#), Lemma 3.7] for the proof.

We recall that the relations  $\leq$  and  $\sim$  are introduced in [Definition 1.5.2](#).

`def:Penv`

**Definition 3.1.2** Given  $\varphi \in \text{PSH}(X, \theta)$ , we define its *P-envelope* as follows:

$$P_\theta[\varphi] := \sup^* \{ \psi \in \text{PSH}(X, \theta) : \psi \leq 0, \psi \leq \varphi \}. \quad (3.1)$$

`{eq:Pthetavarphi}`

Observe that by [Proposition 1.2.1](#), we have  $P_\theta[\varphi] \in \text{PSH}(X, \theta)$ . Moreover, the definition can be equivalently described as

$$P_\theta[\varphi] = \sup_{C \in \mathbb{Z}_{>0}}^* (\varphi + C) \wedge V_\theta. \quad (3.2)$$

`{eq:Penvsups}`

Recall that  $V_\theta$  is introduced in [\(2.9\)](#). Observe that for any  $C \in \mathbb{R}$ , we have  $(\varphi + C) \wedge V_\theta \in \text{PSH}(X, \theta)$  and

$$(\varphi + C) \wedge V_\theta \sim \varphi.$$

`prop:Penvindeptheta`

**Proposition 3.1.1** Let  $\theta' = \theta + \text{dd}^c g$  for some  $g \in C^\infty(X)$ . Then for any  $\varphi \in \text{PSH}(X, \theta)$ , we have  $\varphi - g \in \text{PSH}(X, \theta')$  and

$$P_\theta[\varphi] \sim P_{\theta'}[\varphi'].$$

**Proof** By symmetry, it suffices to show that

$$P_\theta[\varphi] \leq P_{\theta'}[\varphi'].$$

We may assume that  $g \geq 0$ . Then for any  $\psi \in \text{PSH}(X, \theta)$  with  $\psi \leq \varphi$  and  $\psi \leq 0$ , we set  $\psi' := \psi - g \in \text{PSH}(X, \theta')$ . Then  $\psi' \leq \varphi'$  and  $\psi' \leq 0$ , so  $\psi' \leq P_{\theta'}[\varphi']$ . Since  $\psi$  is arbitrary, it follows that

$$P_\theta[\varphi] - g \leq P_{\theta'}[\varphi'].$$

The *P-envelope* preserves the non-pluripolar masses:

`prop:Ppresmass`

**Proposition 3.1.2** Suppose that  $\theta_1, \dots, \theta_n$  be smooth closed real  $(1, 1)$ -forms on  $X$ . Let  $\varphi_i \in \text{PSH}(X, \theta_i)$  for each  $i = 1, \dots, n$ . Then

$$\int_X \theta_{1, P_{\theta_1}[\varphi_1]} \wedge \dots \wedge \theta_{n, P_{\theta_n}[\varphi_n]} = \int_X \theta_{1, \varphi_1} \wedge \dots \wedge \theta_{n, \varphi_n}. \quad (3.3)$$

`{eq:Penvpremass}`

**Proof** For each  $C \in \mathbb{Z}_{>0}$  and each  $i = 1, \dots, n$ , we have

$$(\varphi_i + C) \wedge V_{\theta_i} \sim \varphi_i.$$

It follows from [Theorem 2.3.2](#) that

$$\int_X \theta_{1, (\varphi_1 + C) \wedge V_{\theta_1}} \wedge \dots \wedge \theta_{n, (\varphi_n + C) \wedge V_{\theta_n}} = \int_X \theta_{1, \varphi_1} \wedge \dots \wedge \theta_{n, \varphi_n}.$$

So [\(3.3\)](#) follows from [\(3.2\)](#) and [Corollary 2.3.1](#).  $\square$

Conversely, [Proposition 3.1.2](#) characterizes the  $P$ -envelope:

thm:Pvarphidiffdef

**Theorem 3.1.1** Assume that  $\varphi \in \text{PSH}(X, \theta)_{>0}$ , then

$$P_\theta[\varphi] = \sup \left\{ \psi \in \text{PSH}(X, \theta) : \psi \leq 0, \varphi \leq \psi, \int_X \theta_\varphi^n = \int_X \theta_\psi^n \right\}. \quad (3.4)$$

{eq:Penvdef}

In particular, in this case,

$$P_\theta[P_\theta[\varphi]] = P_\theta[\varphi]. \quad (3.5)$$

{eq:Penvprojop}

We refer to [\[DDNL23, Theorem 3.14\]](#) for the proof. In general, we do not know if [\(3.5\)](#) holds when  $\int_X \theta_\varphi^n > 0$ . We expect it to be wrong. According to our general philosophy, the  $P$ -envelope operator is the correct object only when the non-pluripolar mass is positive. We will avoid using the degenerate case in the whole book.

def:modelpot

**Definition 3.1.3** If  $\varphi = P_\theta[\varphi]$  and  $\int_X \theta_\varphi^n > 0$ , we say  $\varphi$  is a *model potential*.

We remind the readers that the notion of model potentials depends heavily on the choice of  $\theta$ . When there is a risk of confusion, we also say  $\varphi$  is a model potential in  $\text{PSH}(X, \theta)$ .

*Remark 3.1.1* [Definition 3.1.3](#) is different from the common definition in the literature: We impose the extra condition  $\int_X \theta_\varphi^n > 0$ . The author believes that this is the only case where this notion is natural. We sometimes emphasize this point by saying  $\varphi \in \text{PSH}(X, \theta)_{>0}$  is a model potential.

There are plenty of model potentials:

cor:Psendspotentialtomodel

**Corollary 3.1.1** Let  $\varphi \in \text{PSH}(X, \theta)_{>0}$ , then  $P_\theta[\varphi]$  is a model potential in  $\text{PSH}(X, \theta)$ . Moreover,

$$\int_X \theta_{P_\theta[\varphi]}^n = \int_X \theta_\varphi^n.$$

**Proof** This follows immediately from [Theorem 3.1.1](#) and [Proposition 3.1.2](#).  $\square$

prop:landfinitecond1

**Proposition 3.1.3** Assume that  $\varphi, \psi \in \text{PSH}(X, \theta)$  and

$$\int_X \theta_\varphi^n + \int_X \theta_\psi^n > \int_X \theta_{\varphi \vee \psi}^n.$$

Then  $\varphi \wedge \psi \in \text{PSH}(X, \theta)$ .

**Proof** Without loss of generality, we may assume that  $\varphi, \psi \leq 0$ . Take

$$\eta := P_\theta[(1 - \epsilon)\varphi \vee \psi + \epsilon V_\theta]$$

for some small enough  $\epsilon > 0$ , we may guarantee that

$$\int_X \theta_\varphi^n + \int_X \theta_\psi^n > \int_X \theta_\eta^n, \quad \varphi \vee \psi \leq \eta.$$

This is a consequence of [Corollary 3.1.1](#).

Take  $C > 0$  large enough, so that

$$\int_{\{\varphi > \eta - C\}} \theta_{\varphi}^n + \int_{\{\psi > \eta - C\}} \theta_{\psi}^n > \int_X \theta_{\eta}^n.$$

This is possible thanks to [Proposition 2.2.1](#)(4). Fix  $C' > C$ . Using [Proposition 2.2.1](#)(1), we can rewrite this equation as

$$\int_{\{\varphi > \eta - C\}} \theta_{\varphi \vee (\eta - C')}^n + \int_{\{\psi > \eta - C\}} \theta_{\psi \vee (\eta - C')}^n > \int_X \theta_{\eta}^n.$$

Write

$$\gamma_{C'} := (\varphi \vee (\eta - C')) \wedge (\psi \vee (\eta - C')).$$

Then observe that

$$\inf_{C' > C} \gamma_{C'} = \varphi \wedge \psi.$$

Assume by contradiction that  $\varphi \wedge \psi \equiv -\infty$ , then we have

$$\lim_{C' \rightarrow \infty} \sup_X \gamma_{C'} = -\infty.$$

Observe that for each  $C' > C$ ,

$$\sup_X \gamma_{C'} = \sup_{\{\eta \neq -\infty\}} (\gamma_{C'} - \eta)$$

since  $\eta$  is a model potential. It follows that

$$\lim_{C' \rightarrow \infty} \sup_{\{\eta \neq -\infty\}} (\gamma_{C'} - \eta) = -\infty. \quad (3.6)$$

{eq:limsupgammametatemp1}

For each  $C' > C$ , we compute

$$\begin{aligned} \int_{\{\gamma_{C'} \leq \eta - C\}} \theta_{\gamma_{C'}}^n &\leq \int_{\{\varphi \vee (\eta - C') \leq \eta - C\}} \theta_{\varphi \vee (\eta - C')}^n + \int_{\{\psi \vee (\eta - C') \leq \eta - C\}} \theta_{\psi \vee (\eta - C')}^n \\ &= 2 \int_X \theta_{\eta}^n - \int_{\{\varphi > \eta - C\}} \theta_{\varphi}^n - \int_{\{\psi > \eta - C\}} \theta_{\psi}^n \\ &< \int_X \theta_{\eta}^n, \end{aligned}$$

where the first line follows from [Lemma 3.1.1](#). Using (3.6), we can take  $C'$  large enough so that  $\gamma_{C'} \leq \eta - C$ . Then we find

$$\int_X \theta_{\gamma_{C'}}^n < \int_X \theta_{\eta}^n,$$

which contradicts [Theorem 2.3.2](#).  $\square$

### 3.1.2 Properties of the $P$ -envelope

Let  $\theta, \theta_1, \theta_2$  be smooth closed real  $(1, 1)$ -forms on  $X$ .

prop:Penvbimero

**Proposition 3.1.4** *Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism from a Kähler manifold  $Y$  to  $X$ . Then for any  $\varphi \in \text{PSH}(X, \theta)$ , we have*

$$P_{\pi^*\theta}[\pi^*\varphi] = \pi^*P_\theta[\varphi].$$

*In particular, a potential  $\varphi \in \text{PSH}(X, \theta)_{>0}$  is model if and only if  $\pi^*\varphi \in \text{PSH}(Y, \pi^*\theta)_{>0}$  is model.*

**Proof** This follows immediately from [Proposition 1.5.3](#).  $\square$

We have the following concavity property of the  $P$ -envelope.

prop:Pconc

**Proposition 3.1.5**

(1) *Suppose that  $\varphi \in \text{PSH}(X, \theta)$  and  $\lambda \in \mathbb{R}_{>0}$ , then*

$$P_{\lambda\theta}[\lambda\varphi] = \lambda P_\theta[\varphi].$$

(2) *Suppose that  $\varphi_1 \in \text{PSH}(X, \theta_1)$  and  $\varphi_2 \in \text{PSH}(X, \theta_2)$ , then*

$$P_{\theta_1+\theta_2}[\varphi_1 + \varphi_2] \geq P_{\theta_1}[\varphi_1] + P_{\theta_2}[\varphi_2].$$

**Proof** (1) This is obvious by definition.

(2) Suppose that  $\psi_1 \in \text{PSH}(X, \theta_1)$  and  $\psi_2 \in \text{PSH}(X, \theta_2)$  satisfy

$$\psi_i \leq 0, \quad \psi_i \leq \varphi_i$$

for  $i = 1, 2$ . Then

$$\psi_1 + \psi_2 \leq 0, \quad \psi_1 + \psi_2 \leq \varphi_1 + \varphi_2.$$

It follows from [\(3.1\)](#) that

$$\psi_1 + \psi_2 \leq P_{\theta_1+\theta_2}[\varphi_1 + \varphi_2].$$

Since  $\psi_1$  and  $\psi_2$  are arbitrary, we conclude.  $\square$

prop:landpresmodel

**Proposition 3.1.6** *Let  $\varphi, \psi \in \text{PSH}(X, \theta)$ . Assume that*

$$\varphi = P_\theta[\varphi], \quad \psi = P_\theta[\psi], \quad \varphi \wedge \psi \not\equiv -\infty.$$

*Then*

$$P_\theta[\varphi \wedge \psi] = \varphi \wedge \psi.$$

(3.7)

{eq:Pthetaphilandpsi}

**Proof** Observe that we obviously have

$$P_\theta[\varphi \wedge \psi] \leq P_\theta[\varphi] = \varphi, \quad P_\theta[\varphi \wedge \psi] \leq P_\theta[\psi] = \psi.$$

So the  $\leq$  direction in [\(3.7\)](#) holds. The reverse direction is trivial.  $\square$

thm:Pvarphisupport

**Theorem 3.1.2** *Let  $\varphi \in \text{PSH}(X, \theta)$ . Then*

$$\theta_{P_\theta[\varphi]}^n \leq \mathbb{1}_{\{P_\theta[\varphi]=0\}} \theta^n.$$

See [\[DDNL18mono\]](#), [\[DDNL18b\]](#), Theorem 3.8] for the proof.

thm:diamond

**Theorem 3.1.3** *Assume that  $\varphi, \psi \in \text{PSH}(X, \theta)$  and  $\varphi \wedge \psi \in \text{PSH}(X, \theta)$ . Then*

$$\int_X \theta_\varphi^n + \int_X \theta_\psi^n \leq \int_X \theta_{\varphi \vee \psi}^n + \int_X \theta_{\varphi \wedge \psi}^n. \quad (3.8)$$

{eq:diamond}

We refer to [\[DDNLmetric\]](#), [\[DDNL21b\]](#), Theorem 5.4] for the proof.

prop:decseqmodel

**Proposition 3.1.7** *Let  $(\varphi_j)_{j \in I}$  be a decreasing net of potentials in  $\text{PSH}(X, \theta)$  satisfying  $P_\theta[\varphi_j] = \varphi_j$  for each  $j \in I$  and  $\varphi := \inf_j \varphi_j \not\equiv -\infty$ . Then  $P_\theta[\varphi] = \varphi$ .*

**Proof** It follows from [Proposition 1.2.1](#) that  $\varphi \in \text{PSH}(X, \theta)$ . Therefore, for each  $j \in I$ ,

$$\varphi \leq P_\theta[\varphi] \leq P_\theta[\varphi_j] = \varphi_j.$$

Therefore,  $\varphi = P_\theta[\varphi]$ . □

prop:vol\_limit\_model

**Proposition 3.1.8** *Let  $(\epsilon_j)_{j \in I}$  be a decreasing net in  $\mathbb{R}_{\geq 0}$  with limit 0. Take a Kähler form  $\omega$  on  $X$ . Consider a decreasing net  $\varphi_j \in \text{PSH}(X, \theta + \epsilon_j \omega)$  ( $j \in I$ ) satisfying*

$$P_{\theta + \epsilon_j \omega}[\varphi_j] = \varphi_j \quad (3.9)$$

{eq:Palmostmodeltemp}

*with pointwise limit  $\varphi \not\equiv -\infty$ . Then*

$$\lim_{j \in I} \int_X (\theta + \epsilon_j \omega)_{\varphi_j}^n = \int_X \theta_\varphi^n. \quad (3.10)$$

{eq:massmodeldec}

*Moreover, if  $\int_X \theta_\varphi^n > 0$ , then for any prime divisor  $E$  over  $X$ , we have*

$$\lim_{j \in I} \nu(\varphi_j, E) = \nu(\varphi, E). \quad (3.11)$$

{eq:Lelongcontdecseq}

**Proof** Observe that  $\varphi \in \text{PSH}(X, \theta)$ . By [Theorem 2.3.2](#), we have

$$\liminf_{j \in I} \int_X (\theta + \epsilon_j \omega)_{\varphi_j}^n \geq \liminf_{j \in I} \int_X (\theta + \epsilon_j \omega)_\varphi^n = \int_X \theta_\varphi^n.$$

We now argue the reverse inequality.

Fix  $j_0 \in I$ , we have

$$\begin{aligned}
\overline{\lim}_{j \in I} \int_X (\theta + \epsilon_j \omega)_{\varphi_j}^n &= \overline{\lim}_{j \in I} \int_{\{\varphi_j=0\}} (\theta + \epsilon_j \omega)_{\varphi_j}^n \\
&\leq \overline{\lim}_{j \in I} \int_{\{\varphi_j=0\}} (\theta + \epsilon_{j_0} \omega)_{\varphi_j}^n \\
&\leq \int_{\{\varphi=0\}} (\theta + \epsilon_{j_0} \omega)_{\varphi}^n,
\end{aligned}$$

where in the first line we used (3.9) and Theorem 3.1.2, and in the last line we have used the fact that  $\varphi_j \searrow \varphi$  and [DDNL21b, Proposition 4.6] (see also [DDNL23, Lemma 2.11]). Taking limit with respect to  $j_0$ , we arrive at the desired conclusion:

$$\overline{\lim}_{j \in I} \int_X (\theta + \epsilon_j \omega)_{\varphi_j}^n \leq \overline{\lim}_{j_0 \in I} \int_{\{\varphi=0\}} (\theta + \epsilon_{j_0} \omega)_{\varphi}^n = \int_{\{\varphi=0\}} \theta_{\varphi}^n \leq \int_X \theta_{\varphi}^n.$$

This finishes the proof of (3.10).

It remains to argue (3.11). By Lemma 2.3.1 and (3.10), for any  $\epsilon \in (0, 1)$  and  $j$  big enough there exists  $\psi_j \in \text{PSH}(X, \theta + \epsilon_j \omega)$  such that  $(1 - \epsilon)\varphi_j + \epsilon\psi_j \leq \varphi$ . This implies that for  $j$  big enough we have

$$(1 - \epsilon)v(\varphi_j, E) + \epsilon v(\psi_j, E) \geq v(\varphi, E) \geq v(\varphi_j, E).$$

On the other hand, the Lelong numbers  $v(\psi_j, E)$  admit an upper bound for various  $j$  by Proposition 1.5.2. So taking limit with respect to  $j$ , we conclude (3.11).  $\square$

cor:Pprojdec

**Corollary 3.1.2** *Let  $(\varphi_j)_{j \in I}$  be a decreasing net of potentials in  $\text{PSH}(X, \theta)$  with pointwise limit  $\varphi \in \text{PSH}(X, \theta)_{>0}$ . Then*

$$P_{\theta}[\varphi] = \inf_{j \in I} P_{\theta}[\varphi_j].$$

**Proof** We may assume that  $I$  is infinite since otherwise, there is nothing to prove.

Let  $\eta = \inf_{i \in I} P_{\theta}[\varphi_i]$ . We clearly have  $0 \geq \eta \geq P_{\theta}[\varphi]$ .

By Proposition 3.1.8, we have

$$\lim_{i \in I} \int_X \theta_{\varphi_i}^n = \int_X \theta_{\varphi}^n > 0.$$

So by Lemma 2.3.1, we can find a decreasing net  $\epsilon_i \searrow 0$  ( $i \in I$ ) with  $\epsilon_i \in (0, 1)$  and  $\psi_i \in \text{PSH}(X, \theta)$  such that for all  $i \in I$ ,

$$(1 - \epsilon_i)\varphi_i + \epsilon_i\psi_i \leq \varphi, \quad \psi_i \leq \varphi_i.$$

By Proposition 3.1.5, we have

$$\eta + \epsilon_i P_{\theta}[\psi_i] \leq (1 - \epsilon_i)\eta + \epsilon_i P_{\theta}[\psi_i] \leq (1 - \epsilon_i)P_{\theta}[\varphi_i] + \epsilon_i P_{\theta}[\psi_i] \leq P_{\theta}[\varphi].$$

Observe that the  $L^1$ -norms of  $P_\theta[\psi_i]$  (with respect to a fixed volume form) are uniformly bounded by [Proposition 1.5.1](#). Taking limit with respect to  $i \in I$ , we conclude that  $\eta \leq P_\theta[\varphi]$  almost everywhere by [Proposition 1.2.5](#).  $\square$

rmk:limitargument1

*Remark 3.1.2* The arguments like the last sentence in the proof of [Corollary 3.1.2](#) is very common. We will usually omit the details.

**Corollary 3.1.3** *Let  $\varphi \in \text{PSH}(X, \theta)_{>0}$  be a model potential. Let  $\omega$  be a Kähler form on  $X$ . Then*

$$\varphi = \inf_{\epsilon > 0} P_{\theta + \epsilon \omega}[\varphi].$$

**Proof** Clearly, we have the  $\leq$  direction and the right-hand side is non-positive. So by [Theorem 3.1.1](#), it suffices to show that they have the same mass, which follows from [Proposition 3.1.8](#).  $\square$

**Proposition 3.1.9** *Let  $(\varphi_i)_{i \in I}$  be an increasing net of potentials in  $\text{PSH}(X, \theta)_{>0}$  uniformly bounded from above. Let  $\varphi := \sup_{i \in I}^* \varphi_i$ . Then*

$$\sup_{i \in I}^* P_\theta[\varphi_i] = P_\theta[\varphi].$$

*In particular, if  $\varphi_i$  is model for all  $i \in I$ , then so is  $\varphi$ .*

**Proof** We may assume that  $I$  is infinite since otherwise, there is nothing to prove.

We write

$$\eta := \sup_{i \in I}^* P_\theta[\varphi_i].$$

Then it is clear that  $\eta \leq P_\theta[\varphi]$ .

By [Corollary 2.3.1](#), we have

$$\lim_{i \in I} \int_X \theta_{\varphi_i}^n = \int_X \theta_\varphi^n > 0.$$

So by [Lemma 2.3.1](#), we can find a decreasing net  $\epsilon_i \searrow 0$  ( $i \in I$ ) with  $\epsilon \in (0, 1)$  and  $\psi_i \in \text{PSH}(X, \theta)$  ( $i \in I$ ) such that for all  $i \in I$ ,

$$(1 - \epsilon_i)\varphi + \epsilon_i\psi_i \leq \varphi_i.$$

By [Proposition 3.1.5](#), we have

$$P_\theta[\varphi] + \epsilon_i P_\theta[\psi_i] \leq (1 - \epsilon_i)P_\theta[\varphi] + \epsilon_i P_\theta[\varphi_i] \leq \eta.$$

Taking limit with respect to  $i$ , we conclude that  $P_\theta[\varphi] \leq \eta$  (c.f. [Remark 3.1.2](#)).  $\square$

### 3.1.3 Relative full mass classes

subsec:fullmass

Let  $\theta$  be a smooth closed real  $(1, 1)$ -form on  $X$  representing a big cohomology class. Fix a model potential  $\phi \in \text{PSH}(X, \theta)_{>0}$ .



**Definition 3.1.4** We define

$$\begin{aligned}\mathrm{PSH}(X, \theta; \phi) &:= \{\eta \in \mathrm{PSH}(X, \theta) : \eta \leq \phi\}, \\ \mathcal{E}^\infty(X, \theta; \phi) &:= \{\eta \in \mathrm{PSH}(X, \theta) : \eta \sim \phi\}, \\ \mathcal{E}(X, \theta; \phi) &:= \left\{ \eta \in \mathrm{PSH}(X, \theta; \phi) : \int_X \theta_\varphi^n = \int_X \theta_\phi^n \right\}, \\ \mathcal{E}^1(X, \theta; \phi) &:= \left\{ \eta \in \mathcal{E}(X, \theta; \phi) : \int_X |\phi - \eta| \theta_\eta^n < \infty \right\}.\end{aligned}$$

Potentials in the last three classes are said to have *minimal singularities*, *full mass* and *finite energy* relative to  $\phi$  respectively.

We have the following inclusions:

$$\mathcal{E}^\infty(X, \theta; \phi) \subseteq \mathcal{E}^1(X, \theta; \phi) \subseteq \mathcal{E}(X, \theta; \phi) \subseteq \mathrm{PSH}(X, \theta; \phi). \quad (3.12)$$

{eq:energyclassinc}

The only non-trivial part is the first inclusion, which follows from [Theorem 2.3.2](#).

rmk:intwelldef

*Remark 3.1.3* Note that this integral

$$\int_X |\phi - \eta| \theta_\eta^n$$

is defined: The locus where  $\phi - \eta$  is undefined is a pluripolar set, while the product  $\theta_\eta^n$  puts no mass on pluripolar sets ([Proposition 2.2.1](#)).

Similar remarks apply when we talk about similar integrals in the sequel.

When  $\phi = V_\theta$ , we usually write

$$\begin{aligned}\mathcal{E}^\infty(X, \theta; V_\theta) &= \mathcal{E}^\infty(X, \theta), \\ \mathcal{E}(X, \theta; V_\theta) &= \mathcal{E}(X, \theta), \\ \mathcal{E}^1(X, \theta; V_\theta) &= \mathcal{E}^1(X, \theta).\end{aligned}$$

Potentials in the three classes are said to have *minimal singularities*, *full mass* and *finite energy* respectively. The relation (3.12) can be written as

$$\mathcal{E}^\infty(X, \theta; V_\theta) \subseteq \mathcal{E}^1(X, \theta; V_\theta) \subseteq \mathcal{E}(X, \theta; V_\theta)$$

in this case.

The  $P$ -envelope can be used to characterize the full mass classes:

prop:fullmassP

**Proposition 3.1.10** *Let  $\varphi \in \mathrm{PSH}(X, \theta)$ . Then the following are equivalent:*

- (1)  $\varphi \in \mathcal{E}(X, \theta; \phi)$ ;
- (2)  $P_\theta[\varphi] = \phi$ .

**Proof** (2)  $\implies$  (1). This follows from [Proposition 3.1.2](#).

(1)  $\implies$  (2). Note that  $\phi$  is a candidate of  $P_\theta[\varphi]$  as in (3.4). So  $P_\theta[\varphi] = \phi$ .  $\square$

In order to handle the finite energy classes, it is convenient to introduce the following quantity:

`def:MAenergy`

**Definition 3.1.5** We define the *Monge–Ampère energy*  $E_\theta^\phi : \mathcal{E}^\infty(X, \theta; \phi) \rightarrow \mathbb{R}$  as follows

$$E_\theta^\phi(\varphi) := \frac{1}{n+1} \sum_{j=0}^n \int_X (\varphi - \phi) \theta_\varphi^j \wedge \theta_\phi^{n-j}. \quad (3.13)$$

`{eq:Edefbdd}`

More generally, we extend  $E_\theta^\phi$  to a functional  $E_\theta^\phi : \text{PSH}(X, \theta; \phi) \rightarrow [-\infty, \infty)$  as follows

$$E_\theta^\phi(\varphi) := \inf \left\{ E_\theta^\phi(\psi) : \psi \in \mathcal{E}^\infty(X, \theta; \phi), \varphi \leq \psi \right\}. \quad (3.14)$$

`{eq:Eextendgeneral}`

We write  $E_\theta$  instead of  $E_\theta^\phi$  when  $\phi = V_\theta$ .

`prop:cocycE1`

**Proposition 3.1.11** *Let  $\varphi \in \text{PSH}(X, \theta; \phi)$ . The following are equivalent:*

- (1)  $\varphi \in \mathcal{E}^1(X, \theta; \phi)$ ;
- (2)  $E_\theta^\phi(\varphi) > -\infty$ .

When the conditions are satisfied, (3.13) holds.

Given  $\varphi, \psi \in \mathcal{E}^1(X, \theta; \phi)$ , we have the following cocycle equality

$$E_\theta^\phi(\psi) - E_\theta^\phi(\varphi) = \frac{1}{n+1} \sum_{j=0}^n \int_X (\psi - \varphi) \theta_\psi^j \wedge \theta_\varphi^{n-j}. \quad (3.15)$$

`{eq:Ecocyc}`

See [BEGZ10, Proposition 2.11] and [DDNL18big, Proposition 2.5] for the proofs.<sup>1</sup>

`prop:relrooftopclosed`

**Proposition 3.1.12** *Assume that  $\varphi, \psi \in \mathcal{E}(X, \theta; \phi)$  (resp.  $\mathcal{E}^1(X, \theta; \phi)$ ,  $\mathcal{E}^\infty(X, \theta; \phi)$ ), then so is  $\varphi \wedge \psi$ .*

**Proof** The case of  $\mathcal{E}^\infty(X, \theta; \phi)$  is trivial.

We consider the case  $\mathcal{E}(X, \theta; \phi)$ . It follows from Proposition 3.1.3 that  $\varphi \wedge \psi \in \text{PSH}(X, \theta)$ . By Theorem 3.1.3, we have

$$\int_X \theta_{\varphi \wedge \psi}^n \geq \int_X \theta_\phi^n.$$

By Theorem 2.3.2, equality holds. By Theorem 3.1.1, we conclude that

$$P_\theta[\varphi \wedge \psi] = \phi.$$

Finally, the case  $\mathcal{E}^1(X, \theta; \phi)$  is proved in [Xia23Mabuchi, Theorem 4.13] (the arXiv version).  $\square$

`prop:relativeEupperclosed`

**Proposition 3.1.13** *Let  $\varphi, \psi \in \text{PSH}(X, \theta)$  be potentials such that  $\psi \leq \phi$  and  $\varphi \leq \psi$ . Assume that  $\varphi \in \mathcal{E}(X, \theta; \phi)$  (resp.  $\mathcal{E}^1(X, \theta; \phi)$ ,  $\mathcal{E}^\infty(X, \theta; \phi)$ ), then so is  $\psi$ .*

<sup>1</sup> In these references, they took  $\phi = V_\theta$ , but the proof of the general case is almost identical.

**Proof** The case  $\mathcal{E}^\infty(X, \theta; \phi)$  is trivial. The case  $\mathcal{E}(X, \theta; \phi)$  follows from [Theorem 2.3.2](#). The case  $\mathcal{E}^1(X, \theta; \phi)$  follows from [\[Xia23a, Proposition 4.5\]](#) (arXiv version).  $\square$

prop:supsEE1

**Proposition 3.1.14** *Let  $(\varphi_i)_{i \in I}$  be a uniformly bounded from above non-empty family in  $\mathcal{E}(X, \theta; \phi)$  (resp.  $\mathcal{E}^1(X, \theta; \phi)$ ,  $\mathcal{E}^\infty(X, \theta; \phi)$ ), then so is  $\sup^*_i \varphi_i$ .*

**Proof** It suffices to handle the case where  $\varphi_i \in \mathcal{E}(X, \theta; \phi)$  for all  $i \in I$ . The remaining two cases follow from [Proposition 3.1.13](#).

**Step 1.** We first assume that  $I$  is finite. In this case, we can easily further reduce to the case where  $I = \{0, 1\}$ . Assume that  $\varphi_0, \varphi_1 \in \mathcal{E}(X, \theta; \phi)$ . Observe that  $\varphi_0 \leq \phi$  and  $\varphi_1 \leq \phi$ , hence  $\varphi_0 \vee \varphi_1 \leq \phi$ . On the other hand, by [Theorem 2.3.2](#),  $\varphi_0 \vee \varphi_1$  and  $\phi$  have the same mass.

**Step 2.** We come back to the case where  $I$  is infinite.

By [Proposition 1.2.2](#), we may assume that  $I = \mathbb{Z}_{>0}$  as an ordered set. Moreover, by Step 1, we may assume that the sequence  $(\varphi_i)_i$  is increasing. Furthermore, we may assume that  $\varphi_i \leq 0$  for all  $i$ . Then we know that  $\varphi_i \leq \phi$ . Therefore,  $\sup^*_i \varphi_i \leq \phi$ . But they have the same mass as a consequence of [Corollary 2.3.1](#). So we conclude using [Theorem 3.1.1](#).  $\square$

prop:envrelfullmass

**Proposition 3.1.15** *Let  $\varphi, \psi \in \mathcal{E}(X, \theta; \phi)$ . Then*

$$\sup_{C \geq 0}^*(\varphi + C) \wedge \psi = \psi.$$

**Proof** Since for each  $C \geq 0$ ,

$$(\varphi \wedge \psi + C) \wedge \psi \leq (\varphi + C) \wedge \psi \leq \psi,$$

we may replace  $\varphi$  by  $\varphi \wedge \psi$  (c.f. [Proposition 3.1.12](#)) and assume that  $\varphi \leq \psi$ . In this case, the result is proved in [\[DDNL18b, Theorem 3.8, Corollary 3.11\]](#).  $\square$

## 3.2 The $\mathcal{I}$ -envelope

sec:Ienv

From the algebraic point of view, a more natural envelope operator is given by the  $\mathcal{I}$ -envelope.

In this section,  $X$  will denote a connected compact Kähler manifold of dimension  $n$ .

### 3.2.1 $\mathcal{I}$ -equivalence

prop:Iequivchar

**Proposition 3.2.1** *Given  $\varphi, \psi \in \text{QPSH}(X)$ , the following are equivalent:*

(1) *For any  $k \in \mathbb{Z}_{>0}$ , we have*

$$\mathcal{I}(k\varphi) = \mathcal{I}(k\psi);$$

(2) for any  $\lambda \in \mathbb{R}_{>0}$ , we have

$$I(\lambda\varphi) = I(\lambda\psi);$$

(3) for any modification  $\pi: Y \rightarrow X$  and any  $y \in Y$ , we have

$$v(\pi^*\varphi, y) = v(\pi^*\psi, y);$$

(4) for any proper bimeromorphic morphism  $\pi: Y \rightarrow X$  from a Kähler manifold and any  $y \in Y$ , we have

$$v(\pi^*\varphi, y) = v(\pi^*\psi, y);$$

(5) for any prime divisor  $E$  over  $X$ , we have

$$v(\varphi, E) = v(\psi, E).$$

See [Definition B.1.1](#) for the definition of prime divisors over  $X$ . We remind the readers that in the whole book, a *modification* of a compact complex space means a finite composition of blow-ups with smooth centers. This terminology is highly non-standard.

**Proof** (4)  $\iff$  (5). This follows from [Lemma 1.4.1](#).

(3)  $\iff$  (5). This follows from [Corollary B.1.1](#).

(1)  $\implies$  (5). This follows from [Proposition 1.4.4](#).

(5)  $\implies$  (2). This follows from [Theorem 1.4.3](#).

(2)  $\implies$  (1). This is trivial.  $\square$

def:Iequiv

**Definition 3.2.1** Given  $\varphi, \psi \in \text{QPSH}(X)$ , we say they are  $I$ -equivalent and write  $\varphi \sim_I \psi$  if the equivalent conditions in [Proposition 3.2.1](#) are satisfied.

prop:Ienvbimero

**Proposition 3.2.2** Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism from a Kähler manifold  $Y$  to  $X$ . Then for  $\varphi, \psi \in \text{QPSH}(X)$ , we the following are equivalent:

(1)  $\varphi \sim_I \psi$ ;

(2)  $\pi^*\varphi \sim_I \pi^*\psi$ .

**Proof** (1)  $\implies$  (2). This follows from [Proposition 3.2.1](#)(4).

(2)  $\implies$  (1). This follows from the simple fact that

$$I(k\varphi) = \pi_* (\omega_{Y/X} \otimes I(k\pi^*\varphi)), \quad I(k\psi) = \pi_* (\omega_{Y/X} \otimes I(k\pi^*\psi)).$$

prop:Iequivmax

**Proposition 3.2.3** Let  $\varphi, \varphi', \psi, \psi' \in \text{QPSH}(X)$  and  $\lambda > 0$ . Assume that  $\varphi \sim_I \psi$  and  $\varphi' \sim_I \psi'$ , then

$$\varphi \vee \varphi' \sim_I \psi \vee \psi', \quad \varphi + \varphi' \sim_I \psi + \psi', \quad \lambda\varphi \sim_I \lambda\psi.$$

Similarly, if  $(\varphi_i)_{i \in I}, (\psi_i)_{i \in I}$  are two non-empty uniformly bounded from above families in  $\text{PSH}(X, \theta)$  for some closed smooth real  $(1, 1)$ -form  $\theta$  on  $X$  such that  $\varphi_i \sim_I \psi_i$  for all  $i \in I$ , then

$$\sup_{i \in I}^* \varphi_i \sim_I \sup_{i \in I}^* \psi_i.$$

**Proof** This follows from [Proposition 1.4.2](#) and [Corollary 1.4.1](#).  $\square$

### 3.2.2 The definition of the $I$ -envelope

We will fix a smooth closed real  $(1, 1)$ -form  $\theta$  on  $X$ .

`def:Ienv`

**Definition 3.2.2** Given  $\varphi \in \text{PSH}(X, \theta)$ , we define its  $I$ -envelope as follows:

$$P_\theta[\varphi]_I := \sup^* \{\psi \in \text{PSH}(X, \theta) : \psi \leq 0, \psi \sim_I \varphi\}. \quad (3.16)$$

`{eq:Ienvelopedef}`

If  $\varphi = P_\theta[\varphi]_I$ , we say  $\varphi$  is an  $I$ -model potential (in  $\text{PSH}(X, \theta)$ ).

Note that by [Proposition 1.2.1](#),  $P_\theta[\varphi]_I \in \text{PSH}(X, \theta)$ .

`prop:Ienvindeptheta`

**Proposition 3.2.4** Let  $\theta' = \theta + \text{dd}^c g$  for some  $g \in C^\infty(X)$ . Then for any  $\varphi \in \text{PSH}(X, \theta)$ , we have  $\varphi - g \in \text{PSH}(X, \theta')$  and

$$P_\theta[\varphi]_I \sim P_{\theta'}[\varphi']_I.$$

The proof is similar to that of [Proposition 3.1.1](#), so we omit it.

`prop:Ienvelopebimero`

**Proposition 3.2.5** Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism from a connected Kähler manifold  $Y$  to  $X$ . Then for  $\varphi \in \text{PSH}(X, \theta)$ , we have

$$P_{\pi^*\theta}[\pi^*\varphi]_I = \pi^*P_\theta[\varphi]_I.$$

**Proof** The proof is similar to that of [Proposition 3.1.4](#) in view of [Proposition 3.2.2](#).  $\square$

`prop:Ienvprojection`

**Proposition 3.2.6** Let  $\varphi \in \text{PSH}(X, \theta)$ , then

$$\varphi \sim_I P_\theta[\varphi]_I.$$

In particular,

$$P_\theta[P_\theta[\varphi]_I]_I = P_\theta[\varphi]_I$$

and the upper semicontinuous regularization in (3.16) is not necessary.

**Proof** In view of [Proposition 3.2.1](#), it suffices to show that for  $k \in \mathbb{Z}_{>0}$ , we have

$$I(k\varphi) = I(kP_\theta[\varphi]_I). \quad (3.17)$$

`{eq:IenvelopepreservLelong}`

By [Proposition 1.2.2](#), we can find  $\psi_i \in \text{PSH}(X, \theta)$  ( $i \in \mathbb{Z}_{>0}$ ) such that  $\psi_i \leq 0$ ,  $\psi_i \sim_I \varphi$  for all  $i \geq 1$  and

$$\sup_{i>0}^* \psi_i = P_\theta[\varphi]_I.$$

By [Proposition 3.2.3](#), we may replace  $\psi_i$  by  $\psi_1 \vee \cdots \vee \psi_i$  and assume that the sequence  $\psi_i$  is increasing. In this case, it follows from the strong openness theorem [Theorem 1.4.4](#) that for each  $k \in \mathbb{Z}_{>0}$ , we have

$$I(k\varphi) = I(k\psi_j) = I(kP_\theta[\varphi]_I)$$

for  $j$  large enough.  $\square$

`def:volqpsh`

**Definition 3.2.3** Let  $\varphi \in \text{PSH}(X, \theta)$ , we define the *volume*  $\text{vol}(\theta, \varphi)$  as

$$\text{vol}(\theta, \varphi) = \int_X (\theta + \text{dd}^c P_\theta[\varphi]_I)^n.$$

`prop:voldeponlyoncurr`

**Proposition 3.2.7** Let  $\theta' = \theta + \text{dd}^c g$  for some  $g \in C^\infty(X)$ . Then for any  $\varphi \in \text{PSH}(X, \theta)$ , we have  $\varphi - g \in \text{PSH}(X, \theta')$  and

$$\text{vol}(\theta, \varphi) = \text{vol}(\theta', \varphi').$$

*Proof* This follows immediately from [Proposition 3.2.4](#) and [Theorem 2.3.2](#).  $\square$

In view of [Proposition 3.2.7](#), we could write

$$\text{vol } \theta_\varphi = \text{vol}(\theta, \varphi). \quad (3.18) \quad \text{\code{eq:volcurrdef}}$$

The  $I$ -envelope and the  $P$ -envelope are related in a simple manner.

`prop:PandPI`

**Proposition 3.2.8** Let  $\varphi \in \text{PSH}(X, \theta)$ , then

$$P_\theta[\varphi] \leq P_\theta[\varphi]_I, \quad \varphi \sim_I P_\theta[\varphi].$$

*Proof* It suffices to show that  $\varphi \sim_I P_\theta[\varphi]$ . Namely, for each  $k \in \mathbb{Z}_{>0}$ , we have

$$I(k\varphi) = I(kP_\theta[\varphi]). \quad (3.19) \quad \text{\code{eq:IkvarphiIkP}}$$

It follows from [\(3.2\)](#) and the strong openness theorem [Theorem 1.4.4](#) that

$$I(kP_\theta[\varphi]) = I((k\varphi + C) \wedge kV_\theta)$$

when  $C$  is large enough. Since  $(k\varphi + C) \wedge kV_\theta \sim k\varphi$ , we have

$$I((k\varphi + C) \wedge kV_\theta) = I(k\varphi)$$

and [\(3.19\)](#) follows.  $\square$

`cor:compnppmassandvol`

**Corollary 3.2.1** Let  $\varphi \in \text{PSH}(X, \theta)$ , then

$$\int_X \theta_\varphi^n \leq \text{vol } \theta_\varphi.$$

**Proof** This follows from [Proposition 3.2.8](#), [Theorem 2.3.2](#) and [Proposition 3.1.2](#).  $\square$

We note the following special case:

**Proposition 3.2.9** *Let  $\varphi \in \text{PSH}(X, \theta)$ . Assume that  $\varphi$  has analytic singularities, then*

$$\varphi \sim P_\theta[\varphi] \sim_P P_\theta[\varphi]_I.$$

**Proof** In view of [Proposition 3.2.8](#), it suffices to show that

$$P_\theta[\varphi]_I \leq \varphi. \quad (3.20)$$

By [Proposition 3.2.5](#) and [Theorem 1.6.1](#), we may assume that  $\varphi$  has log singularities along an effective  $\mathbb{Q}$ -divisor  $D$ . By rescaling using [Proposition 3.2.10](#), we may assume that  $D$  is a divisor. Take quasi-equisingular approximations  $(\eta_j)_j$  and  $(\varphi_j)_j$  of  $P_\theta[\varphi]_I$  and of  $\varphi$  respectively. Recall that by [Theorem 1.6.2](#), we can guarantee that  $\eta_j$  and  $\varphi_j$  both have the singularity type  $(2^{-j}, \mathcal{I}(2^j \varphi))$  and hence  $\eta_j \sim \varphi_j$  for all  $j \geq 1$ . On the other hand, it is clear that  $\varphi_j \sim \varphi$  for all  $j \geq 1$ . So [\(3.20\)](#) follows.  $\square$

### 3.2.3 Properties of the $\mathcal{I}$ -envelope

Let  $\theta, \theta_1, \theta_2$  be smooth closed real  $(1, 1)$ -forms on  $X$ .

We have the following concavity property of the  $\mathcal{I}$ -envelope.

**Proposition 3.2.10**

(1) *Suppose that  $\varphi \in \text{PSH}(X, \theta)$  and  $\lambda \in \mathbb{R}_{>0}$ , then*

$$P_{\lambda\theta}[\lambda\varphi]_I = \lambda P_\theta[\varphi]_I.$$

(2) *Suppose that  $\varphi_1 \in \text{PSH}(X, \theta_1)$  and  $\varphi_2 \in \text{PSH}(X, \theta_2)$ , then*

$$P_{\theta_1+\theta_2}[\varphi_1 + \varphi_2]_I \geq P_{\theta_1}[\varphi_1]_I + P_{\theta_2}[\varphi_2]_I.$$

(3) *Suppose that  $\varphi_1 \in \text{PSH}(X, \theta_1)$  and  $\varphi_2 \in \text{PSH}(X, \theta_2)$ , then*

$$P_{\theta_1+\theta_2}[\varphi_1 + \varphi_2]_I \sim_I P_{\theta_1}[\varphi_1]_I + P_{\theta_2}[\varphi_2]_I.$$

(4) *Suppose that  $\varphi_1, \varphi_2 \in \text{PSH}(X, \theta)$ , then*

$$P_\theta[\varphi_1 \vee \varphi_2]_I \sim_I P_\theta[\varphi_1]_I \vee P_\theta[\varphi_2]_I.$$

**Proof** (1) This is obvious by definition.

(2) Suppose that  $\psi_1 \in \text{PSH}(X, \theta_1)$  and  $\psi_2 \in \text{PSH}(X, \theta_2)$  satisfy

$$\psi_i \leq 0, \quad \psi_i \sim_I \varphi_i$$

for  $i = 1, 2$ . Then thanks to [Proposition 3.2.3](#),

$$\psi_1 + \psi_2 \leq 0, \quad \psi_1 + \psi_2 \sim_I \varphi_1 + \varphi_2.$$

It follows that

$$\psi_1 + \psi_2 \leq P_{\theta_1 + \theta_2}[\varphi_1 + \varphi_2]_I.$$

Since  $\psi_1$  and  $\psi_2$  are arbitrary, we conclude.

(3) This follows easily from [Proposition 3.2.6](#) and [Proposition 3.2.3](#).

(4) The proof is similar to that of (3). We omit the details.  $\square$

lma:PIenvmono1

**Lemma 3.2.1** *Let  $\varphi, \psi \in \text{QPSH}(X)$ . Assume that  $\varphi \leq \psi$ , then*

$$P_\theta[\varphi]_I \leq P_\theta[\psi]_I.$$

**Proof** It suffices to observe that  $P_\theta[\varphi]_I \vee \psi \sim_I \psi$  as a consequence of [Proposition 1.4.2](#) and [Proposition 3.2.6](#).  $\square$

prop:decnetmodelPI

**Proposition 3.2.11** *Consider a decreasing net  $(\varphi_i)_{i \in I}$  of model potentials in  $\text{PSH}(X, \theta)_{>0}$ . Suppose that  $\varphi := \inf_{i \in I} \varphi_i \not\equiv -\infty$  and  $\int_X \theta_\varphi^n > 0$ . Then*

$$\inf_{i \in I} P_\theta[\varphi_i]_I = P_\theta[\varphi]_I.$$

**Proof** Let  $\eta = \inf_{i \in I} P_\theta[\varphi_i]_I$ . We clearly have  $\eta \geq P_\theta[\varphi]_I$  as a consequence of [Lemma 3.2.1](#).

By [Proposition 3.1.8](#), we have

$$\lim_{i \in I} \int_X \theta_{\varphi_i}^n = \int_X \theta_\varphi^n > 0.$$

So by [Lemma 2.3.1](#), we can find a decreasing net  $\epsilon_i \searrow 0$  ( $i \in I$ ) with  $\epsilon_i \in (0, 1)$  and  $\psi_i \in \text{PSH}(X, \theta)$  such that for all  $i \in I$ ,

$$(1 - \epsilon_i)\varphi_i + \epsilon_i\psi_i \leq \varphi.$$

By [Proposition 3.2.10](#), we have

$$(1 - \epsilon_i)\eta + \epsilon_i P_\theta[\psi_i]_I \leq (1 - \epsilon_i)P_\theta[\varphi_i]_I + \epsilon_i P_\theta[\psi_i]_I \leq P_\theta[\varphi]_I.$$

Taking limit with respect to  $i$ , we conclude that  $\eta \leq P_\theta[\varphi]_I$  (c.f. [Remark 3.1.2](#)).  $\square$

prop:incnetmodelPI

**Proposition 3.2.12** *Let  $(\varphi_i)_{i \in I}$  be an increasing net in  $\text{PSH}(X, \theta)_{>0}$  uniformly bounded from above. Let  $\varphi := \sup^*_{i \in I} \varphi_i$ . Then*

$$\sup^*_{i \in I} P_\theta[\varphi_i]_I = P_\theta[\varphi]_I.$$

**Proof** Let  $\eta = \sup^*_{i \in I} P_\theta[\varphi_i]_I$ . Then  $\eta \leq P_\theta[\varphi]_I$  as a consequence of [Lemma 3.2.1](#).

By [Corollary 2.3.1](#), we have



$$\lim_{i \in I} \int_X \theta_{\varphi_i}^n = \int_X \theta_{\varphi}^n > 0.$$

So by [Lemma 2.3.1](#), we can find a decreasing net  $\epsilon_i \searrow 0$  ( $i \in I$ ) with  $\epsilon_i \in (0, 1)$  and  $\psi_i \in \text{PSH}(X, \theta)$  such that for all  $i \in I$ ,

$$(1 - \epsilon_i)\varphi + \epsilon_i\psi_i \leq \varphi_i.$$

By [Proposition 3.2.10](#), we have

$$P_{\theta}[\varphi]_I + \epsilon_i P_{\theta}[\psi_i]_I \leq (1 - \epsilon_i)P_{\theta}[\varphi]_I + \epsilon_i P_{\theta}[\psi_i]_I \leq P_{\theta}[\varphi_i]_I \leq \eta.$$

Taking limit with respect to  $i$ , we conclude that  $\eta \geq P_{\theta}[\varphi]_I$  (c.f. [Remark 3.1.2](#)).  $\square$

*Remark 3.2.1* One could also define the following interpolation between the  $\mathcal{I}$ -envelope and the  $P$ -envelope: Suppose  $\varphi \in \text{PSH}(X, \theta)_{>0}$ ,  $k \in \{0, \dots, n\}$ . Then we let

$$\begin{aligned} P_{\theta,j}[\varphi] &:= \sup^* \left\{ \psi \in \text{PSH}(X, \theta) : \psi \leq 0, \varphi \leq \psi, \int_X \theta_{\varphi}^j \wedge \theta_{P_{\theta}[\varphi]_I}^{n-j} \right. \\ &\quad \left. = \int_X \theta_{\psi}^j \wedge \theta_{P_{\theta}[\psi]_I}^{n-j} \right\}. \end{aligned}$$

Based on the techniques developed in [Chapter 6](#), one could show that  $P_{\theta,j}[\bullet]$  is a projection operator. When  $j = 0$ , this operator reduces to the  $P$ -envelope, while when  $j = n$ , this operator reduces to the  $\mathcal{I}$ -envelope.



## Chapter 4

### Geodesic rays in the space of potentials

chap:rays

In this chapter, we study subgeodesics and geodesics in the space of quasi-plurisubharmonic functions. Unlike what one usually finds in the literature, here we are carrying out the constructions in the space of Kähler potentials with prescribed singularities. The usual regularization techniques break down in this setup.

The results in [Section 4.2](#) seem to be new, although they have been applied without proofs in the literature.

#### 4.1 Subgeodesics

Let  $X$  be a connected compact Kähler manifold of dimension  $n$  and  $\theta$  be a smooth closed real  $(1, 1)$ -form on  $X$  representing a big cohomology class.

def:subgeod

**Definition 4.1.1** Let us fix  $\varphi_0, \varphi_1 \in \text{PSH}(X, \theta)$ . A *subgeodesic* from  $\varphi_0$  to  $\varphi_1$  is a family  $(\varphi_t)_{t \in (0,1)}$  in  $\text{PSH}(X, \theta)$  such that

(1) if we define

$$\Phi: X \times \{z \in \mathbb{C} : e^{-1} < |z| < 1\} \rightarrow [-\infty, \infty), \quad (x, z) \mapsto \varphi_{-\log |z|}(x),$$

then  $\Phi$  is  $p_1^* \theta$ -psh, where  $p_1: X \times \{z \in \mathbb{C} : e^{-1} < |z| < 1\} \rightarrow X$  is the natural projection;

(2) when  $t \rightarrow 0+$  (resp. to  $1-$ ),  $\varphi_t$  converges to  $\varphi_0$  (resp.  $\varphi_1$ ) with respect to the  $L^1$ -topology.

We also say  $(\varphi_t)_{t \in [0,1]}$  is a subgeodesic.

We say  $\Phi$  is the *complexification* of the subgeodesic  $(\varphi_t)_t$ .

When we do not want to specify  $\varphi_0$  and  $\varphi_1$ , we shall say  $(\varphi_t)_{t \in (0,1)}$  is a subgeodesic.

In general, there are no subgeodesics from  $\varphi_0$  to  $\varphi_1$ .

prop:convexsubgeod

**Proposition 4.1.1** Let  $\varphi_0, \varphi_1 \in \text{PSH}(X, \theta)$  and  $(\varphi_t)_{t \in (0,1)}$  be a subgeodesic from  $\varphi_0$  to  $\varphi_1$ . Then for each  $x \in X$ ,  $[0, 1] \ni t \mapsto \varphi_t(x)$  is a convex function.

**Proof** For each  $x \in X$ , the map

$$\{z \in \mathbb{C} : e^{-1} < |z| < 1\} \rightarrow [-\infty, \infty), \quad z \mapsto \Phi(x, z)$$

is either subharmonic or constantly  $-\infty$ , as follows from [Definition 4.1.1](#) (1) and [Proposition 1.1.4](#). In the latter case, the convexity of  $[0, 1] \ni t \mapsto \varphi_t(x)$  is trivial. In the former case, the convexity on the interval  $(0, 1)$  follows from [Proposition 1.1.3](#).

In order to verify the convexity at the boundary, let us fix  $s \in (0, 1)$ . We need to show that

$$\varphi_s(x) \leq s\varphi_1(x) + (1-s)\varphi_0(x) \quad (4.1)$$

{eq:varphisconvtempl}

for all  $x \in X$ . Thanks to [Proposition 1.2.5](#), it suffices to prove this for almost all  $x$ .

Take a set  $Z \subseteq X$  with zero Lebesgue measure such that for all  $x \in X \setminus Z$ , we have

- (1)  $\varphi_t(x) \neq -\infty$  for all  $t \in [0, 1] \cap \mathbb{Q}$ ;
- (2)  $\varphi_t(x) \rightarrow \varphi_0(x)$  as  $t \rightarrow 0+$  and  $\varphi_t(x) \rightarrow \varphi_1(x)$  as  $t \rightarrow 1-$ .

For all such  $x$ , the convexity of  $\varphi$  guarantees that  $\varphi_t(x) \neq -\infty$  for all  $t \in [0, 1]$  and  $t \mapsto \varphi_t(x)$  is convex for  $t \in [0, 1]$ . In particular, (4.1) holds.  $\square$

prop:maxsubgeod

**Proposition 4.1.2** *Let  $(\varphi_0^i)_{i \in I}$ ,  $(\varphi_1^i)_{i \in I}$  be two non-empty uniformly bounded from above families in  $\text{PSH}(X, \theta)$ . Let  $(\varphi_t^i)_{t \in (0,1)}$  be subgeodesics from  $\varphi_0^i$  to  $\varphi_1^i$  for each  $i \in I$ . Then*

$$\left( \sup_{i \in I}^* \varphi_t^i \right)_{t \in (0,1)}$$

*is a subgeodesic from  $\sup_{i \in I}^* \varphi_0^i$  to  $\sup_{i \in I}^* \varphi_1^i$ .*

**Proof** We may assume that  $\varphi_0^i, \varphi_1^i \leq 0$  for all  $i \in I$ . Then it follows that  $\varphi_t^i \leq 0$  for all  $t \in (0, 1)$  and all  $i \in I$  by [Proposition 4.1.1](#).

We define

$$\varphi_t := \sup_{i \in I}^* \varphi_t^i \in \mathcal{E}(X, \theta; \phi)$$

for all  $t \in [0, 1]$ . Observe that  $[0, 1] \ni t \mapsto \varphi_t$  is convex by the same argument leading to (4.1).

Let  $(\psi_t)_{t \in (0,1)}$  be the subgeodesic whose complexification  $\Phi_\psi$  corresponds to  $\sup_{i \in I}^* \Phi_{\varphi_t^i}$ , where  $\Phi_{\varphi_t^i}$  is the complexification of  $(\varphi_t^i)_{t \in (0,1)}$ . Then clearly,  $\varphi_t \leq \psi_t$  for each  $t \in (0, 1)$ . On the other hand, by [Proposition 1.2.3](#),

$$\psi_t = \sup_{i \in I} \varphi_t^i = \varphi_t \quad \text{almost everywhere}$$

for almost all  $t \in (0, 1)$ . Therefore, using [Proposition 1.2.5](#), we find  $\psi_t = \varphi_t$  for almost all  $t \in (0, 1)$ . Since both functions are convex in  $t$ , we conclude that  $\psi_t = \varphi_t$  for all  $t \in (0, 1)$ .

It remains to argue that  $\varphi_t \xrightarrow{L^1} \varphi_0$  as  $t \rightarrow 0+$  and  $\varphi_t \xrightarrow{L^1} \varphi_1$  as  $t \rightarrow 1-$ . By symmetry, it suffices to argue the former. In fact, we know that for any  $t \in (0, 1)$  and any  $j \in I$ ,

$$\varphi_t^j \leq \varphi_t \leq t\varphi_1 + (1-t)\varphi_0,$$

where the latter inequality follows from [Proposition 4.1.1](#). Letting  $t \rightarrow 0+$  and then taking limit with respect to  $j$ , we conclude.  $\square$

## 4.2 Geodesics in the space of potentials

sec:relativeray

Let  $X$  be a connected compact Kähler manifold of dimension  $n$  and  $\theta$  be a smooth closed real  $(1, 1)$ -form on  $X$  representing a big cohomology class. Fix a model potential  $\phi \in \text{PSH}(X, \theta)_{>0}$ . See [Definition 3.1.3](#) for the definition.

**Definition 4.2.1** Let  $\varphi_0, \varphi_1 \in \mathcal{E}(X, \theta; \phi)$ . The *geodesic*  $(\varphi_t)_{t \in (0,1)}$  from  $\varphi_0$  to  $\varphi_1$  is a family of potentials  $\varphi_t \in \text{PSH}(X, \theta)$  such that

$$\varphi_t = \sup^* \{ \eta_t : (\eta_s)_s \text{ is a subgeodesic from } \psi_0 \text{ to } \psi_1, \psi_0, \psi_1 \in \text{PSH}(X, \theta), \psi_0 \leq \varphi_0, \psi_1 \leq \varphi_1 \}. \quad (4.2)$$

{eq:Perron2}

We refer to [Section 3.1.3](#) for the definition of  $\mathcal{E}(X, \theta; \phi)$ .

def:geod2

**Definition 4.2.2** Let  $(\varphi_t)_{t \in [a,b]}$  ( $a, b \in \mathbb{R}, a \leq b$ ) be a curve in  $\mathcal{E}(X, \theta; \phi)$ . We say  $(\varphi_t)_{t \in [a,b]}$  is a *geodesic* if the curve  $(\varphi_{t(b-a)+a})_{t \in (0,1)}$  is a geodesic from  $\varphi_a$  to  $\varphi_b$ .

We also say  $(\varphi_t)_{t \in [a,b]}$  is a geodesic in  $\mathcal{E}(X, \theta; \phi)$  from  $\varphi_a$  to  $\varphi_b$ .

prop:perronenvissubgeod

**Proposition 4.2.1** Given  $\varphi_0, \varphi_1 \in \mathcal{E}(X, \theta; \phi)$ , the geodesic  $(\varphi_t)_{t \in (0,1)}$  from  $\varphi_0$  to  $\varphi_1$  is a subgeodesic from  $\varphi_0$  to  $\varphi_1$  and  $\varphi_t \in \mathcal{E}(X, \theta; \phi)$  for each  $t \in (0, 1)$ .

Moreover, for any  $0 \leq a \leq b \leq 1$ , the restriction  $(\varphi_t)_{t \in [a,b]}$  is a geodesic.

If furthermore  $\varphi_0, \varphi_1 \in \mathcal{E}^1(X, \theta; \phi)$  (resp.  $\mathcal{E}^\infty(X, \theta; \phi)$ ), then  $\varphi_t \in \mathcal{E}^1(X, \theta; \phi)$  (resp.  $\mathcal{E}^\infty(X, \theta; \phi)$ ) for all  $t \in (0, 1)$ .

**Proof** Without loss of generality, we may assume that  $\varphi_0, \varphi_1 \leq \phi$ . It follows from [Proposition 4.1.1](#) that  $\varphi_t \leq \phi$  for all  $t \in (0, 1)$ . In fact, we have the stronger estimate

$$\varphi_t \leq t\varphi_1 + (1-t)\varphi_0, \quad t \in (0, 1). \quad (4.3)$$

{eq:geodesicconvextemp1}

We first observe that when  $\varphi_0, \varphi_1 \in \mathcal{E}(X, \theta; \phi)$ , so is  $\varphi_0 \wedge \varphi_1$ , see [Proposition 3.1.12](#). In particular, the constant subgeodesic  $t \mapsto \varphi_0 \wedge \varphi_1$  is a candidate in (4.2). So

$$\varphi_t \geq \varphi_0 \wedge \varphi_1, \quad t \in (0, 1). \quad (4.4)$$

{eq:varphitgeqlandtemp1}

By [Proposition 4.1.2](#),  $(\varphi_t)_{t \in (0,1)}$  is a subgeodesic. It follows from [Proposition 3.1.13](#) that  $\varphi_t \in \mathcal{E}(X, \theta; \phi)$  for all  $t \in (0, 1)$ .

Next, we show that as  $t \rightarrow 0+$ , we have  $\varphi_t \xrightarrow{L^1} \varphi_0$ . The corresponding result at  $t = 1$  is similar.

We first argue the special case where  $\varphi_0 \leq \varphi_1$ . Take a constant  $C > 0$  such that

$$\varphi_0 - C \leq \varphi_1.$$

Then  $(\varphi_0 - Ct)_{t \in (0,1)}$  is clearly a candidate in (4.2). Therefore, for all  $t \in (0, 1)$ ,

$$\varphi_0 - Ct \leq \varphi_t \leq t\varphi_1 + (1-t)\varphi_0. \quad (4.5)$$

{eq:varphi0andvarphit}

It follows that  $\varphi_t \xrightarrow{L^1} \varphi_0$  as  $t \rightarrow 0+$ .

Let us come back to the general case. By (4.3), we know that for all  $t \in (0, 1)$ ,

$$\sup_X \varphi_t \leq (\sup_X \varphi_0) \vee (\sup_X \varphi_1)$$

On the other hand,  $\sup_X \varphi_t \geq \sup_X \varphi_0 \wedge \varphi_1$ . It follows from [Proposition 1.5.1](#) that  $\{\varphi_t : t \in (0, 1)\}$  is a relatively compact subset of  $\text{PSH}(X, \theta)$  with respect to the  $L^1$ -topology.

Let  $\psi$  be an  $L^1$ -cluster point of  $\varphi_t$  as  $t \searrow 0$ , it suffices to show that  $\psi = \varphi_0$ .

For each  $M \in \mathbb{N}$ , we write

$$\varphi_0^M = \varphi_0 \wedge (\varphi_1 + M).$$

Observe that  $\varphi_0^M \in \mathcal{E}(X, \theta; \phi)$  by [Proposition 3.1.12](#). Let  $(\varphi_t^M)_{t \in (0,1)}$  be the geodesic from  $\varphi_0^M$  to  $\varphi_1$ . Then it is clear that  $\varphi_t^M \leq \varphi_t$  for all  $t \in (0, 1)$ . Therefore,

$$\psi \geq \varphi_0 \wedge (\varphi_1 + M)$$

almost everywhere hence everywhere by [Proposition 1.2.5](#). On the other hand, by (4.3),  $\psi \leq \varphi_0$ . So it suffices to show that

$$\varphi_0 \wedge (\varphi_1 + M) \xrightarrow{L^1} \varphi_0$$

as  $M \rightarrow \infty$ . This is shown in [Proposition 3.1.15](#).

Next, take  $0 \leq a \leq b \leq 1$ . We want to show that the restriction  $(\varphi_t)_{t \in [a,b]}$  is the geodesic from  $\varphi_a$  to  $\varphi_b$ . We may assume that  $a < b$ . The argument is the standard *balayage* argument.

Let  $(\psi_t)_{t \in (a,b)}$  be the (reparameterized) geodesic from  $\varphi_a$  to  $\varphi_b$ . It is easy to see that the curve  $(\eta_t)_{t \in (0,1)}$  defined by  $\eta_t = \psi_t$  for  $t \in (a, b)$  and  $\eta_t = \varphi_t$  otherwise is a candidate in (4.2). See [\[GZ17, Proposition 1.30\]](#). So we conclude that  $\eta_t = \varphi_t = \psi_t$  for  $t \in (a, b)$ .

Finally, assume furthermore that  $\varphi_0, \varphi_1 \in \mathcal{E}^1(X, \theta; \phi)$  (resp.  $\mathcal{E}^\infty(X, \theta; \phi)$ ). Thanks to (4.4), [Proposition 3.1.12](#) and [Proposition 3.1.13](#), we find  $\varphi_t \in \mathcal{E}^1(X, \theta; \phi)$  (resp.  $\mathcal{E}^\infty(X, \theta; \phi)$ ) for all  $t \in (0, 1)$ .  $\square$

prop:geodsupsublinear

**Proposition 4.2.2** *Let  $\varphi_1, \varphi_0 \in \mathcal{E}(X, \theta; \phi)$  with  $\varphi_1 \leq \varphi_0$ . Let  $(\varphi_t)_{t \in (0,1)}$  be the geodesic from  $\varphi_0$  to  $\varphi_1$ . Then*

$$t \sup_{\{\varphi_0 \neq -\infty\}} (\varphi_1 - \varphi_0) = \sup_{\{\varphi_0 \neq -\infty\}} (\varphi_t - \varphi_0) \quad (4.6)$$

{eq:tsupsuptemp1}

for all  $t \in (0, 1]$ .

**Proof** After replacing  $\varphi_t$  by  $\varphi_t - C't$  for some large enough  $C' > 0$ , we may assume that  $\varphi_1 \leq \varphi_0$ . It follows that  $\varphi_1 \leq \varphi_t \leq 0$  for all  $t \in [0, 1]$ . Similarly,  $[0, 1] \ni t \mapsto \varphi_t$  is decreasing.

Let

$$C = \sup_{\{\varphi_1 \neq -\infty\}} (\varphi_1 - \varphi_0).$$

Then by [Proposition 1.2.5](#), we have

$$\varphi_1 \leq \varphi_0 + C.$$

So  $\varphi_1 - C(1 - t)$  is a candidate in [\(4.2\)](#) and hence

$$\varphi_1 - C(1 - t) \leq \varphi_t, \quad t \in (0, 1). \quad (4.7)$$

$$\{\text{eq:varphi1leqvarphittemp}\}$$

By [Proposition 4.2.1](#), we have  $\varphi_t \xrightarrow{L^1} \varphi_1$  as  $t \rightarrow 1-$ . Since  $\varphi_t$  is decreasing in  $t \in (0, 1)$ . It follows that  $\varphi_1 = \inf_{t \in (0, 1)} \varphi_t$ . Therefore, we can find a pluripolar set  $Z \subseteq X$  such that  $\varphi_t(x) \rightarrow \varphi_1(x) > -\infty$  as  $t \rightarrow 1-$  for all  $x \in X \setminus Z$ .

Similarly, since  $\varphi_0 = \sup_{t \in (0, 1)}^* \varphi_t$ , after enlarging  $Z$ , we may also guarantee that  $\varphi_t(x) \rightarrow \varphi_0(x) > -\infty$  as  $t \rightarrow 0+$  for all  $x \in X \setminus Z$  by [Proposition 1.2.3](#).

For any such  $x \in X \setminus Z$ , the function  $t \mapsto \varphi_t(x)$  is a real-valued continuous convex function on  $[0, 1]$ . Hence,

$$\varphi_1(x) - \varphi_0(x) = \int_0^1 \frac{d}{dt} \varphi_t(x) dt \leq \lim_{t \rightarrow 1-} \frac{\varphi_1(x) - \varphi_t(x)}{1 - t} \leq C,$$

where the second inequality follows from [\(4.7\)](#).

Fix an arbitrary pluripolar set  $Z' \supseteq Z$ . Taking supremum, we find that

$$\begin{aligned} \sup_{x \in X \setminus Z'} \varphi_1(x) - \varphi_0(x) &= \sup_{x \in X, \varphi_1(x) \neq -\infty} \varphi_1(x) - \varphi_0(x) \\ &= \sup_{x \in X \setminus Z'} \lim_{t \rightarrow 1-} \frac{\varphi_1(x) - \varphi_t(x)}{1 - t} = C. \end{aligned}$$

Here we have applied [Corollary 1.3.5](#).

Fix  $s \in (0, 1)$ . The same argument shows that after enlarging  $Z'$ , we may guarantee that

$$\sup_{\{\varphi_1 \neq -\infty\}} (\varphi_1 - \varphi_0) = \sup_{x \in X \setminus Z'} \lim_{t \rightarrow 1-} \frac{\varphi_1(x) - \varphi_t(x)}{1 - t} = \sup_{\{\varphi_1 \neq -\infty\}} \frac{\varphi_1 - \varphi_s}{1 - s}. \quad (4.8)$$

$$\{\text{eq:supvarphi1mivarphi@temp1}\}$$

On the other hand,

$$\sup_{\{\varphi_1 \neq -\infty\}} (\varphi_1 - \varphi_0) \leq s \sup_{\{\varphi_1 \neq -\infty\}} \frac{\varphi_s - \varphi_0}{s} + (1 - s) \sup_{\{\varphi_1 \neq -\infty\}} \frac{\varphi_1 - \varphi_s}{1 - s}.$$

Together with [\(4.8\)](#), we find that

$$\sup_{\{\varphi_1 \neq -\infty\}} (\varphi_1 - \varphi_0) \leq \sup_{\{\varphi_1 \neq -\infty\}} \frac{\varphi_s - \varphi_0}{s}.$$

Using the convexity, we find that equality holds,

$$\sup_{\{\varphi_1 \neq -\infty\}} \frac{\varphi_s - \varphi_0}{s} = \sup_{\{\varphi_1 \neq -\infty\}} (\varphi_1 - \varphi_0).$$

Using [Corollary 1.3.5](#), we conclude (4.6).  $\square$

With an almost identical proof, we find

prop:geodinfsublinear

**Proposition 4.2.3** Let  $\varphi_1, \varphi_0 \in \mathcal{E}^\infty(X, \theta; \phi)$ . Let  $(\varphi_t)_{t \in (0,1)}$  be the geodesic from  $\varphi_0$  to  $\varphi_1$ . Then

$$t \inf_{\{\phi \neq -\infty\}} (\varphi_1 - \varphi_0) = \inf_{\{\phi \neq -\infty\}} (\varphi_t - \varphi_0)$$

for all  $t \in (0, 1]$ .

**Definition 4.2.3** Let  $\ell = (\ell_t)_{t \geq 0}$  be a curve in  $\mathcal{E}(X, \theta; \phi)$ . We say  $\ell$  is a *geodesic ray* in  $\mathcal{E}(X, \theta; \phi)$  emanating from  $\ell_0$  if for each  $0 \leq a \leq b$ , the restriction  $(\ell_t)_{t \in [a,b]}$  is a geodesic.

The set of geodesic rays in  $\mathcal{E}(X, \theta; \phi)$  emanating from  $\phi$  is denoted by  $\mathcal{R}(X, \theta; \phi)$ .

We say a geodesic ray  $\ell \in \mathcal{R}(X, \theta; \phi)$  has *finite energy* if  $\ell_t \in \mathcal{E}^1(X, \theta; \phi)$  for all  $t > 0$ . The set of geodesic rays with finite energy is denoted by  $\mathcal{R}^1(X, \theta; \phi)$ .

We say a geodesic ray  $\ell \in \mathcal{R}(X, \theta; \phi)$  is *bounded* if  $\ell_t \in \mathcal{E}^\infty(X, \theta; \phi)$  for all  $t \geq 0$ . The set of bounded geodesic rays is denoted by  $\mathcal{R}^\infty(X, \theta; \phi)$ .

Given  $\ell, \ell' \in \mathcal{R}(X, \theta; \phi)$ , we write  $\ell \leq \ell'$  if  $\ell_t \leq \ell'_t$  for each  $t \geq 0$ .

When  $\phi = V_\theta$ , we usually omit it from the notations and write  $\mathcal{R}(X, \theta)$ ,  $\mathcal{R}^1(X, \theta)$  and  $\mathcal{R}^\infty(X, \theta)$ ,

prop:raysuplinear

**Proposition 4.2.4** Let  $\ell \in \mathcal{R}(X, \theta; \phi)$ . Then there is a constant  $C > 0$  such that

$$\sup_X \ell_t \leq Ct, \quad t \geq 0.$$

In fact, more precisely, we have

$$\ell_t \leq \phi + Ct.$$

**Proof** Let  $Z = \{\phi = -\infty\}$ . It follows from [Proposition 4.2.2](#) that

$$\ell_t \leq \phi + t \sup_{X \setminus Z} (\ell_1 - \phi), \quad t \geq 0.$$

Since  $\ell_1 \in \mathcal{E}(X, \theta; \phi)$ , we have  $\ell_1 \leq \phi + C$  for some constant  $C$  and our conclusion follows.  $\square$

def:radialMAenergy2

**Definition 4.2.4** We define the *radial Monge–Ampère energy*  $\mathbf{E}^\phi: \mathcal{R}^1(X, \theta; \phi) \rightarrow \mathbb{R}$  as follows:



$$\mathbf{E}^\phi(\ell) := \overline{\lim}_{t \rightarrow \infty} \frac{E_\theta^\phi(\ell_t)}{t}.$$

When  $\phi = V_\theta$ , we write  $\mathbf{E}$  instead of  $\mathbf{E}^{V_\theta}$ .

Thanks to [Proposition 4.2.2](#),  $\mathbf{E}^\phi(\ell) < \infty$ .

def:d1onE12

**Definition 4.2.5** Let  $\varphi, \psi \in \mathcal{E}^1(X, \theta; \phi)$ , we define

$$d_1(\varphi, \psi) = E_\theta^\phi(\varphi) + E_\theta^\phi(\psi) - 2E_\theta^\phi(\varphi \wedge \psi).$$

In particular, if  $\varphi \leq \psi$ , we have

$$d_1(\varphi, \psi) = E_\theta^\phi(\psi) - E_\theta^\phi(\varphi). \quad (4.9)$$

{eq:d1asEdiff}

thm:d1complete

**Theorem 4.2.1** The function  $d_1$  defined in [Definition 4.2.5](#) is a complete metric on  $\mathcal{E}^1(X, \theta; \phi)$ .

The function  $E_\theta^\phi : \mathcal{E}^1(X, \theta; \phi) \rightarrow \mathbb{R}$  is continuous with respect to  $d_1$ .

Moreover, given a decreasing (resp. increasing) sequence  $(\varphi_j)_{j \in \mathbb{Z}_{>0}}$  in  $\mathcal{E}^1(X, \theta; \phi)$  converging (resp. converging almost everywhere) to  $\varphi \in \mathcal{E}^1(X, \theta; \phi)$ , then  $\varphi_j \xrightarrow{d_1} \varphi$ .

See [\[DDNL18big\]](#), Theorem 1.1, Proposition 2.9, Proposition 2.7]. The readers should have no difficulty in generalizing all arguments to the current setting.

thm:d1lor

**Theorem 4.2.2** Let  $\varphi, \psi, \eta \in \mathcal{E}^1(X, \theta; \phi)$ . Then

$$d_1(\varphi \vee \eta, \psi \vee \eta) \leq d_1(\varphi, \psi).$$

See [\[Xia23Mabuchi\]](#), Proposition 4.12] (Proposition 6.8 in the arXiv version).

Next we recall a few particular properties when  $\phi = V_\theta$ .

prop:energylinear

**Proposition 4.2.5** Let  $(\varphi_t)_{t \in [a, b]}$  be a geodesic in  $\mathcal{E}^1(X, \theta)$ , then  $t \mapsto E_\theta(\varphi_t)$  is a linear function of  $t \in [a, b]$ .

We expect that  $t \mapsto E_\theta^\phi(\varphi_t)$  is linear in general. The author does not know how to prove this.

**Proof** This follows from [\[DDNL18fullmass\]](#), Theorem 3.12]. □

prop:d1geod\_diff\_E

**Proposition 4.2.6** Let  $\ell, \ell' \in \mathcal{R}^1(X, \theta)$  and  $\ell \leq \ell'$ . Then

$$d_1(\ell, \ell') = \mathbf{E}(\ell') - \mathbf{E}(\ell). \quad (4.10)$$

{eq:d1rayscompa}

**Proof** This is a direct consequence of [\(4.9\)](#). □

**Proposition 4.2.7** Let  $\ell, \ell' \in \mathcal{R}^1(X, \theta)$ . Then the map

$$t \mapsto d_1(\ell_t, \ell'_t)$$

is convex.

See [\[DDNLmetric, Proposition 2.10\]](#) for the proof. In particular, we can introduce

`def:dirays`

**Definition 4.2.6** Let  $\ell, \ell' \in \mathcal{R}^1(X, \theta)$ . We define

$$d_1(\ell, \ell') := \lim_{t \rightarrow \infty} \frac{1}{t} d_1(\ell_t, \ell'_t).$$

`thm:diraycomplete`

**Theorem 4.2.3** The function  $d_1$  defined in [Definition 4.2.6](#) is a metric and  $(\mathcal{R}^1(X, \theta), d_1)$  is a complete metric space.

See [\[DDNLmetric, Theorem 2.14\]](#) for the proof.

`prop:supsgeod`

**Proposition 4.2.8** Let  $(\varphi_0^i)_{i \in I}, (\varphi_1^i)_{i \in I}$  be two uniformly bounded from above increasing nets in  $\mathcal{E}^\infty(X, \theta)$ . Let  $(\varphi_t^i)_{t \in (0,1)}$  be the geodesic from  $\varphi_0^i$  to  $\varphi_1^i$  for each  $i \in I$ . Then

$$\left( \sup_{i \in I}^* \varphi_t^i \right)_{t \in (0,1)}$$

is the geodesic from  $\sup_{i \in I}^* \varphi_0^i$  to  $\sup_{i \in I}^* \varphi_1^i$ .

**Proof** By [Proposition 1.2.2](#) and [Proposition 4.1.2](#), we may assume that  $I$  is countable. In this case, the assertion follows from [\[DDNL8fullmass, Proposition 3.3\]](#) and [Theorem 2.1.1](#).  $\square$

Next we recall that  $\vee$  operator at the level of geodesic rays.

`def:lorray1`

**Definition 4.2.7** Let  $\ell, \ell' \in \mathcal{R}(X, \theta)$ . We define  $\ell \vee \ell'$  as the minimal ray in  $\mathcal{R}(X, \theta)$  lying above both  $\ell$  and  $\ell'$ .

`prop:lorrays`

**Proposition 4.2.9** Given  $\ell, \ell' \in \mathcal{R}(X, \theta)$ . Then  $\ell \vee \ell' \in \mathcal{R}(X, \theta)$  exists. Moreover, if  $\ell, \ell' \in \mathcal{R}^1(X, \theta)$ , then so is  $\ell \vee \ell'$  and

$$\mathbf{E}(\ell \vee \ell') = \lim_{t \rightarrow \infty} \frac{1}{t} E_\theta(\ell_t \vee \ell'_t). \quad (4.11)$$

`{eq:Elor}`

Furthermore, if both  $\ell, \ell' \in \mathcal{R}^\infty(X, \theta)$ , then so is  $\ell \vee \ell'$ .

**Proof** For each  $t > 0$ , let  $(\ell_s'')_{s \in [0,t]}$  be the geodesic from  $V_\theta$  to  $\ell_t \vee \ell'_t$ . Then clearly, for each fixed  $s \geq 0$ ,  $\ell_s''$  is increasing in  $t \in [s, \infty)$ . Moreover, [Proposition 4.2.4](#) guarantees that  $(\sup_X \ell_s'')_t$  is bounded from above for a fixed  $s$ . Let  $(\ell \vee \ell')_s = \sup_{t \geq s}^* \ell_s''$ . Then [Proposition 4.2.8](#) guarantees that  $\ell \vee \ell'$  is a geodesic ray. It is clear that this ray is minimal among all rays dominating  $\ell$  and  $\ell'$ .

Assume that  $\ell, \ell' \in \mathcal{R}^1(X, \theta)$ , it follows from [Proposition 3.1.13](#) that  $\ell \vee \ell' \in \mathcal{R}^1(X, \theta)$ . Next we compute its energy:

$$\mathbf{E}(\ell \vee \ell') = E_\theta(\ell \vee \ell')_1 = \lim_{t \rightarrow \infty} E_\theta(\ell_1'') = \frac{1}{t} E_\theta(\ell_t \vee \ell'_t),$$

where we applied [Proposition 4.2.5](#) and [Theorem 4.2.1](#).

The last assertion is trivial.  $\square$

lma:d1rayineq

**Lemma 4.2.1** For any  $\ell, \ell' \in \mathcal{R}^1(X, \theta)$ , we have

$$d_1(\ell, \ell') \leq d_1(\ell, \ell \vee \ell') + d_1(\ell', \ell \vee \ell') \leq C_n d_1(\ell, \ell'), \quad (4.12)$$

{eq:d1maxineq}

where  $C_n = 3(n+1)2^{n+2}$ .**Proof** The first inequality is trivial. As for the second, we estimate

$$\begin{aligned} d_1(\ell, \ell \vee \ell') &= \mathbf{E}(\ell \vee \ell') - \mathbf{E}(\ell) \\ &= \lim_{t \rightarrow \infty} \frac{1}{t} \mathbf{E}(\ell_t \vee \ell'_t) - \mathbf{E}(\ell) \\ &= \lim_{t \rightarrow \infty} \frac{1}{t} d_1(\ell_t \vee \ell'_t, \ell_t), \end{aligned}$$

where on the first line, we applied [Proposition 4.2.6](#), on the second line, we used [\(4.11\)](#), the first and the third lines follow from [Proposition 4.2.6](#). In all, we find

$$d_1(\ell, \ell \vee \ell') + d_1(\ell', \ell \vee \ell') \leq \lim_{t \rightarrow \infty} \frac{1}{t} (d_1(\ell_t \vee \ell'_t, \ell_t) + d_1(\ell_t \vee \ell'_t, \ell'_t)).$$

By [\[DDNL18big\]](#), [\[DDNL18a\]](#), Theorem 3.7],

$$d_1(\ell_t \vee \ell'_t, \ell_t) + d_1(\ell_t \vee \ell'_t, \ell'_t) \leq 3(n+1)2^{n+2} d_1(\ell_t, \ell'_t).$$

Now [\(4.12\)](#) follows.  $\square$ 

ex:rayasspsh

**Example 4.2.1** Let  $\varphi \in \text{PSH}(X, \theta)$ . For each  $C > 0$ , let  $(\ell_t^{\varphi, C})_{t \in [0, C]}$  be the geodesic from  $V_\theta$  to  $(V_\theta - C) \vee \varphi$ . For each  $t \geq 0$ , the potential  $\ell_t^{\varphi, C}$  is increasing in  $C \in [t, \infty)$ . We let

$$\ell_t^\varphi := \sup_{C \geq t}^* \ell_t^{\varphi, C}. \quad (4.13)$$

{eq:ellvarphiraydef}

Then  $\ell^\varphi \in \mathcal{R}^\infty(X, \theta)$  and

$$\mathbf{E}(\ell^\varphi) = \frac{1}{n+1} \sum_{j=0}^n \left( \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} - \int_X \theta_{V_\theta}^n \right). \quad (4.14)$$

{eq:Elphi}

**Proof** We first show that for each fixed  $t \geq 0$ ,  $\ell_t^{\varphi, C}$  is increasing in  $C \geq t$ .

To see this, choose  $t \leq C_1 < C_2$ . We need to show that

$$\ell_t^{\varphi, C_1} \leq \ell_t^{\varphi, C_2}.$$

Since both sides are geodesics for  $t \in [0, C_1]$ , it suffices to show that

$$(V_\theta - C_1) \vee \varphi \leq \ell_{C_1}^{\varphi, C_2}. \quad (4.15)$$

{eq:VthetaminusC1temp1}

Then  $((V_\theta - t) \vee \varphi)_{t \in [0, C_2]}$  is a subgeodesic from  $V_\theta$  to  $(V_\theta - C_2) \vee \varphi$  by [Proposition 4.1.2](#). At  $t = 0$  and  $t = C_1$ , it is dominated by the geodesic  $\ell_t^{\varphi, C_2}$ , hence by [\(4.2.1\)](#), we conclude that the same holds at  $t = C_1$ , which is exactly [\(4.15\)](#).

From [Proposition 4.1.1](#), we know that for any  $C \geq t > 0$ , we have

$$\ell_t^{\varphi, C} \leq t((V_\theta - C) \vee \varphi) + (1-t)V_\theta \leq 0.$$

So in [\(4.13\)](#),  $\ell_t^\varphi \in \text{PSH}(X, \theta)$  for any  $t > 0$ . Also observe that by [Proposition 4.2.1](#), we have  $\ell_t^\varphi \in \mathcal{E}^\infty(X, \theta)$  for all  $t > 0$ . It follows from [Proposition 4.2.8](#) that  $\ell^\varphi \in \mathcal{R}^1(X, \theta)$ .

It remains to compute the energy of  $\ell^\varphi$ .

We first fix  $C \geq t > 0$  and compute

$$E_\theta(\ell_t^{\varphi, C}) = \frac{t}{C} E_\theta((V_\theta - C) \vee \varphi).$$

Letting  $C \rightarrow \infty$  and applying [Theorem 4.2.1](#), we find that

$$E_\theta(\ell_t^\varphi) = \lim_{C \rightarrow \infty} \frac{t}{C} E_\theta((V_\theta - C) \vee \varphi).$$

It follows that

$$\mathbf{E}(\ell^\varphi) = \lim_{C \rightarrow \infty} \frac{1}{C} E_\theta((V_\theta - C) \vee \varphi).$$

Using the definition of  $E_\theta$ , it suffices to show that for each  $j = 0, \dots, n$ , we have

$$\lim_{C \rightarrow \infty} \int_X \frac{(V_\theta - C) \vee \varphi - V_\theta}{C} \theta_{(V_\theta - C) \vee \varphi}^j \wedge \theta_{V_\theta}^{n-j} = \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} - \int_X \theta_{V_\theta}^n. \quad (4.16)$$

{eq:limCintXtemp1}

For this purpose, for each  $C > 0$ , we decompose  $X$  as  $\{\varphi > V_\theta - C\}$  and  $\{\varphi \leq V_\theta - C\}$ .

We have

$$\begin{aligned} & \int_{\{\varphi > V_\theta - C\}} \frac{(V_\theta - C) \vee \varphi - V_\theta}{C} \theta_{(V_\theta - C) \vee \varphi}^j \wedge \theta_{V_\theta}^{n-j} \\ &= \int_{\{\varphi > V_\theta - C\}} \frac{\varphi - V_\theta}{C} \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j}. \end{aligned}$$

On the other hand,

$$\begin{aligned} & \int_{\{\varphi \leq V_\theta - C\}} \frac{(V_\theta - C) \vee \varphi - V_\theta}{C} \theta_{(V_\theta - C) \vee \varphi}^j \wedge \theta_{V_\theta}^{n-j} \\ &= - \int_{\{\varphi \leq V_\theta - C\}} \theta_{(V_\theta - C) \vee \varphi}^j \wedge \theta_{V_\theta}^{n-j} \\ &= - \int_X \theta_{V_\theta}^n + \int_{\{\varphi > V_\theta - C\}} \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j}. \end{aligned}$$

Observe that for  $C > 0$ , the functions  $\mathbb{1}_{\{\varphi > V_\theta - C\}} C^{-1}(\varphi - V_\theta)$  is defined almost everywhere and is bounded. When  $C \rightarrow \infty$ , these functions converge to 0 almost everywhere. Therefore, [\(4.16\)](#) follows.  $\square$

## Chapter 5

# Toric pluripotential theory on ample line bundles

chap:toric\_ample

In this chapter, we briefly recall the toric pluripotential theory relative to an ample line bundle. The general case of big line bundles will be handled in [Chapter 12](#) after developing the powerful machinery of partial Okounkov bodies in [Chapter 10](#). The main new result is [Theorem 5.3.1](#) computing the  $L^2$ -sections of a Hermitian big line bundle in the toric setting.

### 5.1 Toric setup

sec:toricsetup

Let  $T$  be a complex torus of dimension  $n$  and  $T_c \subset T(\mathbb{C})$  denotes the corresponding compact torus. Write  $M$  for its character lattice, which is a free Abelian group of rank  $n$ . Similarly, let  $N$  be cocharacter lattice of  $T$ . Let  $P \subseteq M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$  be a full-dimensional *smooth*<sup>1</sup> lattice polytope.

Let  $\Sigma$  be the normal fan of  $P$ . The notation  $\Sigma(1)$  denotes the set of rays in  $\Sigma$ . For each  $\rho \in \Sigma(1)$ , let  $u_\rho \in N$  denote the ray generator of  $\rho$ , namely the first non-zero element in  $N \cap \rho$ . We write

$$P = \{m \in M_{\mathbb{R}} : \langle m, u_\rho \rangle \geq -a_\rho \text{ for all } \rho \in \Sigma(1)\}.$$

Let  $\text{Supp}_P : N_{\mathbb{R}} \rightarrow \mathbb{R}$  denote the support function of  $P$ . Recall that the support function ([Example A.1.2](#)) of  $P$  is defined as

$$\text{Supp}_P(n) = \max \{(m, n) : m \in P\}.$$

Our convention differs from [\[CLS11, Proposition 4.2.14\]](#) by a minus sign.

Let  $X = X_\Sigma$  be the smooth projective toric variety corresponding to  $\Sigma$ . There is a canonical embedding  $T \subseteq X$  as a dense Zariski open subset. Let  $D$  be the Cartier

<sup>1</sup> Recall that *smooth* means that for every vertex  $v \in P$ , if we take the first lattice point  $w_E$  apart from  $v$  as one transverses each edge  $E$  of  $P$  containing  $v$  from  $v$ , then  $\{w_E - v\}_E$  forms a basis of  $M$ . See [\[CLS11, Definition 2.4.2\]](#). We also say  $P$  is a *Delzant polytope* in this case.

divisor on  $X$  defined by  $P$ :

$$D = \sum_{\rho \in \Sigma(1)} a_\rho D_\rho,$$

where  $D_\rho$  is the toric prime divisor defined by  $\rho$  under the orbit–cone correspondence. Let  $L$  be the toric line bundle induced by  $P$ , namely  $L = \mathcal{O}_X(D_\rho)$ . Since  $P$  has full dimension,  $L^k$  is very ample for each  $k \geq n - 1$  by [CLS11, Corollary 2.2.19], we actually know that  $L$  is ample.

We will choose the base  $e$  for the logarithm map

$$\mathbb{C}^* \rightarrow \mathbb{R}, \quad z \mapsto \log |z|^2.$$

This choice will be fixed throughout the whole section. Since we have a canonical identification  $T(\mathbb{C}) \cong N \otimes_{\mathbb{Z}} \mathbb{C}^*$ , we obtain an identification  $T(\mathbb{C})/T_c \cong N_{\mathbb{R}}$ . This gives a tropicalization map

$$\text{Trop}: T(\mathbb{C}) \rightarrow N_{\mathbb{R}}. \quad (5.1)$$

## 5.2 Toric plurisubharmonic functions

We continue to use the notations of Section 5.1.

lma:convextopsh

**Lemma 5.2.1** *Let  $F: N_{\mathbb{R}} \rightarrow [-\infty, \infty]$  be a function. Then the following are equivalent:*

- (1)  $F$  is convex and takes values in  $\mathbb{R}$ , and
- (2)  $\text{Trop}^* F$  is plurisubharmonic on  $T(\mathbb{C})$ .

**Proof** We may choose an identification  $N \cong \mathbb{Z}^n$  so that we have an identification  $T(\mathbb{C}) \cong \mathbb{C}^{*n}$ . Then  $\text{Trop}$  is identified with the map

$$\text{Trop}: \mathbb{C}^{*n} \rightarrow \mathbb{R}^n, \quad (z_1, \dots, z_n) \mapsto (\log |z_1|^2, \dots, \log |z_n|^2).$$

(1)  $\implies$  (2). Let  $F_k \in C^\infty(\mathbb{R}^n) \cap \text{Conv}(\mathbb{R}^n)$  be a decreasing sequence with limit  $F$  (see Proposition A.3.3). It follows from a straightforward computation that

$$\text{dd}^c \text{Trop}^* F_k(z_1, \dots, z_n) = \frac{i}{2\pi} \sum_{i,j=1}^n \partial_{i\bar{j}} F_k \left( \log |z_1|^2, \dots, \log |z_n|^2 \right) z_i^{-1} \bar{z}_j^{-1} dz_i \wedge d\bar{z}_j. \quad (5.2)$$

{eq:ddctrop}

So  $\text{Trop}^* F_k$  is plurisubharmonic. It follows from Proposition 1.2.1 that  $\text{Trop}^* F$  is plurisubharmonic.

(2)  $\implies$  (1). It follows from Lemma 1.2.1 that  $F$  is finite. Moreover, take a radial mollifier, we may find a decreasing sequence  $\varphi_k$  of smooth psh functions on  $\mathbb{C}^{*n}$  with

limit  $\text{Trop}^* F$ . Write  $\varphi_k = \text{Trop}^* F_k$  for some function  $F_k : \mathbb{R}^n \rightarrow \mathbb{R}$ , it follows from (5.2) that  $F_k$  is convex for all  $k$ . Therefore,  $F$  is convex by Lemma A.1.2.  $\square$

Let  $G_0 : M_{\mathbb{R}} \rightarrow (-\infty, \infty]$  be defined as

$$G_0(m) := \begin{cases} \frac{1}{2} \sum_{\rho \in \Sigma(1)} (\langle m, u_{\rho} \rangle + a_{\rho}) \log (\langle m, u_{\rho} \rangle + a_{\rho}), & \text{if } m \in P, \\ \infty, & \text{otherwise.} \end{cases} \quad (5.3) \quad \{\text{eq:G0def}\}$$

This is a closed proper convex function and  $G_0 \sim \chi_P$ . Let

$$F_0 = G_0^* \in \mathcal{E}^{\infty}(N_{\mathbb{R}}, P). \quad (5.4) \quad \{\text{eq:F0def}\}$$

By Guillemin's theorem [Gui94, CDG03],  $\text{dd}^c \text{Trop}^* F_0$  can be extended to a unique Kähler form  $\omega$  in  $c_1(L)$ .

Let  $\text{PSH}_{\text{tor}}(X, \omega)$  denote the set of  $T_c$ -invariant  $\omega$ -psh functions.

thm:toricpsh

**Theorem 5.2.1** *There is a canonical bijection between the following three sets:*

- (1) *The set of  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ ,*
- (2) *the set  $\mathcal{P}(N_{\mathbb{R}}, P)$  in Definition A.3.1, namely, the set of convex functions  $F : N_{\mathbb{R}} \rightarrow \mathbb{R}$  satisfying  $F \leq \text{Supp}_P$ , and*
- (3) *the set of closed proper convex functions  $G \in \text{Conv}(M_{\mathbb{R}})$  satisfying*

$$G|_{M_{\mathbb{R}} \setminus P} \equiv \infty.$$

**Proof** The bijection between (2) and (3) is the classical Legendre duality. Given  $F$  as in (2), we construct  $G = F^*$ , see Proposition A.2.4.

The map from (1) to (2) is given as follows: given  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ , since  $\varphi$  is  $T_c$ -invariant, we can find  $f : N_{\mathbb{R}} \rightarrow [-\infty, \infty)$  such that

$$\varphi|_{T(\mathbb{C})} = \text{Trop}^* f.$$

We then define  $F = f + F_0$ . By Lemma 5.2.1,  $F(n)$  is finite for any  $n \in N_{\mathbb{R}}$  and  $F$  is convex. Moreover,  $F \leq \text{Supp}_P$  since this holds for  $F_0$ .

Conversely, given a map  $F \in \mathcal{P}(N_{\mathbb{R}}, P)$ , then

$$\text{Trop}^*(F - F_0) \in \text{PSH}(T(\mathbb{C}), \omega|_{T(\mathbb{C})}).$$

It follows from Theorem 1.2.1 that this function can be extended uniquely to an  $\omega$ -psh function on  $X$ . The uniqueness of the extension guarantees its  $T_c$ -invariance.

The two maps are clearly inverse to each other.  $\square$

Given  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ , we will write  $F_{\varphi}$  and  $G_{\varphi}$  for the convex functions given by Theorem 5.2.1.

**Proposition 5.2.1** *Given  $\varphi, \psi \in \text{PSH}_{\text{tor}}(X, \omega)$ . The following are equivalent:*

- (1)  $\varphi \leq \psi$ ,

- (2)  $F_\varphi \leq F_\psi$ , and  
 (3)  $G_\varphi \geq G_\psi$ .

In particular,  $\varphi \in \mathcal{E}^\infty(X, \theta)$  if and only if  $F_\varphi \in \mathcal{E}^\infty(N_{\mathbb{R}}, P)$ .

prop:toricpluscst

**Proposition 5.2.2** Given  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$  and  $C \in \mathbb{R}$ . We have

$$F_{\varphi+C} = F_\varphi + C, \quad G_{\varphi+C} = G_\varphi - C.$$

Both results follow immediately from the constructions of  $F$  and  $G$ . We leave the details to the readers.

prop:toricrooftop

**Proposition 5.2.3** Given  $\varphi, \psi \in \text{PSH}_{\text{tor}}(X, \omega)$ , then  $\varphi \wedge \psi \in \text{PSH}_{\text{tor}}(X, \omega)$  and

$$F_{\varphi \wedge \psi} = F_\varphi \wedge F_\psi, \quad G_{\varphi \wedge \psi} = G_\varphi \vee G_\psi.$$

**Proof** It is clear that  $\varphi \wedge \psi \in \text{PSH}_{\text{tor}}(X, \omega)$ . The claim for  $G$  is obvious and the claim for  $F$  follows from [Proposition A.2.2](#).  $\square$

prop:toricseq

**Proposition 5.2.4** Let  $\{\varphi_i\}_{i \in I}$  be a family in  $\text{PSH}_{\text{tor}}(X, \omega)$  uniformly bounded from above. Then  $\sup_{i \in I}^* \varphi_i \in \text{PSH}_{\text{tor}}(X, \omega)$  and

$$F_{\sup_{i \in I}^* \varphi_i} = \sup_{i \in I} F_{\varphi_i}, \quad G_{\sup_{i \in I}^* \varphi_i} = \text{cl} \bigwedge_{i \in I} G_{\varphi_i}.$$

Moreover, if  $I$  is finite, then

$$G_{\max_{i \in I} \varphi_i} = \bigwedge_{i \in I} G_{\varphi_i}.$$

Similarly, if  $\{\varphi_i\}_{i \in I}$  is a decreasing net in  $\text{PSH}_{\text{tor}}(X, \omega)$  such that  $\inf_{i \in I} \varphi_i \not\equiv -\infty$ , then  $\inf_{i \in I} \varphi_i \in \text{PSH}_{\text{tor}}(X, \omega)$  and

$$F_{\inf_{i \in I} \varphi_i} = \inf_{i \in I} F_{\varphi_i}, \quad G_{\inf_{i \in I} \varphi_i} = \sup_{i \in I} G_{\varphi_i}.$$

**Proof** In both cases, the statement for  $F$  is clear. The corresponding statement for  $G$  is obtained via [Proposition A.2.2](#).  $\square$

prop:toricMAandrealMA

**Proposition 5.2.5** Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ , then

$$\text{Trop}_* (\omega|_{T(\mathbb{C})} + \text{dd}^c \varphi|_{T(\mathbb{C})})^n = \text{MA}_{\mathbb{R}}(F_\varphi). \quad (5.5)$$

{eq:tropMAmea}

In particular,

$$\int_X \omega_\varphi^n = \int_{N_{\mathbb{R}}} \text{MA}_{\mathbb{R}}(F_\varphi) = n! \text{vol} \{ \overline{G_\varphi} < \infty \}$$

and

$$\int_X \omega^n = n! \text{vol } P.$$



**Proof** We first prove (5.5). By Proposition A.3.3, we can find a decreasing sequence of smooth convex functions  $F_j$  on  $N_{\mathbb{R}}$  with limit  $F_{\varphi}$ . We write  $F_j = F_{\varphi_j}$  for some  $\varphi_j \in \text{PSH}_{\text{tor}}(X, \omega)$ . By Theorem 2.1.1 and Theorem A.4.1, we may reduce to the case where  $F_{\varphi}$  is smooth. Then it suffices to carry out the straightforward computation using (5.2).  $\square$

### 5.3 Toric pluripotential theory

sec:envelopestoric

Let us begin by consider the  $P$ -envelope.

**Definition 5.3.1** Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ . We define its *Newton body* as

$$\Delta(\omega, \varphi) := \overline{\{G_{\varphi} < \infty\}} \subseteq P.$$

By Proposition A.2.1, we have

$$\Delta(\omega, \varphi) = \overline{\nabla F_{\varphi}(N_{\mathbb{R}})}.$$

prop:GPenvelope

**Proposition 5.3.1** Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ . Then  $P_{\omega}[\varphi] \in \text{PSH}_{\text{tor}}(X, \omega)$  and

$$G_{P_{\omega}[\varphi]}(x) = \begin{cases} G_0(x), & \text{if } x \in \Delta(\omega, \varphi); \\ \infty, & \text{otherwise.} \end{cases} \quad (5.6) \quad \{\text{eq:toricPenv}\}$$

**Proof** By (3.2), we have

$$P_{\omega}[\varphi] = \sup_{C \in \mathbb{R}}^* ((\varphi + C) \wedge 0).$$

It follows from Proposition 5.2.2, Proposition 5.2.3 and Proposition 5.2.4 that  $P_{\omega}[\varphi] \in \text{PSH}_{\text{tor}}(X, \omega)$ . Moreover, by the same propositions, we have

$$G_{P_{\omega}[\varphi]} = \inf_{C \in \mathbb{R}} (G_0 \vee (G_{\varphi} - C)),$$

which is clearly equal to the right-hand side of (5.6).

Next we prove a result of Yi Yao claiming that in the toric setting, all potentials are  $I$ -good.

thm:Yao

**Theorem 5.3.1** Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ , then

$$h^0(X, L \otimes I(\varphi)) = \#(\Delta(\omega, \varphi) \cap M).$$

**Proof** It is well-known that  $H^0(X, L)$  can be identified with the vector space generated by  $\chi^m$  for all  $m \in P \cap M$ , see [CLS11, Proposition 4.3.3]. We will show that

$$H^0(X, L \otimes \mathcal{I}(\varphi)) = \bigoplus_{m \in \Delta(\omega, \varphi) \cap M} \mathbb{C} \chi^m. \quad (5.7)$$

{eq:toricL2sec}

It is convenient to use explicit coordinates. We will identify  $N$  with  $\mathbb{Z}^n$  after choosing a basis. In this way, we get an identification  $M = \mathbb{Z}^n$  and  $T(\mathbb{C}) = \mathbb{C}^{*n}$ . In this case, we have

$$\chi^m(z) = z^m$$

with the multi-index notation.

Observe that  $H^0(X, L \otimes \mathcal{I}(\varphi))$  is a  $\mathbb{C}^{*n}$ -invariant subspace of  $H^0(X, L)$ , it follows that  $H^0(X, L \otimes \mathcal{I}(\varphi))$  is the direct sum of suitable  $\chi^m$ 's.

We first show that  $\chi^m \in H^0(X, L \otimes \mathcal{I}(\varphi))$  for each  $m \in \Delta(\omega, \varphi) \cap M$ . We need to show that

$$\int_{\mathbb{C}^{*n}} |\chi^m|^2 \exp(-P_\omega[\varphi]) \omega^n < \infty.$$

Using [Proposition 5.3.1](#) and [Proposition 5.2.5](#), we find that the latter holds if and only if

$$\int_{\mathbb{R}^n} \exp\left(\langle m, n \rangle - \text{Supp}_{\Delta(\omega, \varphi)}(n)\right) \text{MA}_{\mathbb{R}}(F_0)(n) < \infty,$$

which is obvious since

$$\langle m, n \rangle - \text{Supp}_{\Delta(\omega, \varphi)}(n) \leq 0.$$

Next we show that for any  $m \in M \cap (P \setminus \Delta(\omega, \varphi))$ , the function  $\chi^m$  does not lie in  $H^0(X, L \otimes \mathcal{I}(\varphi))$ . Again, this means

$$\int_{\mathbb{R}^n} \exp\left(\langle m, n \rangle - \text{Supp}_{\Delta(\omega, \varphi)}(n)\right) \text{MA}_{\mathbb{R}}(F_0)(n) = \infty.$$

By change of variables, this is equivalent to

$$\int_P \exp\left(\langle m, \nabla G_0(m') \rangle - \text{Supp}_{\Delta(\omega, \varphi)}(\nabla G_0(m'))\right) dm' = \infty.$$

Since  $m$  does not lie in  $\Delta(\omega, \varphi)$ , we can find  $n_0 \in \mathbb{R}^n$  such that

$$\langle m, n_0 \rangle - \text{Supp}_{\Delta(\omega, \varphi)}(n_0) > 0.$$

In particular, there are closed convex cones  $C' \subseteq C$  containing  $n_0$  in their interiors such that there exists  $\epsilon > 0$  such that

$$\langle m, n \rangle - \text{Supp}_{\Delta(\omega, \varphi)}(n) \geq \epsilon |n|$$

for all  $n \in C$  and  $C'$  intersects the boundary of  $C$  only at 0.

Thus, it would suffice to prove

$$\int_{P \cap \{\nabla G_0 \subseteq C'\}} \exp(\epsilon |\nabla G_0(m')|) dm' = \infty. \quad (5.8)$$

{eq:intexpinftempl}

For each  $\rho \in \Sigma(1)$ , we write

$$r_\rho(m') = \log(\langle m', u_\rho \rangle + a_\rho) + 1, \quad m' \in \mathbb{R}^n.$$

It follows from (5.3) that

$$\nabla G_0(m') = \frac{1}{2} \sum_{\rho \in \Sigma(1)} r_\rho(m') u_\rho.$$

Take a cone  $\sigma$  in  $\Sigma$  such that  $n_0 \in -\text{RelInt } \sigma$ . Let  $\rho_1, \dots, \rho_a$  be the rays of  $\sigma$ . We may find rays  $\rho_{a+1}, \dots, \rho_n \in \Sigma(1)$  such that  $u_{\rho_1}, \dots, u_{\rho_n}$  form a basis of  $\mathbb{R}^n$ .

A subset of  $P \cap \{\nabla G_0 \subseteq C\}$  is given by those  $m' \in P$  such that for all  $\rho \in \Sigma(1)$  different from  $\rho_1, \dots, \rho_a$ , the function  $r_\rho(m')$  is uniformly bounded, while  $m'$  is close enough to the faces corresponding to the rays  $\rho_1, \dots, \rho_n$  and  $\sum_{i=1}^a r_{\rho_i}(m') u_{\rho_i} \in C'$ . Replace the domain of integration in (5.8) to this region and the variable  $m'$  to  $r_{\rho_1}(m'), \dots, r_{\rho_n}(m')$ , we find that the Jacobian is a polynomial in  $r_{\rho_1}, \dots, r_{\rho_a}$ , while the integrand diverges exponentially. We conclude.  $\square$

cor:DXmaintoric

**Corollary 5.3.1** *Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ , then*

$$\lim_{k \rightarrow \infty} \frac{n!}{k^n} h^0(X, L^k \otimes I(k\varphi)) = n! \text{vol } \Delta(\omega, \varphi).$$

We interpret the full mass potentials studied in Section 3.1.3 in the toric setting. We have the following straightforward observation in the full mass case.

**Proposition 5.3.2** *Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ . Then the following are equivalent:*

- (1)  $\varphi \in \mathcal{E}^\infty(X, \omega)$ ;
- (2)  $F_\varphi \sim F_0$ ;
- (3)  $G_\varphi \sim G_0$ .

**Proposition 5.3.3** *Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ . Then the following are equivalent:*

- (1)  $\varphi \in \mathcal{E}(X, \omega)$ ;
- (2)  $F_\varphi \in \mathcal{E}(N_{\mathbb{R}}, P)$ ;
- (3)  $\overline{\text{Dom } G_\varphi} = P$ .

**Proof** (1)  $\iff$  (3). By Proposition 5.2.5

$$\int_X \omega_\varphi^n = \int_{T(\mathbb{C})} (\omega|_{T(\mathbb{C})} + \text{dd}^c \varphi|_{T(\mathbb{C})})^n = n! \text{vol } \overline{\text{Dom } G_\varphi}, \quad \int_X \omega^n = n! \text{vol } P.$$

Therefore, (1) and (3) are equivalent.

(2)  $\iff$  (3). This follows from Proposition A.2.1.  $\square$

**Proposition 5.3.4** *Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ , then*

$$E_\omega(\varphi) = n! \int_P (G_0 - G_\varphi) \, d \text{vol}.$$

**Proof** It suffices to consider the case where  $\varphi$  is bounded. In this case, one could apply [BB13, Proposition 2.9].  $\square$

**Corollary 5.3.2** *Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ . Then the following are equivalent:*

- (1)  $\varphi \in \mathcal{E}^1(X, \omega)$ ;
- (2)  $F_\varphi \in \mathcal{E}^1(N_{\mathbb{R}}, P)$ ;
- (3)  $G_\varphi \in L^1(P)$ .

**Definition 5.3.2** We define

$$\begin{aligned}\mathcal{E}_{\text{tor}}^\infty(X, \omega) &= \mathcal{E}^\infty(X, \omega) \cap \text{PSH}_{\text{tor}}(X, \omega), \\ \mathcal{E}_{\text{tor}}^1(X, \omega) &= \mathcal{E}^1(X, \omega) \cap \text{PSH}_{\text{tor}}(X, \omega), \\ \mathcal{E}_{\text{tor}}(X, \omega) &= \mathcal{E}(X, \omega) \cap \text{PSH}_{\text{tor}}(X, \omega).\end{aligned}$$

cor:toricd1

**Corollary 5.3.3** *Let  $\varphi, \psi \in \mathcal{E}_{\text{tor}}^1(X, \omega)$ , then*

$$d_1(\varphi, \psi) = -n! \int_P (G_\varphi + G_\psi - 2G_{\varphi \vee \psi}) \, d\text{vol}.$$

prop:toricgeodseg

**Proposition 5.3.5** *Let  $\varphi_0, \varphi_1 \in \mathcal{E}_{\text{tor}}^1(X, \omega)$ . The geodesic  $(\varphi_t)_{t \in (0,1)}$  from  $\varphi_0$  to  $\varphi_1$  satisfies the following: for each  $t \in (0, 1)$ ,  $\varphi_t \in \mathcal{E}_{\text{tor}}^1(X, \omega)$  and*

$$G_{\varphi_t} = (1-t)G_{\varphi_0} + tG_{\varphi_1}.$$

This will be proved more generally in [Corollary 12.3.2](#).

**Definition 5.3.3** We define

$$\mathcal{R}_{\text{tor}}^1(X, \omega) := \{\ell \in \mathcal{R}^1(X, \omega) : \ell_t \in \text{PSH}_{\text{tor}}(X, \omega) \text{ for all } t \geq 0\}.$$

**Corollary 5.3.4** *Let  $\ell \in \mathcal{R}_{\text{tor}}^1(X, \omega)$ . Then there is an integrable convex function  $G' \in \text{Conv}(N_{\mathbb{R}})$  with  $\overline{\text{Dom } G'} = P$  such that*

$$G_{\ell_t} = G_0 + tG'$$

for all  $t \geq 0$ .

We could also make [Example 4.2.1](#) concrete.

**Proposition 5.3.6** *Suppose that  $\varphi \in \text{PSH}_{\text{tor}}(X, \omega)$ . Then the ray  $\ell^\varphi$  defined in [Example 4.2.1](#) satisfies:*

$$G_{\ell_t} = G_0 + t f_\ell, \quad f_\ell(x) = \min_{\substack{\lambda \in [0,1] \\ x_1 \in P, x_0 \in \Delta(\omega, \varphi) \\ \lambda x_1 + (1-\lambda)x_0 = x}} \lambda$$

for any  $t \geq 0$  and  $x \in M_{\mathbb{R}}$ .

**Proof** Recall that for each  $C > 0$ , we defined  $(\ell_t^{\varphi, C})_t$  as the geodesic from 0 to  $-C \vee \varphi$ . By [Proposition 5.2.2](#), [Proposition 5.2.4](#), we have  $G_{-C \vee \varphi} = (G_0 + C) \wedge G_\varphi$ . So by [Proposition 5.3.5](#), we have

$$G_{\ell_t^{\varphi, C}} = \frac{t}{C} ((G_0 + C) \wedge G_\varphi) + \frac{C-t}{C} G_0$$

for each  $t \in [0, C]$ .

Recall that for all  $t \geq 0$ ,

$$\ell_t = \sup_{C \geq t}^* \ell_t^{\varphi, C}.$$

It follows from [Proposition 5.2.4](#) that

$$G_{\ell_t} = \text{cl} \inf_{C \geq t} \frac{t}{C} ((G_0 + C) \wedge G_\varphi) + \frac{C-t}{C} G_0.$$

Since the infimum is clearly linear, the closure operation is not needed and  $G_{\ell_t}$  is linear in  $t$ . So it suffices to compute the slope  $f$ :

$$f_\ell := \inf_{C > 0} \frac{1}{C} ((G_0 + C) \wedge G_\varphi) - \frac{1}{C} G_0.$$

We compute this limit using [Proposition A.1.2](#): for  $x \in M_{\mathbb{R}}$ , we compute the slope as follows

$$\begin{aligned} f_\ell(x) &= \inf_{C > 0} \inf_{\substack{\lambda \in (0,1) \\ x_1, x_0 \in M_{\mathbb{R}} \\ \lambda x_1 + (1-\lambda)x_0 = x}} \lambda \left( \frac{G_0(x_1)}{C} + 1 \right) + \frac{1-\lambda}{C} G_\varphi(x_0) - \frac{G_0(x)}{C} \\ &= \inf_{\substack{\lambda \in (0,1) \\ x_1, x_0 \in M_{\mathbb{R}} \\ \lambda x_1 + (1-\lambda)x_0 = x}} \inf_{C > 0} \lambda \left( \frac{G_0(x_1)}{C} + 1 \right) + \frac{1-\lambda}{C} G_\varphi(x_0) - \frac{G_0(x)}{C} \\ &= \min_{\substack{\lambda \in [0,1] \\ x_1 \in P, x_0 \in \Delta(\omega, \varphi) \\ \lambda x_1 + (1-\lambda)x_0 = x}} \lambda. \end{aligned}$$



**Part II**  
**The theory of  $\mathcal{I}$ -good singularities**

This part is the technical core of the whole book. We will develop the theory of  $\mathcal{I}$ -good singularities.

We first develop some general techniques to compare the singularities in [Chapter 6](#): The  $P$ -partial order, the  $\mathcal{I}$ -partial order and the  $d_S$ -pseudometric.

The  $P$ -partial order seems to be new. Some basic properties of the  $d_S$ -pseudometric have never appeared in the literature either.

Then in [Chapter 7](#), we introduce the notion of  $\mathcal{I}$ -good singularities and characterize  $\mathcal{I}$ -good singularities in different ways. In the algebraic situation, we establish the asymptotic Riemann–Roch formula.

In [Chapter 8](#), we will develop two key techniques in the inductive study of singularities: The trace operator and the analytic Bertini theorem. Roughly speaking, the latter tells us the behaviour of a quasi-plurisubharmonic function along a general divisor, while the former handles the case of special divisors. We will establish a relative version of the asymptotic Riemann–Roch formula in the algebraic situation.

In [Chapter 9](#), we develop the theory of test curves. These are curves of model potentials. The key technique is the Ross–Witt Nyström correspondence, which relates test curves with geodesic rays. The complete proof of the most general form of this correspondence has never appeared in the literature, so we will give the full details.

In [Chapter 10](#), we develop the theory of partial Okounkov bodies, in both algebraic and transcendental setting. The partial Okounkov bodies can be regarded as non-toric extensions of the Newton bodies. It turns out that even in the toric setting, our techniques give non-trivial new results.

In [Chapter 11](#), we develop the theory of  $\mathbf{b}$ -divisors in the algebraic setting. We formulate the general form of the Chern–Weil formula in terms of  $\mathbf{b}$ -divisors. We also relate the theory of partial Okounkov bodies to  $\mathbf{b}$ -divisors.



## Chapter 6

### Comparison of singularities

chap:comp

In this chapter, we study several ways of comparing the singularities of quasi-plurisubharmonic functions. In [Section 6.1](#), we will introduce the  $P$  and  $\mathcal{I}$ -partial orders, closely related to the  $P$  and  $\mathcal{I}$ -equivalence relations introduced in [Chapter 3](#).

In [Section 6.2](#), we introduce and study the  $d_S$ -pseudometric characterizing the differences between singularities. We will prove that a number of continuity results with respect to  $d_S$ .

#### 6.1 The $P$ and $\mathcal{I}$ -partial orders

sec:PIpartialorder

Let  $X$  be a connected compact Kähler manifold of dimension  $n$ .

Recall that we have defined a (non-strict) partial order on  $\text{QPSH}(X)$  in [Definition 1.5.2](#) to compare the singularity types of quasi-plurisubharmonic functions. The problem with this partial order is that it is too fine. In general, for our interest, it is helpful to consider rougher relations.

##### 6.1.1 The definitions of the partial orders

Recall that the  $P$ -envelope is defined in [Definition 3.1.2](#).

def:Pmoresing

**Definition 6.1.1** Let  $\varphi, \psi \in \text{QPSH}(X)$ , we say  $\varphi$  is  *$P$ -more singular than  $\psi$*  and write  $\varphi \leq_P \psi$  if for some closed smooth real  $(1, 1)$ -form  $\theta$  on  $X$  such that  $\varphi, \psi \in \text{PSH}(X, \theta)_{>0}$ , we have

$$P_\theta[\varphi] \leq P_\theta[\psi].$$

Suppose that  $\varphi \leq_P \psi$  and  $\psi \leq_P \varphi$ , we shall write  $\varphi \sim_P \psi$  and say  $\varphi$  and  $\psi$  have the same  *$P$ -singularity type*.

This definition is independent of the choice of  $\theta$ :

lma:Pproj\_insens\_omega

**Lemma 6.1.1** *Let  $\varphi, \psi \in \text{PSH}(X, \theta)_{>0}$ . For any Kähler form  $\omega$  on  $X$ , the following are equivalent:*

- (1)  $P_\theta[\varphi] \leq P_\theta[\psi]$ ;
- (2)  $P_{\theta+\omega}[\varphi] \leq P_{\theta+\omega}[\psi]$ .

In particular,  $\leq_P$  defines a non-strict partial order on  $\text{QPSH}(X)$ .

**Proof** (1) implies (2). Observe that

$$P_\theta[\varphi] \leq P_{\theta+\omega}[\varphi], \quad \varphi \leq P_\theta[\varphi].$$

It follows from [Theorem 3.1.1](#) that

$$P_{\theta+\omega}[\varphi] = P_{\theta+\omega}[P_\theta[\varphi]]. \quad (6.1) \quad \{\text{eq:doubleP}\}$$

A similar formula holds for  $\psi$ . So we see that (2) holds.

(2) implies (1). By [\(6.1\)](#), we may assume that  $\varphi$  and  $\psi$  are both model potentials in  $\text{PSH}(X, \theta)_{>0}$ .

Observe that  $\varphi \vee \psi \leq P_{\theta+\omega}[\psi]$ . It follows that  $P_{\theta+\omega}[\varphi \vee \psi] \leq P_{\theta+\omega}[\psi]$ . The reverse inequality is trivial, so

$$P_{\theta+\omega}[\varphi \vee \psi] = P_{\theta+\omega}[\psi].$$

From the direction we have proved, for any  $C \geq 1$ ,

$$P_{\theta+C\omega}[\varphi \vee \psi] = P_{\theta+C\omega}[\psi].$$

So by [Proposition 3.1.2](#),

$$\int_X (\theta + C\omega + \text{dd}^c(\varphi \vee \psi))^n = \int_X (\theta + C\omega + \text{dd}^c\psi)^n.$$

Since both sides are polynomials in  $C$ , the equality extends to  $C = 0$ , namely,

$$\int_X \theta_{\varphi \vee \psi}^n = \int_X \theta_\psi^n.$$

In particular,  $\varphi \vee \psi \leq P_\theta[\psi] = \psi$  by [\(3.4\)](#). So (1) follows.  $\square$

As a first example of  $P$ -equivalence, we have:

ex:Pequiv

**Example 6.1.1** Let  $\theta$  be a closed smooth real  $(1, 1)$ -form on  $X$  and  $\varphi \in \text{PSH}(X, \theta)_{>0}$ , then

$$\varphi \sim_P P_\theta[\varphi].$$

This follows immediately from [Theorem 3.1.1](#).

prop:Pequivchar2

**Proposition 6.1.1** *Let  $\varphi, \psi \in \text{PSH}(X, \theta)$  and  $\varphi \leq \psi$ . Then the following are equivalent:*

- (1)  $\varphi \sim_P \psi$ ;

(2) for each  $j = 0, \dots, n$ , we have

$$\int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} = \int_X \theta_\psi^j \wedge \theta_{V_\theta}^{n-j}. \quad (6.2) \quad \{\text{eq:mixedmassequal}\}$$

Assume furthermore that  $\varphi, \psi \in \text{PSH}(X, \theta)_{>0}$ , then these conditions are equivalent to the following:

(3) We have

$$\int_X \theta_\varphi^n = \int_X \theta_\psi^n.$$

Recall that  $V_\theta$  is introduced in (2.9).

**Proof** We first prove the equivalence between (1) and (3) when  $\varphi, \psi \in \text{PSH}(X, \theta)_{>0}$ .

(1)  $\implies$  (3). Assume that  $\varphi \sim_P \psi$ . By Definition 6.1.1, we have

$$P_\theta[\varphi] = P_\theta[\psi].$$

So (3) follows from Proposition 3.1.2.

(3)  $\implies$  (1). It follows from Theorem 3.1.1 that  $P_\theta[\varphi] = P_\theta[\psi]$ , so (1) follows.

Let us come back to the general case.

(1)  $\implies$  (2). Fix  $j \in \{0, \dots, n\}$ , we argue (6.2).

Take a Kähler form  $\omega$  on  $X$ . By Definition 6.1.1, for each  $\epsilon > 0$ , we have

$$P_{\theta+\epsilon\omega}[\varphi] = P_{\theta+\epsilon\omega}[\psi].$$

It follows from Proposition 3.1.2 that

$$\begin{aligned} \int_X (\theta + \epsilon\omega + \text{dd}^c \psi)^j \wedge \theta_{V_\theta}^{n-j} &= \int_X (\theta + \epsilon\omega + \text{dd}^c P_{\theta+\epsilon\omega}[\psi])^j \wedge \theta_{V_\theta}^{n-j} \\ &= \int_X (\theta + \epsilon\omega + \text{dd}^c P_{\theta+\epsilon\omega}[\varphi])^j \wedge \theta_{V_\theta}^{n-j} \\ &= \int_X (\theta + \epsilon\omega + \text{dd}^c \varphi)^j \wedge \theta_{V_\theta}^{n-j}. \end{aligned}$$

Since the two extremes are both polynomials in  $\epsilon$ , we conclude that the same holds when  $\epsilon = 0$ , that is, (6.2) holds.

(2)  $\implies$  (1). Assume (6.2) holds for all  $j = 0, \dots, n$ . For each  $t \in (0, 1)$ , we have

$$\int_X \theta_{t\varphi+(1-t)V_\theta}^n = \int_X \theta_{t\psi+(1-t)V_\theta}^n$$

by the binomial expansion. By the implication (3)  $\implies$  (1), we have

$$t\varphi + (1-t)V_\theta \sim_P t\psi + (1-t)V_\theta$$

for each  $t \in (0, 1)$ .

Fix a Kähler form  $\omega$  on  $X$ . From the implication (1)  $\implies$  (3), we have

$$\int_X (\theta + \omega)_{t\varphi + (1-t)V_\theta}^n = \int_X (\theta + \omega)_{t\psi + (1-t)V_\theta}^n.$$

Since both sides are polynomials in  $t$ , the same holds when  $t = 1$ . From the implication (3)  $\implies$  (1) again, we have  $\varphi \sim_P \psi$ .  $\square$

prop:Iequivchar2

**Proposition 6.1.2** *Given  $\varphi, \psi \in \text{QPSH}(X)$ , the following are equivalent:*

(1) *For any  $k \in \mathbb{Z}_{>0}$ , we have*

$$I(k\varphi) \subseteq I(k\psi);$$

(2) *for any  $\lambda \in \mathbb{R}_{>0}$ , we have*

$$I(\lambda\varphi) \subseteq I(\lambda\psi);$$

(3) *for any modification  $\pi: Y \rightarrow X$  and any  $y \in Y$ , we have*

$$v(\pi^*\varphi, y) \geq v(\pi^*\psi, y);$$

(4) *for any proper bimeromorphic morphism  $\pi: Y \rightarrow X$  from a Kähler manifold and any  $y \in Y$ , we have*

$$v(\pi^*\varphi, y) \geq v(\pi^*\psi, y);$$

(5) *for any prime divisor  $E$  over  $X$ , we have*

$$v(\varphi, E) \geq v(\psi, E).$$

**Proof** The proof is almost identical to that of [Proposition 3.2.1](#), we omit the details.  $\square$

**Definition 6.1.2** Let  $\varphi, \psi \in \text{QPSH}(X)$ , we say  $\varphi$  is  *$I$ -more singular than  $\psi$*  and write  $\varphi \leq_I \psi$  if the equivalent conditions in [Proposition 6.1.2](#) are satisfied.

It is clear that  $\leq_I$  is a non-strict partial order on  $\text{QPSH}(X)$ .

Note that  $\varphi \leq_I \psi$  and  $\psi \leq_I \varphi$  both hold if and only if  $\varphi \sim_I \psi$  in the sense of [Definition 3.2.1](#).

lma:reform\_preceqP

**Lemma 6.1.2** *Let  $\varphi, \psi \in \text{QPSH}(X)$ . Then the following are equivalent:*

(1)  $\varphi \leq_P \psi$  (resp.  $\varphi \leq_I \psi$ );

(2)  $\varphi \vee \psi \sim_P \psi$  (resp.  $\varphi \vee \psi \sim_I \psi$ ).

**Proof** Take a closed real smooth  $(1, 1)$ -form  $\theta$  on  $X$  such that  $\varphi, \psi \in \text{PSH}(X, \theta)_{>0}$ . We only prove the  $P$  case, the  $I$  case is similar.

(2)  $\implies$  (1). By (2) and [Example 6.1.1](#),  $P_\theta[\varphi \vee \psi] = P_\theta[\psi] \sim_P \psi$ . But  $\varphi \leq P_\theta[\varphi \vee \psi]$ , so (1) follows.

(1)  $\implies$  (2). We may assume that  $\varphi, \psi$  are both model in  $\text{PSH}(X, \theta)_{>0}$  as

$$P_\theta[\varphi \vee \psi] = P_\theta[P_\theta[\varphi] \vee P_\theta[\psi]].$$

Then  $\varphi \leq \psi$  and (2) follows.  $\square$

cor:PimpliesI

**Corollary 6.1.1** Let  $\varphi, \psi \in \text{QPSH}(X)$ . Assume that  $\varphi \leq_P \psi$ , then  $\varphi \leq_I \psi$ .

*Proof* This follows from [Lemma 6.1.2](#) and [Proposition 3.2.8](#).  $\square$

cor:Pvarphidef3

**Corollary 6.1.2** Assume that  $\varphi \in \text{PSH}(X, \theta)_{>0}$ , then

$$\begin{aligned} P_\theta[\varphi] &= \sup \{ \psi \in \text{PSH}(X, \theta) : \psi \leq 0, \psi \sim_P \varphi \} \\ &= \sup \{ \psi \in \text{PSH}(X, \theta) : \psi \leq 0, \psi \leq_P \varphi \}. \end{aligned}$$

*Proof* Note that  $\psi \sim_P \varphi$  implies that  $\psi \in \text{PSH}(X, \theta)_{>0}$  by [Proposition 6.1.4](#). We observe that

$$\begin{aligned} &\sup \{ \psi \in \text{PSH}(X, \theta) : \psi \leq 0, \psi \sim_P \varphi \} \\ &= \sup \{ \psi \in \text{PSH}(X, \theta) : \psi \leq 0, \varphi \leq \psi, \psi \sim_P \varphi \} \end{aligned}$$

by [Lemma 6.1.2](#). So the first equality is a direct consequence of [Proposition 6.1.1](#) and [Theorem 3.1.1](#).

Next we prove the second equality. We only need to show that for any  $\psi \in \text{PSH}(X, \theta)$  with  $\psi \leq 0$  and  $\psi \leq_P \varphi$ , we have  $\psi \leq P_\theta[\varphi]$ .

By [Lemma 6.1.2](#) and [Example 6.1.1](#), we know that  $P_\theta[\varphi] \vee \psi \sim_P \varphi$  and  $P_\theta[\varphi] \vee \psi \leq 0$ . It follows from the first equality that  $\psi \leq P_\theta[\varphi]$ .  $\square$

Similarly, we have

cor:Ienvelopedef2

**Corollary 6.1.3** Assume that  $\varphi \in \text{PSH}(X, \theta)$ , then

$$P_\theta[\varphi]_I = \sup \{ \psi \in \text{PSH}(X, \theta) : \psi \leq 0, \psi \leq_I \varphi \}.$$

prop:Icomparandenvelope

**Proposition 6.1.3** Suppose that  $\varphi, \psi \in \text{QPSH}(X)$  and  $\theta$  is a closed real smooth  $(1, 1)$ -form on  $X$  such that  $\varphi, \psi \in \text{PSH}(X, \theta)$ . Then the following are equivalent:

- (1)  $\varphi \leq_I \psi$ ;
- (2)  $P_\theta[\varphi]_I \leq P_\theta[\psi]_I$ .

*Proof* (1)  $\implies$  (2). This follows immediately from [Corollary 6.1.3](#).

(2)  $\implies$  (1). This follows from [Proposition 3.2.6](#).  $\square$

## 6.1.2 Properties of the partial orders

Now we state a more natural version of the monotonicity theorem [Theorem 2.3.2](#).

prop:mono2

**Proposition 6.1.4** Let  $\theta_1, \dots, \theta_n$  be closed real smooth  $(1, 1)$ -forms on  $X$ . Let  $\varphi_i, \psi_i \in \text{PSH}(X, \theta_i)$  for  $i = 1, \dots, n$ . Assume that  $\varphi_i \leq_P \psi_i$  for each  $i$ . Then

$$\int_X \theta_{\varphi_1} \wedge \dots \wedge \theta_{\varphi_n} \leq \int_X \theta_{\psi_1} \wedge \dots \wedge \theta_{\psi_n}.$$

**Proof** Fix a Kähler form  $\omega$  on  $X$ . For each  $i = 1, \dots, n$ , since  $\varphi_i \leq_P \psi_i$ , we have

$$P_{\theta+\epsilon\omega}[\varphi_i] \leq P_{\theta+\epsilon\omega}[\psi_i]$$

for all  $\epsilon > 0$ . Therefore, by [Proposition 3.1.2](#) and [Theorem 2.3.2](#), we have

$$\int_X (\theta + \epsilon\omega)_{\varphi_1} \wedge \dots \wedge (\theta + \epsilon\omega)_{\varphi_n} \leq \int_X (\theta + \epsilon\omega)_{\psi_1} \wedge \dots \wedge (\theta + \epsilon\omega)_{\psi_n}.$$

Since both sides are polynomials in  $\epsilon$ , we find that the same holds at  $\epsilon = 0$ , which is the desired inequality.  $\square$

prop:Partialsum

**Proposition 6.1.5** Let  $\varphi, \psi, \varphi', \psi' \in \text{QPSH}(X)$ . Assume that

$$\varphi \leq_P \psi, \quad \varphi' \leq_P \psi'.$$

Then

$$\varphi + \varphi' \leq_P \psi + \psi'.$$

The same holds with  $\leq_I$  in place of  $\leq_P$ .

**Proof** Take a Kähler form  $\omega$  on  $X$  such that  $\varphi, \psi, \varphi', \psi' \in \text{PSH}(X, \omega)_{>0}$ . The statement for  $\leq_I$  is a simple consequence of [Proposition 1.4.2](#). We only need to handle the case of  $\leq_P$ .

**Step 1.** We first show that

$$P_\omega[\varphi] + P_\omega[\varphi'] \sim_P \varphi + \varphi'.$$

In fact, we clearly have

$$P_\omega[\varphi] + P_\omega[\varphi'] \geq \varphi + \varphi'.$$

So by [Proposition 6.1.1](#), it suffices to show that they have the same volume. We compute

$$\begin{aligned} & \int_X (2\omega + \text{dd}^c P_\omega[\varphi] + \text{dd}^c P_\omega[\varphi'])^n \\ &= \sum_{j=0}^n \binom{n}{j} \int_X (\omega + \text{dd}^c P_\omega[\varphi])^j \wedge (\omega + \text{dd}^c P_\omega[\varphi'])^{n-j} \\ &= \sum_{j=0}^n \binom{n}{j} \int_X \omega_\varphi^j \wedge \omega_{\varphi'}^{n-j} \\ &= \int_X (2\omega + \varphi + \varphi')^n, \end{aligned}$$

where we applied [Proposition 3.1.2](#) on the third line.

**Step 2.** By Step 1, we may assume that  $\varphi, \psi, \varphi', \psi'$  are all model potentials. So  $\varphi \leq \psi$  and  $\varphi' \leq \psi'$ . Our assertion follows.  $\square$

prop:Partialsup

**Proposition 6.1.6** *Let  $(\varphi_i)_{i \in I}$ ,  $(\psi_i)_{i \in I}$  be uniformly bounded from above non-empty families in  $\text{QPSH}(X)$ . Assume that there exists a closed smooth real  $(1, 1)$ -form  $\theta$  such that  $\varphi_i, \psi_i \in \text{PSH}(X, \theta)$  and  $\varphi_i \leq_P \psi_i$  for all  $i \in I$ . Then*

$$\sup_{i \in I}^* \varphi_i \leq_P \sup_{i \in I}^* \psi_i.$$

*The same holds with  $\leq_I$  in place of  $\leq_P$ .*

**Proof** By increasing  $\theta$ , we may assume that  $\varphi_i, \psi_i \in \text{PSH}(X, \theta)_{>0}$  for all  $i \in I$ . The statement for  $\leq_I$  is a simple consequence of [Corollary 1.4.1](#), we only have to consider the statement for  $\leq_P$ .

**Step 1.** We first handle the case where  $I$  is a directed set and  $(\varphi_i)_{i \in I}$  and  $(\psi_i)_{i \in I}$  are increasing nets.

In this case, our assertion follows simply from [Proposition 3.1.9](#).

**Step 2.** We handle the case where  $I$  is finite. We may assume that  $I = \{0, 1\}$ . It suffices to show that

$$P_\theta[\varphi_0] \vee P_\theta[\varphi_1] \sim_P \varphi_0 \vee \varphi_1.$$

For this purpose, it suffices to prove the following:

$$P_\theta[\varphi_0] \vee \varphi_1 \sim_P \varphi_0 \vee \varphi_1.$$

The  $\geq_P$  direction is obvious. So thanks to [Proposition 6.1.1](#), it suffices to argue that they have the same mass. We may assume that  $\varphi_0 \leq 0$ . Thanks to [Lemma 2.3.1](#), for each  $\epsilon \in (0, 1)$ , we can find  $\eta_\epsilon \in \text{PSH}(X, \theta)_{>0}$  such that

$$(1 - \epsilon)P_\theta[\varphi_0] + \epsilon\eta_\epsilon \leq \varphi_0, \quad \eta_\epsilon \leq \varphi_0 \leq P_\theta[\varphi_0].$$

In particular,

$$(1 - \epsilon)(P_\theta[\varphi_0] \vee \varphi_1) + \epsilon\eta_\epsilon \leq \varphi_0 \vee \varphi_1.$$

It follows from [Theorem 2.3.2](#) that

$$(1 - \epsilon)^n \int_X \theta_{P_\theta[\varphi_0] \vee \varphi_1}^n \leq \int_X \theta_{\varphi_0 \vee \varphi_1}^n.$$

Letting  $\epsilon \rightarrow 0+$  and using [Theorem 2.3.2](#) again, we conclude that

$$\int_X \theta_{P_\theta[\varphi_0] \vee \varphi_1}^n = \int_X \theta_{\varphi_0 \vee \varphi_1}^n.$$

Our assertion is proved.

**Step 3.** The general case can be reduced to the two cases handled in Step 1 and Step 2. More precisely, by [Proposition 1.2.2](#), we could find a countable subset  $J \subseteq I$  such that

$$\sup_{j \in J}^* \varphi_j = \sup_{i \in I}^* \varphi_i, \quad \sup_{j \in J}^* \psi_j = \sup_{i \in I}^* \psi_i.$$

We may replace  $I$  by  $J$  and assume that  $I$  is countable. We may assume that  $I$  is infinite, as otherwise, we could apply Step 2 directly. So let us assume that  $J = \mathbb{Z}_{>0}$ . In this case, by Step 2 again, we may assume that both  $(\varphi_i)_i$  and  $(\psi_i)_i$  are increasing, which is the situation of Step 1.

prop:rooftopprePequiv

**Proposition 6.1.7** *Let  $\varphi, \psi, \varphi', \psi' \in \text{PSH}(X, \theta)_{>0}$  for some closed smooth real  $(1, 1)$ -form on  $X$ . Assume that*

$$\varphi \sim_P \varphi', \quad \psi \sim_P \psi'$$

and

$$\varphi' \wedge \psi' \in \text{PSH}(X, \theta)_{>0}.$$

Then

$$\varphi \wedge \psi \in \text{PSH}(X, \theta)_{>0}$$

and

$$\varphi \wedge \psi \sim_P \varphi' \wedge \psi'.$$

**Proof** Without loss of generality, we may assume that  $\psi = \psi'$ . Replacing  $\varphi'$  by  $P_\theta[\varphi']$ , we may also assume that  $\varphi \leq \varphi'$ .

Using [Corollary 2.3.2](#), for each  $\epsilon \in (0, 1)$ , we can find  $\eta \in \text{PSH}(X, \theta)$  such that

$$\int_X \theta_\eta^n = \int_X \theta_\varphi^n, \quad \epsilon\eta + (1 - \epsilon)\varphi' \leq \varphi, \quad \eta \leq \varphi'.$$

Since

$$\int_X \theta_\eta^n + \int_X \theta_{\varphi' \wedge \psi}^n > \int_X \theta_{\varphi'}^n,$$

by [Proposition 3.1.3](#), we find  $\eta \wedge \psi \in \text{PSH}(X, \theta)$ . Now observe that

$$\epsilon(\eta \wedge \psi) + (1 - \epsilon)(\varphi' \wedge \psi) \leq \varphi \wedge \psi.$$

By [Theorem 2.3.2](#), we find that

$$(1 - \epsilon)^n \int_X \theta_{\varphi' \wedge \psi} \leq \int_X \theta_{\varphi \wedge \psi}^n.$$

Letting  $\epsilon \rightarrow 0+$  and applying [Theorem 2.3.2](#), we find that

$$\int_X \theta_{\varphi' \wedge \psi} = \int_X \theta_{\varphi \wedge \psi}.$$

We conclude by [Proposition 6.1.1](#).



## 6.2 The $d_S$ -pseudometric

sec:dsdef

Let  $X$  be a connected compact Kähler manifold of dimension  $n$  and  $\theta$  be a closed real smooth  $(1, 1)$ -form on  $X$  representing a big cohomology class. The goal of this section is to study a pseudometric on the space  $\text{PSH}(X, \theta)$ .

### 6.2.1 The definition of the $d_S$ -pseudometric

Recall that for any  $\varphi \in \text{PSH}(X, \theta)$ , the geodesic ray  $\ell^\varphi \in \mathcal{R}^1(X, \theta)$  is defined in [Example 4.2.1](#).

def:dS

**Definition 6.2.1** For  $\varphi, \psi \in \text{PSH}(X, \theta)$ , we define

$$d_S(\varphi, \psi) := d_1(\ell^\varphi, \ell^\psi).$$

When we want to be more specific, we write  $d_{S, \theta}$  instead of  $d_S$ .

The  $d_1$  distance of geodesic rays is defined in [Definition 4.2.6](#).

**Proposition 6.2.1** *The function  $d_S$  defined in [Definition 6.2.1](#) is a pseudometric on  $\text{PSH}(X, \theta)$ .*

*Proof* This follows immediately from [Theorem 4.2.3](#). □

When studying a pseudometric, the first thing is to understand when the distance between two elements vanishes.

We first prove a preparation:

lma:dSalmostriang

**Lemma 6.2.1** *Let  $\varphi, \psi \in \text{PSH}(X, \theta)$ . Then*

$$d_S(\varphi, \psi) \leq d_S(\varphi, \varphi \vee \psi) + d_S(\psi, \varphi \vee \psi) \leq C_n d_S(\varphi, \psi),$$

where  $C_n = 3(n+1)2^{n+2}$ .

*Proof* Observe that

$$\ell^\varphi \vee \ell^\psi = \ell^{\varphi \vee \psi}.$$

(6.3)

{eq:elllorsingtype}

In fact, it is clear that

$$\ell^\varphi \leq \ell^{\varphi \vee \psi}, \quad \ell^\psi \leq \ell^{\varphi \vee \psi},$$

so the  $\leq$  direction in [\(6.3\)](#) holds.

Conversely, if  $\ell' \in \mathcal{R}^1(X, \theta)$  and  $\ell' \geq \ell^\varphi \vee \ell^\psi$ , then for each  $t \geq 0$ ,

$$\ell'_t \geq ((V_\theta - t) \vee \varphi) \vee ((V_\theta - t) \vee \psi) = (V_\theta - t) \vee (\varphi \vee \psi).$$

It follows that  $\ell' \geq \ell^{\varphi \vee \psi}$ .

So our assertion follows from [Lemma 4.2.1](#). □

prop:ds@char

**Proposition 6.2.2** Let  $\varphi, \psi \in \text{PSH}(X, \theta)$ . Then the following are equivalent:

- (1)  $\varphi \sim_P \psi$ ;
- (2)  $d_S(\varphi, \psi) = 0$ .

In particular,  $d_S(\varphi, P_\theta[\varphi]) = 0$  for all  $\varphi \in \text{PSH}(X, \theta)_{>0}$ .

**Proof** By Lemma 6.1.2, we have  $\varphi \sim_P \psi$  if and only if  $\varphi \sim_P \varphi \vee \psi$  and  $\psi \sim_P \varphi \vee \psi$ . By Lemma 6.2.1,  $d_S(\varphi, \psi) = 0$  if and only if  $d_S(\varphi, \varphi \vee \psi) = 0$  and  $d_S(\psi, \varphi \vee \psi) = 0$ . So it suffices to prove the assertion when  $\varphi \leq \psi$ . Assuming this, by Proposition 4.2.6 we have that 2 holds if and only if

$$\mathbf{E}(\ell^\varphi) = \mathbf{E}(\ell^\psi),$$

But using (4.14), this holds if and only if

$$\sum_{j=0}^n \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} = \sum_{j=0}^n \int_X \theta_\psi^j \wedge \theta_{V_\theta}^{n-j}.$$

But by Theorem 2.3.2, this holds if and only if for all  $j = 0, \dots, n$ ,

$$\int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} = \int_X \theta_\psi^j \wedge \theta_{V_\theta}^{n-j},$$

which is equivalent to 1 by Proposition 6.1.1.  $\square$

lma:varphileqpsi\_metric

**Lemma 6.2.2** Suppose that  $\varphi, \psi \in \text{PSH}(X, \theta)$  and  $\varphi \leq_P \psi$ , then

$$d_S(\varphi, \psi) = \frac{1}{n+1} \sum_{j=0}^n \left( \int_X \theta_\psi^j \wedge \theta_{V_\theta}^{n-j} - \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} \right).$$

**Proof** This follows trivially from (4.14).  $\square$

cor:dsthreeterm

**Corollary 6.2.1** Suppose that  $\varphi, \psi, \eta \in \text{PSH}(X, \theta)$  and  $\varphi \leq_P \psi \leq_P \eta$ . Then

$$d_S(\varphi, \eta) \geq d_S(\varphi, \psi), \quad d_S(\varphi, \eta) \geq d_S(\psi, \eta).$$

**Proof** This is an immediate consequence of Lemma 6.2.2 and Proposition 6.1.4.  $\square$

cor:dsmetricdoubleineq

**Corollary 6.2.2** For any  $\varphi, \psi \in \text{PSH}(X, \theta)$ , we have

$$\begin{aligned} d_S(\varphi, \psi) &\leq \sum_{j=0}^n \left( 2 \int_X \theta_{\varphi \vee \psi}^j \wedge \theta_{V_\theta}^{n-j} - \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} - \int_X \theta_\psi^j \wedge \theta_{V_\theta}^{n-j} \right) \\ &\leq C_n d_S(\varphi, \psi), \end{aligned} \quad (6.4) \quad \{\text{eq:ds\_biineq}\}$$

where  $C_n = 3(n+1)2^{n+2}$ .

In particular, if  $(\varphi_i)_{i \in I}$  is a net in  $\text{PSH}(X, \theta)$  with  $d_S$ -limit  $\varphi$ , then for each  $j = 0, \dots, n$ ,

$$\lim_{i \in I} \int_X \theta_{\varphi_i}^j \wedge \theta_{V_\theta}^{n-j} = \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} = \int_X \theta_{\varphi_i \vee \varphi}^j \wedge \theta_{V_\theta}^{n-j}$$

**Proof** The estimates (6.4) follows from the combination of Lemma 6.2.2 and Lemma 6.2.1.

The last assertion follows from (6.4) and Theorem 2.3.2.  $\square$

cor:incseqdSconv

**Corollary 6.2.3** Suppose that  $\varphi_i \in \text{PSH}(X, \theta)$  ( $i \in I$ ) be an increasing net, uniformly bounded from above. Then

$$\varphi_i \xrightarrow{d_S} \sup_{j \in I}^* \varphi_j.$$

**Proof** Write  $\varphi = \sup_{j \in I}^* \varphi_j$ . Recall that by Proposition 1.2.1,  $\varphi \in \text{PSH}(X, \theta)$ . By Lemma 6.2.2, it suffices to show that for each  $k = 0, \dots, n$ , we have

$$\lim_{j \in I} \int_X \theta_{\varphi_j}^k \wedge \theta_{V_\theta}^{n-k} = \int_X \theta_\varphi^k \wedge \theta_{V_\theta}^{n-k}.$$

The latter follows from Corollary 2.3.1.  $\square$

By contrast, for decreasing nets, the situation is different:

cor:decnetdS

**Corollary 6.2.4** Suppose that  $\varphi_i \in \text{PSH}(X, \theta)$  is a decreasing net such that  $\varphi := \inf_{i \in I} \varphi_i \not\equiv -\infty$ . Then the following are equivalent:

(1) We have

$$\varphi_i \xrightarrow{d_S} \varphi;$$

(2) for each  $k = 0, \dots, n$ , we have

$$\lim_{j \in I} \int_X \theta_{\varphi_j}^k \wedge \theta_{V_\theta}^{n-k} = \int_X \theta_\varphi^k \wedge \theta_{V_\theta}^{n-k}. \quad (6.5)$$

{eq:mixedmasslim}

If we assume furthermore that  $\int_X \theta_\varphi^n > 0$ , then the above conditions are equivalent to the following:

(3) We have

$$\lim_{j \in I} \int_X \theta_{\varphi_j}^n = \int_X \theta_\varphi^n.$$

In the latter case, we also have

$$P_\theta[\varphi] = \inf_{j \in I} P_\theta[\varphi_j]. \quad (6.6)$$

{eq:Pcontdecseq}

**Proof** Recall that by Proposition 1.2.1,  $\varphi \in \text{PSH}(X, \theta)$ .

(1)  $\iff$  (2). This follows immediately from Lemma 6.2.2.

(2)  $\implies$  (3). This is trivial.

(3)  $\implies$  (2). Let  $(b_j)_{j \in I}$  be a net converging to  $\infty$  such that

$$b_j \in \left(1, \left(\frac{\int_X \theta_{\varphi_j}^n}{\int_X \theta_{\varphi_j}^n - \int_X \theta_{\varphi}^n}\right)^{1/n}\right).$$

By [Lemma 2.3.1](#), for each  $j \in I$ , we can find  $\eta_j \in \text{PSH}(X, \theta)$  such that

$$b_j^{-1} \eta_j + (1 - b_j^{-1}) \varphi_j \leq \varphi.$$

It follows from [Theorem 2.3.2](#) that for any  $k = 0, \dots, n$ ,

$$\int_X \theta_{\varphi}^k \wedge \theta_{V_{\theta}}^{n-k} \geq (1 - b_j^{-1})^k \int_X \theta_{\varphi_j}^k \wedge \theta_{V_{\theta}}^{n-k}.$$

Taking the limit, we conclude the  $\leq$  direction in [\(6.5\)](#). The  $\geq$  direction follows from [Theorem 2.3.2](#).

Finally, we argue [\(6.6\)](#).

Let  $\psi_j = P_{\theta}[\varphi_j]$ . It follows from [Corollary 3.1.1](#) that  $\psi_j$  is a model potential. Let

$$\psi = \inf_{j \in I} \psi_j.$$

It follows from [Proposition 3.1.2](#) and [Proposition 3.1.8](#) that

$$\int_X \theta_{\psi}^n = \lim_{j \in I} \int_X \theta_{\psi_j}^n = \lim_{j \in I} \int_X \theta_{\varphi_j}^n = \int_X \theta_{\varphi}^n.$$

By [Proposition 3.1.7](#),  $\psi$  is a model potential. So by [Proposition 6.1.1](#), we have  $\varphi \sim_P \psi$  and hence  $\psi = P_{\theta}[\varphi]$  by [Corollary 6.1.2](#).  $\square$

Having understood the increasing and decreasing cases, we shall handle more general convergent sequences. In fact, since  $d_S$  is a pseudometric, the topology is completely determined by convergent sequences, so we do not need to consider nets in general.

prop:incanddec

**Proposition 6.2.3** *Let  $\varphi_j, \varphi \in \text{PSH}(X, \theta)$  ( $j \geq 1$ ),  $\varphi_j \xrightarrow{d_S} \varphi$ . Assume that there is  $\delta > 0$  such that*

$$\int_X \theta_{\varphi_j}^n \geq \delta, \quad \int_X \theta_{\varphi}^n \geq \delta$$

*for all  $j$  and the  $\varphi_j$ 's and  $\varphi$  are all model potentials. Then up to replacing  $(\varphi_j)_j$  by a subsequence, there is a decreasing sequence  $\psi_j \in \text{PSH}(X, \theta)$  and an increasing sequence  $\eta_j \in \text{PSH}(X, \theta)$  such that*

- (1)  $\psi_j \xrightarrow{d_S} \varphi, \eta_j \xrightarrow{d_S} \varphi$ ;
- (2)  $\psi_j \geq \varphi_j \geq \eta_j$  for all  $j$ .

*In fact, for any  $j \geq 1$ , we will take*

$$\eta_j = \inf_{k \in \mathbb{N}} \varphi_j \wedge \varphi_{j+1} \wedge \dots \wedge \varphi_{j+k}, \quad \psi_j = \sup_{k \geq j}^* \varphi_k.$$

**Proof** We are free to replace  $(\varphi_j)_j$  by a subsequence. So we may assume that

$$d_S(\varphi_j, \varphi_{j+1}) \leq C_n^{-2j}, \quad d_S(\varphi, \varphi_j) \leq \frac{2^{-j-2}}{(n+1)C_n}, \quad (6.7)$$

{eq:conditiononvarphi jtemp1}

where  $C_n$  is the constant in [Corollary 6.2.2](#).

**Step 1.** We handle the  $\psi_j$ 's. For each  $j \geq 1$  and  $k \geq 1$ , by [Corollary 6.2.2](#) we have

$$\begin{aligned} d_S(\varphi_j, \varphi_j \vee \varphi_{j+1} \vee \cdots \vee \varphi_{j+k}) &\leq C_n d_S(\varphi_j, \varphi_{j+1} \vee \cdots \vee \varphi_{j+k}) \\ &\leq C_n d_S(\varphi_j, \varphi_{j+1}) + C_n d_S(\varphi_{j+1}, \varphi_{j+1} \vee \cdots \vee \varphi_{j+k}). \end{aligned}$$

By iteration, we find

$$\begin{aligned} d_S(\varphi_j, \varphi_j \vee \varphi_{j+1} \vee \cdots \vee \varphi_{j+k}) &\leq \sum_{a=j}^{j+k-1} C_n^{a+1-j} d_S(\varphi_a, \varphi_{a+1}) \\ &\leq \sum_{a=j}^{j+k-1} C_n^{a+1-j} C_n^{-2a} = \frac{C_n^{1-2j}}{1 - C_n^{-1}}. \end{aligned}$$

Using [Corollary 6.2.3](#), we have

$$\varphi_j \vee \varphi_{j+1} \vee \cdots \vee \varphi_{j+k} \xrightarrow{d_S} \psi_j$$

as  $k \rightarrow \infty$  and hence when  $j \geq j_0$  for some  $j_0$ , we have

$$d_S(\varphi_j, \psi_j) \leq \frac{C_n^{1-2j}}{1 - C_n^{-1}} \leq \frac{1}{(n+1)C_n 2^{2+j}}. \quad (6.8)$$

{eq:dsvvarphi jpsijesttemp1}

We conclude that  $\psi_j \xrightarrow{d_S} \varphi$ .

Moreover, we observe that

$$\varphi = \inf_j P_\theta[\psi_j] \quad (6.9)$$

{eq:varphiexpressiontemp1}

by [Corollary 6.2.4](#).

**Step 2.** We consider the  $\eta_j$ 's.

For each  $j \geq 1$  and  $k \geq 0$ , we let

$$\eta_j^k := \varphi_j \wedge \cdots \wedge \varphi_{j+k}.$$

Using the assumption (6.7) and [Corollary 6.2.2](#), we have

$$\left| \int_X \theta_{\varphi_j}^n - \int_X \theta_\varphi^n \right| \leq 2^{-j}.$$

Similarly, using (6.8), we have

$$\left| \int_X \theta_{\psi_j}^n - \int_X \theta_{\varphi}^n \right| \leq 2^{-j}.$$

**Step 2.1.** Take  $j_1$  so that for  $j \geq j_1$ ,  $2^{3-j} < \delta$ . We claim that for a fixed  $j \geq j_0 \vee j_1$ , for any  $k \in \mathbb{N}$ , we have  $\eta_j^k \in \text{PSH}(X, \theta)$  and

$$\int_X \theta_{\eta_j^k} \geq \int_X \theta_{\varphi_j}^n - \sum_{a=0}^k 2^{-j-a+2}.$$

We argue by induction on  $k \geq 0$ . The case  $k = 0$  follows from [Theorem 2.3.2](#). When  $k > 0$ , assume that the case  $k - 1$  is known. Then

$$\begin{aligned} \int_X \theta_{\eta_j^{k-1}}^n + \int_X \theta_{\varphi_{j+k}}^n &> \int_X \theta_{\varphi_j}^n - \sum_{a=0}^{k-1} 2^{2-j-a} + \int_X \theta_{\psi_{j+k-1}}^n - 2^{2-j-k} \\ &\geq \int_X \theta_{\varphi_j}^n - 2^{3-j} + \int_X \theta_{\psi_{j+k-1}}^n > \int_X \theta_{\psi_{j+k-1}}^n. \end{aligned}$$

It follows from [Proposition 3.1.3](#) that  $\eta_j^k \in \text{PSH}(X, \theta)$ . By [Theorem 3.1.3](#), we deduce that

$$\int_X \theta_{\varphi_{j+k}}^n + \int_X \theta_{\eta_j^{k-1}}^n \leq \int_X \theta_{\psi_{j+k-1}}^n + \int_X \theta_{\eta_j^k}^n.$$

Our claim therefore follows.

**Step 2.2.** It follows from [Proposition 3.1.6](#) that

$$P_{\theta}[\eta_j^k] = \eta_k^j.$$

By [Proposition 3.1.8](#), we have

$$\lim_{k \rightarrow \infty} \int_X \theta_{\varphi_j^k}^n = \int_X \theta_{\eta_j}^n.$$

By Step 1, for large enough  $j$ , we have

$$\int_X \theta_{\eta_j}^n \geq \int_X \theta_{\varphi_j}^n - 2^{3-j} > 0.$$

Let  $\eta = \sup^*_j \eta^j$ . Observe that we also have

$$\int_X \theta_{\eta_j}^n \leq \int_X \theta_{\psi_j}^n$$

by [Theorem 2.3.2](#). It follows that

$$\int_X \theta_{\eta}^n = \lim_{j \rightarrow \infty} \int_X \theta_{\varphi_j}^n = \lim_{j \rightarrow \infty} \int_X \theta_{\psi_j}^n = \int_X \theta_{\varphi}^n.$$

Since  $\eta_j \leq \varphi_j \leq \psi_j \leq 0$ , we also have that  $\eta_j \leq P_\theta[\psi_j]$ . Therefore, by [Corollary 6.2.4](#), we also have  $\eta \leq \varphi$ . It follows from [Proposition 6.1.1](#) that  $\eta \sim_P \varphi$ . By [Corollary 6.2.3](#) and [Proposition 6.2.2](#), we have  $\eta^j \xrightarrow{d_S} \varphi$ .  $\square$

cor:completenessdS

**Corollary 6.2.5** *Let  $(\varphi_j)_{j \in I}$  be a net in  $\text{PSH}(X, \theta)$ . Assume that there is  $\delta > 0$  such that  $\int_X \theta_{\varphi_j}^n \geq \delta$  for all  $j \in I$ . Then  $(\varphi_j)_{j \in I}$  has a  $d_S$ -convergent subnet. If moreover  $(\varphi_j)_{j \in I}$  is decreasing, then  $(\varphi_j)_{j \in I}$  itself is convergent.*

**Proof** Since the space of  $\varphi \in \text{PSH}(X, \theta)$  with  $\int_X \theta_\varphi^n \geq \delta$  is a pseudometric space, its completeness can be characterized using sequences instead of nets. So we may assume that  $(\varphi_j)_{j \in I}$  is a sequence.

Replacing  $\varphi_j$  by a subsequence, we may assume that (6.7) holds. By the proof of [Proposition 6.2.3](#) Step 1, we may assume that  $\varphi_j$  is a decreasing sequence. In this case, by [Proposition 6.2.2](#) and [Corollary 6.1.2](#), we may assume that each  $\varphi_j$  is a model potential. Then  $\varphi_j$  converges by [Corollary 6.2.4](#) and [Proposition 3.1.8](#).

On the other hand, if  $(\varphi_j)_{j \in I}$  is decreasing, then it is convergent by [Corollary 6.2.4](#) and [Proposition 3.1.8](#).  $\square$

lma:dSsmallmult

**Lemma 6.2.3** *There is a constant  $C > 0$  such that for any  $\varphi \in \text{PSH}(X, \theta)$  satisfying that  $\theta_\varphi$  is a Kähler current, we have*

$$d_{S, \theta}((1 - \epsilon)\varphi, \varphi) \leq C\epsilon$$

for  $\epsilon > 0$  such that  $(1 - \epsilon)\varphi \in \text{PSH}(X, \theta)$ .

**Proof** By [Lemma 6.2.2](#), we can compute

$$\begin{aligned} d_{S, \theta}((1 - \epsilon)\varphi, \varphi) &= \frac{1}{n+1} \sum_{j=0}^n \left( \int_X \theta_{(1-\epsilon)\varphi}^j \wedge \theta_{V_\theta}^{n-j} - \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} \right) \\ &= \frac{1}{n+1} \sum_{j=0}^n \left( \int_X (1 - \epsilon)^j \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} - \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} \right) \\ &\quad + \sum_{j=0}^n \sum_{k=0}^{j-1} \binom{j}{k} (1 - \epsilon)^k \epsilon^{j-k} \int_X \theta_\varphi^{j-k} \wedge \theta_\varphi^k \wedge \theta_{V_\theta}^{n-j}. \end{aligned}$$

Both terms are of the order of  $O(\epsilon)$ .  $\square$

## 6.2.2 Convergence theorems

lma:dsconvperv

**Lemma 6.2.4** *Let  $(\varphi_i)_{i \in I}$  be a net in  $\text{PSH}(X, \theta)$  and  $\varphi \in \text{PSH}(X, \theta)$ . Assume that  $\varphi_i \xrightarrow{d_S} \varphi$ . Then for any  $t \in (0, 1]$ ,*

$$(1 - t)\varphi_i + tV_\theta \xrightarrow{d_S} (1 - t)\varphi + tV_\theta.$$

**Proof** Fix  $t \in (0, 1]$ , we write

$$\varphi_{i,t} = (1-t)\varphi_i + tV_\theta, \quad \varphi_t = (1-t)\varphi + tV_\theta$$

for any  $i \in I$ . By [Corollary 6.2.2](#), it suffices to show that for each  $j = 0, \dots, n$ ,

$$2 \int_X \theta_{\varphi_{i,t} \vee \varphi_t}^j \wedge \theta_{V_\theta}^{n-j} - \int_X \theta_{\varphi_{i,t}}^j \wedge \theta_{V_\theta}^{n-j} - \int_X \theta_{\varphi_t}^j \wedge \theta_{V_\theta}^{n-j} \rightarrow 0. \quad (6.10)$$

{eq:massconvafterpert}

Observe that

$$\varphi_{i,t} \vee \varphi_t = (1-t)(\varphi \vee \varphi_i) + tV_\theta.$$

So after binary expansion, (6.10) follows from [Corollary 6.2.2](#).  $\square$

Similarly,

lma:linearpertbyVtheta

**Lemma 6.2.5** Let  $\varphi \in \text{PSH}(X, \theta)$ . For each  $t \in (0, 1)$ , let  $\varphi_t = (1-t)\varphi + tV_\theta$ . Then

$$\varphi_t \xrightarrow{d_S} \varphi$$

as  $t \rightarrow 0+$ .

**Proof** By [Lemma 6.2.2](#), we need to show that for each  $j = 1, \dots, n$ , we have

$$\lim_{t \rightarrow 0+} \int_X \theta_{\varphi_t}^j \wedge \theta_{V_\theta}^{n-j} = \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j}.$$

For this purpose, we compute

$$\begin{aligned} & \int_X \theta_{\varphi_t}^j \wedge \theta_{V_\theta}^{n-j} - \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} \\ &= \sum_{i=0}^{j-1} \binom{j}{i} (1-t)^i t^{j-i} \theta_\varphi^i \wedge \theta_{V_\theta}^{n-i}. \end{aligned}$$

As  $t \rightarrow 0+$ , the right-hand side clearly tends to 0.  $\square$

The following convergent theorem lies at the heart of the whole theory.

thm:convdS

**Theorem 6.2.1** Let  $\theta_1, \dots, \theta_n$  be smooth closed real  $(1, 1)$ -forms on  $X$  representing big cohomology classes. Suppose that  $(\varphi_j^k)_{k \in I}$  are nets in  $\text{PSH}(X, \theta_j)$  for  $j = 1, \dots, n$  and  $\varphi_1, \dots, \varphi_n \in \text{PSH}(X, \theta)$ . We assume that  $\varphi_j^k \xrightarrow{d_S} \varphi_j$  for each  $j = 1, \dots, n$ . Then

$$\lim_{k \in I} \int_X \theta_{1, \varphi_1^k} \wedge \dots \wedge \theta_{n, \varphi_n^k} = \int_X \theta_{1, \varphi_1} \wedge \dots \wedge \theta_{n, \varphi_n}. \quad (6.11)$$

{eq:convmixedmassds}

**Proof** Since  $d_S$  is a pseudometric, in order to establish the continuity of mixed masses, it suffices to consider sequences instead of nets. So we may assume that  $I = \mathbb{Z}_{>0}$  as ordered sets.



**Step 1.** We reduce to the case where  $\varphi_j^k, \varphi_j$  all have positive masses and there is a constant  $\delta > 0$ , such that for all  $j$  and  $k$ ,

$$\int_X \theta_{j, \varphi_j^k}^n > \delta.$$

Take  $t \in (0, 1)$ . By [Lemma 6.2.4](#), we have

$$(1-t)\varphi_j^k + tV_{\theta_j} \xrightarrow{ds} (1-t)\varphi_j + tV_{\theta_j}$$

for each  $j$ . Assume that we have proved the special case of the theorem, we have

$$\begin{aligned} & \lim_{k \in I} \int_X \theta_{1, (1-t)\varphi_1^k + tV_{\theta_1}} \wedge \cdots \wedge \theta_{n, (1-t)\varphi_n^k + tV_{\theta_n}} \\ &= \int_X \theta_{1, (1-t)\varphi_1 + tV_{\theta_1}} \wedge \cdots \wedge \theta_{n, (1-t)\varphi_n + tV_{\theta_n}}. \end{aligned}$$

Since both sides are polynomials in  $t$ , it follows that the same holds at  $t = 0$ . From this, [\(6.11\)](#) follows.

**Step 2.** Next we may assume that  $\varphi_j^k, \varphi_j$  are model potentials by [Proposition 6.2.2](#) and [Corollary 3.1.1](#).

It suffices to prove that any subsequence of  $\int_X \theta_{1, \varphi_1^k} \wedge \cdots \wedge \theta_{n, \varphi_n^k}$  has a converging subsequence with limit  $\int_X \theta_{1, \varphi_1} \wedge \cdots \wedge \theta_{n, \varphi_n}$ . Thus, by [Proposition 6.2.3](#) and [Theorem 2.3.2](#), we may assume that for each fixed  $i$ ,  $\varphi_i^k$  is either increasing or decreasing. We may assume that for  $i \leq i_0$ , the sequence is decreasing and for  $i > i_0$ , the sequence is increasing.

Recall that in [\(6.11\)](#) the  $\geq$  inequality always holds by [Theorem 2.3.2](#), it suffices to prove

$$\overline{\lim}_{k \in I} \int_X \theta_{1, \varphi_1^k} \wedge \cdots \wedge \theta_{n, \varphi_n^k} \leq \int_X \theta_{1, \varphi_1} \wedge \cdots \wedge \theta_{n, \varphi_n}. \quad (6.12)$$

{eq:limsup}

By [Theorem 2.3.2](#) in order to prove [\(6.12\)](#), we may assume that for  $j > i_0$ , the sequences  $\varphi_j^k$  are constant. Thus, we are reduced to the case where for all  $i$ ,  $\varphi_i^k$  are decreasing.

In this case, for each  $i$  we may take an increasing sequence  $b_i^k > 1$ , tending to  $\infty$ , such that

$$(b_i^k)^n \int_X \theta_{i, \varphi_i}^n \geq ((b_i^k)^n - 1) \int_X \theta_{i, \varphi_i^k}^n.$$

Let  $\psi_i^k$  be the maximal  $\theta_i$ -psh function such that

$$(b_i^k)^{-1} \psi_i^k + (1 - (b_i^k)^{-1}) \varphi_i^k \leq \varphi_i,$$

whose existence is guaranteed by [Lemma 2.3.1](#).

Then by [Theorem 2.3.2](#) again,

$$\prod_{i=1}^n \left(1 - (b_i^k)^{-1}\right) \int_X \theta_{1, \varphi_1^k} \wedge \cdots \wedge \theta_{n, \varphi_n^k} \leq \int_X \theta_{1, \varphi_1} \wedge \cdots \wedge \theta_{n, \varphi_n}.$$

Letting  $k \rightarrow \infty$ , we conclude (6.12).  $\square$

cor:dsconvcrit

**Corollary 6.2.6** Suppose that  $(\varphi_i)_{i \in I}$  is a net in  $\text{PSH}(X, \theta)$  and  $\varphi \in \text{PSH}(X, \theta)$ . Then the following are equivalent:

- (1)  $\varphi_i \xrightarrow{d_S} \varphi$ ;
- (2)  $\varphi_i \vee \varphi \xrightarrow{d_S} \varphi$  and

$$\lim_{i \in I} \int_X \theta_{\varphi_i}^j \wedge \theta_{V_\theta}^{n-j} = \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} \quad (6.13)$$

{eq:massconv\_varphi}

for each  $j = 0, \dots, n$ .

The corollary allows us to reduce a number of convergence problems related to  $d_S$  to the case  $\varphi_i \geq \varphi$ , which is much easier to handle by Lemma 6.2.2. This is the most handy way of establishing  $d_S$ -convergence in practice.

**Proof** (1)  $\implies$  (2).  $\varphi_i \vee \varphi \xrightarrow{d_S} \varphi$  follows from Corollary 6.2.2. While (6.13) follows from Theorem 6.2.1.

(2)  $\implies$  (1). By (6.4), we need to show that for each  $j = 0, \dots, n$ , we have

$$2 \int_X \theta_{\varphi_i \vee \varphi}^j \wedge \theta_{V_\theta}^{n-j} - \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j} - \int_X \theta_{\varphi_i}^j \wedge \theta_{V_\theta}^{n-j} \rightarrow 0.$$

This follows from Theorem 6.2.1 and (6.13).  $\square$

cor:dSconv\_changetheta

**Corollary 6.2.7** Let  $(\varphi_i)_{i \in I}$  be a net in  $\text{PSH}(X, \theta)$  and  $\varphi \in \text{PSH}(X, \theta)$ . Let  $\omega$  be a Kähler form on  $X$ . Then the following are equivalent:

- (1)  $\varphi_i \xrightarrow{d_{S, \theta}} \varphi$ ;
- (2)  $\varphi_i \xrightarrow{d_{S, \theta + \omega}} \varphi$ .

In particular, there is no risk when we simply write  $\varphi_i \xrightarrow{d_S} \varphi$ .

**Proof** (1)  $\implies$  (2). It suffices to show that for each  $j = 0, \dots, n$ , we have

$$\begin{aligned} 2 \int_X (\theta + \omega)_{\varphi_i \vee \varphi}^j \wedge (\theta + \omega)_{V_{\theta + \omega}}^{n-j} - \int_X (\theta + \omega)_\varphi^j \wedge (\theta + \omega)_{V_{\theta + \omega}}^{n-j} \\ - \int_X (\theta + \omega)_{\varphi_i}^j \wedge (\theta + \omega)_{V_{\theta + \omega}}^{n-j} \rightarrow 0. \end{aligned}$$

Note that this quantity is a linear combination of terms of the following form:

$$\begin{aligned} 2 \int_X \theta_{\varphi_i \vee \varphi}^r \wedge \omega^{j-r} \wedge (\theta + \omega)_{V_{\theta + \omega}}^{n-j} - \int_X \theta_{\varphi_i}^r \wedge \omega^{j-r} \wedge (\theta + \omega)_{V_{\theta + \omega}}^{n-j} \\ - \int_X \theta_\varphi^r \wedge \omega^{j-r} \wedge (\theta + \omega)_{V_{\theta + \omega}}^{n-j}, \end{aligned}$$

where  $r = 0, \dots, j$ . By [Theorem 6.2.1](#), it suffices to show that  $\varphi \vee \varphi_i \xrightarrow{d_S} \varphi$ . But this follows from [Corollary 6.2.6](#).

(2)  $\implies$  (1). From the direction we already proved, for each  $C \geq 1$ , we have that

$$\varphi_i \xrightarrow{d_{S, \theta + C\omega}} \varphi.$$

By [Theorem 6.2.1](#), it follows that

$$\lim_{i \in I} \int_X (\theta + C\omega)_{\varphi_i}^j \wedge \theta_{V_\theta}^{n-j} = \int_X (\theta + C\omega)_\varphi^j \wedge \theta_{V_\theta}^{n-j}$$

for all  $j = 0, \dots, n$ . It follows that

$$\lim_{i \in I} \int_X \theta_{\varphi_i}^j \wedge \theta_{V_\theta}^{n-j} = \int_X \theta_\varphi^j \wedge \theta_{V_\theta}^{n-j}. \quad (6.14)$$

{eq:varphi\_jmass\_limit}

By [Corollary 6.2.6](#), it remains to show that  $\varphi_i \vee \varphi \xrightarrow{d_{S, \theta}} \varphi$ . By [Corollary 6.2.6](#) again, we know that  $\varphi_i \vee \varphi \xrightarrow{d_{S, \theta + \omega}} \varphi$ . So it suffices to apply (6.14) to  $\varphi_i \vee \varphi$  instead of  $\varphi_i$ , and we conclude by [Lemma 6.2.2](#).  $\square$

We sometimes need a slightly more general form.

cor:dseivalenceindep

**Corollary 6.2.8** *Let  $(\varphi_j)_{j \in I}$ ,  $(\psi_j)_{j \in I}$  be nets in  $\text{PSH}(X, \theta)$ . Consider a Kähler form  $\omega$  on  $X$ . Then the following are equivalent:*

- (1)  $d_{S, \theta}(\varphi_i, \psi_i) \rightarrow 0$ ;
- (2)  $d_{S, \theta + \omega}(\varphi_i, \psi_i) \rightarrow 0$ .

In particular, we can write  $d_S(\varphi_i, \psi_i) \rightarrow 0$  without ambiguity.

**Proof** The proof is similar to that of [Corollary 6.2.7](#), which is therefore left to the readers.  $\square$

We have the following sandwich criterion:

lma:dsconvupplower

**Corollary 6.2.9** *Let  $(\varphi_i)_{i \in I}$ ,  $(\psi_i)_{i \in I}$ ,  $(\eta_i)_{i \in I}$  be three nets in  $\text{PSH}(X, \theta)$  and  $\varphi \in \text{PSH}(X, \theta)$ . Assume that*

- (1)  $\psi_i \leq_P \varphi_i \leq_P \eta_i$  for each  $i \in I$ ;
- (2)  $\eta_i \xrightarrow{d_S} \varphi$ ,  $\psi_i \xrightarrow{d_S} \varphi$ .

Then  $\varphi_i \xrightarrow{d_S} \varphi$ .

**Proof** By [Corollary 6.2.7](#), we may replace  $\theta$  by  $\theta + \omega$ , where  $\omega$  is a Kähler form on  $X$ . In particular, we may assume that  $\varphi_i, \psi_i, \eta_i \in \text{PSH}(X, \theta)_{>0}$  for all  $i \in I$ . By [Proposition 6.2.2](#), we may assume that  $\varphi_i, \psi_i, \eta_i$  are model potentials for all  $i \in I$  and hence  $\varphi_i \leq \psi_i \leq \eta_i$  for all  $i \in I$ .

It follows from [Theorem 2.3.2](#) that for each  $k = 0, \dots, n$ , we have

$$\int_X \theta_{\psi_i}^k \wedge \theta_{V_\theta}^{n-k} \leq \int_X \theta_{\varphi_i}^k \wedge \theta_{V_\theta}^{n-k} \leq \int_X \theta_{\eta_i}^k \wedge \theta_{V_\theta}^{n-k}$$

for all  $i \in I$ . By [Theorem 6.2.1](#), the limits of the both ends are  $\int_X \theta_\varphi^k \wedge \theta_{V_\theta}^{n-k}$  as  $j \rightarrow \infty$ . It follows that

$$\lim_{i \in I} \int_X \theta_{\varphi_i}^k \wedge \theta_{V_\theta}^{n-k} = \int_X \theta_\varphi^k \wedge \theta_{V_\theta}^{n-k}. \quad (6.15)$$

`{eq:thetak_conv}`

By [Corollary 6.2.6](#), it remains to prove that  $\varphi_i \vee \varphi \xrightarrow{d_S} \varphi$ . By [Corollary 6.2.6](#), up to replacing  $\psi_i$  (resp.  $\varphi_i, \eta_i$ ) by  $\psi_i \vee \varphi$  (resp.  $\varphi_i \vee \varphi, \eta_i \vee \varphi$ ), we may assume from the beginning that  $\psi_i, \varphi_i, \eta_i \geq \varphi$ . Now  $\varphi_i \xrightarrow{d_S} \varphi$  by (6.15) and [Lemma 6.2.2](#).  $\square$

`prop:dconvpresorder`

**Proposition 6.2.4** *Let  $(\varphi_i)_{i \in I}, (\psi_i)_{i \in I}$  be nets in  $\text{PSH}(X, \theta)$  such that  $\varphi_i \xrightarrow{d_S} \varphi \in \text{PSH}(X, \theta)$  and  $\psi_i \xrightarrow{d_S} \psi \in \text{PSH}(X, \theta)$ . Assume that  $\varphi_i \leq_P \psi_i$  for all  $i \in I$ . Then  $\varphi \leq_P \psi$ .*

**Proof** It follows from [Proposition 6.2.5](#) that

$$\varphi_i \vee \psi_i \xrightarrow{d_S} \varphi \vee \psi.$$

By [Lemma 6.1.2](#), we have  $\varphi_i \vee \psi_i \sim_P \psi_i$  for all  $i \in I$ . In particular, by [Proposition 6.2.2](#),

$$\varphi_i \vee \psi_i \xrightarrow{d_S} \psi.$$

By [Proposition 6.2.2](#) again,  $\varphi \vee \psi \sim_P \psi$  and hence  $\varphi \leq_P \psi$  by [Lemma 6.1.2](#).  $\square$

`lma:dsior`

**Lemma 6.2.6** *Let  $\varphi, \psi, \eta \in \text{PSH}(X, \theta)$ , then*

$$d_S(\varphi \vee \eta, \psi \vee \eta) \leq C_n d_S(\varphi, \psi), \quad (6.16)$$

`{eq:dSmax}`

where  $C_n = 3(n+1)2^{n+2}$ .

**Proof** According to [Corollary 6.2.2](#), we may assume that  $\varphi \leq \psi$ .

We will show that for each  $C \geq t \geq 0$ ,

$$d_1(\ell_t^{\varphi \vee \eta, C}, \ell_t^{\psi \vee \eta, C}) \leq d_1(\ell_t^{\varphi, C}, \ell_t^{\psi, C}). \quad (6.17)$$

`{eq:d1maxcomp}`

When  $C \rightarrow \infty$ , by [Corollary 2.3.1](#) and [Theorem 4.2.1](#), it follows that

$$d_1(\ell_t^{\varphi \vee \eta}, \ell_t^{\psi \vee \eta}) \leq d_1(\ell_t^\varphi, \ell_t^\psi),$$

which implies (6.16).

It remains to argue (6.17). As  $\varphi \leq \psi$ , we know that

$$d_1(\ell_t^\varphi, \ell_t^\psi) = \frac{t}{C} d_1(\ell_C^\varphi, \ell_C^\psi), \quad d_1(\ell_t^{\varphi \vee \eta}, \ell_t^{\psi \vee \eta}) = \frac{t}{C} d_1(\ell_C^{\varphi \vee \eta}, \ell_C^{\psi \vee \eta}).$$

It suffices to handle the case  $t = C$ , namely,

$$d_1(\varphi \vee \eta \vee (V_\theta - C), \psi \vee \eta \vee (V_\theta - C)) \leq d_1(\varphi \vee (V_\theta - C), \psi \vee (V_\theta - C)).$$

This is a consequence of [Theorem 4.2.2](#).  $\square$

prop:lor\_dS\_conv

**Proposition 6.2.5** *Let  $(\varphi_i)_{i \in I}$  (resp.  $(\psi_i)_{i \in I}$ ) be a net in  $\text{PSH}(X, \theta)$  such that  $\varphi_i \xrightarrow{d_S} \varphi \in \text{PSH}(X, \theta)$  (resp.  $\psi_i \xrightarrow{d_S} \psi \in \text{PSH}(X, \theta)$ ). Then*

$$\varphi_i \vee \psi_i \xrightarrow{d_S} \varphi \vee \psi.$$

**Proof** We compute

$$\begin{aligned} d_S(\varphi_i \vee \psi_i, \varphi \vee \psi) &\leq d_S(\varphi_i \vee \psi_i, \varphi_i \vee \psi) + d_S(\varphi_i \vee \psi, \varphi \vee \psi) \\ &\leq C_n (d_S(\psi_i, \psi) + d_S(\varphi_i, \varphi)), \end{aligned}$$

where the second inequality follows from [Lemma 6.2.6](#). The right-hand side converges to 0 by our hypothesis.  $\square$

thm:dSadditivity

**Theorem 6.2.2** *Let  $\theta_1, \theta_2$  be smooth real closed  $(1, 1)$ -forms on  $X$  representing big cohomology classes. Suppose that  $(\varphi_i)_{i \in I}$  (resp.  $(\psi_i)_{i \in I}$ ) be a net in  $\text{PSH}(X, \theta_1)$  (resp.  $\text{PSH}(X, \theta_2)$ ) and  $\varphi \in \text{PSH}(X, \theta_1)$  (resp.  $\psi \in \text{PSH}(X, \theta_2)$ ). Consider the following three conditions:*

- (1)  $\varphi_i \xrightarrow{d_S} \varphi$ ;
- (2)  $\psi_i \xrightarrow{d_S} \psi$ ;
- (3)  $\varphi_i + \psi_i \xrightarrow{d_S} \varphi + \psi$ .

*Then any two of these conditions imply the third.*

**Proof** By [Corollary 6.2.7](#), we may assume that  $\theta_1, \theta_2$  are both Kähler forms. We denote them by  $\omega_1, \omega_2$  instead. Let  $\omega = \omega_1 + \omega_2$ .

(1)+(2)  $\implies$  (3). It suffices to show that for each  $r = 0, \dots, n$ ,

$$2 \int_X \omega_{(\varphi_j + \psi_j) \vee (\varphi + \psi)}^r \wedge \omega^{n-r} - \int_X \omega_{\varphi_j + \psi_j}^r \wedge \omega^{n-r} - \int_X \omega_{\varphi + \psi}^r \wedge \omega^{n-r} \rightarrow 0.$$

Observe that for each  $j \in I$ ,

$$(\varphi_j + \psi_j) \vee (\varphi + \psi) \leq \varphi_j \vee \varphi + \psi_j \vee \psi.$$

Thus, it suffices to show that

$$2 \int_X \omega_{\varphi_j \vee \varphi + \psi_j \vee \psi}^r \wedge \omega - \int_X \omega_{\varphi_j + \psi_j}^r \wedge \omega^{n-r} - \int_X \omega_{\varphi + \psi}^r \wedge \omega^{n-r} \rightarrow 0.$$

The left-hand side is a linear combination of

$$2 \int_X \omega_{1, \varphi_j \vee \varphi}^a \wedge \omega_{2, \psi_j \vee \psi}^{r-a} \wedge \omega^{n-r} - \int_X \omega_{1, \varphi_j}^a \wedge \omega_{2, \psi_j}^{r-a} \wedge \omega^{n-r} - \int_X \omega_{1, \varphi}^a \wedge \omega_{2, \psi}^{r-a} \wedge \omega^{n-r}$$

with  $a = 0, \dots, r$ . Observe that  $\varphi_j \vee \varphi \xrightarrow{d_S} \varphi$  and  $\psi_j \vee \psi \xrightarrow{d_S} \psi$  by [Corollary 6.2.2](#), each term tends to 0 by [Theorem 6.2.1](#).

(2)+(3)  $\implies$  (1). This is similar.

(1)+(3)  $\implies$  (2). For each  $C \geq 1$ , from the direction we already proved,

$$C\varphi_i + \psi_i \xrightarrow{d_S} C\varphi + \psi.$$

By [Theorem 6.2.1](#), for each  $j = 0, \dots, n$ ,

$$\begin{aligned} & \lim_{i \in I} \int_X (C\omega_1 + \omega_2 + \text{dd}^c(C\varphi_i + \psi_i))^j \wedge \omega_2^{n-j} \\ &= \int_X (C\omega_1 + \omega_2 + \text{dd}^c(C\varphi + \psi))^j \wedge \omega_2^{n-j}. \end{aligned}$$

It follows that

$$\lim_{i \in I} \int_X \omega_{2, \psi_i}^j \wedge \omega_2^{n-j} = \int_X \omega_{2, \psi}^j \wedge \omega_2^{n-j}. \quad (6.18)$$

{eq:psii\_quant\_conv}

Therefore, 2 follows if  $\psi_i \geq \psi$  for each  $i$  by [Lemma 6.2.2](#).

Next we prove the general case. By the direction that we already proved, we know that  $\varphi_i + \psi \xrightarrow{d_S} \varphi + \psi$ . By [Proposition 6.2.5](#), we have that

$$\varphi_i + \psi_i \vee \psi \xrightarrow{d_S} \varphi + \psi.$$

It follows from the special case above that  $\psi_i \vee \psi \xrightarrow{d_S} \psi$ . It follows from (6.18) and [Corollary 6.2.6](#) that (2) holds.  $\square$

thm:contPI

**Theorem 6.2.3** *The map*

$$P_\theta[\bullet]_I : \text{PSH}(X, \theta)_{>0} \rightarrow \text{PSH}(X, \theta)_{>0}$$

*is continuous with respect to  $d_S$ .*

**Proof** Let  $(\varphi_i)_{i \in \mathbb{Z}_{>0}}$  be a sequence in  $\text{PSH}(X, \theta)_{>0}$  such that  $\varphi_i \xrightarrow{d_S} \varphi \in \text{PSH}(X, \theta)_{>0}$ . We want to show that

$$P[\varphi_i]_I \xrightarrow{d_S} P[\varphi]_I. \quad (6.19)$$

We may assume that the  $\varphi_i$ 's and  $\varphi$  are all model potentials by [Proposition 6.2.2](#).

By [Proposition 6.2.3](#) and [Corollary 6.2.9](#), we may assume that  $(\varphi_i)_i$  is either increasing or decreasing. The two cases are handled by [Proposition 3.2.12](#) and [Proposition 3.2.11](#) respectively.  $\square$

### 6.2.3 Continuity of invariants

thm:Lelongcont

**Theorem 6.2.4** *Let  $(\varphi_j)_{j \in I}$  be a net in  $\text{PSH}(X, \theta)$  and  $\varphi_j \xrightarrow{d_S} \varphi \in \text{PSH}(X, \theta)$ . Then for any prime divisor  $E$  over  $X$ , we have*

$$\lim_{j \in I} \nu(\varphi_j, E) = \nu(\varphi, E). \quad (6.20)$$

{eq:convnu}

**Proof** First observe that since  $d_S$  is a pseudometric, it suffices to prove (6.20) when  $I = \mathbb{Z}_{>0}$  as partially ordered sets.

By **Corollary 6.2.7**, we may assume that the masses of  $\varphi_j$  and of  $\varphi$  are bounded from below by a positive constant.

By **Theorem 6.2.3**, we may assume that  $\varphi_i$  and  $\varphi$  are both  $\mathcal{I}$ -model. When proving (6.20), we are free to pass to subsequences.

By **Proposition 6.2.3**, we may assume that the sequence  $(\varphi_i)$  is either increasing or decreasing. In the increasing case, there is nothing to prove. In the decreasing case, (6.20) follows from **Proposition 3.1.8**.  $\square$

thm:contvolu

**Theorem 6.2.5** *Let  $(\varphi_j)_{j \in I}$  be a net in  $\text{PSH}(X, \theta)$  and  $\varphi \in \text{PSH}(X, \theta)_{>0}$ . Assume that  $\varphi_j \xrightarrow{d_S} \varphi \in \text{PSH}(X, \theta)$ , then*

$$\text{vol } \theta_{\varphi_j} \rightarrow \text{vol } \theta_{\varphi}. \quad (6.21)$$

{eq:Ivolcont}

Recall the volume is defined in **Definition 3.2.3**.

**Proof** It follows from **Theorem 6.2.1** that

$$\int_X \theta_{\varphi_j}^n \rightarrow \int_X \theta_{\varphi}^n.$$

We may therefore assume that  $\int_X \theta_{\varphi_j}^n > 0$  for all  $j \in I$ . Then by **Theorem 6.2.3**, we have

$$P_{\theta}[\varphi_j]_I \xrightarrow{d_S} P_{\theta}[\varphi]_I.$$

Therefore, (6.21) follows from **Theorem 6.2.1**.  $\square$

thm:equising\_cond\_general

**Theorem 6.2.6** *Let  $\varphi_j, \varphi \in \text{PSH}(X, \theta)$  ( $j \in \mathbb{Z}_{>0}$ ). Assume that  $\varphi_j \xrightarrow{d_S} \varphi$ . Then for each  $\lambda' > \lambda > 0$ , there is  $j_0 > 0$  so that for  $j \geq j_0$ ,*

$$\mathcal{I}(\lambda' \varphi_j) \subseteq \mathcal{I}(\lambda \varphi). \quad (6.22)$$

{eq:quasi\_equi\_cond}

**Proof** Fix  $\lambda' > \lambda > 0$ , we want to find  $j_0 > 0$  so that for  $j \geq j_0$ , (6.22) holds.

**Step 1.** We first assume that  $\varphi$  has analytic singularities.

Let  $\pi : Y \rightarrow X$  be a log resolution of  $\varphi$  and let  $E_1, \dots, E_N$  be all prime divisors of the singular part of  $\varphi$  on  $Y$ . Recall that a local holomorphic function  $f$  lies in the right-hand side of (6.22) if and only if

$$\mathrm{ord}_{E_i}(f) > \lambda \mathrm{ord}_{E_i}(\varphi) - \frac{1}{2} A_X(E_i) \quad (6.23)$$

 $\{\mathrm{eq:ordEif}\}$ 

whenever they make sense. Here  $A_X$  denotes the log discrepancy. Similarly,  $f$  lies in the left-hand side of (6.22) implies that there is  $\epsilon > 0$  so that

$$\mathrm{ord}_{E_i}(f) \geq (1 + \epsilon) \lambda' \mathrm{ord}_{E_i}(\varphi_j) - \frac{1}{2} A_X(E_i).$$

As Lelong numbers are continuous with respect to  $d_S$  by [Theorem 6.2.4](#), we can find  $j_0 > 0$  so that when  $j \geq j_0$ ,  $\lambda' \mathrm{ord}_{E_i}(\varphi_j) \geq \lambda \mathrm{ord}_{E_i}(\varphi)$  for all  $i$ . In particular, (6.23) follows.

**Step 2.** We handle the general case.

By [Corollary 6.2.7](#), we are free to increase  $\theta$  and assume that  $\theta_\varphi$  is a Kähler current.

Take a quasi-equisingular approximation  $(\psi_k)_k$  of  $\varphi$ . The existence is guaranteed by [Theorem 1.6.2](#). Take  $\lambda'' \in (\lambda, \lambda')$ , then by definition, we can find  $k > 0$  so that

$$I(\lambda'' \psi_k) \subseteq I(\lambda \varphi).$$

Observe that  $\varphi_j \vee \psi_k \xrightarrow{d_S} \psi_k$  as  $j \rightarrow \infty$  by [Proposition 6.2.5](#). By Step 1, we can find  $j_0 > 0$  so that for  $j \geq j_0$ ,

$$I(\lambda'(\varphi_j \vee \psi_k)) \subseteq I(\lambda'' \psi_k).$$

It follows that for  $j \geq j_0$ ,

$$I(\lambda' \varphi_j) \subseteq I(\lambda \varphi).$$



## Chapter 7

### $\mathcal{I}$ -good singularities

chap:Igood

In this chapter, we study the key notion in the whole theory: the  $\mathcal{I}$ -good singularities. We will give several useful characterizations of  $\mathcal{I}$ -good singularities. The key result is the asymptotic Riemann–Roch formula for Hermitian big line bundles [Theorem 7.3.1](#).

#### 7.1 The notion of $\mathcal{I}$ -good singularities

Let  $X$  be a connected compact Kähler manifold of dimension  $n$ .

thm:charIgoodasclosure

**Theorem 7.1.1** *Let  $\theta$  be a closed real smooth  $(1, 1)$ -form on  $X$  representing a big cohomology class. Let  $\varphi \in \text{PSH}(X, \theta)_{>0}$ . Then the following are equivalent:*

(1) *There exists a sequence  $(\varphi_j)_j$  in  $\text{PSH}(X, \theta)$  with analytic singularities such that*

$$\varphi_j \xrightarrow{ds} \varphi.$$

(2) *We have*

$$\int_X \theta_\varphi^n = \text{vol } \theta_\varphi. \quad (7.1)$$

{eq:nppmassequalvolume}

(3) *We have*

$$P_\theta[\varphi] = P_\theta[\varphi]_{\mathcal{I}}.$$

*In (1), we could in addition require that each  $\theta_{\varphi_j}$  is a Kähler current.*

*Moreover, if  $\theta_\varphi$  is a Kähler current, the sequence in (1) can be taken as any quasi-equisingular approximation of  $\varphi$  in  $\text{PSH}(X, \theta)$ .*

**Proof** (1)  $\implies$  (2). By [Theorem 6.2.1](#), we may assume that  $\int_X \theta_{\varphi_j}^n > 0$  for all  $j$ . It follows from [Proposition 3.2.9](#) that

$$\int_X \theta_{\varphi_j}^n = \text{vol } \theta_{\varphi_j}$$

for any  $j \geq 1$ . Using [Theorem 6.2.5](#) and [Theorem 6.2.1](#), we conclude (7.1).

(2)  $\iff$  (3). This follows from [Theorem 3.1.1](#).

(3)  $\implies$  (1). Note that the condition in (1) characterizes the closure of analytic singularities in  $\text{PSH}(X, \theta)$ .

**Step 1.** We first reduce to the case where  $\theta_\varphi$  is a Kähler current.

By [Lemma 2.3.2](#), we can find  $\psi \in \text{PSH}(X, \theta)$  so that  $\theta_\psi$  is a Kähler current and  $\psi \leq \varphi$ . We let

$$\psi_j = (1 - j^{-1})\varphi + j^{-1}\psi$$

for each  $j \in \mathbb{Z}_{>0}$ . Then  $(\psi_j)_j$  is an increasing sequence converging almost everywhere to  $\varphi$ . Then

$$P_\theta[\psi_j]_I \xrightarrow{d_S} P_\theta[\varphi]_I = P_\theta[\varphi]$$

by [Proposition 3.2.12](#), [Corollary 6.2.3](#). So it suffices to show that  $P_\theta[\psi_j]_I$  lies in the closure of analytic singularities.

**Step 2.** We assume that  $\theta_\varphi$  is a Kähler current. We show that  $P_\theta[\varphi]_I$  lies in the closure of analytic singularities.

Let  $(\varphi_j)_j$  be a quasi-equisingular approximation of  $\varphi$  in  $\text{PSH}(X, \theta)$ . We will show that  $\varphi_j \xrightarrow{d_S} P_\theta[\varphi]_I$ . Let

$$\psi = \inf_{j \in \mathbb{Z}_{>0}} P_\theta[\varphi_j].$$

We know that  $\varphi_j \xrightarrow{d_S} \psi$  by [Proposition 6.2.2](#), [Proposition 3.1.8](#) and [Corollary 6.2.4](#).

Moreover, observe that  $\psi$  is  $\mathcal{I}$ -model by [Proposition 3.2.11](#) and [Example 7.1.1](#). So it suffices to show that  $\varphi \sim_{\mathcal{I}} \psi$ .

It is clear that  $\psi \geq \varphi$ . Conversely, it remains to argue that  $\psi \leq_{\mathcal{I}} \varphi$ . For this purpose, take  $\lambda > 0$ , we need to show that

$$\mathcal{I}(\lambda\psi) \subseteq \mathcal{I}(\lambda\varphi).$$

By the strong openness [Theorem 1.4.4](#), we may take  $\lambda' > \lambda$  such that  $\mathcal{I}(\lambda\psi) = \mathcal{I}(\lambda'\psi)$ , then it follows from the definition of the quasi-equisingular approximation that

$$\mathcal{I}(\lambda'\psi) \subseteq \mathcal{I}(\lambda'\varphi_j) \subseteq \mathcal{I}(\lambda\varphi)$$

for large enough  $j$ . Our assertion follows.  $\square$

def:Igoodpot

**Definition 7.1.1** We say a potential  $\varphi \in \text{QPSH}(X)$  is  $\mathcal{I}$ -good if for some smooth closed real  $(1, 1)$ -form on  $X$  such that  $\varphi \in \text{PSH}(X, \theta)_{>0}$ , we have

$$P_\theta[\varphi] = P_\theta[\varphi]_I. \tag{7.2}$$

{eq:envelopeeq}

An immediate question is to verify that this definition is independent of the choice of  $\theta$ .

lma:Igoodinsenspert

**Lemma 7.1.1** Let  $\varphi \in \text{PSH}(X, \theta)_{>0}$  for some smooth closed real  $(1, 1)$ -form  $\theta$  on  $X$ . Take a Kähler form  $\omega$  on  $X$ . Then the following are equivalent:

- (1)  $P_\theta[\varphi] = P_\theta[\varphi]_I$ ;
- (2)  $P_{\theta+\omega}[\varphi] = P_\theta[\varphi + \omega]_I$ .

**Proof** (1)  $\implies$  (2). By [Theorem 7.1.1](#), we can find  $\varphi_j \in \text{PSH}(X, \theta)$  with analytic singularities such that  $\varphi_j \xrightarrow{d_{S, \theta}} \varphi$ . By [Corollary 6.2.7](#), we have  $\varphi_j \xrightarrow{d_{S, \theta + \omega}} \varphi$ . Therefore, by [Theorem 7.1.1](#) again, 2 holds.

(2)  $\implies$  (1). Suppose that (1) fails, so that

$$\int_X (\theta + \text{dd}^c \varphi)^n < \int_X (\theta + \text{dd}^c P_\theta[\varphi]_I)^n.$$

It follows that

$$\begin{aligned} \int_X (\theta + \omega + \text{dd}^c \varphi)^n &= \sum_{i=0}^n \binom{n}{i} \int_X \theta_\varphi^i \wedge \omega^{n-i} \\ &< \sum_{i=0}^n \binom{n}{i} \int_X \theta_{P_\theta[\varphi]_I}^i \wedge \omega^{n-i} \\ &= \int_X (\theta + \omega + \text{dd}^c P_\theta[\varphi]_I)^n \\ &\leq \int_X (\theta + \omega + \text{dd}^c P_{\theta+\omega}[\varphi]_I)^n. \end{aligned}$$

So (2) fails as well.  $\square$

cor:Igoodclosed

**Corollary 7.1.1** *Let  $\theta$  be a closed real smooth  $(1, 1)$ -form on  $X$  representing a big cohomology class. Let  $(\varphi_j)_{j \in I}$  be a net of  $\mathcal{I}$ -good potentials in  $\text{PSH}(X, \theta)$  such that  $\varphi_j \xrightarrow{d_S} \varphi$ . Then  $\varphi$  is  $\mathcal{I}$ -good.*

**Proof** By [Corollary 6.2.7](#), we may assume that  $\varphi_j, \varphi \in \text{PSH}(X, \theta)_{>0}$  for all  $j \in I$ . It follows from [Theorem 7.1.1](#) that

$$\int_X \theta_{\varphi_j}^n = \text{vol } \theta_{\varphi_j}$$

for all  $j \in I$ . Taking limit with respect to  $j$  with the help of [Theorem 6.2.5](#) and [Theorem 6.2.1](#), we conclude that

$$\int_X \theta_\varphi^n = \text{vol } \theta_\varphi.$$

Therefore, by [Theorem 7.1.1](#) again, we find that  $\varphi$  is  $\mathcal{I}$ -good.  $\square$

ex:analyIgood

**Example 7.1.1** Assume that  $\varphi \in \text{QPSH}(X)$  has analytic singularities. Then  $\varphi$  is  $\mathcal{I}$ -good. This is proved in [Proposition 3.2.9](#).

ex:ImodelIgood

**Example 7.1.2** Assume that  $\varphi \in \text{PSH}(X, \theta)_{>0}$  is an  $\mathcal{I}$ -model potential for some closed real smooth  $(1, 1)$ -form  $\theta$  on  $X$ . Then  $\varphi$  is  $\mathcal{I}$ -good.

cor:quasi-equichar

**Corollary 7.1.2** *Let  $\varphi \in \text{PSH}(X, \theta)_{>0}$  and  $(\epsilon_j)_j$  be a decreasing sequence in  $\mathbb{R}_{\geq 0}$  with limit 0. Fix a Kähler form  $\omega$  on  $X$ . Consider a decreasing sequence  $\varphi_j \in$*

$\text{PSH}(X, \theta + \epsilon_j \omega)$  of potentials with analytic singularities for each  $j \geq 1$ . Assume that  $\varphi = \inf_j \varphi_j$ . Then the following are equivalent:

- (1)  $\varphi_j \xrightarrow{d_S} P_\theta[\varphi]_I$ , and
- (2)  $(\varphi_j)_j$  is a quasi-equisingular approximation of  $\varphi$ .

**Proof** By [Corollary 6.2.7](#) and [Example 7.1.2](#), we may replace  $\theta$  by  $\theta + C\omega$  for some large constant  $C > 0$  and assume that  $\varphi, \varphi_j \in \text{PSH}(X, \theta - \omega)$  for all  $j \geq 1$ .

(2)  $\implies$  (1). This is already proved in the proof of [Theorem 7.1.1](#).

(1)  $\implies$  (2). This follows from [Theorem 6.2.6](#).  $\square$

ex:BBJ

**Example 7.1.3** Let  $X = \mathbb{P}^1$  and  $\omega$  be the Fubini–Study metric. Let  $K \subseteq \mathbb{P}^1$  be a polar Cantor sets carrying an atom free probability measure  $\mu$  supported on  $K$  (see [\[Car83, Page 31\]](#)). Write  $\mu = \omega + \Delta\varphi$  for some  $\varphi \in \text{SH}(X, \omega)$ . Since  $\mu$  is atom free, we know that all Lelong numbers of  $\varphi$  are 0. On the other hand,  $\varphi$  has 0 non-pluripolar mass since  $K$  is pluripolar. In particular,  $c\varphi$  for  $c \in (0, 1)$  is not  $\mathcal{I}$ -good.

## 7.2 Properties of $\mathcal{I}$ -good singularities

Let  $X$  be a connected compact Kähler manifold.

prop:Igoodlinear

**Proposition 7.2.1** Let  $\varphi, \psi \in \text{QPSH}(X)$  be  $\mathcal{I}$ -good and  $\lambda > 0$ . Then the following potentials are all  $\mathcal{I}$ -good.

- (1)  $\varphi + \psi$ ;
- (2)  $\varphi \vee \psi$ ;
- (3)  $\lambda\varphi$ .

**Proof** Take a closed real smooth  $(1, 1)$ -form  $\theta$  on  $X$  such that  $\varphi, \psi \in \text{PSH}(X, \theta)_{>0}$ . It follows from [Theorem 7.1.1](#) that there are sequences  $\varphi_j, \psi_j$  in  $\text{PSH}(X, \theta)$  with analytic singularities such that  $\varphi_j \xrightarrow{d_S} \varphi$  and  $\psi_j \xrightarrow{d_S} \psi$ .

By [Theorem 6.2.2](#), [Proposition 6.2.5](#), we have

$$\varphi_j + \psi_j \xrightarrow{d_S} \varphi + \psi, \quad \varphi_j \vee \psi_j \xrightarrow{d_S} \varphi \vee \psi.$$

On the other hand, it is clear that

$$\lambda\varphi_j \xrightarrow{d_S} \lambda\varphi.$$

Therefore, our assertions follow from [Theorem 7.1.1](#).  $\square$

prop:Igoodsup

**Proposition 7.2.2** Let  $\{\varphi_j\}_{j \in I}$  be a non-empty family of  $\mathcal{I}$ -good potentials. Assume that the family is uniformly bounded from above and there exists a closed real smooth  $(1, 1)$ -form  $\theta$  on  $X$  such that  $\varphi_j \in \text{PSH}(X, \theta)$  for all  $j \in I$ . Then  $\sup_{j \in I}^* \varphi_j$  is  $\mathcal{I}$ -good.

**Proof** Without loss of generality, we may assume that  $\varphi_j \in \text{PSH}(X, \theta)_{>0}$  for all  $j \in I$ .

When  $I$  is finite, this result follows from [Proposition 7.2.1](#). When  $I$  is infinite, we may assume that  $I = \mathbb{Z}_{>0}$  by [Proposition 1.2.2](#). By [Proposition 7.2.1](#), we may assume that the sequence  $(\varphi_j)_j$  is increasing. In this case, as shown in [Corollary 6.2.3](#),

$$\varphi_j \xrightarrow{ds} \sup_{i \in \mathbb{Z}_{>0}}^* \varphi_i.$$

Therefore,  $\sup_{i \in \mathbb{Z}_{>0}}^* \varphi_i$  is  $\mathcal{I}$ -good by [Theorem 7.1.1](#).  $\square$

thm:contvolu2

**Theorem 7.2.1** Let  $(\varphi_j)_{j \in I}$  be a net in  $\text{PSH}(X, \theta)$  such that  $\varphi_j \xrightarrow{ds} \varphi \in \text{PSH}(X, \theta)$ . Assume that  $\varphi$  is  $\mathcal{I}$ -good, then we have

$$\text{vol } \theta_{\varphi_j} \rightarrow \text{vol } \theta_{\varphi}. \quad (7.3)$$

{eq:Ivolcont2}

**Proof** Fix a Kähler form  $\omega$  on  $X$ . Then for any  $\epsilon > 0$ , we have

$$\begin{aligned} \text{vol}(\theta + \epsilon\omega)_{\varphi} &= \int_X (\theta + \epsilon\omega + \text{dd}^c P_{\theta + \epsilon\omega}[\varphi]_I)^n \\ &= \int_X (\theta + \epsilon\omega + \text{dd}^c \varphi)^n. \end{aligned}$$

On the other hand,

$$\begin{aligned} \int_X (\theta + \epsilon\omega + \text{dd}^c P_{\theta + \epsilon\omega}[\varphi]_I)^n &\geq \int_X (\theta + \epsilon\omega + \text{dd}^c P_{\theta}[\varphi]_I)^n \\ &\geq \int_X (\theta + \text{dd}^c P_{\theta}[\varphi]_I)^n \\ &\geq \int_X \theta_{\varphi}^n. \end{aligned}$$

Therefore,

$$\text{vol}(\theta + \epsilon\omega)_{\varphi} - \text{vol } \theta_{\varphi} \leq \int_X (\theta + \epsilon\omega + \text{dd}^c \varphi)^n - \int_X \theta_{\varphi}^n.$$

The difference can be controlled by a polynomial in  $\epsilon$  without constant term independent of the choice of  $\varphi$ . We have a similar estimate for  $\varphi_j$  as well. So our assertion follows from [Theorem 6.2.5](#).  $\square$

prop:vollinearlimit

**Proposition 7.2.3** Let  $\varphi, \psi \in \text{PSH}(X, \theta)_{>0}$ . Then

(1) we have

$$\lim_{\epsilon \rightarrow 0+} \text{vol}(\theta, (1 - \epsilon)\varphi + \epsilon\psi) = \text{vol}(\theta, \varphi).$$

(2) Let  $\omega$  be a Kähler form on  $X$ , then

$$\mathrm{vol} \theta_\varphi = \lim_{\epsilon \rightarrow 0+} \mathrm{vol}(\theta + \epsilon\omega)_\varphi.$$

(3) Consider a prime divisor  $E$  on  $X$ . Then

$$\mathrm{vol} \theta_\varphi = \mathrm{vol}(\theta_\varphi - \nu(\varphi, E)[E]).$$

**Proof** (1) We need to show that

$$\lim_{\epsilon \rightarrow 0+} \int_X (\theta + \mathrm{dd}^c P_\theta[(1 - \epsilon)\varphi + \epsilon\psi]_I)^n = \int_X (\theta + \mathrm{dd}^c P_\theta[\varphi]_I)^n.$$

By [Proposition 3.2.10](#), for any  $\epsilon \in (0, 1)$ ,

$$(1 - \epsilon)\varphi + \epsilon\psi \sim_I (1 - \epsilon)P_\theta[\varphi]_I + \epsilon P_\theta[\psi]_I.$$

In particular, we may replace  $\varphi$  and  $\psi$  by  $P_\theta[\varphi]_I$  and  $P_\theta[\psi]_I$  respectively. By [Proposition 7.2.1](#), it remains to show that

$$\lim_{\epsilon \rightarrow 0+} \int_X (\theta + \mathrm{dd}^c((1 - \epsilon)\varphi + \epsilon\psi))^n = \int_X (\theta + \mathrm{dd}^c \varphi)^n,$$

which is obvious.

(2) For each  $\epsilon > 0$ ,

$$\begin{aligned} \mathrm{vol}(\theta + \epsilon\omega)_\varphi &= \int_X (\theta + \epsilon\omega + \mathrm{dd}^c P_{\theta+\epsilon\omega}[\varphi]_I)^n \\ &= \int_X (\theta + \epsilon\omega + \mathrm{dd}^c P_{\theta+\epsilon\omega}[P_\theta[\varphi]_I])^n \\ &= \int_X (\theta + \epsilon\omega + \mathrm{dd}^c P_\theta[\varphi]_I)^n, \end{aligned}$$

where the third equality follows from [Example 7.1.2](#). Letting  $\epsilon \rightarrow 0+$ , we conclude.

(3) By (2), we may assume that  $\theta_\varphi$  is a Kähler current. Take a quasi-equisingular approximation  $(S_j)_j$  of  $\theta_\varphi - \nu(\varphi, E)[E]$ . By [Theorem 6.2.2](#),

$$S_j + \nu(\varphi, E)[E] \xrightarrow{d_S} \theta_\varphi.$$

For each  $j \geq 1$ , the currents  $S_j + \nu(\varphi, E)[E]$  and  $S_j$  are  $\mathcal{I}$ -good as follows from [Proposition 7.2.1](#), we have

$$\mathrm{vol}(S_j + \nu(\varphi, E)[E]) = \int_X (S_j + \nu(\varphi, E)[E])^n = \int_X S_j^n = \mathrm{vol} S_j.$$

Letting  $j \rightarrow \infty$ , we conclude by [Theorem 6.2.6](#).  $\square$

### 7.3 The volume of Hermitian big line bundles

sec:volHermitianbig

Let  $X$  be a connected compact Kähler manifold of dimension  $n$ .

**Definition 7.3.1** A *Hermitian pseudoeffective line bundle*  $(L, h)$  on  $X$  consists of a pseudoeffective line bundle  $L$  on  $X$  together with a plurisubharmonic metric  $h$  on  $L$ .

A *Hermitian big line bundle*  $(L, h)$  on  $X$  is a big line bundle  $L$  on  $X$  together with a plurisubharmonic metric  $h$  on  $L$  such that  $\text{vol}(\text{dd}^c h) > 0$ .

When  $X$  admits a big line bundle, it is necessarily projective. See [MM07, Theorem 2.2.26].

thm:DXmain1

**Theorem 7.3.1** Let  $(L, h)$  be a Hermitian big line bundle and  $T$  be a holomorphic line bundle on  $X$ . We have

$$\lim_{k \rightarrow \infty} \frac{n!}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(h^k)) = \text{vol}(\text{dd}^c h). \quad (7.4)$$

{eq:DXmain1}

In particular, the limit exists.

*Remark 7.3.1* This theorem also holds for a general Hermitian pseudoeffective line bundle. The proof is more involved. We would have to apply the singular holomorphic Morse inequality of Bonavero [Bon98]. See [DX21, Theorem 1.1].

For the proof, let us fix a smooth Hermitian metric  $h_0$  on  $L$  with  $\theta = c_1(L, h_0)$ . We identify  $h$  with  $h_0 \exp(-\varphi)$  for some  $\varphi \in \text{PSH}(X, \theta)$ .

We first handle the case where  $\varphi$  has analytic singularities.

prop:DXmainanalytic

**Proposition 7.3.1** Under the assumptions of [Theorem 7.3.1](#), assume furthermore that  $\varphi$  has analytic singularities, then [\(7.4\)](#) holds.

**Proof Step 1.** Reduce to the case of log singularities.

Let  $\pi: Y \rightarrow X$  be a modification such that  $\pi^*\varphi$  has log singularities. In this case, for each  $k \in \mathbb{Z}_{>0}$ , we have

$$h^0(X, T \otimes L^k \otimes \mathcal{I}(kh)) = h^0(Y, K_{Y/X} \otimes \pi^*T \otimes \pi^*L^k \otimes \mathcal{I}(k\pi^*h)).$$

By [Proposition 3.2.5](#), we have

$$\text{vol}(\text{dd}^c h) = \text{vol}(\text{dd}^c \pi^*h).$$

Therefore, it suffices to argue [\(7.4\)](#) with  $K_{Y/X} \otimes \pi^*T$ ,  $\pi^*L$  and  $\pi^*h$  in place of  $T$ ,  $L$  and  $h$ .

**Step 2.** Assume that  $D$  has log singularities along an effective  $\mathbb{Q}$ -divisor  $D$ , we decompose  $D$  into irreducible components, say

$$D = \sum_{i=1}^N a_i D_i.$$

In this case, we can easily compute

$$\mathcal{I}(k\varphi) = \mathcal{O}_X \left( - \sum_{i=1}^N [ka_i] D_i \right)$$

for each  $k \in \mathbb{Z}_{>0}$ . Observe that  $L - D$  is nef (see [Lemma 1.6.1](#)), so we could apply the asymptotic Riemann–Roch theorem to conclude that

$$\lim_{k \rightarrow \infty} \frac{n!}{k^n} h^0 \left( X, T \otimes L^k \otimes \mathcal{O}_X \left( - \sum_{i=1}^N [ka_i] D_i \right) \right) = (L - D)^n.$$

Observe that by [Proposition 1.8.1](#),

$$\theta_\varphi = [D] + T,$$

where  $T$  is a closed positive  $(1, 1)$ -current with bounded potential. Therefore,

$$(L - D)^n = \int_X T^n = \int_X \theta_\varphi^n.$$

By [Example 7.1.1](#), we know that the right-hand side is exactly  $\text{vol } \theta_\varphi$ .  $\square$

**Proof (Proof of [Theorem 7.3.1](#)) Step 1.** We first handle the case where  $\theta_\varphi$  is a Kähler current. Fix a Kähler form  $\omega \geq \theta$  on  $X$  such that  $\theta_\varphi \geq 2\delta\omega$  for some  $\delta \in (0, 1)$ .

Let  $(\varphi_j)_j$  be a quasi-equisingular approximation of  $\varphi$  in  $\text{PSH}(X, \theta)$ . We may assume that  $\theta_{\varphi_j} \geq \delta\omega$  for all  $j$ . From [Proposition 7.3.1](#), we know that for each  $j \geq 1$ ,

$$\overline{\lim}_{k \rightarrow \infty} \frac{n!}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi)) \leq \lim_{k \rightarrow \infty} \frac{n!}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi_j)) = \text{vol } \theta_{\varphi_j}.$$

It follows from [Theorem 7.1.1](#) and [Theorem 6.2.5](#) that the right-hand side converges to  $\text{vol } \theta_\varphi$  as  $j \rightarrow \infty$ . Therefore,

$$\overline{\lim}_{k \rightarrow \infty} \frac{n!}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi)) \leq \text{vol } \theta_\varphi.$$

Conversely, fix an integer  $N > \delta^{-1}$ . From [Theorem 7.1.1](#) and [Theorem 6.2.1](#), we know that

$$\lim_{j \rightarrow \infty} \int_X \theta_{\varphi_j}^n = \int_X \theta_{P_\theta[\varphi]_I}^n > 0. \quad (7.5)$$

{eq:quasiequassconvtempl}

Therefore, by [Lemma 2.3.1](#), we can find  $j_0 > 0$  such that for  $j \geq j_0$ , there is  $\psi \in \text{PSH}(X, \theta)_{>0}$  with

$$(1 - N^{-1})\varphi_j + N^{-1}\psi \leq P_\theta[\varphi]_I. \quad (7.6)$$

{eq:linearlowerbdPItemp1}



For each  $k > 0$ , we write  $k = k'N - r$ , where  $k' \in \mathbb{N}$  and  $r \in \{0, 1, \dots, N-1\}$ . Then we compute for  $j > j_0$  and large enough  $k$  that

$$\begin{aligned} & h^0(X, T \otimes L^k \otimes I(k\varphi)) \\ & \geq h^0(X, T \otimes L^{-r} \otimes L^{k'N} \otimes I(k'N\varphi)) \\ & \geq h^0\left(X, T \otimes L^{-r} \otimes L^{k'N} \otimes I\left(k'(\psi + (N-1)\varphi_j)\right)\right) \\ & \geq h^0\left(X, T \otimes L^{-r} \otimes L^{k'N} \otimes L^{k'(N-1)} \otimes I\left(k'N\varphi_j\right)\right), \end{aligned}$$

where the third line follows from (7.6), the fourth line can be argued as follows: for large enough  $k$ , there is a non-zero section  $s \in H^0(X, L^{k'} \otimes I(k'\psi))$  by Lemma 2.3.3; It follows from Lemma 1.6.3 that for large enough  $k$ ,

$$I(k'N\varphi_j) \subseteq I_\infty(k'(N-1)\varphi_j).$$

It follows that multiplication by  $s$  gives an injective map

$$\begin{aligned} & H^0\left(X, T \otimes L^{-r} \otimes L^{k'(N-1)} \otimes I(k'N\varphi_j)\right) \hookrightarrow \\ & H^0\left(X, T \otimes L^{-r} \otimes L^{k'N} \otimes I(k'\psi + k'(N-1)\varphi_j)\right). \end{aligned}$$

Next observe that

$$(N-1)\theta + N\mathrm{dd}^c\varphi_j \geq 0.$$

So Proposition 7.3.1 is applicable. We let  $k \rightarrow \infty$  to conclude that

$$\begin{aligned} \lim_{k \rightarrow \infty} h^0(X, T \otimes L^k \otimes I(k\varphi)) & \geq \frac{1}{n! \cdot N^{-n}} \int_X ((N-1)\theta + N\mathrm{dd}^c\varphi_j)^n \\ & = \frac{1}{n!} \int_X \left((1 - N^{-1})\theta + \mathrm{dd}^c\varphi_j\right)^n. \end{aligned}$$

Letting  $j \rightarrow \infty$  and then  $N \rightarrow \infty$  and using (7.5), we find that

$$\lim_{k \rightarrow \infty} h^0(X, T \otimes L^k \otimes I(k\varphi)) \geq \int_X \theta_{P_\theta[\varphi]_I}^n.$$

**Step 2.** We handle the general case. We may assume that  $\varphi$  is  $I$ -model.

Take an ample line bundle  $A$  on  $X$  and a Kähler form  $\omega$  in  $c_1(A)$ . Then for any fixed  $N \in \mathbb{Z}_{>0}$ , we apply Step 1 to  $L^N \otimes A$  in place of  $L$  and  $T \otimes L^i$  with  $i = 0, \dots, N-1$  in place of  $T$ , we have

$$\lim_{k \rightarrow \infty} \frac{n!}{k^n} h^0(X, T \otimes L^k \otimes I(k\varphi)) \leq \int_X \left(N^{-1}\omega + \theta + \mathrm{dd}^c P_{\theta+N^{-1}\omega}[\varphi]_I\right)^n.$$

On the other hand, since  $\varphi$  is  $I$ -good by Example 7.1.2, we have

$$P_{\theta+N^{-1}\omega}[\varphi]_I = P_{\theta+N^{-1}\omega}[\varphi].$$

It follows from [Proposition 3.1.2](#) that

$$\overline{\lim}_{k \rightarrow \infty} \frac{n!}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi)) \leq \int_X \left( \theta + N^{-1}\omega + \text{dd}^c \varphi \right)^n.$$

Letting  $N \rightarrow \infty$ , we conclude

$$\overline{\lim}_{k \rightarrow \infty} \frac{n!}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi)) \leq \int_X \theta_\varphi^n.$$

It remains to argue the reverse inequality.

Choose  $\psi \in \text{PSH}(X, \theta)$  such that  $\theta_\psi$  is a Kähler current and  $\psi \leq \varphi$ . The existence of  $\psi$  is guaranteed by [Lemma 2.3.2](#). Then for any  $t \in (0, 1)$ , we set

$$\varphi_t = (1-t)\varphi + t\psi.$$

It follows again from Step 1 that

$$\lim_{k \rightarrow \infty} \frac{n!}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi)) \geq \lim_{k \rightarrow \infty} \frac{n!}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi_t)) = \text{vol } \theta_{\varphi_t}.$$

On the other hand, by [Corollary 6.2.3](#), we have  $\varphi_t \xrightarrow{d_S} \varphi$  as  $t \rightarrow 0+$ . It follows from [Theorem 6.2.5](#) that

$$\lim_{t \rightarrow 0+} \text{vol } \theta_{\varphi_t} = \text{vol } \theta_\varphi.$$

So we find

$$\lim_{k \rightarrow \infty} \frac{n!}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi)) \geq \text{vol } \theta_\varphi.$$

ex:toricIgood

*Example 7.3.1* If  $X$  is a toric smooth projective variety and  $\theta$  is invariant under the action of the compact torus. Suppose that  $\varphi \in \text{PSH}(X, \theta)_{>0}$  is also invariant under the action of the compact torus, then  $\varphi$  is  $\mathcal{I}$ -good.

**Proof** Thanks to [Lemma 7.1.1](#), we may assume that  $\theta \in c_1(L)$  for some toric invariant ample line bundle  $L$ . In this case, the result follows from [Theorem 7.1.1](#), [Theorem 7.3.1](#) and [Theorem 5.3.1](#).  $\square$

cor:volbigL

**Corollary 7.3.1** *We have*

$$\lim_{k \rightarrow \infty} \frac{n!}{k^n} h^0(X, L^k) = \int_X \theta_{V_\theta}^n. \quad (7.7) \quad \{\text{eq:volbig}\}$$

This common quantity is the *volume* of  $L$ , usually denoted by  $\text{vol } L$ .

## Chapter 8

### The trace operator

chap:trace

In this chapter, we develop the theory of trace operators and prove the analytic Bertini theorem. These techniques allow us to make induction on the dimension while studying the singularities.

#### 8.1 The definition of the trace operator

Let  $X$  be a connected compact Kähler manifold and  $Y \subseteq X$  be an irreducible analytic subset. The trace operator gives a way to restrict a quasi-plurisubharmonic function on  $X$  to  $\tilde{Y}$ , the normalization of  $Y$ . It follows from [GK20, Proposition 3.5] that  $\tilde{Y}$  is a normal Kähler space. We refer to Appendix B for the pluripotential theory on unibranch Kähler spaces.

For later applications, we need this generality even if initially we are only interested in the smooth case.

We first observe that given  $\varphi \in \text{QPSH}(X)$  with analytic singularities such that  $\nu(\varphi, Y) = 0$ , then  $\varphi|_Y \not\equiv -\infty$ . This observation will be crucial in the sequel.

op:traceindquasiequisingapp

**Proposition 8.1.1** *Let  $\varphi \in \text{QPSH}(X)$ . Consider a smooth closed real  $(1, 1)$ -form on  $X$  and  $\varphi \in \text{PSH}(X, \theta)$  such that  $\nu(\varphi, Y) = 0$ . Let  $(\varphi_i)_i, (\psi_i)_i$  be quasi-equisingular approximations of  $\varphi$ . Then*

$$\lim_{i \rightarrow \infty} d_S(\varphi_i|_{\tilde{Y}}, \psi_i|_{\tilde{Y}}) = 0. \quad (8.1)$$

{eq:dsequivtemp1}

The meaning of (8.1) is explained in Corollary 6.2.8.

**Proof** Take a Kähler form  $\omega$  on  $X$ . By Corollary 6.2.8, we may assume that  $\varphi, \varphi_i, \psi_i \in \text{PSH}(X, \theta - \omega)$  for all  $i \geq 1$ . Replacing  $\varphi$  by  $P_\theta[\varphi]_I$ , we may assume that  $\varphi$  is  $I$ -good. It follows from Corollary 7.1.2 and Proposition 6.2.5 that we can assume  $\varphi_i \leq \psi_i$  for all  $i \geq 1$ .

Take a decreasing sequence  $(\epsilon_j)_j$  in  $\mathbb{R}_{>0}$  with limit 0 such that  $(1 - \epsilon_j)\varphi_j \in \text{PSH}(X, \theta)$ . We first observe that

$$\lim_{i \rightarrow \infty} d_S(\varphi_i|_{\tilde{Y}}, (1 - \epsilon_i)\varphi_i|_{\tilde{Y}}) = 0.$$

This is a consequence of [Lemma 6.2.3](#).

Next by [Proposition 1.6.3](#), we could find a subsequence  $(\psi_{j_i})_{i \in \mathbb{Z}_{>0}}$  of  $(\psi_j)_j$  such that for each  $i \geq 1$ ,

$$\varphi_{j_i} \leq \psi_{j_i} \leq (1 - \epsilon_i)\varphi_i.$$

Therefore, [\(8.1\)](#) follows from [Corollary 6.2.1](#).  $\square$

def:traceop

**Definition 8.1.1** Let  $\varphi \in \text{QPSH}(X)$  such that  $\nu(\varphi, Y) = 0$ . We say a potential  $\psi \in \text{QPSH}(\tilde{Y})$  is a *trace operator* of  $\varphi$  along  $Y$  if there is a smooth closed real  $(1, 1)$ -form  $\theta$  on  $X$  such that  $\varphi \in \text{PSH}(X, \theta)$  and a quasi-equisingular approximation  $(\varphi_j)_j$  of  $\varphi$  such that

$$\varphi_j|_{\tilde{Y}} \xrightarrow{d_S} \psi. \quad (8.2)$$

{eq:deftrace}

By [Corollary 6.2.5](#), the trace operator is always defined. Observe that by [Proposition 8.1.1](#), the condition [\(8.2\)](#) is independent of the choice of  $(\varphi_j)_j$ . It is also independent of the choice of  $\theta$  by [Corollary 6.2.7](#).

prop:traceunique

**Proposition 8.1.2** Let  $\varphi \in \text{QPSH}(X)$  such that  $\nu(\varphi, Y) = 0$ . Suppose that  $\psi$  and  $\psi'$  are trace operators of  $\varphi$  along  $Y$ . Then  $\psi$  and  $\psi'$  are  $\mathcal{I}$ -good and  $\psi \sim_P \psi'$ .

*Proof* That  $\psi$  and  $\psi'$  are  $\mathcal{I}$ -good follows from [Theorem 7.1.1](#). The fact that  $\psi \sim_P \psi'$  follows from [Proposition 8.1.1](#) and [Proposition 6.2.2](#).  $\square$

**Definition 8.1.2** Let  $\varphi \in \text{QPSH}(X)$  such that  $\nu(\varphi, Y) = 0$ . We write  $\text{Tr}_Y(\varphi)$  for any trace operator of  $\varphi$  along  $Y$ .

Given a closed smooth real  $(1, 1)$ -form  $\theta$  on  $X$ . When  $\text{Tr}_Y(\varphi)$  can be chosen to lie in  $\text{PSH}(\tilde{Y}, \theta|_{\tilde{Y}})_{>0}$ , we write

$$\text{Tr}_Y^\theta(\varphi) := P_{\theta|_{\tilde{Y}}} [\text{Tr}_Y(\varphi)] = P_{\theta|_{\tilde{Y}}} [\text{Tr}_Y(\varphi)]_I.$$

The trace operator  $\text{Tr}_Y(\varphi)$  is therefore well-defined only up to  $P$ -equivalence by [Proposition 8.1.2](#).

rmk:tracecurrent

*Remark 8.1.1* As in [Remark 1.7.1](#), the trace operator could also be applied to closed positive  $(1, 1)$ -currents on  $X$ . If  $T \in \mathcal{Z}_+(X, \alpha)$  (see [Definition 1.7.3](#)) and  $\beta \in H^{1,1}(\tilde{Y}, \mathbb{R})$ , then we write

$$\text{Tr}_Y^\beta(T)$$

for any closed positive  $(1, 1)$ -current in  $\beta$  representing  $\text{Tr}_Y(T)$  when  $\nu(T, Y) = 0$ .

prop:Trdominarest

**Proposition 8.1.3** Let  $\varphi \in \text{QPSH}(X)$  such that  $\nu(\varphi, Y) = 0$ . Assume that  $\varphi|_Y \not\equiv -\infty$ . Then

$$\varphi|_{\tilde{Y}} \leq_P \text{Tr}_Y(\varphi).$$

*Proof* Take a Kähler form  $\omega$  such that  $\omega_\varphi$  is a Kähler current. Let  $(\varphi_j)_j$  be a quasi-equisingular approximation of  $\varphi$  in  $\text{PSH}(X, \omega)$ . We may assume that  $\varphi_j \leq 0$  for all  $j \geq 1$ .

Then

$$\varphi_j|_{\bar{Y}} \leq P_{\theta|_{\bar{Y}}} [\varphi_j|_{\bar{Y}}] \quad (8.3) \quad \{\text{eq:varphijrestrleqPtemp}\}$$

for all  $j \geq 1$ .

Thanks to [Corollary 6.2.4](#),

$$\text{Tr}_Y(\varphi) \sim_P \inf_{j \geq 1} P_{\theta|_{\bar{Y}}} [\varphi_j|_{\bar{Y}}]. \quad (8.4) \quad \{\text{eq:TrYnewexpression}\}$$

Letting  $j \rightarrow \infty$  in (8.3), we conclude our assertion.  $\square$

ex:resanalyt

*Example 8.1.1* Let  $\varphi \in \text{QPSH}(X)$  such that  $v(\varphi, Y) = 0$ . Assume that  $\varphi$  has analytic singularities, then

$$\text{Tr}_Y(\varphi) \sim_P \varphi|_{\bar{Y}}.$$

*Example 8.1.2* Let  $\varphi \in \text{QPSH}(X)$ . Take a closed real smooth  $(1, 1)$ -form  $\theta$  on  $X$  such that  $\varphi \in \text{PSH}(X, \theta)_{>0}$ , then

$$\text{Tr}_X(\varphi) \sim_P P_\theta[\varphi]_I, \quad \text{Tr}_X^\theta(\varphi) = P_\theta[\varphi]_I.$$

In particular, the trace operator can be regarded as a generalization of the  $I$ -envelope.

ex:tracedefinedposmass

*Example 8.1.3* Assume that  $\varphi \in \text{PSH}(X, \theta)$  for some closed smooth real  $(1, 1)$ -form  $\theta$  on  $X$  and

$$\lim_{\epsilon \searrow 0} \int_Y \left( \theta|_Y + \epsilon \omega|_Y + \text{dd}^c \text{Tr}_Y^{\theta+\epsilon}(\varphi) \right)^m > 0 \quad (8.5) \quad \{\text{eq:traceposmasscond}\}$$

for any arbitrary choice of a Kähler form  $\omega$  on  $X$ . Then it follows from [Proposition 3.1.8](#) that  $\text{Tr}_Y^\theta(\varphi)$  is defined, and its mass is exact the above limit.

In particular, if  $\theta_\varphi$  is a Kähler current,  $\text{Tr}_Y^\theta(\varphi)$  is always defined.

*Remark 8.1.2* The trace operator allows us to introduce the following extension of the moving Seshadri constant: Let  $T \in \mathcal{Z}_+(X, \alpha)$  and  $x \in X$ , we define

$$\epsilon(T, x) := \inf_{V \ni x} \left( \frac{\text{vol Tr}_V^{\alpha|_{\bar{V}}} T}{\text{mult}_x V} \right)^{\frac{1}{\dim V}},$$

where  $\text{vol Tr}_V^{\alpha|_{\bar{V}}} T = 0$  if  $\text{Tr}_V^{\alpha|_{\bar{V}}} T$  is not defined. Here  $V$  runs over all positive-dimensional closed irreducible analytic subsets of  $X$  containing  $x$ .

These moving Seshadri constants seem to be new.

## 8.2 Properties of the trace operator

Let  $X$  be a connected compact Kähler manifold and  $Y \subseteq X$  be an irreducible analytic subset.

prop:tracelinear

**Proposition 8.2.1** Let  $\varphi, \psi \in \text{QPSH}(X)$ ,  $\lambda > 0$ . Assume that  $\nu(\varphi, Y) = \nu(\psi, Y) = 0$ . Then we have the following:

- (1) Suppose that  $\varphi \leq_I \psi$ , then  $\text{Tr}_Y(\varphi) \leq_P \text{Tr}_Y(\psi)$ .  
 (2) We have

$$\text{Tr}_Y(\varphi + \psi) \sim_P \text{Tr}_Y(\varphi) + \text{Tr}_Y(\psi).$$

- (3) We have

$$\text{Tr}_Y(\lambda\varphi) \sim_P \lambda \text{Tr}_Y(\varphi).$$

- (4) We have

$$\text{Tr}_Y(\varphi \vee \psi) \sim_P \text{Tr}_Y(\varphi) \vee \text{Tr}_Y(\psi).$$

**Proof** Take a closed smooth real  $(1, 1)$ -form  $\theta$  on  $X$  such that  $\theta_\varphi, \theta_\psi$  are both Kähler currents. Let  $(\varphi_j)_j$  and  $(\psi_j)_j$  be quasi-equisingular approximations of  $\varphi$  and  $\psi$  in  $\text{PSH}(X, \theta)$  respectively.

(1) By [Corollary 7.1.2](#) and [Proposition 6.2.5](#), we may assume that  $\varphi_j \leq \psi_j$  for all  $j$ . Then our assertion follows from [Proposition 6.2.4](#).

(2) It follows from [Theorem 6.2.2](#) that  $\varphi_j + \psi_j \xrightarrow{ds} P_\theta[\varphi]_I + P_\theta[\psi]_I$ . However, by [Proposition 3.2.10](#) and [Proposition 7.2.1](#), we have

$$P_\theta[\varphi]_I + P_\theta[\psi]_I \sim_P P_\theta[\varphi + \psi]_I.$$

Therefore, by [Proposition 6.2.2](#), [Corollary 7.1.2](#) and [Proposition 1.6.1](#),  $\varphi_j + \psi_j$  is a quasi-equisingular approximation of  $\varphi + \psi$ . We conclude using [Theorem 6.2.2](#).

(3) Let  $(\lambda_j)_j$  be an increasing sequence of positive rational numbers with limit  $\lambda$ . Then  $(\lambda_j \varphi_j)_j$  is a quasi-equisingular approximation of  $\varphi$ . Our assertion follows [Lemma 6.2.3](#).

(4) By [Proposition 6.2.5](#), we have

$$\varphi_j \vee \psi_j \xrightarrow{ds} P_\theta[\varphi]_I \vee P_\theta[\psi]_I.$$

By [Proposition 3.2.10](#) and [Proposition 7.2.1](#), we have

$$P_\theta[\varphi]_I \vee P_\theta[\psi]_I \sim_P P_\theta[\varphi \vee \psi]_I.$$

Therefore, our assertion follows exactly as in the proof of (2).  $\square$

prop:tracedeclimit

**Proposition 8.2.2** Let  $(\varphi_j)_{j \in I}$  be a decreasing net in  $\text{QPSH}(X)$ . Assume that there exists a closed real smooth  $(1, 1)$ -form  $\theta$  such that  $\varphi_j \in \text{PSH}(X, \theta)$  for each  $j \in I$ .

Assume that  $\varphi_j \xrightarrow{ds} \varphi \in \text{QPSH}(X)$  and  $\nu(\varphi, Y) = 0$ . Then

$$\text{Tr}_Y(\varphi_j) \xrightarrow{ds} \text{Tr}_Y(\varphi).$$

**Proof** By [Corollary 6.2.7](#), we may assume that there is a Kähler form  $\omega$  on  $X$  such that  $\varphi, \varphi_j \in \text{PSH}(X, \theta - \omega)$  for all  $j \in I$ . Note that for each  $j \geq 1$ ,

$$\text{Tr}_Y(\varphi_{j+1}) \leq_P \text{Tr}_Y(\varphi_j).$$

It follows from [Proposition 8.2.1](#) and [Corollary 6.2.5](#) that there exists  $\psi \in \text{PSH}(\tilde{Y}, \theta|_{\tilde{Y}})$  such that  $\text{Tr}_Y(\varphi_j) \xrightarrow{ds} \psi$ .

For each  $j$ , we take a quasi-equisingular approximation  $(\varphi_j^k)_k$  in  $\text{PSH}(X, \theta)$  of  $\varphi_j$ . Using [Theorem 1.6.2](#), we may guarantee that

$$\varphi_{j+1}^k \leq \varphi_j^k$$

for each  $j, k \geq 1$ . In particular,  $(\varphi_j^j)_j$  is a quasi-equisingular approximation of  $\varphi$ . By [Proposition 6.2.4](#), we have  $\psi \leq_P \text{Tr}_Y(\varphi)$ .

Conversely, by [Proposition 8.2.1](#),  $\text{Tr}_Y(\varphi_j) \geq_P \text{Tr}_Y(\varphi)$ . It follows again from [Proposition 6.2.4](#) that  $\text{Tr}_Y(\varphi) \leq_P \psi$ .  $\square$

*Example 8.2.1* The trace operator is not continuous along increasing sequences. Let us consider the case  $X = \mathbb{P}^2$  with coordinates  $(z_1, z_2)$ . Let  $\omega_{\text{FS}}$  denote the Fubini–Study metric. The subvariety  $Y \cong \mathbb{P}^1$  is defined by  $z_2 = 0$ . Consider an increasing sequence  $(\varphi_j)_j$  in  $\text{PSH}(X, \omega_{\text{FS}})$ , whose potentials near  $(0, 0)$  are given by

$$\log |z_1|^2 \vee \left( k^{-1} \log |z_2|^2 \right) + O(1).$$

The pointwise restriction of these potentials to  $Y$  are given locally by

$$\log |z_1|^2 + O(1).$$

On the other hand, locally

$$\log |z_1|^2 \vee \left( k^{-1} \log |z_2|^2 \right) \rightarrow 0$$

almost everywhere as  $k \rightarrow \infty$ . So the trace operator is not continuous along the sequence  $(\varphi_j)_j$ .

`lma:rescommpullback`

**Lemma 8.2.1** *Let  $\pi: Z \rightarrow X$  be a proper bimeromorphic morphism with  $Z$  being a connected Kähler manifold. Assume that  $W$  (resp.  $Y$ ) be analytic subsets in  $Z$  (resp.  $X$ ) of codimension 1 such that the restriction  $\Pi: W \rightarrow Y$  of  $\pi$  is defined and is bimeromorphic, so that we have the following commutative diagram*

$$\begin{array}{ccccc} \tilde{W} & \longrightarrow & W & \hookrightarrow & Z \\ \downarrow \tilde{\Pi} & & \downarrow \Pi & & \downarrow \pi \\ \tilde{Y} & \longrightarrow & Y & \hookrightarrow & X. \end{array}$$

Then for any  $\varphi \in \text{QPSH}(X)$  with  $v(\varphi, Y) = 0$ , we have

$$\tilde{\Pi}^* \text{Tr}_Y(\varphi) \sim_P \text{Tr}_W(\pi^* \varphi). \quad (8.6)$$

`{eq:rescommpullback}`

**Proof** We first observe that by Zariski’s main theorem,  $v(\pi^* \varphi, W) = 0$ . So the right-hand side of (8.6) makes sense.

**Step 1.** Assume that  $T$  has analytic singularities. It suffices to apply [Example 8.1.1](#) to reformulate (8.6) as

$$\tilde{\Pi}^*(\varphi|_{\tilde{Y}}) \sim_P (\pi^*\varphi)|_{\tilde{W}}.$$

In fact, the strict equality holds, which is nothing but the functoriality of pullbacks.

**Step 2.** Next we handle the general case. Up to replacing  $\theta$  by  $\theta + \omega$  for some Kähler form  $\omega$  on  $X$ , we may assume that  $T$  is a Kähler current. Take a quasi-equisingular approximation  $(\varphi_j)_j$  of  $\varphi$  in  $\text{PSH}(X, \theta)$ . By [Corollary 7.1.2](#),  $(\pi^*\varphi_j)_j$  is a quasi-equisingular approximation of  $\pi^*\varphi$ . From Step 1, we know that for each  $j$ ,

$$\tilde{\Pi}^* \text{Tr}_Y(\varphi_j) \sim_P \text{Tr}_W(\pi^*\varphi_j).$$

Letting  $j \rightarrow \infty$ , we conclude (8.6) using [Proposition 8.2.2](#).  $\square$

prop:OT2

**Proposition 8.2.3** *Let  $\varphi \in \text{QPSH}(X)$  with  $v(\varphi, Y) = 0$ . Assume that  $Y$  is smooth. Then for any  $\lambda > 0$ , we have*

$$I(\lambda \text{Tr}_Y(\varphi)) \subseteq \text{Res}_Y I(\lambda\varphi). \quad (8.7)$$

{eq:OT}

**Proof** Take a Kähler form  $\omega$  on  $X$  such that  $\omega_\varphi$  is a Kähler current.

Let  $(\varphi_j)_j$  be a quasi-equisingular approximation of  $\varphi$  in  $\text{PSH}(X, \omega)$ .

By definition, for each  $j \geq 1$ , we get that

$$\text{Tr}_Y(\varphi) \leq_P \varphi_j|_Y.$$

For any  $\lambda' > \lambda > 0$ , we can find  $j > 0$  so that

$$I(\lambda' \varphi_j) \subseteq I(\lambda\varphi).$$

By [Theorem 1.4.5](#), we have

$$I(\lambda' \text{Tr}_Y(\varphi)) \subseteq I(\lambda' \varphi_j|_Y) \subseteq \text{Res}_Y I(\lambda' \varphi_j) \subseteq \text{Res}_Y I(\lambda\varphi).$$

Thanks to [Theorem 1.4.4](#), we conclude (8.7).  $\square$

Lastly, we turn our attention to global sections. For this we will need the following global Ohsawa–Takegoshi extension theorem for the trace operator:

thm: OT\_ext\_global

**Theorem 8.2.1** *Let  $L$  be a big line bundle on  $X$  and  $\theta$  is a closed real smooth  $(1, 1)$ -form on  $X$  representing  $c_1(L)$ . Suppose that  $\varphi \in \text{PSH}(X, \theta)$  and  $\theta_\varphi$  is a Kähler current. Assume that  $v(\varphi, Y) = 0$ . Let  $T$  be a holomorphic line bundle on  $X$ . Then there exists  $k_0$  such that for all  $k \geq k_0$  and  $s \in H^0(Y, T|_Y \otimes L|_Y^k \otimes I(k \text{Tr}_Y^\theta(\varphi)))$ , there exists an extension  $\tilde{s} \in H^0(X, T \otimes L^k \otimes I(k\varphi))$ .*

It is of interest to know if one could control the  $L^2$ -norm of  $\tilde{s}$  in the above result.

**Proof** Fix a Kähler form  $\omega$  on  $X$ . We may assume that  $Y \neq X$  and that  $\theta_\varphi \geq 3\delta\omega$  for some  $\delta > 0$ . Let  $(\varphi_j)_j$  be the decreasing quasi-equisingular approximation of  $\varphi$  in  $\text{PSH}(X, \theta)$ . We can assume that  $\theta_{\varphi_j} \geq 2\delta\omega$  for all  $j \geq 1$ . Also, there exists  $\epsilon_0 > 0$  such that  $\theta_{(1+\epsilon)\varphi_j} \geq \delta\omega$  for any  $\epsilon \in (0, \epsilon_0)$ . Take  $k_0 = k_0(\delta)$  as in [Theorem 1.8.1](#).



We fix  $k \geq k_0$  and  $s \in H^0(Y, T|_Y \otimes L|_Y^k \otimes \mathcal{I}(k \operatorname{Tr}_Y^\theta(\varphi)))$ . By [Theorem 1.4.4](#), there exists  $\epsilon \in (0, \epsilon_0)$  such that  $s \in H^0(Y, T|_Y \otimes L|_Y^k \otimes \mathcal{I}(k(1+\epsilon) \operatorname{Tr}_Y^\theta(\varphi)))$ .

Since  $\operatorname{Tr}_Y^\theta(\varphi) \leq \varphi_j|_Y$ , we obtain that  $s \in H^0(Y, T|_Y \otimes L|_Y^k \otimes \mathcal{I}(k(1+\epsilon)\varphi_j|_Y))$ . Due to [Theorem 1.8.1](#) there exists  $\tilde{s}_j \in H^0(X, T \otimes L^k \otimes \mathcal{I}(k(1+\epsilon)\varphi_j))$  such that  $\tilde{s}_j|_Y = s$ , for all  $j$ .

But by definition of quasi-equisingular approximation, we obtain that for high enough  $j$  the inclusion  $\mathcal{I}(k(1+\epsilon)\varphi_j) \subseteq \mathcal{I}(k\varphi)$  holds. As a result,  $\tilde{s}_j \in H^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi))$  for high enough  $j$ , finishing the argument.  $\square$

### 8.3 Restricted volumes

Let  $X$  be a connected projective manifold of dimension  $n$  and  $Y \subseteq X$  be a connected submanifold of dimension  $m$ . Consider a big line bundle  $L$  on  $X$ , a Hermitian metric  $h_0$  on  $L$  with  $\theta = c_1(L, h_0)$ . Let  $A$  be a very ample line bundle on  $X$ . Take a Hermitian metric  $h_A$  on  $A$  such that  $\omega = \operatorname{dd}^c h_A$  is a Kähler form.

Using the trace operator, one could prove the following generalization of [Theorem 7.3.1](#).

thm: rest\_volume

**Theorem 8.3.1** *Let  $h$  be a singular plurisubharmonic metric on  $L$  with  $v(\operatorname{dd}^c h, Y) = 0$ . Assume that*

$$\lim_{\epsilon \searrow 0} \left( \operatorname{Tr}_Y^{c_1(L|_Y) + \epsilon \omega} (c_1(L, h)) \right)^m > 0. \quad (8.8)$$

{eq: traceposmasscond2}

*Then for any holomorphic line bundle  $T$  on  $X$  we have that*

$$\int_Y \left( \operatorname{Tr}_Y^{c_1(L|_Y)} (c_1(L, h)) \right)^m = \lim_{k \rightarrow \infty} \frac{m!}{k^m} h^0 \left( Y, T|_Y \otimes L|_Y^k \otimes \operatorname{Res}_Y(\mathcal{I}(h^k)) \right). \quad (8.9)$$

{eq: DXmainrelative}

Recall that  $\operatorname{Res}_Y$  is defined in [Definition 1.4.5](#). Observe that by [Example 8.1.3](#), (8.8) implies that  $\operatorname{Tr}_Y^{c_1(L|_Y)}(c_1(L, h))$  is defined. So (8.9) is defined.

We will identify  $h$  with  $\varphi \in \operatorname{PSH}(X, \theta)$  as in (1.10).

We only need to consider the case  $Y \neq X$ , since otherwise, the result is proved in [Theorem 7.3.1](#). We will always assume  $Y \neq X$  in the sequel.

**Lemma 8.3.1** *There is  $\psi_Y \in \operatorname{QPSH}(X)$  with neat analytic singularities such that  $\{\psi_Y = -\infty\} = Y$  and in an open neighbourhood of  $Y$ , we have*

$$\psi_Y(x) = 2(n-m) \log \operatorname{dist}(x, Y) \quad (8.10)$$

{eq: Psi\_Y\_def}

*for some Riemannian distance function  $\operatorname{dist}(\cdot, Y)$ .*

See [Definition 1.6.1](#) for the definition of neat analytic singularities.

See [Fin22](#), Lemma 2.3] for the proof.

lma: IpsiY

**Lemma 8.3.2** *The multiplier ideal sheaf of  $\psi_Y$  can be calculated as*

$$\mathcal{I}(\psi_Y) = \mathcal{I}_Y. \quad (8.11) \quad \text{\texttt{\{eq:mis_psi\}}}$$

Moreover, given  $y \in Y$  and  $\epsilon > 0$ , for any germ  $f \in \mathcal{I}_{Y,y}$  we have

$$\int_U |f|^\epsilon e^{-\psi_Y} \omega^n < \infty, \quad (8.12) \quad \text{\texttt{\{eq:integrabilitypsiY\}}}$$

where  $U$  is an open neighbourhood of  $y$  in  $X$ .

In other words,  $\psi_Y$  has *log canonical singularities*.

**Proof** Since  $\psi_Y$  is locally bounded away from  $Y$ , it suffices to prove (8.11) along  $Y$ . Fix  $y \in Y$ , and we will verify (8.11) germ-wise at  $y$ .

Take an open neighbourhood  $U \subset X$  of  $y$  and a biholomorphic map  $F: U \rightarrow V \times W$ , where  $V$  is an open neighbourhood of  $y$  in  $Y$  and  $W$  is a connected open subset in  $\mathbb{C}^{n-m}$  containing  $0$ , such that  $F(Y \cap U) = V \times \{0\}$ . For any  $x \in U$ , write  $x_V, x_W$  for the two components of  $F(x)$  in  $V$  and  $W$  respectively. We denote the coordinates in  $\mathbb{C}^{n-m}$  as  $w_1, \dots, w_{n-m}$ .

Due to (8.10), after possibly shrinking  $U$ , we may assume that

$$\exp(-\psi_Y(x)) = |x_W|^{2m-2n} + \mathcal{O}(1)$$

for any  $x \in U \setminus Y$ .

Given  $f \in \mathcal{I}_{Y,y}$ , after shrinking  $U$ , we may assume that there exists  $g_1, \dots, g_{n-m} \in H^0(V \times W, \mathcal{O}_{V \times W})$  such that

$$f = \sum_{i=1}^{n-m} w_i g_i.$$

In order to verify  $f \in \mathcal{I}(\psi_Y)_y$ , it suffices to show  $w_i g_i \in \mathcal{I}((\sum_{i=1}^{n-m} |w_i|^2)^{m-n})_{F(y)}$ , which follows from Fubini's theorem. The proof of (8.12) is similar.

Conversely, take  $f \in \mathcal{I}(\psi_Y)$ , the similar application of Fubini's theorem shows that after possible shrinking  $U$ , we have  $f|_Y = 0$ . By Rückert's Nullstellensatz [GR84, Page 67], it follows that  $f \in \mathcal{I}_Y$ .  $\square$

\texttt{lem: analytic\_formula}

**Lemma 8.3.3** Assume that  $\varphi$  has analytic singularity type and  $\theta_u$  is a Kähler current. Suppose that  $\varphi|_Y \not\equiv -\infty$ . Then

$$\int_Y (\theta|_Y + \text{dd}^c \varphi|_Y)^m = \lim_{k \rightarrow \infty} \frac{m!}{k^m} \dim_{\mathbb{C}} \{s|_Y : s \in H^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi))\}. \quad (8.13) \quad \text{\texttt{\{eq:asymanasing\}}}$$

Recall that  $\mathcal{I}_\infty$  is defined in Definition 1.6.5.

**Proof** Suppose that  $\epsilon \in (0, 1)$  is small enough so that  $(1 - \epsilon)u \in \text{PSH}(X, \theta)$ .

Using Theorem 7.3.1 we can start to write the following sequence of inequalities:

$$\begin{aligned}
& \frac{1}{m!} \int_Y (\theta|_Y + \text{dd}^c \varphi|_Y)^m \\
&= \lim_{k \rightarrow \infty} \frac{1}{k^m} h^0(Y, T|_Y \otimes L|_Y^k \otimes \mathcal{I}(k\varphi|_Y)) \\
&\leq \lim_{k \rightarrow \infty} \frac{1}{k^m} \dim \{s|_Y : s \in H^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi))\} \quad \text{by Theorem 1.8.1} \\
&\leq \lim_{k \rightarrow \infty} \frac{1}{k^m} \dim \{s|_Y : s \in H^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi))\} \\
&\leq \lim_{k \rightarrow \infty} \frac{1}{k^m} \dim \{s|_Y : s \in H^0(X, T \otimes L^k \otimes \mathcal{I}_\infty((1-\epsilon)k\varphi))\} \quad \text{by Lemma 1.6.3} \\
&\leq \lim_{k \rightarrow \infty} \frac{1}{k^m} \dim_{\mathbb{C}} \{s \in H^0(Y, T|_Y \otimes L|_Y^k) : \log h^k(s, s) \leq (1-\epsilon)k\varphi|_Y\} \\
&\leq \lim_{k \rightarrow \infty} \frac{1}{k^m} h^0(Y, T|_Y \otimes L|_Y^k \otimes \mathcal{I}((1-\epsilon)k\varphi|_Y)) \\
&= \frac{1}{m!} \int_Y (\theta|_Y + (1-\epsilon)\text{dd}^c \varphi|_Y)^m \quad \text{by Theorem 7.3.1.}
\end{aligned}$$

Letting  $\epsilon \rightarrow 0$ , (8.13) follows from multi-linearity of the non-pluripolar product.  $\square$

prop: rest\_volume

**Proposition 8.3.1** *In the setting of Theorem 8.3.1, assume that  $\text{dd}^c h$  is a Kähler current. Then (8.9) holds.*

**Proof** Let  $(\varphi_j)_j$  a quasi-equisingular approximation of  $\varphi$  in  $\text{PSH}(X, \theta)$ . After possibly replacing  $(\varphi_j)_j$  by a subsequence, there exists  $\epsilon_0 \in (0, 1) \cap \mathbb{Q}$  such that  $\theta_{(1-\epsilon)^2 \varphi_j}$  and  $\theta_{(1-\epsilon) \varphi_j}$  are also Kähler currents for any  $\epsilon \in (0, \epsilon_0)$ .

We claim that for any  $j \geq 1$  and  $k \in \mathbb{N}$ , we have

$$\mathcal{I}_\infty((1-\epsilon)k\varphi_j) \cap \mathcal{I}(\psi_Y) \subseteq \mathcal{I}((1-\epsilon)^2 k\varphi_j + \psi_Y). \quad (8.14)$$

{eq: JcapI}

Take  $x \in X$ , and it suffices to argue (8.14) along the germ of  $x$ . Since  $\psi_Y$  is locally bounded outside  $Y$ , we may assume that  $x \in Y$ . Recall that by Lemma 8.3.2,  $\mathcal{I}(\psi_Y) = \mathcal{I}_Y$ .

Let  $f \in \mathcal{I}_\infty((1-\epsilon)k\varphi_j)_x \cap \mathcal{I}(\psi_Y)_x$ . Then there is an open neighbourhood  $U$  of  $x$  in  $X$  such that  $|f|^{2(1-\epsilon)} e^{-k(1-\epsilon)^2 \varphi_j} \leq C$  holds on  $U \setminus \{\varphi_j = -\infty\}$  for some  $C > 0$ , hence

$$\begin{aligned}
\int_U |f|^2 e^{-k(1-\epsilon)^2 \varphi_j - \psi_Y} \omega^n &= \int_U |f|^{2(1-\epsilon)} e^{-k(1-\epsilon)^2 \varphi_j} |f|^{2\epsilon} e^{-\psi_Y} \omega^n \\
&\leq C \int_U |f|^{2\epsilon} e^{-\psi_Y} \omega^n < \infty,
\end{aligned}$$

where the last inequality follows from Lemma 8.3.2. We have proved the claim (8.14).

Next we consider the following composition morphism of coherent sheaves on  $Y$ :

$$\text{Res}_Y \mathcal{I}_\infty((1-\epsilon)k\varphi_j) \hookrightarrow \frac{\mathcal{I}((1-\epsilon)^2 k\varphi_j)}{\mathcal{I}_\infty((1-\epsilon)k\varphi_j) \cap \mathcal{I}_Y} \rightarrow \frac{\mathcal{I}((1-\epsilon)^2 k\varphi_j)}{\mathcal{I}((1-\epsilon)^2 k\varphi_j + \psi_Y)}. \quad (8.15)$$

{eq: sheaf\_injection}

Here we have identified the coherent  $\mathcal{O}_X$ -modules supported on  $Y$  with coherent  $\mathcal{O}_Y$ -modules. Note that the target of (8.15) is also supported on  $Y$  as  $\psi_Y$  is locally bounded outside  $Y$ . We denote the coherent  $\mathcal{O}_Y$ -module whose pushforward to  $X$  gives  $\frac{\mathcal{I}((1-\epsilon)^2 k \varphi_j)}{\mathcal{I}((1-\epsilon)^2 k \varphi_j + \psi_Y)}$  by  $\mathcal{I}_{k,j}$ .

In (8.15), the first map is the inclusion and the second one is the obvious projection induced by (8.14). Although in general the second map fails to be injective, we observe that the composition is still injective as  $\mathcal{I}((1-\epsilon)^2 k \varphi_j + \psi_Y) \subseteq \mathcal{I}(\psi_Y) = \mathcal{I}_Y$ . Therefore, for any  $k \in \mathbb{N}$ , we have an injective morphism of coherent  $\mathcal{O}_Y$ -modules:

$$L_Y^k \otimes T|_Y \otimes \text{Res}_Y \mathcal{I}_\infty((1-\epsilon)k\varphi_j) \hookrightarrow L_Y^k \otimes T|_Y \otimes \mathcal{I}_{k,j}. \quad (8.16)$$

{eq:injLkTideal}

Using Theorem 7.3.1 we can start the following inequalities:

$$\begin{aligned} & \frac{1}{m!} \int_Y \left( \theta|_Y + \text{dd}^c \text{Tr}_Y^\theta(\varphi) \right)^m \\ &= \lim_{k \rightarrow \infty} \frac{1}{k^m} h^0(Y, T|_Y \otimes L_Y^k \otimes \mathcal{I}(k \text{Tr}_Y^\theta(\varphi))) \quad \text{by Theorem 7.3.1} \\ &\leq \lim_{k \rightarrow \infty} \frac{1}{k^m} h^0(Y, T|_Y \otimes L_Y^k \otimes \text{Res}_Y(\mathcal{I}(k\varphi))) \quad \text{by Theorem 1.4.5} \\ &\leq \overline{\lim}_{k \rightarrow \infty} \frac{1}{k^m} h^0(Y, T|_Y \otimes L_Y^k \otimes \text{Res}_Y(\mathcal{I}(k\varphi))) \\ &\leq \overline{\lim}_{k \rightarrow \infty} \frac{1}{k^m} h^0(Y, T|_Y \otimes L_Y^k \otimes \mathcal{I}(k\varphi_j)|_Y) \\ &\leq \overline{\lim}_{k \rightarrow \infty} \frac{1}{k^m} h^0(Y, T|_Y \otimes L_Y^k \otimes \mathcal{I}_\infty((1-\epsilon)k\varphi_j)|_Y) \quad \text{by Lemma 1.6.3} \\ &\leq \overline{\lim}_{k \rightarrow \infty} \frac{1}{k^m} h^0(Y, T|_Y \otimes L_Y^k \otimes \mathcal{I}_{k,j}) \quad \text{by (8.16)} \\ &\leq \overline{\lim}_{k \rightarrow \infty} \frac{1}{k^m} \dim_{\mathbb{C}} \left\{ s|_Y : s \in H^0 \left( X, T \otimes L^k \otimes \frac{\mathcal{I}((1-\epsilon)^2 k \varphi_j)}{\mathcal{I}((1-\epsilon)^2 k \varphi_j + \psi_Y)} \right) \right\} \\ &= \overline{\lim}_{k \rightarrow \infty} \frac{1}{k^m} \dim_{\mathbb{C}} \{ s|_Y : s \in H^0(X, T \otimes L^k \otimes \mathcal{I}((1-\epsilon)^2 k \varphi_j)) \} \quad (\text{see below}) \\ &= \frac{1}{m!} \int_Y \left( \theta|_Y + (1-\epsilon)^2 \text{dd}^c \varphi_j|_Y \right)^m \quad \text{by Lemma 8.3.3,} \end{aligned}$$

where in the penultimate line we used <sup>CDM17</sup>[CDM17, Theorem 1.1(6)] for  $q = 0$ . Letting  $\epsilon \rightarrow \infty$  and then  $j \rightarrow \infty$  the result follows.  $\square$

**Proof (Proof of Theorem 8.3.1)** Using Proposition 8.2.3 and Theorem 7.3.1 we obtain that

$$\begin{aligned} \int_Y \left( \theta|_Y + \text{dd}^c \text{Tr}_Y^\theta(\varphi) \right)^m &= \lim_{k \rightarrow \infty} \frac{m!}{k^m} h^0(Y, T|_Y \otimes L|_Y^k \otimes \mathcal{I}(k \text{Tr}_Y^\theta(\varphi))) \\ &\leq \lim_{k \rightarrow \infty} \frac{m!}{k^m} h^0(Y, T|_Y \otimes L|_Y^k \otimes \text{Res}_Y(\mathcal{I}(k\varphi))). \end{aligned}$$

{eq:DX\_cor}

Now we address the other direction in (8.9). Let  $\phi \in H^0(X, A)$  be a section that does not vanish identically on  $Y$ . Such  $\phi$  exists since  $A$  is very ample.

We fix  $k_0 \in \mathbb{N}$ . For any  $k \geq 0$ , we have that  $k = qk_0 + r$  with  $q, r \in \mathbb{N}$  and  $r \in \{0, \dots, k_0 - 1\}$ . Also, we have an injective linear map

$$H^0(Y, T|_Y \otimes L|_Y^k \otimes \mathcal{I}(k\varphi|_Y)) \xrightarrow{\cdot \phi^{\otimes q}} H^0(Y, T|_Y \otimes L|_Y^k \otimes A|_Y^q \otimes \mathcal{I}(k\varphi|_Y)).$$

Therefore,

$$\begin{aligned} &\overline{\lim}_{k \rightarrow \infty} \frac{m!}{k^m} h^0(Y, T|_Y \otimes L|_Y^k \otimes \mathcal{I}(k\varphi|_Y)) \\ &\leq \overline{\lim}_{k \rightarrow \infty} \frac{m!}{k^m} h^0(Y, T|_Y \otimes L|_Y^k \otimes A|_Y^q \otimes \mathcal{I}(k\varphi|_Y)) \\ &= \frac{1}{k_0^m} \overline{\lim}_{q \rightarrow \infty} \frac{m!}{q^m} h^0(Y, T|_Y \otimes L|_Y^{qk_0} \otimes A|_Y^q \otimes L|_Y^r \otimes \mathcal{I}(k\varphi|_Y)) \\ &\leq \frac{1}{k_0^m} \overline{\lim}_{q \rightarrow \infty} \frac{m!}{q^m} h^0(Y, T|_Y \otimes L|_Y^{qk_0} \otimes A|_Y^q \otimes L|_Y^r \otimes \mathcal{I}(k_0 q \varphi|_Y)) \\ &= \int_Y \left( \theta|_Y + k_0^{-1} \omega|_Y + \text{dd}^c \text{Tr}_Y^{\theta + k_0^{-1} \omega}(\varphi) \right)^m \\ &= \int_Y \left( \theta|_Y + k_0^{-1} \omega|_Y + \text{dd}^c \text{Tr}_Y^\theta(\varphi) \right)^m, \end{aligned}$$

where in the fourth line we have used that  $k_0 q \leq k$  and in the last line we have used [Proposition 8.3.1](#) for the big line bundle  $L^{k_0} \otimes A$ , the Kähler current  $k_0 \theta_u - \text{dd}^c \log g = k_0 \theta_u + \omega$ , and twisting bundle  $T \otimes L^r$ . Letting  $k_0 \rightarrow \infty$ , we conclude that

$$\overline{\lim}_{k \rightarrow \infty} \frac{m!}{k^m} h^0(Y, T|_Y \otimes L|_Y^k \otimes \mathcal{I}(k\varphi|_Y)) \leq \int_Y \left( \theta|_Y + \text{dd}^c \text{Tr}_Y^\theta(\varphi) \right)^m.$$

thm: rest\_volume\_2

**Theorem 8.3.2** *Let  $\varphi \in \text{PSH}(X, \theta)$  such that  $v(\varphi, Y) = 0$ . Assume that  $\theta_\varphi$  is a Kähler current. Then*

$$\int_Y \left( \theta|_Y + \text{dd}^c \text{Tr}_Y^\theta(\varphi) \right)^m = \lim_{k \rightarrow \infty} \frac{m!}{k^m} \dim_{\mathbb{C}} \{s|_Y : s \in H^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi))\}.$$

**Proof** This is a consequence of [Theorem 7.3.1](#), [Theorem 8.2.1](#) and [Theorem 8.3.1](#):

$$\begin{aligned}
\int_Y \left( \theta|_Y + \text{dd}^c \text{Tr}_Y^\theta(\varphi) \right)^m &= \lim_{k \rightarrow \infty} \frac{m!}{k^m} h^0(Y, T|_Y \otimes L|_Y^k \otimes I(k \text{Tr}_Y^\theta(\varphi))) \\
&\leq \lim_{k \rightarrow \infty} \frac{m!}{k^m} \dim_{\mathbb{C}} \{s|_Y : s \in H^0(X, T \otimes L^k \otimes I(k\varphi))\} \\
&\leq \lim_{k \rightarrow \infty} \frac{m!}{k^m} \dim_{\mathbb{C}} \{s|_Y : s \in H^0(X, T \otimes L^k \otimes I(k\varphi))\} \\
&\leq \lim_{k \rightarrow \infty} \frac{m!}{k^m} h^0(Y, T|_Y \otimes L|_Y^k \otimes I(k\varphi)|_Y) \\
&= \int_Y \left( \theta|_Y + \text{dd}^c \text{Tr}_Y^\theta(\varphi) \right)^m.
\end{aligned}$$

*Remark 8.3.1* One could also show that when (8.8) fails, the right-hand side of (8.9) is 0. See [DX24].

## 8.4 Analytic Bertini theorems

Let  $X$  be a connected projective manifold of dimension  $n \geq 1$ .

The analytic Bertini theorem handles the restriction along a generic subvariety.

thm:Bert

**Theorem 8.4.1** *Let  $\varphi \in \text{QPSH}(X)$ . Let  $p: X \rightarrow \mathbb{P}^N$  be a morphism ( $N \geq 1$ ). Define*

$$\mathcal{G} := \{H \in |\mathcal{O}_{\mathbb{P}^N}(1)| : H' := H \cap X \text{ is smooth and } I(\varphi|_{H'}) = \text{Res}_{H'}(I(\varphi))\}.$$

*Then  $\mathcal{G} \subseteq |\mathcal{O}_{\mathbb{P}^N}(1)|$  is co-pluripolar.*

Recall that co-pluripolar sets are defined in Definition 1.1.4. We adopt the convention that  $I(-\infty) = 0$ .

*Remark 8.4.1* Here and in the sequel, we slightly abuse the notation by writing  $H \cap X$  for  $p^{-1}H$ , the scheme-theoretic inverse image of  $H$ . In other words,  $H \cap X := H \times_{\mathbb{P}^N} X$ .

By definition, any  $H \in |\mathcal{O}_{\mathbb{P}^N}(1)|$  such that  $p^{-1}H = \emptyset$  lies in  $\mathcal{G}$ .

**Proof** Take an ample line bundle  $L$  with a smooth Hermitian metric  $h$  such that  $c_1(L, h) + \text{dd}^c \varphi \geq 0$ , where  $c_1(L, h)$  is the first Chern form of  $(L, h)$ , namely the curvature form of  $h$ . We introduce  $\Lambda := |\mathcal{O}_{\mathbb{P}^N}(1)|$  to simplify our notations.

**Step 1.** We prove that the following set is co-pluripolar:

$$\begin{aligned}
\mathcal{G}_L := \{H \in \Lambda : H \cap X \text{ is smooth and } H^0(H \cap X, \omega_{H \cap X} \otimes L|_{H \cap X} \otimes I(\varphi|_{H \cap X})) = \\
H^0(H \cap X, \omega_{H \cap X} \otimes L|_{H \cap X} \otimes \text{Res}_{H \cap X}(I(\varphi)))\}.
\end{aligned}$$

Here  $\omega_{H \cap X}$  denotes the dualizing sheaf of  $H \cap X$ .

Let  $U \subseteq \Lambda \times X$  be the closed subvariety whose  $\mathbb{C}$ -points correspond to pairs  $(H, x) \in \Lambda \times X$  with  $p(x) \in H$ . Let  $\pi_1: U \rightarrow \Lambda$  be the natural projection. We may assume that  $\pi_1$  is surjective, as otherwise there is nothing to prove.

Observe that  $U$  is a local complete intersection scheme by *Krulls Hauptidealsatz* and *a fortiori* a Cohen–Macaulay scheme. It follows from miracle flatness [Mat89, Theorem 23.1] that the natural projection  $\pi_2: U \rightarrow X$  is flat. As the fibers of  $\pi_2$  over closed points of  $X$  are isomorphic to  $\mathbb{P}^{N-1}$ , it follows that  $\pi_2$  is smooth. Thus,  $U$  is smooth as well. Moreover, observe that

$$I(\pi_2^* \varphi) = \pi_2^* I(\varphi) \quad (8.17)$$

{eq:pi2pullvarphiItem1}

by [Proposition 1.4.5](#).

In the following, we will construct pluripolar sets  $\Sigma_1 \subseteq \Sigma_2 \subseteq \Sigma_3 \subseteq \Sigma_4 \subseteq \Lambda$  such that the behaviour of  $\pi_1$  is improved successively on the complement of  $\Sigma_i$ .

**Step 1.1.** The usual Bertini theorem shows that there is a proper Zariski closed set  $\Sigma_1 \subseteq \Lambda$  such that  $\pi_1$  has smooth fibres outside  $\Sigma_1$ . Enlarging  $\Sigma_1$ , we could guarantee that  $\pi_1$  and  $I(\pi_2^* \varphi)$  are both flat outside  $\Sigma_1$ . See [EGAIV-2, Théorème 6.9.1]. Then after further enlarging  $\Sigma_1$  so that  $H$  avoids all associated points of  $\mathcal{O}_X/I(\varphi)$ , for all  $H \in \Lambda \setminus \Sigma_1$ . Let  $\pi_{1,H}$  denote the fibre of  $\pi_1$  at  $H$  and write  $i_H: \pi_{1,H} \rightarrow U$  for the inclusion morphism. We arrive at

$$\text{Res}_{\pi_{1,H}}(I(\pi_2^* \varphi)) = i_H^* I(\pi_2^* \varphi)$$

for all  $H \in \Lambda \setminus \Sigma_1$ .<sup>1</sup>

**Step 1.2.** By Grauert’s coherence theorem,

$$\mathcal{F}^i := R^i \pi_{1*} (\omega_{U/\Lambda} \otimes \pi_2^* L \otimes I(\pi_2^* \varphi))$$

is coherent for all  $i$ . Here  $\omega_{U/\Lambda}$  denotes the relative dualizing sheaf of the morphism  $U \rightarrow \Lambda$ . Thus, there is a proper Zariski closed set  $\Sigma_2 \subseteq \Lambda$  such that

- (1)  $\Sigma_2 \supseteq \Sigma_1$ .
- (2) The  $\mathcal{F}^i$ ’s are locally free outside  $\Sigma_2$ .

We write  $\mathcal{F} = \mathcal{F}^0$ . By cohomology and base change [Har77, Theorem III.12.11], for any  $H \in \Lambda \setminus \Sigma_2$ , the fibre  $\mathcal{F}|_H$  of  $\mathcal{F}$  is given by

$$\mathcal{F}|_H = H^0(\pi_{1,H}, \omega_{U/\Lambda}|_{\pi_{1,H}} \otimes \pi_2^* L|_{\pi_{1,H}} \otimes \text{Res}_{\pi_{1,H}}(I(\pi_2^* \varphi))).$$

**Step 1.3.** In order to proceed, we need to make use of the Hodge metric  $h_{\mathcal{H}}$  on  $\mathcal{F}$  defined in [HPS18]. We briefly recall its definition in our setting. By [HPS18, Section 22], we can find a proper Zariski closed set  $\Sigma_3 \subseteq \Lambda$  such that

- (1)  $\Sigma_3 \supseteq \Sigma_2$ ,
- (2)  $\pi_1$  is smooth outside  $\Sigma_3$ ,
- (3) both  $\mathcal{F}$  and  $\pi_{1*} (\omega_{U/\Lambda} \otimes \pi_2^* L) / \mathcal{F}$  are locally free outside  $\Sigma_3$ , and
- (4) for each  $i$ ,

$$R^i \pi_{1*} (\omega_{U/\Lambda} \otimes \pi_2^* L)$$

is locally free outside  $\Sigma_3$ .

---

<sup>1</sup> This subtle point was overlooked in the proof of [Xia22a].

Then for any  $H \in \Lambda \setminus \Sigma_3$ ,

$$H^0(H \cap X, \omega_{H \cap X} \otimes L|_{H \cap X} \otimes \mathcal{I}(\varphi|_{H \cap X})) \subseteq \mathcal{F}|_H \subseteq H^0(H \cap X, \omega_{H \cap X} \otimes L|_{H \cap X}).$$

See [HPS18, Lemma 22.1].

Now we can give the definition of the Hodge metric on  $\Lambda \setminus \Sigma_3$ . Given any  $H \in \Lambda \setminus \Sigma_3$ , any  $\alpha \in \mathcal{F}|_H$ , the Hodge metric is defined as

$$h_{\mathcal{H}}(\alpha, \alpha) := \int_{X \cap H} |\alpha|_h^2 e^{-\varphi} \in [0, \infty].$$

Observe that  $h_{\mathcal{H}}(\alpha, \alpha) < \infty$  if and only if  $\alpha \in H^0(H \cap X, \omega_{H \cap X} \otimes L|_{H \cap X} \otimes \mathcal{I}(\varphi|_{H \cap X}))$ . Moreover,  $h_{\mathcal{H}}(\alpha, \alpha) > 0$  if  $\alpha \neq 0$ . It is shown in [HPS18] (c.f. [PT18, Theorem 3.3.5]) that  $h_{\mathcal{H}}$  is indeed a singular Hermitian metric, and it extends to a positive metric on  $\mathcal{F}$ .

**Step 1.4.** The determinant  $\det h_{\mathcal{H}}$  is singular at all  $H \in \Lambda \setminus \Sigma_3$  such that

$$H^0(H \cap X, \omega_{H \cap X} \otimes L|_{H \cap X} \otimes \mathcal{I}(\varphi|_{H \cap X})) \neq \mathcal{F}|_H.$$

As the map  $\pi_2$  is smooth, we have  $\pi_2^* \mathcal{I}(\varphi) = \mathcal{I}(\pi_2^* \varphi)$  by Proposition 1.4.5. Under the identification  $\pi_{1,H} \cong H \cap X$ , we have

$$\text{Res}_{\pi_{1,H}} (\pi_2^* \mathcal{I}(\varphi)) \cong \text{Res}_{H \cap X} (\mathcal{I}(\varphi)).$$

Thus, we have the following inclusions:

$$\begin{aligned} & H^0(H \cap X, \omega_{H \cap X} \otimes L|_{H \cap X} \otimes \mathcal{I}(\varphi|_{H \cap X})) \\ & \subseteq H^0(H \cap X, \omega_{H \cap X} \otimes L|_{H \cap X} \otimes \text{Res}_{H \cap X}(\mathcal{I}(\varphi))), \end{aligned}$$

the right-hand side being  $\mathcal{F}|_H$ .

Recall that the first inclusion follows from Theorem 1.4.5. Hence,  $\det h_{\mathcal{H}}$  is singular at all  $H \in |\mathcal{O}_{\mathbb{P}^N}(1)| \setminus \Sigma_3$  such that

$$\begin{aligned} & H^0(H \cap X, \omega_{H \cap X} \otimes L|_{H \cap X} \otimes \mathcal{I}(\varphi|_{H \cap X})) \\ & \neq H^0(H \cap X, \omega_{H \cap X} \otimes L|_{H \cap X} \otimes \text{Res}_{H \cap X}(\mathcal{I}(\varphi))). \end{aligned}$$

Let  $\Sigma_4$  be the union of  $\Sigma_3$  and the set of all such  $H$ . Since the Hodge metric  $h_{\mathcal{H}}$  is positive ([PT18, Theorem 3.3.5] and [HPS18, Theorem 21.1]), its determinant  $\det h_{\mathcal{H}}$  is also positive ([Rau15, Proposition 1.3] and [HPS18, Proposition 25.1]), it follows that  $\Sigma_4$  is pluripolar. As a consequence,  $\mathcal{G}_L$  is co-pluripolar.

**Step 2.**

Fix an ample invertible sheaf  $S$  on  $X$ . The same result holds with  $L \otimes S^{\otimes a}$  in place of  $L$ . Thus, the set

$$A := \bigcap_{a=0}^{\infty} \mathcal{G}_{L \otimes S^{\otimes a}}$$



is co-pluripolar. For each  $H \in W$  such that  $X \cap H$  is smooth and  $\mathcal{I}(\varphi|_{X \cap H}) \neq \text{Res}_{H \cap X}(\mathcal{I}(\varphi))$ , let  $\mathcal{K}$  be the following cokernel:

$$0 \rightarrow \mathcal{I}(\varphi|_{X \cap H}) \rightarrow \text{Res}_{H \cap X}(\mathcal{I}(\varphi)) \rightarrow \mathcal{K} \rightarrow 0.$$

By Serre vanishing theorem, taking  $a$  large enough, we may guarantee that

$$H^1(X \cap H, \omega_{X \cap H} \otimes (L \otimes S^{\otimes a})|_{X \cap H} \otimes \mathcal{I}(\varphi|_{X \cap H})) = 0$$

and

$$H^0(X \cap H, \omega_{X \cap H} \otimes (L \otimes S^{\otimes a})|_{X \cap H} \otimes \mathcal{K}) \neq 0.$$

Then

$$\begin{aligned} H^0(X \cap H, \omega_{X \cap H} \otimes (L \otimes S^{\otimes a})|_{X \cap H} \otimes \mathcal{I}(\varphi|_{X \cap H})) &\neq \\ H^0(X \cap H, \omega_{X \cap H} \otimes (L \otimes S^{\otimes a})|_{X \cap H} \otimes \text{Res}_{H \cap X}(\mathcal{I}(\varphi))) &. \end{aligned}$$

Thus,  $H \notin A$ . We conclude that  $\mathcal{G}$  is co-pluripolar.  $\square$

In the sequel of this section, we fix a base-point free linear system  $\Lambda$  on  $X$ .

cor:qpshtgeneralres

**Corollary 8.4.1** *Let  $\varphi \in \text{QPSH}(X)$ . Then there is a co-pluripolar subset  $\Lambda' \subseteq \Lambda$  such that  $\varphi|_H \not\equiv -\infty$  for any  $H \in \Lambda'$ .*

**Proof** This follows immediately from [Theorem 8.4.1](#).  $\square$

cor:ABTfortrace

**Corollary 8.4.2** *Assume that  $n \geq 2$ . Let  $\varphi \in \text{QPSH}(X)$ . Then there is a co-pluripolar set  $\Lambda' \subseteq \Lambda$  such that any  $H \in \Lambda'$  is connected and smooth,  $\nu(\varphi, H) = 0$  and we have*

$$\text{Tr}_H(\varphi) \sim_{\mathcal{I}} \varphi|_H.$$

The assumption  $n \geq 2$  is only to guarantee that a general element  $H \in \Lambda$  is connected, since we developed most of our theories only in this case.

**Proof** First observe that the set  $\{x \in X : \nu(\varphi, x) > 0\}$  is a countable union of proper analytic subsets by [Theorem 1.4.1](#). It follows that a very general element in  $\Lambda$  is not contained in this set.

Fix an ample line bundle  $L$  so that there is a smooth psh metric  $h_L$  such that  $c_1(L, h_L) + \text{dd}^c \varphi$  is a Kähler current. Thanks to [Theorem 8.4.1](#), we can find a co-pluripolar set  $\Lambda' \subseteq \Lambda$  such that each  $H \in \Lambda'$  satisfies the following:

- (1)  $H$  is smooth;
- (2)  $\nu(\varphi, H) = 0$ ;
- (3)  $\mathcal{I}(k\varphi|_H) = \text{Res}_H(\mathcal{I}(\varphi))$  for all  $k > 0$ .

It follows from [Theorem 8.3.1](#) and [Theorem 7.3.1](#) that

$$\int_H \left( c_1(L, h_L)|_H + \text{dd}^c \text{Tr}_Y^{c_1(L, h_L)}(\varphi) \right)^{n-1} = \int_H (c_1(L, h_L)|_H + \text{dd}^c \varphi|_H)^{n-1}.$$

Since  $\varphi|_H \leq \text{Tr}_Y(\varphi)$  by [Proposition 8.1.3](#), our assertion follows.  $\square$

lma:posmasscurrres

**Lemma 8.4.1** Assume that  $n \geq 2$ . Let  $T$  be a closed positive  $(1, 1)$ -current on  $X$  with  $\int_X T^n > 0$ . Then there is a co-pluripolar set  $\Lambda' \subseteq \Lambda$  such that any  $H \in \Lambda'$  is connected and smooth,  $T|_H$  is well-defined and satisfies

$$\int_H T|_H^{n-1} > 0.$$

**Proof** Write  $T = \theta_\varphi$  for some smooth closed real  $(1, 1)$ -form  $\theta$  on  $X$  and  $\varphi \in \text{PSH}(X, \theta)_{>0}$ . Thanks to [Lemma 2.3.2](#), we can find  $\psi \in \text{PSH}(X, \theta)$  such that  $\theta_\psi$  is a Kähler current and  $\psi \leq \varphi$ . By [Corollary 8.4.1](#), we can find a co-pluripolar set  $\Lambda' \subseteq \Lambda$  such that each  $H \in \Lambda'$  satisfies:

- (1)  $H$  is smooth and connected;
- (2) the restriction  $\psi|_H$  is not identically  $-\infty$ .

Therefore,  $\psi|_H \leq \varphi|_H$  are two potentials in  $\text{PSH}(H, \theta|_H)$  for any  $H \in \Lambda'$ . Our assertion follows from [Theorem 2.3.2](#).  $\square$

cor:tracegeneralwelldef

**Corollary 8.4.3** Assume that  $n \geq 2$ . Let  $T$  be a closed positive  $(1, 1)$ -current on  $X$  with  $\text{vol } T > 0$ . Then there is a co-pluripolar set  $\Lambda' \subseteq \Lambda$  such that any  $H \in \Lambda'$  is connected and smooth, and  $\text{Tr}_H^{[T]|_H}(T)$  is well-defined.

**Proof** This follows from [Example 8.1.3](#), [Corollary 8.4.2](#) and [Lemma 8.4.1](#).  $\square$

**Proposition 8.4.1** Assume that  $n \geq 2$ . Let  $\varphi, \psi \in \text{QPSH}(X)$ . Assume that  $\varphi \leq_P \psi$ . Then there is a co-pluripolar set  $\Lambda' \subseteq \Lambda$  such that any  $H \in \Lambda'$  is connected and smooth, and  $\varphi|_H \leq_P \psi|_H$ .

**Proof** Thanks to [Lemma 6.1.2](#), we may replace  $\varphi$  by  $\varphi \vee \psi$  and assume that  $\varphi \sim_P \psi$ . It suffices to show that  $\varphi|_H \sim \psi|_H$ .

Take a smooth closed real  $(1, 1)$ -form  $\theta$  on  $X$  so that  $\varphi, \psi \in \text{PSH}(X, \theta)_{>0}$ . It suffices to compare  $\varphi$  and  $\psi$  with  $P_\theta[\varphi]$ , so without loss of generality, we may assume that  $\psi$  is a model potential in  $\text{PSH}(X, \theta)_{>0}$ . Up to adding a constant to  $\varphi$ , we may then assume that  $\varphi \leq \psi$ . It follows from [Lemma 2.3.1](#) that we can find a sequence  $(\eta_j)_j$  in  $\text{PSH}(X, \theta)_{>0}$  such that

$$j^{-1}\eta_j + (1 - j^{-1})\psi \leq \varphi$$

for all  $j \geq 2$ . By [Corollary 8.4.1](#), [Lemma 8.4.1](#), we can find a co-pluripolar set  $\Lambda' \subseteq \Lambda$  such that any  $H \in \Lambda'$  satisfies:

- (1)  $H$  is smooth and connected;
- (2)  $\eta_j|_H \in \text{PSH}(H, \theta|_H)_{>0}$  for all  $j \geq 2$  and  $\psi|_H \in \text{PSH}(H, \theta|_H)_{>0}$ .

Therefore, taking [Proposition 3.1.5](#) into account, we arrive at

$$j^{-1}P_{\theta|_H}[\eta_j|_H] + (1 - j^{-1})P_{\theta|_H}[\psi|_H] \leq P_{\theta|_H}[\varphi|_H]$$

for all  $j \geq 2$ . Letting  $j \rightarrow \infty$ , we conclude that

$$P_{\theta|_H}[\psi|_H] \leq P_{\theta|_H}[\varphi|_H]$$

and hence  $\psi|_H \leq_P \varphi|_H$ .  $\square$

lma:Igoodrest

**Lemma 8.4.2** Assume that  $n \geq 2$ . Let  $\theta$  be a closed smooth  $(1, 1)$ -form on  $X$  representing a big cohomology class and  $(\varphi_j)_j$  be a decreasing sequence in  $\text{PSH}(X, \theta)$ . Assume that  $\varphi \in \text{PSH}(X, \theta)$  and  $\varphi_j \xrightarrow{d_S} \varphi$ . Then there is a co-pluripolar set  $\Lambda' \subseteq \Lambda$  such that any  $H \in \Lambda'$  is connected and smooth,  $\varphi_j|_H \not\equiv -\infty$  for all  $j \geq 1$ ,  $\varphi|_H \not\equiv -\infty$ , and

$$\varphi_j|_H \xrightarrow{d_S} \varphi|_H.$$

**Proof** By [Corollary 6.2.7](#), we may assume that  $\varphi \in \text{PSH}(X, \theta)_{>0}$ . Using [Lemma 2.3.1](#), we could find a decreasing sequence  $(\epsilon_j)_j$  in  $(0, 1)$  with limit 0 and  $\eta_j \in \text{PSH}(X, \theta)_{>0}$  such that  $\eta_j \leq \varphi_j$  and

$$\epsilon_j \eta_j + (1 - \epsilon_j) \varphi_j \leq \varphi.$$

By [Corollary 8.4.1](#), [Lemma 8.4.1](#), we can find a co-pluripolar set  $\Lambda' \subseteq \Lambda$  such that any  $H \in \Lambda'$  satisfies:

- (1)  $H$  is smooth and connected;
- (2)  $\eta_j|_H \in \text{PSH}(H, \theta|_H)_{>0}$  for all  $j \geq 1$  and  $\varphi|_H \in \text{PSH}(H, \theta|_H)_{>0}$ .

Therefore, taking [Proposition 3.1.5](#) into account, we arrive at

$$\epsilon_j P_{\theta|_H}[\eta_j|_H] + (1 - \epsilon_j) P_{\theta|_H}[\varphi_j|_H] \leq P_{\theta|_H}[\varphi|_H].$$

Letting  $j \rightarrow \infty$ , we get

$$\lim_{j \rightarrow \infty} P_{\theta|_H}[\varphi_j|_H] \leq P_{\theta|_H}[\varphi|_H].$$

By [Theorem 2.3.2](#) and [Proposition 3.1.8](#), we conclude that

$$\lim_{j \rightarrow \infty} \int_H (\theta|_H + \text{dd}^c \varphi_j|_H)^{n-1} = \int_H (\theta|_H + \text{dd}^c \varphi|_H)^{n-1}.$$

Therefore, using [Corollary 6.2.4](#), we conclude that  $\varphi_j|_H \xrightarrow{d_S} \varphi|_H$ .  $\square$

**Corollary 8.4.4** Assume that  $n \geq 2$ . Let  $\varphi \in \text{QPSH}(X)$  be an  $\mathcal{I}$ -good potential. Then there is a co-pluripolar set  $\Lambda' \subseteq \Lambda$  such that any  $H \in \Lambda'$  satisfies:

- (1)  $H$  is connected and smooth;
- (2)  $\varphi|_H \in \text{PSH}(X, \theta|_H)$  is  $\mathcal{I}$ -good;
- (3)  $v(\varphi, H) = 0$ ;
- (4)  $\text{Tr}_H \varphi \sim_P \varphi|_H$ .

Furthermore, if  $\theta$  is a closed smooth real  $(1, 1)$ -form on  $X$  such that  $\varphi \in \text{PSH}(X, \theta)_{>0}$ , then we could further guarantee that  $\text{Tr}_H(\varphi)$  has a representative  $\text{Tr}_H(\varphi) \in \text{PSH}(H, \theta|_H)_{>0}$  for all  $H \in \Lambda'$ .

***Proof*** This is a consequence of Lemma 8.4.2, Theorem 7.1.1, Corollary 8.4.2 and Corollary 8.4.3.  $\square$

## Chapter 9

### Test curves

chap:testcurve

In this chapter, we develop the theory of test curves. Roughly speaking, a test curve is a concave curve of model potentials. In [Section 9.2](#), we will prove the Ross–Witt Nyström correspondence, through which the test curves are related to geodesic rays in the space of quasi-plurisubharmonic functions. In [Section 9.4](#), we define operations on test curves, anticipating applications in non-Archimedean pluripotential theory in [Chapter 13](#).

#### 9.1 The notion of test curves

Let  $X$  be a connected compact Kähler manifold of dimension  $n$  and  $\theta$  be a smooth closed real  $(1, 1)$ -form on  $X$  representing a big cohomology class.

def:testcur

**Definition 9.1.1** A *test curve*  $\Gamma$  in  $\text{PSH}(X, \theta)$  consists of a real number  $\Gamma_{\max}$  together with a map  $(-\infty, \Gamma_{\max}) \rightarrow \text{PSH}(X, \theta)$  denoted by  $\tau \mapsto \Gamma_\tau$  satisfying the following conditions:

- (1) The map  $\tau \mapsto \Gamma_\tau$  is concave and decreasing;
- (2) each  $\Gamma_\tau$  is a model potential;
- (3) the potential

$$\Gamma_{-\infty} := \sup_{\tau < \Gamma_{\max}}^* \Gamma_\tau \quad (9.1)$$

{eq:Gammaminf}

satisfies

$$\int_X (\theta + \text{dd}^c \Gamma_{-\infty})^n > 0.$$

Let  $\phi \in \text{PSH}(X, \theta)_{>0}$  be a model potential. The set of test curves  $\Gamma$  with  $\Gamma_{-\infty} = \phi$  is denoted by  $\text{TC}(X, \theta; \phi)$ .

The union of all  $\text{TC}(X, \theta; \phi)$ 's for various model potentials  $\phi \in \text{PSH}(X, \theta)_{>0}$  is denoted by  $\text{TC}(X, \theta)_{>0}$ .

By (2),  $\sup_X \Gamma_\tau = 0$  for each  $\tau < \Gamma_{\max}$ . So  $\Gamma_{-\infty} \in \text{PSH}(X, \theta)$  by [Proposition 1.2.1](#). Moreover,  $\Gamma_{-\infty}$  is a model potential by [Proposition 3.1.9](#).

rmk:extendtestcur

*Remark 9.1.1* Sometimes it is convenient to extend  $\Gamma_\tau$  to  $\tau \geq \Gamma_{\max}$  as well. This can be done as follows: for  $\tau > \Gamma_{\max}$ , we set  $\Gamma_\tau \equiv -\infty$ . For  $\tau = \Gamma_{\max}$ , we set

$$\Gamma_\tau := \inf_{\tau' < \Gamma_{\max}} \Gamma_{\tau'} \in \text{PSH}(X, \theta).$$

We will always make this extension in the sequel.

Recall that according to our general principle, we only talk about model potentials when a potential has positive mass. Fortunately, this principle is not violated in the above definition, as shown below:

lma:testcurvposmass

**Lemma 9.1.1** *Assume that  $\Gamma \in \text{TC}(X, \theta)_{>0}$ . Then for each  $\tau < \Gamma_{\max}$ , we have*

$$\int_X (\theta + \text{dd}^c \Gamma_\tau)^n > 0. \quad (9.2) \quad \{\text{eq:dalethtauposmass}\}$$

**Proof** Fix  $\tau \in (-\infty, \Gamma_{\max})$ .

By assumption,  $\Gamma_{-\infty}$  has positive mass. By [Corollary 2.3.1](#), we have

$$\int_X \theta_{\Gamma_{-\infty}}^n = \lim_{\tau \rightarrow -\infty} \int_X \theta_{\Gamma_\tau}^n.$$

In particular, for a sufficiently small  $\tau_0 < \tau$ , we have

$$\int_X \theta_{\Gamma_{\tau_0}}^n > 0.$$

Now take  $\tau' \in (\tau, \Gamma_{\max})$  and  $t \in (0, 1)$  so that

$$\tau = (1 - t)\tau' + t\tau_0.$$

From the concavity of  $\Gamma$ , we find that

$$\Gamma_\tau \geq (1 - t)\Gamma_{\tau'} + t\Gamma_{\tau_0}.$$

By [Theorem 2.3.2](#),

$$\int_X \theta_{\Gamma_\tau}^n \geq \int_X \theta_{(1-t)\Gamma_{\tau'} + t\Gamma_{\tau_0}}^n \geq t^n \int_X \theta_{\Gamma_{\tau_0}}^n > 0$$

and (9.2) follows.  $\square$

prop:testcurvmasslogconc

**Proposition 9.1.1** *Let  $\Gamma \in \text{TC}(X, \theta)_{>0}$ . Then the map*

$$[-\infty, \Gamma_{\max}) \rightarrow \mathbb{R}, \quad \tau \mapsto \log \int_X \theta_{\Gamma_\tau}^n$$

*is concave and continuous.*

**Proof** The concavity of this function follows from [Theorem 2.3.3](#) and [Theorem 2.3.2](#). The continuity at  $-\infty$  is a consequence of [Corollary 2.3.1](#).  $\square$

def:relatetestcurv

**Definition 9.1.2** Let  $\phi \in \text{PSH}(X, \theta)_{>0}$  be a model potential.

A test curve  $\Gamma \in \text{TC}(X, \theta; \phi)$  is said to be *bounded* if for  $\tau$  small enough,  $\Gamma_\tau = \phi$ . The subset of bounded test curves is denoted by  $\text{TC}^\infty(X, \theta; \phi)$ . In this case, we write

$$\Gamma_{\min} := \{\tau \in \mathbb{R} : \Gamma_\tau = \phi\}.$$

A test curve  $\Gamma \in \text{TC}(X, \theta; \phi)$  is said to have *finite energy* if

$$\mathbf{E}^\phi(\Gamma) := \Gamma_{\max} \int_X \theta_\phi^n + \int_{-\infty}^{\Gamma_{\max}} \left( \int_X \theta_{\Gamma_\tau}^n - \int_X \theta_\phi^n \right) d\tau > -\infty. \quad (9.3)$$

{eq:tcfiniteenergy}

The subset of test curves with finite energy is denoted by  $\text{TC}^1(X, \theta; \phi)$ .

We first observe that the notion of test curves does not really depend on the choice of  $\theta$  within its cohomology class.

prop:testcurveindeptheta

**Proposition 9.1.2** Let  $\theta'$  be another smooth closed real  $(1, 1)$ -form on  $X$  representing the same cohomology class as  $\theta$ . Let  $\phi \in \text{PSH}(X, \theta)_{>0}$  be a model potential. Let  $\phi' \in \text{PSH}(X, \theta')_{>0}$  be the unique model potential satisfying  $\phi \sim \phi'$ .

Then there is a canonical bijection

$$\text{TC}(X, \theta; \phi) \xrightarrow{\sim} \text{TC}(X, \theta'; \phi').$$

This bijection induces the following bijections:

$$\text{TC}^1(X, \theta; \phi) \xrightarrow{\sim} \text{TC}^1(X, \theta'; \phi'), \quad \text{TC}^\infty(X, \theta; \phi) \xrightarrow{\sim} \text{TC}^\infty(X, \theta'; \phi').$$

These bijections satisfy the obvious cocycle conditions.

**Proof** Choose  $g \in C^\infty(X)$  such that  $\theta' = \theta + \text{dd}^c g$ . Given any  $\Gamma \in \text{TC}(X, \theta; \phi)$ , we observe that  $\Gamma' : (-\infty, \Gamma_{\max}) \rightarrow \text{PSH}(X, \theta')$  defined as

$$\tau \mapsto P_{\theta'}[\Gamma_\tau - g]$$

lies in  $\text{TC}(X, \theta'; \phi')$ . Moreover, the choice of  $g$  is irrelevant since for any other choice of  $g$ , say  $g'$ , we have

$$\Gamma_\tau - g \sim \Gamma_\tau - g'.$$

All assertions follow directly from the definition.  $\square$

prop:ETCbimero

**Proposition 9.1.3** Let  $\pi : Y \rightarrow X$  be a proper bimeromorphic morphism from a compact Kähler manifold. Then the pointwise pull-back induces a bijection

$$\pi^* : \text{TC}(X, \theta; \phi) \xrightarrow{\sim} \text{TC}(Y, \pi^*\theta; \pi^*\phi).$$

**Proof** This follows immediately from [Proposition 3.1.4](#).  $\square$

prop:Gammaclosed

**Proposition 9.1.4** Let  $\Gamma$  be a test curve in  $\text{PSH}(X, \theta)$ . For each  $x \in X$ , the map  $\mathbb{R} \ni \tau \mapsto \Gamma_\tau(x)$  is a closed concave function. Moreover, the map is proper as long as  $\Gamma_{\Gamma_{\max}}(x) \neq -\infty$ .

The notion of closedness is recalled in [Definition A.1.6](#).

**Proof** We argue the closedness. Fix  $x \in X$ . Assume that  $\Gamma_\tau(x) \neq -\infty$  for some  $\tau \in \mathbb{R}$ . We only need to argue the upper-semicontinuity of  $\tau \mapsto \Gamma_\tau(x)$ . The upper semi-continuity is clear at  $\tau \geq \Gamma_{\max}$ , so we are reduced to prove the following:

$$\Gamma_\tau = \inf_{\tau' < \tau} \Gamma_{\tau'} \quad (9.4)$$

{eq:Gammatautempl}

for any  $\tau < \Gamma_{\max}$ . Take  $\tau'' \in (\tau, \Gamma_{\max})$ . Outside the polar locus of  $\Gamma_{\tau''}$ , we know that (9.4) holds by continuity. So (9.4) holds everywhere by [Proposition 1.2.5](#).

The final assertion is trivial.  $\square$

def:Ptestcurve

**Definition 9.1.3** Let  $\Gamma \in \text{TC}(X, \theta)_{>0}$  and  $\omega$  be a smooth closed real positive  $(1, 1)$ -form. Then we define  $P_{\theta+\omega}[\Gamma] \in \text{TC}(X, \theta + \omega)_{>0}$  as follows:

(1) Define

$$P_{\theta+\omega}[\Gamma]_{\max} = \Gamma_{\max};$$

(2) for each  $\tau < \Gamma_{\max}$ , define

$$P_{\theta+\omega}[\Gamma]_\tau = P_{\theta+\omega}[\Gamma_\tau].$$

It follows from [Proposition 3.1.5](#) that  $P_{\theta+\omega}[\Gamma] \in \text{TC}(X, \theta + \omega)_{>0}$ .

## 9.2 Ross–Witt Nyström correspondence

sec:RWN

Let  $X$  be a connected compact Kähler manifold of dimension  $n$  and  $\theta$  be a smooth closed real  $(1, 1)$ -form on  $X$  representing a big cohomology class. Fix a model potential  $\phi \in \text{PSH}(X, \theta)_{>0}$ .

[Proposition 9.1.4](#) allows us to talk about the Legendre transforms in the expected way.

The general definition of the Legendre transform [Definition A.2.1](#) can be translated as follows:

def:Legtrans

**Definition 9.2.1** Let  $\Gamma \in \text{TC}(X, \theta; \phi)$ . We define its *Legendre transform* as  $\Gamma^*: [0, \infty) \rightarrow \text{PSH}(X, \theta)$  given by

$$\Gamma_t^* = \sup_{\tau \in \mathbb{R}} (t\tau + \Gamma_\tau). \quad (9.5)$$

{eq:testcurveLegtran}

rmk:negativeray

**Remark 9.2.1** Here we do not talk about the case  $t < 0$  because its behaviour is pretty trivial: take  $x \in X$ , if  $\Gamma_\tau(x) = -\infty$  for all  $\tau$ , then  $\Gamma_t^* = -\infty$ ; otherwise,  $\Gamma_t^* = \infty$ .

As we will see later on, the information about  $t \geq 0$  suffices to characterize  $\Gamma$ .

We have made a non-trivial claim that  $\Gamma_t^* \in \text{PSH}(X, \theta)$  for all  $t \geq 0$ . Let us prove this.



lma:testcurvelegusc

**Lemma 9.2.1** *Let  $\Gamma \in \text{TC}(X, \theta; \phi)$ . Then  $\Gamma_t^* \in \text{PSH}(X, \theta)$  for all  $t \geq 0$ . In fact,  $\Gamma$  is upper semicontinuous as a function of  $X \times (0, \infty)$ .*

**Proof** We first observe that for each  $x \in X$ , we have

$$\Gamma_t^*(x) \leq t\Gamma_{\max} < \infty.$$

Let  $R = \{a + ib \in \mathbb{C} : a > 0\}$ . We consider

$$F : X \times R \rightarrow [-\infty, \infty), \quad (x, a + ib) \mapsto \Gamma_a^*(x).$$

Let  $\pi : X \times R \rightarrow X$  be the natural projection. Observe that the upper semicontinuous envelope  $G$  of  $F$  is  $\pi^*\theta$ -psh by [Proposition 1.2.1](#). It suffices to show that  $F = G$ . We let

$$E := \{(x, z) \in X \times R : F(x, z) < G(x, z)\}.$$

We want to argue that  $E = \emptyset$ . Clearly,  $E$  can be written as  $B \times i\mathbb{R}$  for some set  $B \subseteq X \times (0, \infty)$ . Since  $E$  is a pluripolar set by [Proposition 1.2.3](#), it has zero Lebesgue measure. Hence,  $B$  has zero Lebesgue measure. For each  $x \in X$ , write

$$B_x = \{t \in (0, \infty) : (t, x) \in B\}.$$

By Fubini theorem,  $B_x$  has zero 1-dimensional Lebesgue measure for all  $x \in X \setminus Z$ , where  $Z \subseteq X$  is a subset of measure 0. We may assume that  $Z \supseteq \{\Gamma_{-\infty} = 0\}$  so that for  $x \in X \setminus Z$ ,  $\Gamma_t(x) \neq -\infty$  for all  $t > 0$ .

For any  $x \in X \setminus Z$ , both  $t \mapsto F(x, t)$  and  $G(x, t)$  are convex functions with values in  $\mathbb{R}$  on  $(0, \infty)$ . They agree almost everywhere, hence everywhere by their continuity. It follows that for  $x \in X \setminus Z$ , we have  $B_x = \emptyset$ .

By [Theorem A.2.1](#), for any  $x \in X$ , we have

$$\Gamma_\tau(x) = \inf_{t>0} (F(t, x) - t\tau), \quad \tau < \Gamma_{\max}.$$

On the other hand, let

$$\chi_\tau(x) = \inf_{t>0} (G(t, x) - t\tau), \quad \tau < \Gamma_{\max}, x \in X.$$

By Kiselman's principle [Proposition 1.2.6](#),  $\chi_\tau \in \text{PSH}(X, \theta)$ . But on  $X \setminus Z$ , we already know that  $\Gamma_\tau = \chi_\tau$  for all  $\tau < \Gamma_{\max}$ . By [Proposition 1.2.5](#), they are equal everywhere. By [Theorem A.2.1](#) again, we find that  $F = G$ .  $\square$

lma:suplegenlinear

**Lemma 9.2.2** *Let  $\Gamma \in \text{TC}(X, \theta; \phi)$ , then*

$$\sup_X \Gamma_t^* = t\Gamma_{\max}$$

for all  $t \geq 0$ .

In particular,  $t \mapsto \Gamma_t^* - t\Gamma_{\max}$  is a decreasing function in  $t \geq 0$ .

**Proof** Choose  $x \in X$  such that  $\Gamma_{\max}(x) = 0$ . Then

$$\Gamma_t^*(x) = t\Gamma_{\max}$$

by definition. On the other hand, since  $\Gamma_\tau \leq 0$  for all  $\tau < \Gamma_{\max}$ , we have

$$\sup_X \Gamma_t^* \leq t\Gamma_{\max}.$$

lma:LegsendsTCtoR

**Lemma 9.2.3** *Given  $\Gamma \in \text{TC}(X, \theta; \phi)$ , we have  $\Gamma^* \in \mathcal{R}(X, \theta; \phi)$ .*

**Proof** It follows from [Lemma 9.2.1](#), [\(9.5\)](#) and [Proposition 1.2.1](#) that  $\Gamma^*$  is a subgeodesic (in the sense that for each  $0 \leq a \leq b$ , the restriction  $(\Gamma_t^*)_{t \in (a,b)}$  is a subgeodesic from  $\Gamma_a^*$  to  $\Gamma_b^*$ ).

First observe that as  $t \rightarrow 0+$ , we have

$$\Gamma_t^* \xrightarrow{L^1} \phi. \quad (9.6)$$

{eq:Gammatophil1temp}

To see this, first observe that by [\(9.5\)](#), for any fixed  $t > 0$  and any  $x \in X$  with  $\phi(x) \neq -\infty$ , we have

$$\Gamma_t^*(x) \leq t\Gamma_{\max} + \phi(x).$$

By [Proposition 1.2.5](#), the same holds everywhere. Therefore, any  $L^1$ -cluster point  $\psi$  of  $\Gamma_t^*$  as  $t \rightarrow 0$  satisfies  $\psi \leq \phi$ . On the other hand, for any fixed  $\tau < \Gamma_{\max}$ , by [\(9.5\)](#), we have

$$\Gamma_t^* \geq \Gamma_\tau + t\tau$$

for any  $t > 0$ . So  $\psi \geq \Gamma_\tau$  almost everywhere and hence everywhere by [Proposition 1.2.5](#). It follows that  $\psi \geq \phi$ . Therefore,  $\psi = \phi$ . On the other hand, from the above estimates and [Proposition 1.5.1](#) that  $(\Gamma_t^*)_{t \in (0,1)}$  is a relative compact subset in  $\text{PSH}(X, \theta)$  with respect to the  $L^1$ -topology. We therefore conclude [\(9.6\)](#).

Assume that  $\Gamma^*$  is not a geodesic ray. Then we can find  $0 \leq a < b$  such that  $(\Gamma_t^*)_{t \in (a,b)}$  differs from the geodesic  $(\eta_t)_{t \in (a,b)}$  from  $\Gamma_a^*$  to  $\Gamma_b^*$ . We consider the subgeodesic  $(\ell_t)_{t > 0}$  given by  $\ell_t = \eta_t$  for  $t \in (a, b)$  and  $\ell_t = \Gamma_t^*$  otherwise. Consider the Legendre transform

$$\Gamma'_\tau = \inf_{t > 0} (\ell_t - t\tau), \quad \tau \in \mathbb{R}.$$

Then  $\Gamma'_\tau \geq \Gamma_\tau$  and  $\Gamma'_\tau \in \text{PSH}(X, \theta) \cup \{-\infty\}$  by [Proposition 1.2.6](#) for all  $\tau \in \mathbb{R}$ .

We claim that

$$\Gamma'_\tau \leq \Gamma_\tau + (b - a)(\Gamma_{\max} - \tau), \quad \tau \in \mathbb{R}.$$

Observe that  $\Gamma'_\tau \equiv -\infty$  when  $\tau > \Gamma_{\max}$  by [Lemma 9.2.2](#). So it suffices to consider  $\tau \leq \Gamma_{\max}$ . In this case, we compute

$$\inf_{t \in [a,b]} (\ell_t - t\tau) \leq \Gamma_b^* - b\tau \leq (b - a)(\Gamma_{\max} - \tau) \inf_{t \in [a,b]} (\Gamma_t^* - t\tau),$$

where we applied [Lemma 9.2.2](#). In particular, for any  $\tau < \Gamma_{\max}$ , we have

$$\Gamma'_\tau \leq \Gamma_\tau.$$

On the other hand, by definition of  $\Gamma'_\tau$ , we clearly have  $\Gamma'_\tau \leq 0$  for all  $\tau < \Gamma_{\max}$ . It follows from the fact that  $\Gamma_\tau$  is a model potential that  $\Gamma_\tau = \Gamma'_\tau$  for all  $\tau < \Gamma_{\max}$ . Therefore, by [Theorem A.2.1](#), we have  $\Gamma_t^* = \ell'_t$  for all  $t > 0$ , which is a contradiction.  $\square$

thm:Legendbij

**Theorem 9.2.1** *The Legendre transform in [Definition 9.2.1](#) is a bijection*

$$\mathrm{TC}(X, \theta; \phi) \xrightarrow{\sim} \mathcal{R}(X, \theta; \phi).$$

Moreover, this bijection restricts to the following bijections:

$$\mathrm{TC}^1(X, \theta; \phi) \xrightarrow{\sim} \mathcal{R}^1(X, \theta; \phi), \quad \mathrm{TC}^\infty(X, \theta; \phi) \xrightarrow{\sim} \mathcal{R}^\infty(X, \theta; \phi).$$

For any  $\Gamma \in \mathrm{TC}^1(X, \theta; \phi)$ , we have

$$\mathbf{E}^\phi(\Gamma) = \mathbf{E}^\phi(\Gamma^*). \quad (9.7)$$

{eq:RWNenergy}

**Proof** It follows from [Lemma 9.2.3](#) that the forward map is well-defined.

The inverse map is of course also given by the Legendre transform: given  $\ell \in \mathcal{R}(X, \theta; \phi)$ , its Legendre transform is given by

$$\ell_\tau^* := \inf_{t > 0} (\ell_t - t\tau), \quad \tau \in \mathbb{R}. \quad (9.8)$$

{eq:invLeg}

By [Proposition 4.2.4](#), there is a constant  $C > 0$  such that  $\ell_t \leq Ct$ .

Note that it follows from [Proposition 1.2.6](#) that  $\ell_\tau^* \in \mathrm{PSH}(X, \theta) \cup \{-\infty\}$  for all  $\tau \in \mathbb{R}$ .

We need to argue for any  $\tau \in \mathbb{R}$  such that  $\ell_\tau^* \neq -\infty$ , we have  $P_\theta[\ell_\tau^*] = \ell_\tau^*$ . Fix such  $\tau$  and some  $C > 0$ . It suffices to show that

$$(\ell_\tau^* + C) \wedge \phi \leq \ell_\tau^*. \quad (9.9)$$

{eq:ellstarleqetemp1}

For this purpose, let us consider the following geodesics: for any  $M > 0$  and  $t \in [0, 1]$ , let

$$\ell_t^{1,M} = \ell_{tM} - tM\tau, \quad \ell_t^{2,M} = (\ell_\tau^* + C) \wedge \phi - Ct.$$

It is clear that at  $t = 0, 1$ , we have  $\ell_t^{2,M} \leq \ell_t^{1,M}$ . Hence, the same holds for all  $t \in [0, 1]$ . In particular, for any fixed  $s \in [0, 1]$ , we have

$$(\ell_\tau^* + C) \wedge \phi - Cs \leq \ell_{sM} - sM.$$

Take infimum with respect to  $M \geq 1$  and then the supremum with respect to  $s$ , we conclude [\(9.9\)](#).

The two operations are inverse to each other thanks to [Theorem A.2.1](#).

Next we consider the bounded situation. Suppose that  $\Gamma \in \mathrm{TC}^\infty(X, \theta; \phi)$ . Take  $\tau_0 \in \mathbb{R}$  so that  $\Gamma_\tau = \phi$  for all  $\tau \leq \tau_0$ . It follows from that

$$\Gamma_t^* \geq \phi + t\tau_0$$

for all  $t > 0$ . Therefore,  $\Gamma_t^* \sim \phi$  for all  $t > 0$  and hence  $\Gamma^* \in \mathcal{R}^\infty(X, \theta; \phi)$ .

Conversely, suppose that  $\ell \in \mathcal{R}^\infty(X, \theta; \phi)$ . Thanks to [Proposition 4.2.3](#), there is a constant  $C > 0$  such that

$$\ell_t \geq \phi - Ct.$$

Therefore, according to [\(9.8\)](#), we have

$$\ell_\tau^* \geq \inf_{t>0} \phi - (C + \tau)t = \phi$$

if  $\tau \leq -C$ . Therefore,  $\ell_\tau^* = \phi$  for all  $\tau \leq -C$ .

Finally, it remains to handle [\(9.7\)](#). Take  $\Gamma \in \text{TC}^\infty(X, \theta; \phi)$ . We may assume that  $\Gamma_{\max} = 0$  after a translation.

For  $N \in \mathbb{Z}_{>0}$ ,  $M \in \mathbb{Z}$ , we introduce the following:

$$\Gamma_t^{*,N,M} := \max_{\substack{k \in \mathbb{Z} \\ k \leq M}} \left( \Gamma_{k/2^N} + tk/2^N \right) \in \mathcal{E}^\infty(X, \theta; \phi), \quad t > 0.$$

Moreover, we now argue that

$$\frac{t}{2^N} \int_X \theta_{\Gamma_{(M+1)/2^N}}^n \leq E_\theta^\phi(\Gamma_t^{*,N,M+1}) - E_\theta^\phi(\Gamma_t^{*,N,M}) \leq \frac{t}{2^N} \int_X \theta_{\Gamma_{M/2^N}}^n. \quad (9.10)$$

{eq: diff\_eq\_I}

Indeed, for elementary reasons:

$$\begin{aligned} \int_X \left( \Gamma_t^{*,N,M+1} - \Gamma_t^{*,N,M} \right) \theta_{\Gamma_t^{*,N,M+1}}^n &\leq E_\theta^\phi(\Gamma_t^{*,N,M+1}) - E_\theta^\phi(\Gamma_t^{*,N,M}) \\ &\leq \int_X \left( \Gamma_t^{*,N,M+1} - \Gamma_t^{*,N,M} \right) \theta_{\Gamma_t^{*,N,M}}^n. \end{aligned} \quad (9.11)$$

{eq: first\_I\_ineq}

Clearly  $\Gamma_t^{*,N,M+1} \geq \Gamma_t^{*,N,M}$ , and using  $\tau$ -concavity, we notice that

$$U_t := \left\{ \Gamma_t^{*,N,M+1} - \Gamma_t^{*,N,M} > 0 \right\} = \left\{ \Gamma_{(M+1)/2^N} + 2^{-N}t - \Gamma_{M/2^N} > 0 \right\}.$$

Moreover, on  $U_t$  we have

$$\Gamma_t^{*,N,M+1} = \Gamma_{(M+1)/2^N} + t(M+1)/2^N, \quad \Gamma_t^{*,N,M} = \Gamma_{M/2^N} + tM/2^N.$$

We also note that  $U_t$  is an open set in the plurifine topology, implying that

$$\begin{aligned} \theta_{\Gamma_{(M+1)/2^N}}^n|_{U_t} &= \theta_{\Gamma_t^{*,N,M+1}}^n|_{U_t}, \\ \theta_{\Gamma_{M/2^N}}^n|_{U_t} &= \theta_{\Gamma_t^{*,N,M}}^n|_{U_t}. \end{aligned}$$

Recall that  $\theta_{\Gamma_{M/2^N}}^n$  and  $\theta_{\Gamma_{(M+1)/2^N}}^n$  are supported on the sets  $\{\Gamma_{M/2^N} = 0\}$  and  $\{\Gamma_{(M+1)/2^N} = 0\}$  respectively, see [Theorem 3.1.2](#). Since  $\{\Gamma_{(M+1)/2^N} = 0\} \subseteq U_t$  and  $\{\Gamma_{(M+1)/2^N} = 0\} \subseteq \{\Gamma_{M/2^N} = 0\}$ , applying the above to [\(9.11\)](#), we arrive at [\(9.10\)](#).

Fixing  $N$ , let  $M = \lfloor 2^N \Gamma_{\min} \rfloor$ . Then repeated application of [\(9.10\)](#) yields

$$\sum_{M+1 \leq j \leq 0} \frac{t}{2^N} \int_X \theta_{\Gamma_{j/2^N}}^n \leq E_\theta^\phi(\Gamma_t^{*,N,0}) - E_\theta^\phi(E_t^{*,N,M}) \leq \sum_{M \leq j \leq -1} \frac{t}{2^N} \int_X \theta_{\Gamma_{j/2^N}}^n.$$

Since  $M \leq 2^N \Gamma_{\min}$ , we have that

$$\Gamma_t^{*,N,M} = \Gamma_{M/2^N} + tM/2^N = \phi + tM/2^N,$$

we can continue to write

$$\sum_{j=M+1}^0 \frac{t}{2^N} \left( \int_X \theta_{\Gamma_{j/2^N}}^n - \int_X \theta_\phi^n \right) \leq E_\theta^\phi(\Gamma_t^{*,N,0}) \leq \sum_{j=M}^{-1} \frac{t}{2^N} \left( \int_X \theta_{\Gamma_{j/2^N}}^n - \int_X \theta_\phi^n \right).$$

We now notice that we have Riemann sums on both the left and right of the above inequality. Using [Proposition 9.1.1](#), it is possible to let  $N \rightarrow \infty$  and obtain

$$E_\theta^\phi(\Gamma_t^*) = tE^\phi(\Gamma)$$

So (9.7) follows as desired. Note that we have furthermore shown that  $t \mapsto E_\theta^\phi(\Gamma_t^*)$  is linear.

Finally, let us come back to the general case. Let  $\Gamma \in \text{TC}(X, \theta; \phi)$ . Again, we may assume that  $\Gamma_{\max} = 0$ . For each  $\epsilon > 0$ , we introduce  $\Gamma^\epsilon \in \text{TC}^\infty(X, \theta; \phi)$  as follows:

- (1) Let  $\Gamma_{\max}^\epsilon = 0$ , and
- (2) for each  $\tau < 0$ , we set

$$\Gamma_\tau^\epsilon = P_\theta [(1 + \epsilon\tau) \vee 0) \Gamma_\tau + (1 - (1 + \epsilon\tau) \vee 0)) \phi].$$

It follows from [Corollary 3.1.2](#) that for each  $\tau < 0$ , the sequence  $\Gamma_\tau^\epsilon$  is a decreasing sequence with limit  $\Gamma_\tau$  as  $\epsilon \searrow 0$ . Therefore, by [Proposition 3.1.8](#), we have

$$\lim_{\epsilon \rightarrow 0+} \int_X (\theta + \text{dd}^c \Gamma_\tau^\epsilon)^n = \int_X (\theta + \text{dd}^c \Gamma_\tau)^n$$

for all  $\tau < 0$ . Hence, by the monotone convergence theorem, we find

$$E^\phi(\Gamma) = \lim_{\epsilon \rightarrow 0+} E^\phi(\Gamma^\epsilon) = \lim_{\epsilon \rightarrow 0+} E^\phi(\Gamma^{\epsilon,*}). \quad (9.12)$$

{eq:EphiGammatempl}

Furthermore, according to [Proposition A.2.2](#), we have

$$\Gamma_t^* = \inf_{\epsilon > 0} \Gamma_t^{\epsilon,*}$$

for all  $t > 0$ .

Now suppose that  $\Gamma \in \text{TC}^1(X, \theta; \phi)$ . Then it follows from [Theorem 4.2.1](#) that for each  $t > 0$ ,

$$E_\theta^\phi(\Gamma_t^*) = \lim_{\epsilon \rightarrow 0+} E_\theta^\phi(\Gamma_t^{\epsilon,*}) = tE^\phi(\Gamma).$$

Hence,  $\Gamma^* \in \mathcal{E}^1(X, \theta; \phi)$ .

Conversely, suppose that  $\Gamma^* \in \mathcal{E}^1(X, \theta; \phi)$ . Then (9.12) implies that  $\Gamma \in \text{TC}^1(X, \theta; \phi)$ .  $\square$

As an immediate consequence of the proof, we have

**Corollary 9.2.1** *Let  $\ell \in \mathcal{R}^1(X, \theta; \phi)$ , then  $[0, \infty) \ni t \mapsto E_\theta^\phi(\ell_t)$  is linear.*

**Corollary 9.2.2** *Let  $\ell \in \mathcal{R}(X, \theta; \phi)$ . Then  $\sup_X \ell_t = \ell_{\max}^* t$ .*

*Proof* This follows from Lemma 9.2.2 and Theorem 9.2.1.  $\square$

### 9.3 $\mathcal{I}$ -model test curves

Let  $X$  be a connected compact Kähler manifold of dimension  $n$  and  $\theta$  be a smooth closed real  $(1, 1)$ -form on  $X$  representing a big cohomology class. Fix a model potential  $\phi \in \text{PSH}(X, \theta)_{>0}$ .

**Definition 9.3.1** A test curve  $\Gamma \in \text{TC}(X, \theta; \phi)$  is  $\mathcal{I}$ -model if for any  $\tau < \Gamma_{\max}$ , the potential  $\Gamma_\tau$  is  $\mathcal{I}$ -model.

The subset of  $\mathcal{I}$ -model test curves in  $\text{TC}(X, \theta; \phi)$  is denoted by  $\text{PSH}^{\text{NA}}(X, \theta; \phi)$ .

The set of  $\mathcal{I}$ -model test curves in  $\text{PSH}(X, \theta)$  for any model potential  $\phi \in \text{PSH}(X, \theta)_{>0}$  is denoted by  $\text{PSH}^{\text{NA}}(X, \theta)_{>0}$ .

**Proposition 9.3.1** *Let  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ . Then  $\Gamma_{-\infty}$  is an  $\mathcal{I}$ -model potential.*

*Proof* This follows from Proposition 3.2.12.  $\square$

**Proposition 9.3.2** *Let  $\theta'$  be another smooth closed real  $(1, 1)$ -form on  $X$  representing the same cohomology class as  $\theta$ . Then there is a canonical bijection*

$$\text{PSH}^{\text{NA}}(X, \theta)_{>0} \xrightarrow{\sim} \text{PSH}^{\text{NA}}(X, \theta')_{>0}.$$

*This bijection satisfies the obvious cocycle condition.*

*Proof* This is an immediate consequence of Proposition 9.1.2 and Example 7.1.2.  $\square$

**Proposition 9.3.3** *Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism from a compact Kähler manifold. Then the pointwise pull-back induces a bijection*

$$\pi^*: \text{PSH}^{\text{NA}}(X, \theta; \phi) \xrightarrow{\sim} \text{PSH}^{\text{NA}}(Y, \pi^*\theta; \pi^*\phi).$$

*Proof* This is an immediate consequence of Proposition 9.1.3 and Proposition 3.2.5.  $\square$

**Definition 9.3.2** Given  $\Gamma \in \text{TC}(X, \theta; \phi)$ , we define its  $\mathcal{I}$ -envelope  $P_\theta[\Gamma]_{\mathcal{I}}$  as the map  $(-\infty, \Gamma_{\max}) \rightarrow \text{PSH}(X, \theta)$  given by

$$\tau \mapsto P_\theta[\Gamma_\tau]_{\mathcal{I}}.$$

prop:transitionPI

**Proposition 9.3.4** *Let  $\Gamma \in \text{TC}(X, \theta; \phi)$ , then*

$$P_\theta[\Gamma]_I \in \text{PSH}^{\text{NA}}(X, \theta; P_\theta[\phi]_I).$$

*More generally, for any closed real smooth positive  $(1, 1)$ -form  $\omega$  on  $X$ , we have*

$$P_{\theta+\omega}[\Gamma]_I \in \text{PSH}^{\text{NA}}(X, \theta + \omega; P_{\theta+\omega}[\phi]_I).$$

**Proof** The only non-trivial point is to show that

$$\sup_{\tau < \Gamma_{\max}}^* P_\theta[\Gamma_\tau]_I = P_\theta[\phi]_I, \quad \sup_{\tau < \Gamma_{\max}}^* P_{\theta+\omega}[\Gamma_\tau]_I = P_{\theta+\omega}[\phi]_I.$$

This follows from [Proposition 3.2.12](#).  $\square$

## 9.4 Operations on test curves

sec:operationontc

Let  $X$  be a connected compact Kähler manifold of dimension  $n$  and  $\theta, \theta', \theta''$  be smooth closed real  $(1, 1)$ -forms on  $X$  representing big cohomology classes.

def:potestcurve

**Definition 9.4.1** Given  $\Gamma, \Gamma' \in \text{TC}(X, \theta)_{>0}$ , we say  $\Gamma \leq \Gamma'$  if for all  $\Gamma_{\max} \leq \Gamma'_{\max}$  and for all  $\tau < \Gamma_{\max}$ , we have

$$\Gamma_\tau \leq \Gamma'_\tau. \quad (9.13)$$

{eq:GammatauGammap}

Observe that (9.13) actually holds for all  $\tau \in \mathbb{R}$ . It is easy to verify that for all  $\leq$  defines a partial order on  $\text{TC}(X, \theta)_{>0}$ .

lma:testcurord1

**Lemma 9.4.1** *Let  $\Gamma, \Gamma' \in \text{TC}(X, \theta)_{>0}$  and  $\omega$  be a closed real smooth positive  $(1, 1)$ -form on  $X$ . Then the following are equivalent:*

- (1)  $\Gamma \leq \Gamma'$ ;
- (2)  $P_{\theta+\omega}[\Gamma] = P_{\theta+\omega}[\Gamma']$ .

**Proof** It suffices to observe that we could rewrite (9.13) as

$$\Gamma_\tau \leq_P \Gamma'_\tau,$$

since both potentials are model.  $\square$

def:sumtestcur

**Definition 9.4.2** Let  $\Gamma \in \text{TC}(X, \theta)_{>0}$  and  $\Gamma' \in \text{TC}(X, \theta')_{>0}$ , then we define  $\Gamma + \Gamma' \in \text{TC}(X, \theta + \theta')_{>0}$  as follows:

- (1) we set

$$(\Gamma + \Gamma')_{\max} := \Gamma_{\max} + \Gamma'_{\max};$$

- (2) for any  $\tau < (\Gamma + \Gamma')_{\max}$ , we define

$$(\Gamma + \Gamma')_\tau := P_\theta \left[ \sup_{t \in \mathbb{R}} (\Gamma_t + \Gamma'_{\tau-t}) \right]. \quad (9.14)$$

{eq:GammaGammagsum}

lma:testcurvplus

**Lemma 9.4.2** Let  $\Gamma \in \text{TC}(X, \theta)_{>0}$  and  $\Gamma' \in \text{TC}(X, \theta')_{>0}$ , then for any  $\tau < (\Gamma + \Gamma')_{\max}$ , we have

$$\sup_{t \in \mathbb{R}} (\Gamma_t + \Gamma'_{\tau-t}) \in \text{PSH}(X, \theta).$$

This potential is  $\mathcal{I}$ -good if  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$  and  $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')_{>0}$ .

In particular, (9.14) in [Definition 9.4.2](#) makes sense.

**Proof** Let

$$\eta_\tau = \sup_{t \in \mathbb{R}} (\Gamma_t + \Gamma'_{\tau-t}) = \sup_{t < \Gamma_{\max}, \tau-t < \Gamma'_{\max}} (\Gamma_t + \Gamma'_{\tau-t})$$

for all  $\tau \in \mathbb{R}$ . Set

$$Z = \{x \in X : \Gamma_{-\infty}(x) = -\infty \text{ or } \Gamma'_{-\infty}(x) = -\infty\}.$$

It follows from [Proposition A.2.3](#) that for any  $x \in X \setminus Z$ , we have

$$\eta_t^*(x) = \Gamma_t^*(x) + \Gamma_t'^*(x)$$

for all  $t > 0$ . The same trivially holds when  $x \in Z$ , so the equation holds everywhere.

In particular, by [Theorem A.2.1](#) and [Proposition 1.2.6](#), we have

$$\eta_\tau = (\Gamma^* + \Gamma'^*)_\tau^* \in \text{PSH}(X, \theta + \theta') \cup \{-\infty\}.$$

Next, assume that  $\Gamma$  and  $\Gamma'$  are  $\mathcal{I}$ -model. We need to argue that so is  $\Gamma + \Gamma'$ . Fix  $\tau < \Gamma_{\max} + \Gamma'_{\max}$ . Then for each  $t \in \mathbb{R}$  such that  $t < \Gamma_{\max}$  and  $\tau - t < \Gamma'_{\max}$ , we know that  $\Gamma_t \in \text{PSH}(X, \theta)_{>0}$  and  $\Gamma'_{\tau-t} \in \text{PSH}(X, \theta')_{>0}$  by [Lemma 9.1.1](#). It follows from [Example 7.1.2](#) that  $\Gamma_t$  and  $\Gamma'_{\tau-t}$  are both  $\mathcal{I}$ -good, hence so is  $\Gamma_t + \Gamma'_{\tau-t} \in \text{PSH}(X, \theta + \theta')_{>0}$  by [Proposition 7.2.1](#). Therefore,  $\eta_\tau$  is  $\mathcal{I}$ -good by [Proposition 7.2.2](#). Therefore,  $\Gamma + \Gamma'$  is  $\mathcal{I}$ -model.  $\square$

prop:testcurvesumproperty

**Proposition 9.4.1** Let  $\Gamma \in \text{TC}(X, \theta)_{>0}$  and  $\Gamma' \in \text{TC}(X, \theta')_{>0}$ , then  $\Gamma + \Gamma' \in \text{TC}(X, \theta + \theta')_{>0}$ . Moreover,

$$(\Gamma + \Gamma')_{-\infty} = P_{\theta+\theta'}[\Gamma_{-\infty} + \Gamma'_{-\infty}]. \quad (9.15)$$

{eq:sumGammaGammap}

When  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$  and  $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')_{>0}$ , we have  $\Gamma + \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta + \theta')_{>0}$ .

The operation  $+$  is commutative and associative.

**Proof** It follows immediately from [Lemma 9.4.2](#) that  $\Gamma + \Gamma' \in \text{TC}(X, \theta + \theta')_{>0}$ , and it lies in  $\text{PSH}^{\text{NA}}(X, \theta + \theta')_{>0}$  if  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$  and  $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')_{>0}$ .

We argue (9.15). By definition, for any small enough  $\tau$ , we have

$$(\Gamma + \Gamma')_{-\infty} \geq (\Gamma + \Gamma')_{2\tau} \geq_P \Gamma_\tau + \Gamma'_\tau.$$

Letting  $\tau \rightarrow -\infty$  and applying [Proposition 6.2.4](#) and [Theorem 6.2.2](#), we find that

$$(\Gamma + \Gamma')_{-\infty} \geq_P \Gamma_{-\infty} + \Gamma'_{-\infty}.$$



On the other hand, for each small enough  $\tau$ , we have

$$(\Gamma + \Gamma')_\tau \sim_P \sup_{t \in \mathbb{R}} (\Gamma_t + \Gamma'_{\tau-t}) \leq_P \Gamma_{-\infty} + \Gamma'_{-\infty}$$

by [Proposition 6.1.5](#) and [Proposition 6.2.4](#). We apply [Proposition 6.2.4](#) again, we conclude that

$$(\Gamma + \Gamma')_{-\infty} \leq_P \Gamma_{-\infty} + \Gamma'_{-\infty}.$$

So [\(9.15\)](#) follows.

Finally, let us show that  $+$  is commutative and associative. Commutativity is obvious. Let  $\Gamma'' \in \text{TC}(X, \theta'')_{>0}$ . Then we want to show that

$$(\Gamma + \Gamma') + \Gamma'' = \Gamma + (\Gamma' + \Gamma'').$$

First observe that

$$((\Gamma + \Gamma') + \Gamma'')_{\max} = (\Gamma + (\Gamma' + \Gamma''))_{\max}.$$

Fix  $\tau$  less than this common value. We observe that

$$\begin{aligned} & ((\Gamma + \Gamma') + \Gamma'')_\tau \\ &= P_\theta \left[ \sup_{t_1 \in \mathbb{R}} ((\Gamma + \Gamma')_{t_1} + \Gamma''_{\tau-t_1}) \right] \\ &\sim_P \sup_{t_1 \in \mathbb{R}} ((\Gamma + \Gamma')_{t_1} + \Gamma''_{\tau-t_1}) \\ &\sim_P \sup_{t_1, t_2 \in \mathbb{R}} (\Gamma_{t_2} + \Gamma'_{t_1-t_2} + \Gamma''_{\tau-t_1}), \end{aligned}$$

where in the last line, we applied [Proposition 6.2.4](#) and [Proposition 6.1.5](#). Similarly, for  $(\Gamma + (\Gamma' + \Gamma''))_\tau$ , we get the same expression. The associativity follows.  $\square$

lma:testcursumcomp

**Lemma 9.4.3** *Let  $\Gamma \in \text{TC}(X, \theta)_{>0}$  and  $\Gamma' \in \text{TC}(X, \theta')_{>0}$ , then for any closed smooth positive  $(1, 1)$ -forms  $\omega$  and  $\omega'$  on  $X$ , we have*

$$P_{\theta+\omega+\theta'+\omega'}[\Gamma + \Gamma'] = P_{\theta+\omega}[\Gamma] + P_{\theta'+\omega'}[\Gamma].$$

**Proof** Observe that

$$P_{\theta+\omega+\theta'+\omega'}[\Gamma + \Gamma']_{\max} = (P_{\theta+\omega}[\Gamma] + P_{\theta'+\omega'}[\Gamma])_{\max} = \Gamma_{\max} + \Gamma'_{\max}.$$

Take  $\tau \in \mathbb{R}$  less than this common value, we need to verify that

$$(\Gamma + \Gamma')_\tau \sim_P (P_{\theta+\omega}[\Gamma] + P_{\theta'+\omega'}[\Gamma])_\tau.$$

By definition, this means that

$$\sup_{t \in \mathbb{R}} (\Gamma_t + \Gamma'_{\tau-t}) \sim_P \sup_{t \in \mathbb{R}} (P_{\theta+\omega}[\Gamma_t] + P_{\theta'+\omega'}[\Gamma'_{\tau-t}]).$$

This is a consequence of [Proposition 6.1.5](#) and [Proposition 6.1.6](#).  $\square$

`def:testcurvplusC`

**Definition 9.4.3** Let  $\Gamma \in \text{TC}(X, \theta)_{>0}$  and  $C \in \mathbb{R}$ , we define  $\Gamma + C \in \text{TC}(X, \theta)_{>0}$  as follows:

(1) We set

$$(\Gamma + C)_{\max} := \Gamma_{\max} + C;$$

(2) for any  $\tau < (\Gamma + C)_{\max}$ , we set

$$\Gamma_{\tau} := \Gamma_{\tau-C}.$$

It is obvious that if  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ , then so is  $\Gamma + C$ .

`prop:testcurveplusC`

**Proposition 9.4.2** Let  $\Gamma \in \text{TC}(X, \theta)_{>0}$ ,  $\Gamma' \in \text{TC}(X, \theta')_{>0}$  and  $C, C' \in \mathbb{R}$ , then

- (1)  $(\Gamma + \Gamma') + C = \Gamma + (\Gamma' + C) = (\Gamma + C) + \Gamma'$ ;
- (2)  $\Gamma + (C + C') = (\Gamma + C) + C'$ .

**Proof** (1) We first observe that

$$((\Gamma + \Gamma') + C)_{\max} = (\Gamma + (\Gamma' + C))_{\max} = ((\Gamma + C) + \Gamma')_{\max} = \Gamma_{\max} + \Gamma'_{\max} + C.$$

Take any  $\tau \in \mathbb{R}$  less than this common value. We compute

$$\begin{aligned} ((\Gamma + \Gamma') + C)_{\tau} &= (\Gamma + \Gamma')_{\tau-C} = P_{\theta+\theta'} \left[ \sup_{t \in \mathbb{R}} (\Gamma_t + \Gamma'_{\tau-C-t}) \right], \\ (\Gamma + (\Gamma' + C))_{\tau} &= P_{\theta+\theta'} \left[ \sup_{t \in \mathbb{R}} (\Gamma_t + (\Gamma' + C)_{\tau-t}) \right] = P_{\theta+\theta'} \left[ \sup_{t \in \mathbb{R}} (\Gamma_t + \Gamma'_{\tau-C-t}) \right], \\ ((\Gamma + C) + \Gamma')_{\tau} &= P_{\theta+\theta'} \left[ \sup_{t \in \mathbb{R}} ((\Gamma + C)_{\tau-t} + \Gamma'_t) \right] \\ &= P_{\theta+\theta'} \left[ \sup_{t \in \mathbb{R}} (\Gamma_t + \Gamma'_{\tau-C-t}) \right]. \end{aligned}$$

(2) Observe that

$$(\Gamma + (C + C'))_{\max} = ((\Gamma + C) + C')_{\max} = \Gamma_{\max} + C + C'.$$

For any  $\tau \in \mathbb{R}$  less than this value, we have

$$(\Gamma + (C + C'))_{\tau} = \Gamma_{\tau-C-C'} = ((\Gamma + C) + C')_{\tau}.$$

`def:testcurlor`

**Definition 9.4.4** Let  $\Gamma, \Gamma' \in \text{TC}(X, \theta)_{>0}$ . We define  $\Gamma \vee \Gamma' \in \text{TC}(X, \theta)_{>0}$  as follows:

(1) We set

$$(\Gamma \vee \Gamma')_{\max} := \Gamma_{\max} \vee \Gamma'_{\max},$$

and

(2) for any  $\tau < (\Gamma \vee \Gamma')_{\max}$ , we define

$$(\Gamma \vee \Gamma')_{\tau} := P_{\theta} \left[ \text{CE} \left( \rho \mapsto \Gamma_{\rho} \vee \Gamma'_{\rho} \right) \right]. \quad (9.16)$$

{eq:testcurlordef}

Recall that the upper convex hull CE is defined in [Definition A.1.4](#). Trivially, we have  $\Gamma \vee \Gamma' \geq \Gamma$  and  $\Gamma \vee \Gamma' \geq \Gamma'$ .

lma:testcurlor

**Lemma 9.4.4** *Let  $\Gamma, \Gamma' \in \text{TC}(X, \theta)_{>0}$ . Then for any  $\tau < \Gamma_{\max} \vee \Gamma'_{\max}$ , we have*

$$\text{CE} \left( \rho \mapsto \Gamma_{\rho} \vee \Gamma'_{\rho} \right)_{\tau} \in \text{PSH}(X, \theta).$$

*This potential is  $\mathcal{I}$ -good if  $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ .*

*In particular, (9.16) in [Definition 9.4.4](#) makes sense.*

**Proof** To simply the notations, we write

$$\psi_{\tau} = \text{CE} \left( \rho \mapsto \Gamma_{\rho} \vee \Gamma'_{\rho} \right)_{\tau}$$

for all  $\tau \in \mathbb{R}$ . Thanks to [Proposition A.2.2](#), we have

$$\psi_t^*(x) = \Gamma_t^*(x) \vee \Gamma_t'^*(x) \quad (9.17)$$

{eq:psistartempl}

for all  $t > 0$  as long as  $\Gamma_{\tau}(x) \neq -\infty$  and  $\Gamma'_{\tau}(x) \neq -\infty$  for some  $\tau \in \mathbb{R}$ . Otherwise, assume that  $x \in X$  is such that  $\Gamma_{\tau} = -\infty$  for all  $\tau \in \mathbb{R}$ , then by definition,  $\psi_{\tau}(x) = \Gamma'_{\tau}(x)$  for all  $\tau \in \mathbb{R}$ . Therefore,  $\Gamma_t^*(x) = -\infty$  for all  $t > 0$  and hence (9.17) continues to hold. Therefore, we have shown that

$$\psi_t^* = \Gamma_t^* \vee \Gamma_t'^* \in \text{PSH}(X, \theta).$$

It follows from [Proposition 4.1.2](#) that  $(\psi_t^*)_{t \in [a, b]}$  is a subgeodesic for any  $0 < a < b$ .

Next we observe that  $\psi_{\bullet}$  is closed by definition. So it follows from [Proposition A.2.2](#) and [Proposition 1.2.6](#) that

$$\psi_{\tau} = (\psi_{\bullet}^*)_{\tau}^* \in \text{PSH}(X, \theta) \cup \{-\infty\}.$$

Due to [Proposition 9.1.4](#) and [Proposition A.1.2](#), there is a pluripolar set  $Z \subseteq X$  such that for  $x \in X \setminus Z$ , we have

$$\psi_{\tau}(x) = \sup \left\{ \lambda \Gamma_{\rho}(x) + (1 - \lambda) \Gamma'_{\rho'}(x) : \lambda \in (0, 1), \rho, \rho' \in \mathbb{R}, \lambda \rho + (1 - \lambda) \rho' = \tau \right\}$$

for all  $\tau < \Gamma_{\max} \vee \Gamma'_{\max}$ . It follows from [Proposition 1.2.5](#) that

$$\psi_{\tau} = \sup^* \left\{ \lambda \Gamma_{\rho} + (1 - \lambda) \Gamma'_{\rho'} : \lambda \in (0, 1), \rho, \rho' \in \mathbb{R}, \lambda \rho + (1 - \lambda) \rho' = \tau \right\} \quad (9.18)$$

{eq:psitausupslinartemp}

for all  $\tau < \Gamma_{\max} \vee \Gamma'_{\max}$ .

It follows from (9.18) that  $\psi_{\tau}$  is  $\mathcal{I}$ -good if  $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ , thanks to [Proposition 7.2.1](#) and [Proposition 7.2.2](#).  $\square$

cor:testcurvlorprop

**Corollary 9.4.1** *Let  $\Gamma, \Gamma' \in \text{TC}(X, \theta)_{>0}$ . Then  $\Gamma \vee \Gamma' \in \text{TC}(X, \theta)_{>0}$  and*

$$(\Gamma \vee \Gamma')_{-\infty} = P_\theta [\Gamma_{-\infty} \vee \Gamma'_{-\infty}]. \quad (9.19)$$

{eq:GammalorGammaminfty}

*If  $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ , then  $\Gamma \vee \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ .*

*For each  $\Gamma'' \in \text{TC}(X, \theta)_{>0}$  and each  $\Gamma'' \geq \Gamma$  and  $\Gamma'' \geq \Gamma'$ , we have  $\Gamma'' \geq \Gamma \vee \Gamma'$ .*

*Moreover, the operation  $\vee$  is associative and commutative.*

**Proof** It follows immediately from [Lemma 9.4.4](#) that  $\Gamma \vee \Gamma' \in \text{TC}(X, \theta)_{>0}$ , and it lies in  $\text{PSH}^{\text{NA}}(X, \theta)_{>0}$  if  $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ .

The argument of [\(9.19\)](#) is very similar to that of [\(9.15\)](#), which we leave to the readers.

Take  $\Gamma''$  as in the statement of the proposition. First observe that

$$\Gamma''_{\max} \geq \Gamma_{\max} \vee \Gamma'_{\max} = (\Gamma \vee \Gamma')_{\max}.$$

Take  $\tau < (\Gamma \vee \Gamma')_{\max}$ , we argue that

$$\Gamma''_{\tau} \geq (\Gamma \vee \Gamma')_{\tau}.$$

By the concavity of  $\Gamma''$ , this is equivalent to

$$\Gamma''_{\tau} \geq \Gamma_{\tau} \vee \Gamma'_{\tau}.$$

Therefore,

$$\Gamma'' \geq \Gamma \vee \Gamma'.$$

The commutativity and associativity of  $\vee$  are trivial.  $\square$

lma:testcurlorPthetapomega

**Lemma 9.4.5** *Let  $\Gamma, \Gamma' \in \text{TC}(X, \theta)_{>0}$  and  $\omega$  be a closed smooth positive  $(1, 1)$ -form on  $X$ . Then*

$$P_{\theta+\omega}[\Gamma \vee \Gamma'] = P_{\theta+\omega}[\Gamma] \vee P_{\theta+\omega}[\Gamma'].$$

**Proof** We first observe that

$$(P_{\theta+\omega}[\Gamma \vee \Gamma'])_{\max} = (P_{\theta+\omega}[\Gamma] \vee P_{\theta+\omega}[\Gamma'])_{\max} = \Gamma_{\max} \vee \Gamma'_{\max}.$$

Let  $\tau \in \mathbb{R}$  be less than this common value. We need to show that

$$(\Gamma \vee \Gamma')_{\tau} \sim_P (P_{\theta+\omega}[\Gamma] \vee P_{\theta+\omega}[\Gamma'])_{\tau}.$$

We need the formula [\(9.18\)](#) proved in the proof of [Lemma 9.4.4](#):

$$(\Gamma \vee \Gamma')_{\tau} = \sup^* \left\{ \lambda \Gamma_{\rho} + (1 - \lambda) \Gamma'_{\rho'} : \lambda \in (0, 1), \rho, \rho' \in \mathbb{R}, \lambda \rho + (1 - \lambda) \rho' = \tau \right\}.$$

A similar result holds with  $P_{\theta+\omega}[\Gamma]$  and  $P_{\theta+\omega}[\Gamma']$  in place of  $\Gamma$  and  $\Gamma'$ . So our assertion is a direct consequence of [Proposition 6.1.5](#) and [Proposition 6.1.6](#).  $\square$

def:testcursup

**Definition 9.4.5** Let  $(\Gamma^i)_{i \in I}$  be an increasing net in  $\text{TC}(X, \theta)_{>0}$ . Assume that

$$\sup_{i \in I} \Gamma_{\max}^i < \infty.$$

(9.20)

{eq:Gammaisupfinite}

Then we define  $\sup_{i \in I}^* \Gamma^i \in \text{TC}(X, \theta)_{>0}$  as follows:

(1) We set

$$\left( \sup_{i \in I}^* \Gamma^i \right)_{\max} = \sup_{i \in I} \Gamma_{\max}^i;$$

(2) for any  $\tau < \sup_{i \in I} \Gamma_{\max}^i$ , we let

$$\left( \sup_{i \in I}^* \Gamma^i \right)_{\tau} := \sup_{i \in I} \Gamma_{\tau}^i.$$

prop:supsincnetteestcur

**Proposition 9.4.3** *Let  $(\Gamma^i)_{i \in I}$  be an increasing net in  $\text{TC}(X, \theta)_{>0}$  satisfying (9.20). Then  $\sup_{i \in I}^* \Gamma^i$  as defined in Definition 9.4.5 lies in  $\sup_{i \in I}^* \Gamma^i \in \text{TC}(X, \theta)_{>0}$ . Moreover, if  $\Gamma^i \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$  for all  $i \in I$ , then  $\sup_{i \in I}^* \Gamma^i$  lies in  $\text{PSH}^{\text{NA}}(X, \theta)_{>0}$  as well.*

Moreover, we have

$$\left( \sup_{i \in I}^* \Gamma^i \right)_{-\infty} = \sup_{i \in I} \Gamma_{-\infty}^i.$$

(9.21)

{eq:Gammiminf}

**Proof** The first assertion follows easily from Proposition 3.1.9, while the second follows from Proposition 3.2.12.

It remains to argue (9.21). Without loss of generality, we may assume that  $I$  contains a minimal element  $i_0$ .

By Proposition 1.2.3, there is a pluripolar set  $Z \subseteq X$  such that for any  $x \in X \setminus Z$ ,

$$\left( \sup_{i \in I}^* \Gamma^i \right)_{-\infty}(x) = \sup_{\tau < \Gamma_{\max}^{i_0}} \left( \sup_{i \in I}^* \Gamma_{\tau}^i \right)(x) = \sup_{\tau < \Gamma_{\max}^{i_0}, i \in I} \Gamma_{\tau}^i(x) = \sup_{i \in I} \Gamma_{-\infty}^i(x).$$

So they are equal everywhere by Proposition 1.2.5.  $\square$

lma:suptestcurvcompatible

**Lemma 9.4.6** *Let  $(\Gamma^i)_{i \in I}$  be an increasing net in  $\text{TC}(X, \theta)_{>0}$  satisfying (9.20). Assume that  $\omega$  is a closed smooth positive  $(1, 1)$ -form on  $X$ . Then*

$$P_{\theta+\omega} \left[ \sup_{i \in I}^* \Gamma^i \right] = \sup_{i \in I}^* P_{\theta+\omega} [\Gamma^i].$$

**Proof** Observe that

$$\left( P_{\theta+\omega} \left[ \sup_{i \in I}^* \Gamma^i \right] \right)_{\max} = \left( \sup_{i \in I}^* P_{\theta+\omega} [\Gamma^i] \right)_{\max} = \sup_{i \in I} \Gamma_{\max}^i.$$

Fix  $\tau \in \mathbb{R}$  less than this common value.

It suffices to show that

$$\left( \sup_{i \in I}^* \Gamma^i \right)_{\tau} = \left( \sup_{i \in I}^* P_{\theta+\omega} [\Gamma^i] \right)_{\tau}.$$

This is an immediate consequence of [Proposition 6.1.6](#).  $\square$

**Definition 9.4.6** Let  $(\Gamma^i)_{i \in I}$  be a non-empty family in  $\text{TC}(X, \theta)_{>0}$  satisfying (9.20). Then we define

$$\sup_{i \in I}^* \Gamma^i := \sup_{J \in \text{Fin}(I)}^* \left( \bigvee_{j \in J} \Gamma^j \right). \quad (9.22)$$

Observe that by [Definition 9.4.4](#), we have

$$\sup_{J \in \text{Fin}(I)} \left( \bigvee_{j \in J} \Gamma^j \right)_{\max} = \sup_{i \in I} \Gamma_{\max}^i < \infty.$$

So (9.22) makes sense. In particular,

$$\left( \sup_{i \in I} \Gamma^i \right)_{\max} = \sup_{i \in I} \Gamma_{\max}^i. \quad (9.23)$$

It is clear that [Definition 9.4.6](#) extends both [Definition 9.4.5](#) and [Definition 9.4.4](#).

**Proposition 9.4.4** Let  $(\Gamma^i)_{i \in I}$  be a non-empty family in  $\text{TC}(X, \theta)_{>0}$  satisfying (9.20). Then  $\sup_{i \in I}^* \Gamma^i \in \text{TC}(X, \theta)_{>0}$ . Moreover, if  $\Gamma^i \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ , then so is  $\sup_{i \in I}^* \Gamma^i$ .

Finally, we have

$$\left( \sup_{i \in I}^* \Gamma^i \right)_{-\infty} = P_{\theta} \left[ \sup_{i \in I}^* \Gamma_{-\infty}^i \right]. \quad (9.24)$$

**Proof** The first assertion and the second follow from [Proposition 9.4.3](#) and [Corollary 9.4.1](#).

It remains to argue (9.24). For this purpose, it suffices to show that

$$\left( \sup_{i \in I}^* \Gamma^i \right)_{-\infty} \sim_P \sup_{i \in I}^* \Gamma_{-\infty}^i.$$

For any  $J \in \text{Fin}(I)$ , it follows from [Corollary 9.4.1](#) and [Proposition 6.1.6](#) that

$$\left( \bigvee_{j \in J} \Gamma^j \right)_{-\infty} \sim_P \bigvee_{j \in J} \Gamma_{-\infty}^j.$$

From this, applying [Proposition 6.1.6](#) and [Proposition 9.4.3](#), we conclude our assertion.  $\square$

**Lemma 9.4.7** Let  $(\Gamma^i)_{i \in I}$  be a non-empty family in  $\text{TC}(X, \theta)_{>0}$  satisfying (9.20). Assume that  $\omega$  is a closed smooth positive  $(1, 1)$ -form on  $X$ . Then

$$P_{\theta+\omega} \left[ \sup_{i \in I}^* \Gamma^i \right] = \sup_{i \in I}^* P_{\theta+\omega} [\Gamma^i].$$

**Proof** This is a direct consequence of [Lemma 9.4.6](#) and [Lemma 9.4.5](#).  $\square$

prop:testcurvChoquet

**Proposition 9.4.5** Let  $(\Gamma^i)_{i \in I}$  be a non-empty family in  $\text{TC}(X, \theta)_{>0}$  satisfying (9.20). Then there is a countable subset  $I' \subseteq I$  such that

$$\sup_{i \in I}^* \Gamma^i = \sup_{i \in I'}^* \Gamma^i.$$

**Proof** We may assume that  $I$  is infinite.

It follows from [Proposition 1.2.2](#) that we can find a countable subset  $I' \subseteq I$  such that for each

$$\tau \in \left( -\infty, \sup_{i \in I}^* \Gamma_{\max}^i \right) \cap \mathbb{Q},$$

we have

$$\sup_{i \in I}^* \Gamma_{\tau}^i = \sup_{i \in I'}^* \Gamma_{\tau}^i.$$

Let  $\Gamma' = \sup_{i \in I'}^* \Gamma^i$ . Then clearly,  $\Gamma' \leq \Gamma$ . We claim that they are actually equal. For this purpose, it suffices to show that for any  $\tau < \sup_{i \in I}^* \Gamma_{\max}^i$ , we have

$$\int_X (\theta + \text{dd}^c \Gamma'_{\tau})^n = \int_X (\theta + \text{dd}^c \Gamma_{\tau})^n.$$

Since we know that this holds on a dense subset of  $\tau$ , this holds everywhere by [Theorem 2.3.3](#).  $\square$

prop:supGammiOtherprop

**Proposition 9.4.6** Let  $(\Gamma^i)_{i \in I}$  be a non-empty family in  $\text{TC}(X, \theta)_{>0}$  satisfying (9.20). Let  $C \in \mathbb{R}$ . Then

$$\sup_{i \in I}^* (\Gamma^i + C) = \sup_{i \in I}^* \Gamma^i + C.$$

Suppose that  $(\Gamma'^i)_{i \in I}$  is another family in  $\text{TC}(X, \theta)_{>0}$  satisfying (9.20). Suppose that  $\Gamma^i \leq \Gamma'^i$  for all  $i \in I$ , then

$$\sup_{i \in I}^* \Gamma^i \leq \sup_{i \in I}^* \Gamma'^i.$$

**Proof** This is immediate by definition.  $\square$

def:res

**Definition 9.4.7** Let  $\Gamma \in \text{TC}(X, \theta)_{>0}$  and  $\lambda > 0$ , we define  $\lambda\Gamma \in \text{TC}(X, \lambda\theta)_{>0}$  as follows:

(1) We set

$$(\lambda\Gamma)_{\max} = \lambda\Gamma_{\max};$$

(2) for any  $\tau < \lambda\Gamma_{\max}$ , we set

$$(\lambda\Gamma)_{\tau} = \lambda\Gamma_{\lambda^{-1}\tau}.$$

prop:testcurrecaling

**Proposition 9.4.7** Let  $\Gamma \in \text{TC}(X, \theta)_{>0}$  and  $\lambda > 0$ , then  $\lambda\Gamma$  as defined in [Definition 9.4.7](#) lies in  $\text{TC}(X, \lambda\theta)_{>0}$ . Moreover, if  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ , then  $\lambda\Gamma \in \text{PSH}^{\text{NA}}(X, \lambda\theta)_{>0}$ .

We have

$$(\lambda\Gamma)_{-\infty} = \lambda\Gamma_{-\infty}. \quad (9.25)$$

prop:resclcompat

**Proposition 9.4.8** *Let  $\Gamma \in \text{TC}(X, \theta)_{>0}$ ,  $\Gamma' \in \text{TC}(X, \theta')_{>0}$ ,  $C \in \mathbb{R}$  and  $\lambda, \lambda' > 0$ , we have*

$$\begin{aligned}\lambda(\Gamma + \Gamma') &= \lambda\Gamma + \lambda\Gamma', \\ (\lambda\lambda')\Gamma &= \lambda(\lambda'\Gamma), \\ \lambda(\Gamma + C) &= \lambda\Gamma + \lambda C.\end{aligned}$$

*Suppose that  $(\Gamma^i)_{i \in I}$  is a non-empty family in  $\text{TC}(X, \theta)_{>0}$  satisfying (9.20), then*

$$\lambda \left( \sup_{i \in I}^* \Gamma^i \right) = \sup_{i \in I}^* (\lambda \Gamma^i).$$

lma:testcurvrescompatible

**Lemma 9.4.8** *Let  $\Gamma \in \text{TC}(X, \theta)_{>0}$  and  $\lambda > 0$ . Then for any closed smooth positive  $(1, 1)$ -form  $\omega$  on  $X$ , we have*

$$P_{\lambda(\theta+\omega)}[\lambda\Gamma] = \lambda P_{\theta+\omega}[\Gamma].$$

**Proof** This is clear by definition. □



# Chapter 10

## The theory of Okounkov bodies

chap:Okou

In this chapter, we apply our theory of singularities to the study of Okounkov bodies. We establish the theory of partial Okounkov bodies, which are convex bodies constructed from a given plurisubharmonic singularity. These objects allow us to reduce many problems in pluripotential theory to problems in convex geometry, which are usually simpler.

We will establish two related theories. One in the algebraic setting in [Section 10.2](#) and one in the transcendental setting in [Section 10.3](#).

### 10.1 Flags and valuations

#### 10.1.1 The algebraic setting

subsec:flagvalalgebraic

Let  $X$  be an irreducible normal projective variety of dimension  $n$ .

def:admf1

**Definition 10.1.1** An *admissible flag*  $Y_\bullet$  on  $X$  is a flag of subvarieties

$$X = Y_0 \supseteq Y_1 \supseteq \cdots \supseteq Y_n$$

such that  $Y_i$  is irreducible of codimension  $i$  and is smooth at the point  $Y_n$ .

Given any admissible flag  $Y_\bullet$ , we can define a rank  $n$  valuation  $v_{Y_\bullet} : \mathbb{C}(X)^\times \rightarrow \mathbb{Z}^n$ . Here we consider  $\mathbb{Z}^n$  as a totally ordered Abelian group with the lexicographic order. We sometimes write  $\mathbb{Z}_{\text{lex}}^n$  to emphasize this point.

The automorphism group  $\text{Aut}(\mathbb{Z}_{\text{lex}}^n)$  of  $\mathbb{Z}_{\text{lex}}^n$  is then identified with the subgroup of  $\text{GL}(n, \mathbb{Z})$  consisting of matrices of the form  $I + U$ , where  $I$  is the identity matrix and  $U$  is a strictly upper triangular matrix with elements in  $\mathbb{Z}$ .

We recall the definition: Let  $s \in \mathbb{C}(X)^\times$ . Let  $v(s)_1 = \text{ord}_{Y_1} s$ . After localization around  $Y_n$ , we can take a local defining equation  $t^1$  of  $Y_1$ , set  $s_1 = (s(t^1)^{-v_1(s)})|_{Y_1}$ . Then  $s_1 \in \mathbb{C}(Y_1)^\times$ . We can repeat this construction with  $Y_2$  in place of  $Y_1$  to get  $v(s)_2$  and  $s_2$ . Repeating this construction  $n$  times, we get

$$\nu_{Y_\bullet}(s) = \nu(s) = (\nu(s)_1, \nu(s)_2, \dots, \nu(s)_n) \in \mathbb{Z}^n.$$

It is easy to verify that  $\nu$  is indeed a rank  $n$  valuation.

The same construction can be applied to define  $\nu_{Y_\bullet}(s)$  when  $s \in H^0(X, L)$  or  $\nu_{Y_\bullet}(D)$  when  $D$  is an effective divisor on  $X$ .

rmk:Abhyankar

*Remark 10.1.1* Conversely, by a theorem of Abhyankar, any valuation of  $\mathbb{C}(X)$  with Noetherian valuation ring of rank  $n$  is equivalent to a valuation taking value in  $\mathbb{Z}^n$ , see [FK18, Chapter 0, Theorem 6.5.2]. As shown in [CFKLS17, Theorem 2.9], any such valuation is equivalent<sup>1</sup> to (but not necessarily equal to) a valuation induced by an admissible flag on a modification of  $X$ .

### 10.1.2 The transcendental setting

Let  $X$  be a connected compact Kähler manifold of dimension  $n$ .

**Definition 10.1.2** A *smooth flag*  $Y_\bullet$  on  $X$  consists of a flag of connected submanifolds of  $X$ :

$$X = Y_0 \supseteq Y_1 \supseteq \dots \supseteq Y_n,$$

where  $Y_i$  has dimension  $n - i$ .

In this section, we will fix a smooth flag  $Y_\bullet$  on  $X$ .

def:valcurr

**Definition 10.1.3** Let  $T$  be a closed positive  $(1, 1)$ -current on  $X$ . We define the *valuation* of  $T$  along  $Y_\bullet$  as

$$\nu_{Y_\bullet}(T) = (\nu_{Y_\bullet}(T)_1, \dots, \nu_{Y_\bullet}(T)_n) \in \mathbb{R}_{\geq 0}^n$$

by induction on  $n$ . When  $n = 0$ , we define  $\nu_{Y_\bullet}(T)$  as the unique point in  $\mathbb{R}^0$ . When  $n > 1$ , we define

$$\nu_{Y_\bullet}(T)_1(T) = \nu(T, Y_1);$$

Then for  $i = 2, \dots, n$ , we define

$$\nu_{Y_\bullet}(T)_i = \nu_{Y_1 \supseteq \dots \supseteq Y_n}(\text{Tr}_{Y_1}(T - \nu(T, Y_1)[Y_1]))_{i-1}.$$

**Proposition 10.1.1** Let  $T$  be a closed positive  $(1, 1)$ -current on  $X$ . Then  $\nu_{Y_\bullet}(T) \in \mathbb{R}_{\geq 0}^n$  defined in *Definition 10.1.3* is independent of the choices of the trace operators in the definition. Moreover,  $\nu_{Y_\bullet}(T)$  depends only on the  $\mathcal{I}$ -equivalence class of  $T$ .

**Proof** We will prove both statements at the same time by induction on  $n \geq 0$ . The case  $n = 0$  is trivial.

<sup>1</sup> Two valuations  $\nu, \nu'$  with value in  $\mathbb{Z}^n$  are equivalent if one can find a matrix  $G$  of the form  $I + N$ , where  $N$  is strictly upper triangular with integral entries, such that  $\nu' = \nu G$ .

Let us consider the case  $n > 0$  and assume that the result is known in dimension  $n - 1$ . We first observe that  $\nu_{Y_\bullet}(T)$  is independent of the choice of the trace operator: different choices of  $\text{Tr}_{Y_1}(T - \nu(T, Y_1)[Y_1])$  are  $\mathcal{I}$ -equivalent by [Proposition 8.1.2](#). Therefore, by induction, its valuation is well-defined.

Next, let  $T'$  be another closed positive  $(1, 1)$ -current such that  $T \sim_{\mathcal{I}} T'$ . Using [Proposition 3.2.1](#), we know that  $\nu(T, Y_1) = \nu(T', Y_1)$ . Therefore,

$$T - \nu(T, Y_1)[Y_1] \sim_{\mathcal{I}} T' - \nu(T', Y_1)[Y_1].$$

It follows by induction that

$$\nu_{Y_1 \supseteq \dots \supseteq Y_n}(\text{Tr}_{Y_1}(T - \nu(T, Y_1)[Y_1])) = \nu_{Y_1 \supseteq \dots \supseteq Y_n}(\text{Tr}_{Y_1}(T' - \nu(T', Y_1)[Y_1])).$$

ex:valuationdivcompatible

*Example 10.1.1* When  $X$  is projective, we have

$$\nu_{Y_\bullet}([D]) = \nu_{Y_\bullet}(D),$$

where the right-hand side is defined in [Section 10.1.1](#).

prop:nuvaluationlinear

**Proposition 10.1.2** *Let  $T, S$  be closed positive  $(1, 1)$ -currents on  $X$ ,  $\lambda \in \mathbb{R}_{\geq 0}$ . Then*

(1) *if  $T \leq_{\mathcal{I}} S$ , we have*

$$\nu_{Y_\bullet}(T) \geq_{\text{lex}} \nu_{Y_\bullet}(S). \quad (10.1)$$

{eq:nuTS}

(2) *We have the following additivity property:*

$$\nu_{Y_\bullet}(T + S) = \nu_{Y_\bullet}(T) + \nu_{Y_\bullet}(S), \quad \nu_{Y_\bullet}(\lambda T) = \lambda \nu_{Y_\bullet}(T). \quad (10.2)$$

{eq:nuvaluationlinear}

**Proof** (1) We make an induction on  $n \geq 0$ . The case  $n = 0, 1$  is trivial. Assume that  $n \geq 2$  and the case  $n - 1$  is known. Observe that  $\nu(T, Y_1) \geq \nu(S, Y_1)$ , if the inequality is strict, we are done. So let us assume that  $\nu(T, Y_1) = \nu(S, Y_1)$ . By [Proposition 8.2.1](#), we find that

$$\text{Tr}_{Y_1}(T - \nu(T, Y_1)[Y_1]) \leq_{\mathcal{I}} \text{Tr}_{Y_1}(S - \nu(T, Y_1)[Y_1]).$$

By the inductive hypothesis, we conclude [\(10.1\)](#).

(2) We make an induction on  $n \geq 0$ . The cases  $n = 0, 1$  are trivial. Assume that  $n \geq 2$  and the case  $n - 1$  is known. By [Proposition 1.4.2](#), we have

$$\nu(T + S, Y_1) = \nu(T, Y_1) + \nu(S, Y_1), \quad \nu(\lambda T, Y_1) = \lambda \nu(T, Y_1).$$

By [Proposition 8.2.1](#), we have

$$\begin{aligned} \text{Tr}_{Y_1}(T + S - \nu(T + S, Y_1)[Y_1]) &\sim_P \text{Tr}_{Y_1}(T - \nu(T, Y_1)[Y_1]) + \text{Tr}_{Y_1}(S - \nu(S, Y_1)[Y_1]), \\ \text{Tr}_{Y_1}(\lambda T - \nu(\lambda T, Y_1)[Y_1]) &\sim_P \lambda \text{Tr}_{Y_1}(T - \nu(T, Y_1)[Y_1]). \end{aligned}$$

By the inductive hypothesis, we conclude [\(10.2\)](#).

**Definition 10.1.4** Let  $\pi: Z \rightarrow X$  be a proper bimeromorphic morphism with  $Z$  being a Kähler manifold. We say that a smooth flag  $W_\bullet$  on  $Z$  is a *lifting* of  $Y_\bullet$  to  $Z$  if the restriction of  $\pi$  to  $W_i \rightarrow Y_i$  is defined and bimeromorphic for each  $i = 0, \dots, n$ .

In this case, we define  $\text{cor}(Y_\bullet, \pi) \in \text{Aut}(\mathbb{Z}_{\text{lex}}^n)$  inductively as follows:

$$\text{cor}(Y_\bullet, \pi) := \begin{bmatrix} 1 & -\nu_{W_1 \supseteq \dots \supseteq W_n}((\pi^*[Y_1] - [W_1])|_{W_1}) \\ 0 & \text{cor}(Y_1 \supseteq \dots \supseteq Y_n, \pi|_{W_1}: W_1 \rightarrow Y_1) \end{bmatrix}. \quad (10.3)$$

{eq:correcur}

We observe that a lifting  $W_\bullet$  of  $Y_\bullet$  on  $Z$  is unique if it exists. For each  $i = 0, \dots, n-1$ , the component  $W_{i+1}$  is necessarily the strict transform of  $Y_{i+1}$  with respect to the bimeromorphic morphism  $W_i \rightarrow Y_i$ . We shall also say that  $(W_\bullet, \text{cor}(Y_\bullet, \pi))$  is the *lifting* of  $Y_\bullet$  to  $Z$ .

prop:cormult

**Proposition 10.1.3** Let  $\pi: Z \rightarrow X, p: Z' \rightarrow Z$  be proper bimeromorphic morphisms with  $Z$  and  $Z'$  being Kähler manifolds. Assume that  $Y_\bullet$  admits a lifting  $W_\bullet$  (resp.  $W'_\bullet$ ) to  $Z$  (resp.  $Z'$ ). Then

$$\text{cor}(Y_\bullet, \pi \circ p) = \text{cor}(Y_\bullet, \pi) \text{cor}(W_\bullet, p). \quad (10.4)$$

{eq:cormul}

**Proof** We let  $\pi' = \pi \circ p$ :

$$\begin{array}{ccc} Z' & \xrightarrow{p} & Z \\ & \searrow \pi' & \swarrow \pi \\ & X & \end{array}.$$

We make induction on  $n \geq 1$ . The case  $n = 1$  is trivial. Assume that  $n \geq 2$  and the case  $n - 1$  has been solved. Then by (10.3), the desired formula (10.4) can be reformulated as

$$\begin{aligned} & \begin{bmatrix} 1 & -\nu_{W'_1 \supseteq \dots \supseteq W'_n}((\pi'^*[Y_1] - [W'_1])|_{W'_1}) \\ 0 & \text{cor}(Y_1 \supseteq \dots \supseteq Y_n, \pi'|_{W'_1}: W'_1 \rightarrow Y_1) \end{bmatrix} = \\ & \begin{bmatrix} 1 & -\nu_{W_1 \supseteq \dots \supseteq W_n}((\pi^*[Y_1] - [W_1])|_{W_1}) \\ 0 & \text{cor}(Y_1 \supseteq \dots \supseteq Y_n, \pi|_{W_1}: W_1 \rightarrow Y_1) \end{bmatrix} \cdot \\ & \begin{bmatrix} 1 & -\nu_{W'_1 \supseteq \dots \supseteq W'_n}((p^*[W_1] - [W'_1])|_{W'_1}) \\ 0 & \text{cor}(W_1 \supseteq \dots \supseteq W_n, p|_{W'_1}: W'_1 \rightarrow W_1) \end{bmatrix} \end{aligned}$$

By the inductive hypothesis, this is equivalent to

$$\begin{aligned} & \nu_{W'_1 \supseteq \dots \supseteq W'_n}((\pi'^*[Y_1] - [W'_1])|_{W'_1}) = \nu_{W'_1 \supseteq \dots \supseteq W'_n}((p^*[W_1] - [W'_1])|_{W'_1}) + \\ & \nu_{W_1 \supseteq \dots \supseteq W_n}((\pi^*[Y_1] - [W_1])|_{W_1}) \text{cor}(W_1 \supseteq \dots \supseteq W_n, p|_{W'_1}: W'_1 \rightarrow W_1), \end{aligned}$$

which can be further rewritten as

$$\begin{aligned} \nu_{W'_1 \supseteq \dots \supseteq W'_n}((\pi^*[Y_1] - [W'_1])|_{W'_1}) &= \nu_{W'_1 \supseteq \dots \supseteq W'_n}((p^*[W_1] - [W'_1])|_{W'_1}) + \\ &\quad \nu_{W'_1 \supseteq \dots \supseteq W'_n}(p|_{W'_1}^*(\pi^*[Y_1] - [W_1])|_{W_1}). \end{aligned}$$

This follows from [Proposition 10.1.2](#).  $\square$

prop:cormatrix

**Proposition 10.1.4** *Let  $\pi: Z \rightarrow X$  be a proper bimeromorphic morphism with  $Z$  being a Kähler manifold. Let  $W_\bullet$  be a lifting of  $Y_\bullet$ , then for any closed positive  $(1, 1)$ -current  $T$  on  $X$ , we have*

$$\nu_{W_\bullet}(\pi^*T) = \nu_{Y_\bullet}(T) \operatorname{cor}(Y_\bullet, \pi). \quad (10.5)$$

**Proof** We make induction on  $n \geq 0$ . The case  $n = 0$  is trivial. In general, assume that  $n \geq 1$  and the result is proved in dimension  $n - 1$ .

For simplicity, we write  $\nu = \nu_{Y_\bullet}$  and  $\nu' = \nu_{W_\bullet}$ . Let  $\mu$  (resp.  $\mu'$ ) be the valuation of currents defined by the truncated flag  $Y_1 \supseteq \dots \supseteq Y_n$  (resp.  $W_1 \supseteq \dots \supseteq W_n$ ). Then we need to show that

$$\begin{aligned} &[\nu'(\pi^*T)_1 \mu'(\operatorname{Tr}_{W_1}(\pi^*T - \nu'(\pi^*T)_1[W_1]))] \\ &= [\nu(T)_1 \mu(\operatorname{Tr}_{Y_1}(T - \nu(T)_1[Y_1]))] \operatorname{cor}(Y_\bullet, \pi). \end{aligned} \quad (10.6)$$

{eq:mubiration}

By Zariski's main theorem,

$$\nu'(\pi^*T)_1 = \nu(T)_1 =: c.$$

By the inductive hypothesis, we have

$$\mu'(\Pi^* \operatorname{Tr}_{Y_1}(T - c[Y_1])) = \mu(\operatorname{Tr}_{Y_1}(T - c[Y_1])) \operatorname{cor}(Y_1 \supseteq \dots \supseteq Y_n, \Pi), \quad (10.7)$$

{eq: ind\_hypos}

where  $\Pi: W_1 \rightarrow Y_1$  is the restriction of  $\pi$ . By [Lemma 8.2.1](#) and [Proposition 8.2.1](#),

$$\begin{aligned} \Pi^* \operatorname{Tr}_{Y_1}(T - c[Y_1]) &\sim_P \operatorname{Tr}_{W_1}(\pi^*(T - c[Y_1])) \\ &\sim_P \operatorname{Tr}_{W_1}(\pi^*T - c[W_1]) + c \operatorname{Tr}_{W_1}(\pi^*[Y_1] - [W_1]). \end{aligned}$$

So

$$\mu'(\Pi^* \operatorname{Tr}_{Y_1}(T - c[Y_1])) = \mu'(\operatorname{Tr}_{W_1}(\pi^*T - c[W_1])) + c\mu'(\operatorname{Tr}_{W_1}(\pi^*[Y_1] - [W_1])).$$

Combining the above with (10.7), we see that (10.6) follows.  $\square$

thm:lifttableflag

**Theorem 10.1.1** *Let  $\pi: Z \rightarrow X$  be a proper bimeromorphic morphism from a reduced complex space  $Z$ . Then there is a modification  $W \rightarrow X$  dominating  $Z \rightarrow X$  such that  $Y_\bullet$  admits a lifting to  $W$ .*

**Proof** By Hironaka's Chow lemma, we may assume that  $\pi$  is a modification.

We begin by setting  $W_0 = Z$ . We will construct  $W_i$  inductively for each  $i$ . Assume that for  $0 \leq i < n$  a smooth partial flag  $W_0 \supset \dots \supset W_i$  has been constructed on a modification  $\pi_i: Z_i \rightarrow Z$  so that  $\pi \circ \pi_i$  restricts to bimeromorphic morphisms  $W_j \rightarrow Y_j$  for each  $j = 0, \dots, i$ .

By Zariski's main theorem,  $W_i \rightarrow Y_i$  is an isomorphism outside a codimension 2 subset of  $Y_i$ . We let  $W_{i+1}$  be the strict transform of  $Y_{i+1}$  in  $W_i$ . The problem is that  $W_{i+1}$  is not necessarily smooth.

We will further modify  $Z_i$  and lift  $W_1, \dots, W_{i+1}$  in order to make the flag smooth. Take the embedded resolution of  $(W_j, W_{i+1})$ , say  $W'_j \rightarrow W_j$  for each  $j = 0, \dots, i$ .

We have canonical embeddings  $W'_i \hookrightarrow W'_{i-1} \hookrightarrow \dots \hookrightarrow W'_0$  making the following diagram commutative:

$$\begin{array}{ccccccc} W'_i & \hookrightarrow & W'_{i-1} & \hookrightarrow & \dots & \hookrightarrow & W'_0 \\ \downarrow & & \downarrow & & \vdots & & \downarrow \\ W_i & \hookrightarrow & W_{i-1} & \hookrightarrow & \dots & \hookrightarrow & W_0 \end{array}$$

Let  $W'_{i+1}$  be the strict transform of  $W_{i+1}$  in  $W'_i$ . It suffices to define  $\pi_{i+1}$  as the morphism  $W'_0 \rightarrow Z_i \rightarrow Z$  and replace  $W_0 \supset \dots \supset W_{i+1}$  by  $W'_0 \supset \dots \supset W'_{i+1}$ .  $\square$

## 10.2 Algebraic partial Okounkov bodies

sec:PoB

Let  $X$  be a connected smooth complex projective variety of dimension  $n$  and  $(L, h)$  be a Hermitian big line bundle on  $X$ .

Let  $h_0$  be a smooth Hermitian metric on  $L$ . Let  $\theta = c_1(L, h_0)$ . Then we can identify  $h$  with a function  $\varphi \in \text{PSH}(X, \theta)$ . We will use interchangeably the notations  $(\theta, \varphi)$  and  $(L, h)$ .

Fix a rank  $n$  valuation  $\nu: \mathbb{C}(X)^\times \rightarrow \mathbb{Z}^n$ , which without loss of generality can be assumed to be surjective.

We will adopt the notations of [Appendix C.2](#).

### 10.2.1 The spaces of sections

**Definition 10.2.1** We will write

$$\begin{aligned} \Gamma(\theta, \varphi) &:= \{(\nu(s), k) : k \in \mathbb{N}, s \in H^0(X, L^k \otimes \mathcal{I}(k\varphi))^\times\}, \\ \Delta_k(\theta, \varphi) &:= \text{Conv} \{k^{-1}\nu(f) : f \in H^0(X, L^k \otimes \mathcal{I}(k\varphi))^\times\} \subseteq \mathbb{R}^n, \quad k \geq 0. \end{aligned}$$

When  $\theta = V_\theta$ , we simply write  $\Gamma(L)$  and  $\Delta_k(L)$  instead.

Here  $\text{Conv}$  denotes the convex hull. For large enough  $k$ ,  $\Delta_k(\theta, \varphi)$  is non-empty thanks to [Theorem 7.3.1](#).

**Definition 10.2.2** Assume that  $\varphi$  has analytic singularities. We define

$$\Gamma^\infty(\theta, \varphi) := \{(\nu(s), k) : k \in \mathbb{N}, s \in H^0(X, L^k \otimes \mathcal{I}_\infty(k\varphi))^\times\}. \quad (10.8)$$

{eq:Weps1}

For later use, we introduce a twisted version as well.

**Definition 10.2.3** If  $T$  is a holomorphic line bundle on  $X$ , we introduce

$$\begin{aligned}\Delta_{k,T}(\theta, \varphi) &:= \text{Conv} \left\{ k^{-1} \nu(f) : f \in H^0(X, T \otimes L^k \otimes I(k\varphi))^\times \right\} \subseteq \mathbb{R}^n, \\ \Delta_{k,T}(L) &:= \text{Conv} \left\{ k^{-1} \nu(f) : f \in H^0(X, T \otimes L^k)^\times \right\} \subseteq \mathbb{R}^n.\end{aligned}$$

### 10.2.2 Algebraic Okounkov bodies

prop:Okounbiglbd1

**Proposition 10.2.1** *There is a convex body  $\Delta \in \mathcal{K}_n$  such that  $\Gamma(L) \in \mathcal{S}'(\Delta)$ .*

**Proof Step 1.** We first show that there is  $\Delta \in \mathcal{K}_n$  such that  $\Delta_k(L) \subseteq \Delta$ . For this purpose, using [Remark 10.1.1](#), we may assume that  $\nu$  is induced by an admissible flag  $Y_\bullet$  on  $X$ .

Fix  $s \in H^0(X, L^k)^\times$  for some  $k \in \mathbb{Z}_{>0}$ . Assume that  $s \neq 0$ . We need to show that for each  $i = 1, \dots, n$ ,  $\nu(s)_i \leq Ck$  for some constant  $C > 0$ , independent of the choices of  $k$  and  $s$ .

Fix an ample divisor  $H$  on  $X$ . Take a large enough integer  $b_1 > 0$  such that

$$(L - b_1 Y_1) \cdot H^{n-1} < 0.$$

Then  $\nu(s)_1 \leq b_1 k$ . Next take a large enough integer  $b_2$  such that

$$((L - aY_1)|_{Y_1} - b_2 Y_2) \cdot H^{n-2} < 0.$$

It follows that  $\nu(s)_2 \leq b_2 k$ . Continue in this manner, we conclude that  $\nu(s)_i/k$  is bounded for each  $i$ .

**Step 2.** Observe that  $\Gamma(L)$  is clearly a semigroup. It remains to show that  $\Gamma(L)$  generates  $\mathbb{Z}^{n+1}$  as an Abelian group.

For this purpose, take two very ample divisors  $A$  and  $B$  so that  $L = \mathcal{O}_X(A - B)$ . After choosing  $A$  and  $B$  ample enough, we may guarantee that there exist sections  $s_0 \in H^0(X, A)$ ,  $t_i \in H^0(X, B)$  for  $i = 0, \dots, n$  such that

$$\nu(s_0) = \nu(t_0) = 0$$

and  $\nu(t_i)$  is the  $i$ -th unit vector  $e_i \in \mathbb{R}^n$  for  $i = 1, \dots, n$ .

Since  $L$  is big, we can find  $m_0 > 0$  such that for any  $m \geq m_0$  we can find an effective divisor  $F_m$  on  $X$  linearly equivalent to  $mL - B$ . Let  $f_m = \nu([F_m])$ . Then we find that

$$(f_m, m), (f_m + e_1, m), \dots, (f_m + e_n, m) \in \Gamma(L).$$

Since  $(m+1)L$  is linearly equivalent to  $A + F_m$ , so

$$(f_m, m+1) \in \Gamma(L).$$

It follows that  $\Gamma(L)$  generates  $\mathbb{Z}^{n+1}$ .  $\square$

Thanks to [Proposition 10.2.1](#), we can introduce the next definition.

**Definition 10.2.4** We define the *Okounkov body* of  $L$  with respect to the valuation  $v$  as

$$\Delta_v(L) := \Delta(\Gamma(L)).$$

prop:Okounonlydepnum

**Proposition 10.2.2** *The Okounkov body  $\Delta_v(L)$  depends only on the numerical class of  $L$ .*

See [\[LM09, Proposition 4.1\]](#) for the elegant proof.

cor:Okounvol

**Corollary 10.2.1** *We have*

$$\text{vol } \Delta_v(L) = \frac{1}{n!} \text{vol } L. \quad (10.9)$$

**Proof** This follows immediately from [Proposition 10.2.1](#) and [Theorem C.2.1](#).  $\square$

prop:GammaepsSp

**Proposition 10.2.3** *Assume that  $\varphi$  has analytic singularities and  $\theta_\varphi$  is a Kähler current. Then we have*

$$\Gamma^\infty(\theta, \varphi) \in \mathcal{S}'(X, \theta)$$

and

$$\text{vol } \Gamma^\infty(\theta, \varphi) = \frac{1}{n!} \int_X \theta_\varphi^n.$$

**Proof** Replacing  $X$  by a modification, we may assume that  $\varphi$  has log singularities along an effective  $\mathbb{Q}$ -divisor  $D$ . See [Theorem 1.6.1](#).

In this case,

$$\Gamma^\infty(\theta, \varphi) = \{(\nu(s), k) : k \in \mathbb{N}, s \in H^0(X, L^k \otimes \mathcal{O}_X(-\lfloor kD \rfloor))\}.$$

Since  $L - D$  is ample by [Lemma 1.6.1](#), our assertion follows from the same argument as [Proposition 10.2.1](#).  $\square$

We first extend [Theorem C.2.1](#) to the twisted case.

prop-Deltaconvtwisted

**Proposition 10.2.4** *For any holomorphic line bundle  $T$  on  $X$ , as  $k \rightarrow \infty$*

$$\Delta_{k,T}(L) \xrightarrow{d_{\text{Haus}}} \Delta_v(L).$$

**Proof** As  $L$  is big, we can take  $k_0 \in \mathbb{Z}_{>0}$  so that

- (1)  $T^{-1} \otimes L^{k_0}$  admits a non-zero global holomorphic section  $s_0$ , and
- (2)  $T \otimes L^{k_0}$  admits a non-zero global holomorphic section  $s_1$ .



Then for  $k \in \mathbb{Z}_{>k_0}$ , we have injective linear maps

$$H^0(X, L^{k-k_0}) \xrightarrow{\times s_1} H^0(X, T \otimes L^k) \xrightarrow{\times s_0} H^0(X, L^{k+k_0}).$$

It follows that

$$(k - k_0)\Delta_{k-k_0}(L) + \nu(s_1) \subseteq k\Delta_{k,T}(L) \subseteq (k + k_0)\Delta_{k+k_0}(L) - \nu(s_0).$$

Using [Theorem C.2.1](#), we conclude.  $\square$

prop:subaddOkoun

**Proposition 10.2.5** *Let  $L'$  be another big line bundle on  $X$ . Then*

$$\Delta_\nu(L) + \Delta_\nu(L') \subseteq \Delta_\nu(L \otimes L').$$

**Proof** Observe that for each  $k \in \mathbb{N}$ , we have

$$\Delta_k(L) + \Delta_k(L') \subseteq \Delta_k(L \otimes L').$$

So our assertion follows immediately from [Theorem C.2.1](#).  $\square$

prop:Okourescaling

**Proposition 10.2.6** *For any  $a \in \mathbb{Z}_{>0}$ , we have*

$$\Delta_\nu(L^a) = a\Delta_\nu(L).$$

**Proof** This is an immediate consequence of [Theorem C.2.1](#).  $\square$

### 10.2.3 Construction of partial Okounkov bodies

thm:Gammaasg

**Theorem 10.2.1** *We have*

$$\Gamma(\theta, \varphi) \in \overline{\mathcal{S}'(\Delta_\nu(L))}_{>0}.$$

This theorem allows us to give the following definition:

**Definition 10.2.5** The *partial Okounkov body* of  $(L, h)$  is defined as

$$\Delta_\nu(L, h) = \Delta_\nu(\theta, \varphi) := \Delta(\Gamma(\theta, \varphi)). \quad (10.10)$$

{eq:Deltalbdef}

When  $\nu$  is induced by an admissible flag  $Y_\bullet$  on  $X$  (see [Definition 10.1.1](#)), we also say that  $\Delta_\nu(\theta, \varphi)$  the *partial Okounkov body* of  $(L, h)$  or of  $(\theta, \varphi)$  with respect to  $Y_\bullet$ . In this case, we also write  $\Delta_{Y_\bullet}$  instead of  $\Delta_\nu$ .

cor:POBvolume

**Corollary 10.2.2** *We have*

$$\text{vol } \Delta_\nu(\theta, \varphi) = \frac{1}{n!} \text{vol } \theta_\varphi. \quad (10.11)$$

{eq:Okov}

**Proof** This follows immediately from [Theorem 10.2.1](#), [Theorem 7.3.1](#) and [Theorem C.2.2](#).  $\square$

We will prove [Theorem 10.2.1](#) and [Corollary 10.2.2](#) at the same time.

**Proof Step 1.** We first assume that  $\varphi$  has analytic singularities and  $\theta_\varphi$  is a Kähler current.

We claim that

$$d_{\text{sg}}(\Gamma^\infty(\theta, \varphi), \Gamma(\theta, \varphi)) = 0. \quad (10.12)$$

{eq:Gamma0Gammaanalytic}

Observe that for each  $\epsilon \in \mathbb{Q}_{>0}$ , we have

$$H^0(X, L^k \otimes \mathcal{I}_\infty(k\varphi)) \subseteq H^0(X, L^k \otimes \mathcal{I}(k\varphi)) \subseteq H^0(X, L^k \otimes \mathcal{I}_\infty(k(1-\epsilon)\varphi))$$

for all large enough  $k$ . This is a consequence of [Lemma 1.6.3](#). Therefore, it suffices to show that

$$\lim_{\mathbb{Q} \ni \epsilon \rightarrow 0+} \text{vol } \Gamma^\infty(\theta, (1-\epsilon)\varphi) = \text{vol } \Gamma^\infty(\theta, \varphi).$$

This follows from the explicit formula in [Proposition 10.2.3](#).

**Step 2.** We next handle the case where  $\theta_\varphi$  is a Kähler current.

Let  $(\varphi_j)_j$  be a quasi-equisingular approximation of  $\varphi$  in  $\text{PSH}(X, \theta)$ . Then  $\varphi_j \xrightarrow{d_S} P_\theta[\varphi]_I$  by [Corollary 7.1.2](#).

In this case, it suffices to prove that

$$\Gamma(\theta, \varphi_j) \xrightarrow{d_{\text{sg}}} \Gamma(\theta, \varphi). \quad (10.13)$$

{eq:WtoWclaim}

In fact, by [Theorem 7.3.1](#), we have

$$\begin{aligned} & d_{\text{sg}}(\Gamma(\theta, \varphi_j), \Gamma(\theta, \varphi)) \\ &= \overline{\lim}_{k \rightarrow \infty} k^{-n} \left( H^0(X, L^k \otimes \mathcal{I}(k\varphi_j)) - H^0(X, L^k \otimes \mathcal{I}(k\varphi)) \right) \\ &= \lim_{k \rightarrow \infty} k^{-n} H^0(X, L^k \otimes \mathcal{I}(k\varphi_j)) - \lim_{k \rightarrow \infty} k^{-n} H^0(X, L^k \otimes \mathcal{I}(k\varphi)) \\ &= \frac{1}{n!} \text{vol } \theta_{\varphi_j} - \frac{1}{n!} \text{vol } \theta_\varphi. \end{aligned}$$

Letting  $j \rightarrow \infty$ , we conclude (10.13) by [Theorem 6.2.5](#).

**Step 3.** Now we only assume that  $\text{vol } \theta_\varphi > 0$ . We may replace  $\varphi$  with  $P_\theta[\varphi]_I$  and then assume that  $\varphi \in \text{PSH}(X, \theta)_{>0}$ .

Take a potential  $\psi \in \text{PSH}(X, \theta)$  such that  $\psi \leq \varphi$  and  $\theta_\psi$  is a Kähler current. The existence of  $\psi$  is proved in [Lemma 2.3.2](#). For each  $\epsilon \in (0, 1)$ , let  $\varphi_\epsilon = (1-\epsilon)\varphi + \epsilon\psi$ . It suffices to show that

$$\Gamma(\theta, \varphi_\epsilon) \xrightarrow{d_{\text{sg}}} \Gamma(\theta, \varphi)$$

as  $\epsilon \rightarrow 0+$ . We compute using [Theorem 7.3.1](#):

$$\begin{aligned}
& d_{\text{sg}}(\Gamma(\theta, \varphi_\epsilon), \Gamma(\theta, \varphi)) \\
&= \overline{\lim}_{k \rightarrow \infty} k^{-n} \left( H^0(X, L^k \otimes I(k\varphi)) - H^0(X, L^k \otimes I(k\varphi_\epsilon)) \right) \\
&= \lim_{k \rightarrow \infty} k^{-n} H^0(X, L^k \otimes I(k\varphi)) - \lim_{k \rightarrow \infty} k^{-n} H^0(X, L^k \otimes I(k\varphi_\epsilon)) \\
&= \frac{1}{n!} \text{vol } \theta_\varphi - \frac{1}{n!} \text{vol } \theta_{\varphi_\epsilon} \\
&\rightarrow 0
\end{aligned}$$

by [Theorem 6.2.5](#), as  $\epsilon \rightarrow 0+$ .  $\square$

rmk:DeltaanaW0

*Remark 10.2.1* It follows from the proof that if  $\varphi$  has analytic singularities and  $\theta_\varphi$  is a Kähler current, then (10.12) holds.

If we take a modification  $\pi: Y \rightarrow X$  such that  $\pi^*\varphi$  has log singularities along an effective  $\mathbb{Q}$ -divisor  $D$  on  $Y$ , then

$$\Delta_\nu(\theta, \varphi) = \Delta_\nu(\pi^*L - D) + \nu(D).$$

#### 10.2.4 Basic properties of partial Okounkov bodies

cor:Okocurrent

**Proposition 10.2.7** *The partial Okounkov body  $\Delta_\nu(L, h)$  depends only on  $\text{dd}^c h$ , not on the explicit choices of  $L, h_0, h$ .*

Thanks to this result, given a closed positive  $(1, 1)$ -current  $T \in c_1(L)$  on  $X$  with  $\int_X T^n > 0$ , we can write

$$\Delta_\nu(T) := \Delta_\nu(\theta, \varphi)$$

if  $T = \theta + \text{dd}^c \varphi$  for some  $\varphi \in \text{PSH}(X, \theta)$ .

**Proof** There are two different claims to prove, as detailed in the two steps below.

**Step 1.** Let  $h'_0$  be another Hermitian metric on  $L$ . Set  $\theta' = c_1(L, h'_0)$ . Write  $\text{dd}^c f = \theta - \theta'$ . Let  $\varphi' = \varphi + f \in \text{PSH}(X, \theta')$ . Then

$$\Delta_\nu(\theta, \varphi) = \Delta_\nu(\theta', \varphi'). \quad (10.14)$$

{eq:DeltaDelta1}

This is obvious since  $\Gamma(\theta, \varphi) = \Gamma(\theta', \varphi')$ .

**Step 2.** Let  $L'$  be another big line bundle on  $X$ . By Step 1, we may assume that the reference Hermitian metric  $h'_0$  on  $L'$  is such that  $c_1(L', h'_0) = \theta$ .

Let  $h'$  be a plurisubharmonic metric on  $L'$  with  $c_1(L, h) = c_1(L', h')$ . Then

$$\Delta_\nu(L, h) = \Delta_\nu(L', h').$$

From our construction, we may assume that  $c_1(L, h)$  has analytic singularities. After taking a birational resolution, it suffices to deal with the case where  $c_1(L, h)$  has analytic singularities along an effective  $\mathbb{Q}$ -divisors  $D$ . By rescaling, we may also

assume that  $D$  is a divisor. By [Remark 10.2.1](#), we further reduce to the case where  $c_1(L, h)$  is not singular.

In this case, the assertion is proved in [Proposition 10.2.2](#).  $\square$

prop:IcompimplyDeltacomp

**Proposition 10.2.8** *Let  $\varphi, \psi \in \text{PSH}(X, \theta)_{>0}$ . Assume that  $\varphi \leq_I \psi$ , then*

$$\Delta_\nu(\theta, \varphi) \subseteq \Delta_\nu(\theta, \psi). \quad (10.15)$$

{eq:Deltacomp}

**Proof** This follows from [Corollary C.2.2](#).  $\square$

thm:Okoucont

**Theorem 10.2.2** *The Okounkov body map*

$$\Delta_\nu(\theta, \bullet) : (\text{PSH}(X, \theta)_{>0}, d_S) \rightarrow (\mathcal{K}_n, d_{\text{Haus}})$$

*is continuous.*

**Proof** Let  $\varphi_j \rightarrow \varphi$  be a  $d_S$ -convergent sequence in  $\text{PSH}(X, \theta)_{>0}$ . We want to show that

$$\Delta_\nu(\theta, \varphi_j) \xrightarrow{d_{\text{Haus}}} \Delta_\nu(\theta, \varphi). \quad (10.16)$$

{eq:Deltavjv}

By [Proposition 10.2.8](#), we may assume that all  $\varphi_j$ 's and  $\varphi$  are model potentials.

By [Theorem C.1.1](#) and [Proposition 6.2.3](#), we may assume that  $(\varphi_j)_j$  is either decreasing or increasing. By [Theorem 6.2.3](#), we may further assume that the  $\varphi_j$ 's are  $I$ -model. In both cases, we claim that

$$\Gamma(\theta, \varphi_j) \xrightarrow{d_{\text{sg}}} \Gamma(\theta, \varphi)$$

as  $j \rightarrow \infty$ . In fact, using [Theorem 7.3.1](#), we can compute

$$\begin{aligned} d_{\text{sg}}(\Gamma(\theta, \varphi_j), \Gamma(\theta, \varphi)) &= \overline{\lim}_{k \rightarrow \infty} k^{-n} |\mathbf{H}^0(X, L^k \otimes I(k\varphi_j)) - \mathbf{H}^0(X, L^k \otimes I(k\varphi))| \\ &= \frac{1}{n!} |\text{vol } \theta_{\varphi_j} - \text{vol } \theta_\varphi|, \end{aligned}$$

which converges to 0 by [Theorem 6.2.5](#).  $\square$

prop:birinv0

**Proposition 10.2.9** *Let  $\pi: Y \rightarrow X$  be a modification. Then*

$$\Delta_\nu(\pi^*L, \pi^*h) = \Delta_\nu(L, h).$$

**Proof** Thanks to [Proposition 3.2.5](#), we may assume that  $\varphi$  is  $I$ -model. By [Theorem 7.1.1](#), we can find a sequence  $(\varphi_j)_j$  with analytic singularities in  $\text{PSH}(X, \theta)$  such that  $\varphi_j \xrightarrow{d_S} \varphi$ . It is clear that  $\pi^*\varphi_j \xrightarrow{d_S} \pi^*\varphi$ . By [Theorem 10.2.2](#), we may then reduce to the case where  $\varphi$  has analytic singularities. In this case, it suffices to apply [Remark 10.2.1](#).  $\square$

prop:suba

**Proposition 10.2.10** *Let  $(L', h')$  be another Hermitian big line bundle on  $X$ . Then*

$$\Delta_\nu(L, h) + \Delta_\nu(L', h') \subseteq \Delta_\nu(L \otimes L', h \otimes h').$$

**Proof** Take a smooth metric  $h'_0$  on  $L'$  and let  $\theta' = c_1(L', h'_0)$ . We identify  $h'$  with  $\varphi' \in \text{PSH}(X, \theta')$ . Then we need to show

$$\Delta_\nu(\theta, \varphi) + \Delta_\nu(\theta', \varphi') \subseteq \Delta_\nu(\theta + \theta', \varphi + \varphi'). \quad (10.17)$$

{eq:suba}

By **Theorem 7.1.1**, we can find sequences  $(\varphi_j)_j$  and  $(\varphi'_j)_j$  in  $\text{PSH}(X, \theta)_{>0}$  and  $\text{PSH}(X, \theta')_{>0}$  respectively such that

- (1)  $\varphi_j$  and  $\varphi'_j$  both have analytic singularities for all  $j \geq 1$ , and
- (2)  $\varphi_j \xrightarrow{ds} \varphi, \varphi'_j \xrightarrow{ds} \varphi'$ .

Then  $\varphi_j + \varphi'_j \in \text{PSH}(X, \theta + \theta')_{>0}$  and  $\varphi_j + \varphi'_j \xrightarrow{ds} \varphi + \varphi'$  by **Theorem 6.2.2**. Thus, by **Theorem 10.2.2**, we may assume that  $\varphi$  and  $\psi$  both have analytic singularities. Taking a birational resolution, we may further assume that they have log singularities. By **Remark 10.2.1**, we reduce to the case without singularities, in which case the result is just **Proposition 10.2.5**.  $\square$

thm:concOkO

**Theorem 10.2.3** Let  $\varphi, \psi \in \text{PSH}(X, \theta)_{>0}$ . Then for any  $t \in (0, 1)$ ,

$$\Delta_\nu(\theta, t\varphi + (1-t)\psi) \supseteq t\Delta_\nu(\theta, \varphi) + (1-t)\Delta_\nu(\theta, \psi). \quad (10.18)$$

{eq:Deltaconcave}

**Proof** We may assume that  $t$  is rational as a consequence of **Theorem 10.2.2**. Similarly, as in the proof of **Proposition 10.2.10**, we could reduce to the case where both  $\varphi$  and  $\psi$  have analytic singularities. In this case, let  $N > 0$  be an integer such that  $Nt$  is an integer. Then for any  $s \in H^0(X, L^k \otimes \mathcal{I}_\infty(k\varphi))$  and  $r \in H^0(X, L^k \otimes \mathcal{I}_\infty(k\psi))$ , we have

$$s^{tN} \otimes r^{N-tN} \in H^0(X, L^{kN} \otimes \mathcal{I}_\infty(Nt\varphi + (N-Nt)\psi)).$$

By **Theorem C.2.1** and **Remark 10.2.1**, (10.18) follows.  $\square$

prop:res

**Proposition 10.2.11** For any  $a \in \mathbb{Z}_{>0}$ ,

$$\Delta_\nu(a\theta, a\varphi) = a\Delta_\nu(\theta, \varphi).$$

**Proof** As in the proof of **Proposition 10.2.10**, we may assume that  $\varphi$  has log singularities. Using **Remark 10.2.1**, we reduce to the case without the singularity  $\varphi$ , which is proved in **Proposition 10.2.6**.  $\square$

In particular, if  $T$  is a closed positive  $(1, 1)$ -current on  $X$  with  $\int_X T^n > 0$  and such that

$$[T] \in \text{NS}^1(X)_{\mathbb{Q}},$$

we can define

$$\Delta_\nu(T) := a^{-1}\Delta_\nu(aT) \quad (10.19)$$

{eq:DeltaTanuTalgebraic1}

for a sufficiently divisible positive integer  $a$ .

We also need the following perturbation. Let  $A$  be an ample line bundle on  $X$ . Fix a Hermitian metric  $h_A$  on  $A$  such that  $\omega := c_1(A, h_A)$  is a Kähler form on  $X$ .

prop:Deltapert

**Proposition 10.2.12** *As  $\delta \searrow 0$ , the convex bodies  $\Delta_\nu(\theta + \delta\omega + \text{dd}^c\varphi)$  are decreasing and*

$$\Delta_\nu(\theta + \delta\omega + \text{dd}^c\varphi) \xrightarrow{d_{\text{Haus}}} \Delta_\nu(\theta_\varphi).$$

**Proof** Let  $0 \leq \delta < \delta'$  be two rational numbers. Take  $C \in \mathbb{N}_{>0}$  divisible enough, so that  $C\delta$  and  $C\delta'$  are both integers. Then by [Proposition 10.2.10](#),

$$\Delta_\nu(C\theta + C\delta\omega + C\text{dd}^c\varphi) \subseteq \Delta_\nu(C\theta + C\delta'\omega + C\text{dd}^c\varphi).$$

It follows that

$$\Delta_\nu(\theta + \delta\omega + \text{dd}^c\varphi) \subseteq \Delta_\nu(\theta + \delta'\omega + \text{dd}^c\varphi).$$

On the other hand,

$$\text{vol } \Delta_\nu(\theta + \delta\omega + \text{dd}^c\varphi) = \frac{1}{n!} \text{vol}(\theta + \delta\omega)_\varphi = \frac{1}{n!} \int_X (\theta + \delta\omega)_{P_\theta[\varphi]_I}^n,$$

where we applied [Example 7.1.2](#). As  $\delta \rightarrow 0+$ , the right-hand side converges to

$$\text{vol } \Delta_\nu(\theta, \varphi) = \frac{1}{n!} \text{vol } \theta_\varphi.$$

Our assertion therefore follows.  $\square$

### 10.2.5 The Hausdorff convergence property of partial Okounkov bodies

Let  $T$  be a holomorphic line bundle on  $X$ .

thm:HCP

**Theorem 10.2.4** *As  $k \rightarrow \infty$ , we have  $\Delta_{k,T}(\theta, \varphi) \xrightarrow{d_{\text{Haus}}} \Delta_\nu(\theta, \varphi)$ .*

Although we are only interested in the untwisted case, the proof given below requires twisted case.

lma:twistedHcp

**Lemma 10.2.1** *Assume that  $\varphi$  has analytic singularities and  $\theta_\varphi$  is a Kähler current, then as  $k \rightarrow \infty$ ,*

$$\Delta_{k,T}(\theta, \varphi) \xrightarrow{d_{\text{Haus}}} \Delta_\nu(\theta, \varphi).$$

**Proof** Up to replacing  $X$  by a birational model and twisting  $T$  accordingly, we may assume that  $\varphi$  has log singularities along an effective  $\mathbb{Q}$ -divisor  $D$ , see [Proposition 10.2.9](#) and [Theorem 1.6.1](#).

Take a small enough  $\epsilon \in \mathbb{Q}_{>0}$ . In this case, for large enough  $k \in \mathbb{Z}_{>0}$  we have

$$H^0(X, T \otimes L^k \otimes I_\infty(k\varphi)) \subseteq H^0(X, T \otimes L^k \otimes I(k\varphi)) \subseteq H^0(X, T \otimes L^k \otimes I_\infty(k(1-\epsilon)\varphi)).$$

Take an integer  $N \in \mathbb{Z}_{>0}$  so that  $ND$  is a divisor and  $N\epsilon$  is an integer.

Let  $\Delta'$  be the limit of a subsequence of  $(\Delta_{k,T}(\theta, \varphi))_k$ , say the sequence defined by the indices  $k_1, k_2, \dots$ . We want to show that  $\Delta' = \Delta(\theta, \varphi)$ .

There exists  $t \in \{0, 1, \dots, N-1\}$  such that  $k_i \equiv t$  modulo  $N$  for infinitely many  $i$ , up to replacing  $k_i$  by a subsequence, we may assume that  $k_i \equiv t$  modulo  $N$  for all  $i$ . Write  $k_i = Ng_i + t$ . Then for large enough  $i$ , we have

$$\begin{aligned} H^0(X, T \otimes L^{-N+t} \otimes L^{N(g_i+1)} \otimes \mathcal{I}_\infty(N(g_i+1)\varphi)) &\subseteq H^0(X, T \otimes L^{k_i} \otimes \mathcal{I}(k_i\varphi)) \\ &\subseteq H^0(X, T \otimes L^t \otimes L^{Ng_i} \otimes \mathcal{I}_\infty(g_iN(1-\epsilon)\varphi)). \end{aligned}$$

So

$$\begin{aligned} (g_i+1)\Delta_{g_i+1, T \otimes L^{-N+t}}(NL - ND) + N(g_i+1)v(D) &\subseteq (Ng_i+t)\Delta_{k, T}(\theta, \varphi) \\ &\subseteq g_i\Delta_{g_i, T \otimes L^t}(NL - N(1-\epsilon)D) + Ng_i(1-\epsilon)v(D). \end{aligned}$$

Letting  $i \rightarrow \infty$ , by [Proposition 10.2.4](#),

$$\Delta_v(L - D) + v(D) \subseteq \Delta' \subseteq \Delta_v(L - (1-\epsilon)D) + (1-\epsilon)v(D).$$

Letting  $\epsilon \rightarrow 0+$ , we find that

$$\Delta_v(L - D) + v(D) = \Delta'.$$

It follows from [Theorem C.1.1](#) that

$$\Delta_{k, T}(\theta, \varphi) \xrightarrow{d_{\text{Haus}}} \Delta_v(L - D) + v(D) = \Delta_v(\theta, \varphi)$$

as  $k \rightarrow \infty$ . □

lma-Hausconvbetato0

**Lemma 10.2.2** Assume that  $\theta_\varphi$  is a Kähler current, then as  $\mathbb{Q} \ni \beta \rightarrow 0+$ , we have

$$\Delta_v((1-\beta)\theta, \varphi) \xrightarrow{d_{\text{Haus}}} \Delta_v(\theta, \varphi).$$

Here and in the sequel,  $\Delta_v((1-\beta)\theta, \varphi) = \Delta_v((1-\beta)\theta + dd^c\varphi)$ .

**Proof** By [Proposition 10.2.10](#), we have

$$\Delta_v((1-\beta)\theta, \varphi) + \beta\Delta_v(L) \subseteq \Delta_v(\theta, \varphi).$$

In particular, if  $\Delta'$  is the Hausdorff limit of a subsequence of  $(\Delta_v((1-\beta)\theta, \varphi))_\beta$ , then  $\Delta' \subseteq \Delta_v(\theta, \varphi)$ . But

$$\begin{aligned} \text{vol } \Delta' &= \lim_{\beta \rightarrow 0+} \Delta_v((1-\beta)\theta, \varphi) = \lim_{\beta \rightarrow 0+} \int_X ((1-\beta)\theta + dd^c P_{(1-\beta)\theta}[\varphi]_I)^n \\ &= \int_X (\theta + dd^c P_\theta[\varphi]_I)^n, \end{aligned}$$

where the last step follows easily from [Theorem 11.2.1](#). It follows that  $\Delta' = \Delta_v(\theta, \varphi)$ . We conclude by [Theorem C.1.1](#). □

**Proof (Proof of Theorem 10.2.4)** Fix a Kähler form  $\omega \geq \theta$  on  $X$ .

**Step 1.** We first handle the case where  $\theta_\varphi$  is a Kähler current, say  $\theta_\varphi \geq 2\delta\omega$  for some  $\delta \in (0, 1)$ . Take a quasi-equisingular approximation  $(\varphi_j)_j$  of  $\varphi$  in  $\text{PSH}(X, \theta)$ . We may assume that  $\theta_{\varphi_j} \geq \delta\omega$  for all  $j \geq 1$ .

Let  $\Delta'$  be a limit of a subsequence of  $(\Delta_{k,T}(\theta, \varphi))_k$ . Let us say the indices of the subsequence are  $k_1 < k_2 < \dots$ . By [Theorem C.1.1](#), it suffices to show that  $\Delta' = \Delta_\nu(\theta, \varphi)$ .

Observe that for each  $j \geq 1$ , we have  $\Delta' \subseteq \Delta_\nu(\theta, \varphi_j)$  by [Lemma 10.2.1](#). Letting  $j \rightarrow \infty$ , we find  $\Delta' \subseteq \Delta_\nu(\theta, \varphi)$ . Therefore, it suffices to prove that

$$\text{vol } \Delta' \geq \text{vol } \Delta_\nu(\theta, \varphi). \quad (10.20)$$

Fix an integer  $N > \delta^{-1}$ . Observe that for any  $j \geq 1$ , we have  $\varphi_j \in \text{PSH}(X, (1-N^{-1})\theta)$ . Similarly,  $\varphi \in \text{PSH}(X, (1-N^{-1})\theta)$ . By [Lemma 10.2.2](#), it suffices to argue that

$$\text{vol } \Delta' \geq \text{vol } \Delta_\nu((1-N^{-1})\theta, \varphi). \quad (10.21)$$

{eq:volDeltatoprove}

For this purpose, we are free to replace  $k_i$ 's by a subsequence, so we may assume that  $k_i \equiv a$  modulo  $q$  for all  $i \geq 1$ , where  $a \in \{0, 1, \dots, q-1\}$ . We write  $k_i = g_i q + a$ . Observe that for each  $i \geq 1$ ,

$$H^0(X, T \otimes L^{k_i} \otimes I(k_i \varphi)) \supseteq H^0(X, T \otimes L^{-q+a} \otimes L^{g_i q + a} \otimes I((g_i q + a)\varphi)).$$

Up to replacing  $T$  by  $T \otimes L^{-q+a}$ , we may therefore assume that  $a = 0$ .

By [Lemma 2.3.1](#), we can find  $k' \in \mathbb{Z}_{>0}$  such that for all  $k \geq k'$ , there is  $\psi \in \text{PSH}(X, \theta)_{>0}$  satisfying

$$P_\theta[\varphi]_I \geq (1-N^{-1})\varphi_k + N^{-1}\psi_k.$$

Fix  $k \geq k'$ . It suffices to show that

$$\Delta_\nu((1-N^{-1})\theta, \varphi_k) + \nu' \subseteq \Delta' \quad (10.22)$$

{eq:DeltatransinDeltaprime}

for some  $\nu' \in \mathbb{R}^n$ . In fact, if this is true, we have

$$\text{vol } \Delta' \geq \text{vol } \Delta((1-N^{-1})\theta, \varphi_k).$$

Letting  $k \rightarrow \infty$  and applying [Theorem 10.2.2](#), we conclude (10.21).

It remains to prove (10.22). By the proof of [Theorem 7.3.1](#), there is  $j_0 > 0$  such that for any  $j \geq j_0$ , we can find a non-zero section  $s_j \in H^0(X, L^j \otimes I(j\psi_k))$  such that we get an injective linear map

$$H^0(X, T \otimes L^{(N-1)j} \otimes I(jN\varphi_k)) \xrightarrow{\times s_j} H^0(X, T \otimes L^{jN} \otimes I(jN\varphi)).$$

In particular, when  $j = k_i$  for some  $i$  large enough, we then find

$$\Delta_{k_i, T}((N-1)\theta, N\varphi_k) + (k_i)^{-1}\nu(s_{k_i}) \subseteq N\Delta_{k_i, T}(\theta, \varphi).$$



We observe that  $(k_i)^{-1}v(s_{k_i})$  is bounded as both convex bodies appearing in this equation are bounded when  $i$  varies. Then by [Lemma 10.2.1](#), there is a vector  $v' \in \mathbb{R}^n$  such that [\(10.22\)](#) holds.

**Step 2.** Next we handle the general case.

Let  $\Delta'$  be the Hausdorff limit of a subsequence of  $(\Delta_{k,T}(\theta, \varphi))_k$ , say the subsequence with indices  $k_1 < k_2 < \dots$ . By [Theorem C.1.1](#), it suffices to prove that  $\Delta' = \Delta_v(\theta, \varphi)$ .

Take  $\psi \in \text{PSH}(X, \theta)$  such that  $\theta_\psi$  is a Kähler current and  $\psi \leq \varphi$ . The existence of  $\psi$  follows from [Lemma 2.3.2](#).

Then for any  $\epsilon \in \mathbb{Q} \cap (0, 1)$ ,

$$\Delta_{k,T}(\theta, \varphi) \supseteq \Delta_{k,T}(\theta, (1 - \epsilon)\varphi + \epsilon\psi)$$

for all  $k \geq 1$ . It follows from Step 1 that

$$\Delta' \supseteq \Delta_v(\theta, (1 - \epsilon)\varphi + \epsilon\psi).$$

Letting  $\epsilon \rightarrow 0$  and applying [Theorem 10.2.2](#), we have  $\Delta' \supseteq \Delta_v(\theta, \varphi)$ . It remains to establish that

$$\text{vol } \Delta' \leq \text{vol } \Delta_v(\theta, \varphi). \quad (10.23)$$

{eq:Deltapvolumeupp}

For this purpose, we are free to replace  $k_1 < k_2 < \dots$  by a subsequence. Fix  $q > 0$ , we may then assume that  $k_i \equiv a$  modulo  $q$  for all  $i \geq 1$  for some  $a \in \{0, 1, \dots, q-1\}$ . We write  $k_i = g_i q + a$ . Observe that

$$H^0(X, T \otimes L^{k_i} \otimes I(k_i \varphi)) \subseteq H^0(X, T \otimes L^a \otimes L^{g_i q} \otimes I(g_i q \varphi)).$$

Up to replacing  $T$  by  $T \otimes L^a$ , we may assume that  $a = 0$ .

Take a very ample line bundle  $H$  on  $X$  and fix a Kähler form  $\omega \in c_1(H)$ , take a non-zero section  $s \in H^0(X, H)$ .

We have an injective linear map

$$H^0(X, T \otimes L^{jq} \otimes I(jq\varphi)) \xrightarrow{\times s^j} H^0(X, T \otimes H^j \otimes L^{jq} \otimes I(jq\varphi))$$

for each  $j \geq 1$ . In particular, for each  $i \geq 1$ ,

$$k_i \Delta_{k_i, T}(q\theta, q\varphi) + k_i v(s) \subseteq k_i \Delta_{k_i, T}(\omega + q\theta, q\varphi).$$

Letting  $i \rightarrow \infty$ , by Step 1, we have

$$q\Delta' + v(s) \subseteq \Delta_v(\omega + q\theta, q\varphi).$$

So

$$\text{vol } \Delta' \leq \text{vol } \Delta_v(q^{-1}\omega + \theta, \varphi) = \int_X (q^{-1}\omega + \theta + \text{dd}^c P_{q^{-1}\omega + \theta}[\varphi])^n.$$

By [Example 7.1.2](#),

$$\text{vol } \Delta' \leq \int_X (q^{-1}\omega + \theta + \text{dd}^c P_\theta[\varphi]_I)^n.$$

Letting  $q \rightarrow \infty$ , we conclude (10.23).  $\square$

## 10.2.6 Recover Lelong numbers from partial Okounkov bodies

thm:nuOk

**Theorem 10.2.5** *Let  $E$  be a prime divisor on  $X$ . Let  $Y_\bullet$  be an admissible flag with  $E = Y_1$ . Then*

$$v(\varphi, E) = \min_{x \in \Delta_{Y_\bullet}(\theta, \varphi)} x_1. \quad (10.24)$$

{eq:numinOk}

Here  $x_1$  denotes the first component of  $x$ .

**Proof** Replacing  $\varphi$  by  $P_\theta[\varphi]_I$ , we may assume that  $\varphi$  is  $I$ -good.

**Step 1.** We first reduce to the case where  $\varphi$  has analytic singularities.

By [Theorem 7.1.1](#), we can find a sequence  $(\varphi_j)_j$  in  $\text{PSH}(X, \theta)_{>0}$  with analytic singularities such that  $\varphi_j \xrightarrow{d_S} \varphi$ . It follows from [Theorem 10.2.2](#) that

$$\Delta_{Y_\bullet}(\theta, \varphi_j) \xrightarrow{d_{\text{Haus}}} \Delta_{Y_\bullet}(\theta, \varphi).$$

Therefore,

$$\lim_{j \rightarrow \infty} \min_{x \in \Delta_{Y_\bullet}(\theta, \varphi_j)} x_1 = \min_{x \in \Delta_{Y_\bullet}(\theta, \varphi)} x_1.$$

In view of [Theorem 6.2.4](#), it suffices to prove (10.24) with  $\varphi_j$  in place of  $\varphi$ .

**Step 2.** Assume that  $\varphi$  has analytic singularities. In view of [Proposition 10.2.9](#) and [Theorem 1.6.1](#), after replacing  $X$  by a birational model, we may assume that  $\varphi$  has log singularities along an effective  $\mathbb{Q}$ -divisor  $F$ .

Perturbing  $L$  by an ample  $\mathbb{Q}$ -line bundle by [Proposition 10.2.12](#), we may assume that  $\theta_\varphi$  is a Kähler current. Therefore,  $L - F$  is ample by [Lemma 1.6.1](#). Finally, by rescaling, we may assume that  $F$  is a divisor and  $L$  is a line bundle.

By [Theorem 10.2.4](#), we know that

$$\min_{x \in \Delta_{Y_\bullet}(\theta, \varphi)} x_1 = \lim_{k \rightarrow \infty} \min_{x \in \Delta_k(\theta, \varphi)} x_1.$$

By definition,

$$\min_{x \in \Delta_k(\theta, \varphi)} x_1 = k^{-1} \text{ord}_E H^0(X, L^k \otimes I(k\varphi)).$$

It remains to show that

$$\lim_{k \rightarrow \infty} k^{-1} \text{ord}_E H^0(X, L^k \otimes I(k\varphi)) = \lim_{k \rightarrow \infty} k^{-1} \text{ord}_E I(k\varphi). \quad (10.25)$$

{eq:temp1}

The  $\geq$  direction is trivial, we prove the converse. Observe that

$$H^0(X, L^k \otimes I(k\varphi)) = H^0(X, L^k \otimes \mathcal{O}_X(-kF)), \quad I(k\varphi) = \mathcal{O}_X(-kF).$$

As  $L - F$  is ample, for large enough  $k$ , we have

$$\mathrm{ord}_E H^0(X, L^k \otimes \mathcal{O}_X(-kF)) = \mathrm{ord}_E(kF).$$

Thus, (10.25) is clear.  $\square$

cor:Deltacontimplyvarphi

**Corollary 10.2.3** *Let  $\varphi, \psi \in \mathrm{PSH}(X, \theta)_{>0}$ . If*

$$\Delta_{W_\bullet}(\pi^*\theta, \pi^*\varphi) \subseteq \Delta_{W_\bullet}(\pi^*\theta, \pi^*\psi)$$

*for all birational models  $\pi : Y \rightarrow X$  and all admissible flags  $W_\bullet$  on  $Y$ , then  $\varphi \leq_I \psi$ .*

**Proof** This follows immediately from Theorem 10.2.5.  $\square$

cor:numin

**Corollary 10.2.4** *Let  $E$  be a prime divisor over  $X$ . Then*

$$v(V_\theta, E) = \lim_{k \rightarrow \infty} \frac{1}{k} \mathrm{ord}_E H^0(X, L^k). \quad (10.26)$$

**Proof** This follows from Theorem 10.2.5 and the fact that  $\Delta_{Y_\bullet}(\theta, V_\theta) = \Delta_{Y_\bullet}(L)$  for any admissible flag  $Y_\bullet$  on  $X$ .  $\square$

### 10.3 Transcendental partial Okounkov bodies

sec:tpob

Let  $X$  be a connected compact Kähler manifold of dimension  $n$ . Fix a smooth flag  $Y_\bullet$  on  $X$ .

#### 10.3.1 The traditional approach to the Okounkov body problem

**Definition 10.3.1** Let  $\alpha$  be a big cohomology class on  $X$ . We define the *Okounkov body* of  $\alpha$  as

$$\Delta_{Y_\bullet}(\alpha) := \overline{\{v_{Y_\bullet}(S) : S \in \mathcal{Z}_+(X, \alpha), S \text{ has gentle analytic singularities}\}}. \quad (10.27)$$

{eq:twodefspob}

See Definition 1.6.4 for the definition of gentle analytic singularities.

The results of [DRWN<sup>+</sup>23] can be summarized as follows:

thm:Okounkovtranmain

**Theorem 10.3.1** *For any big cohomology class  $\alpha$  on  $X$ , the set  $\Delta_{Y_\bullet}(\alpha) \subseteq \mathbb{R}^n$  is a convex body satisfying the following properties:*

(1) *we have*

$$\mathrm{vol} \Delta_{Y_\bullet}(\alpha) = \frac{1}{n!} \mathrm{vol} \alpha;$$

(2) Given another big cohomology class  $\alpha'$  on  $X$ , we have

$$\Delta_{Y_\bullet}(\alpha) + \Delta_{Y_\bullet}(\alpha') \subseteq \Delta_{Y_\bullet}(\alpha + \alpha');$$

(3) Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism with  $Y$  being a Kähler manifold. Assume that  $(W_\bullet, g)$  is the lifting of  $Y_\bullet$  to  $Y$ , then

$$\Delta_{W_\bullet}(\pi^* \alpha) = \Delta_{Y_\bullet}(\alpha)g;$$

(4) The map  $\alpha \mapsto \Delta_{Y_\bullet}(\alpha)$  is continuous in the big cone with respect to the Hausdorff metric;

(5) For any small enough  $t > 0$ , we have

$$\{y \in \mathbb{R}^{n-1} : (t, y) \in \Delta_{Y_\bullet}(\beta)\} = \Delta_{Y_1 \geq \dots \geq Y_n}((\beta - t[Y_1])|_{Y_1}).$$

### 10.3.2 Definitions of partial Okounkov bodies

Let  $\theta$  be a closed real smooth  $(1, 1)$ -form on  $X$  representing a big cohomology class  $\alpha$ .

Let  $T = \theta_\varphi \in \mathcal{Z}_+(X, \alpha)$ . We shall define a convex body  $\Delta_{Y_\bullet}(T) \subseteq \mathbb{R}^n$ , which is also written as  $\Delta_{Y_\bullet}(\theta, \varphi)$ . This convex body is called the *partial Okounkov body* of  $T$  with respect to the flag  $Y_\bullet$ .

#### 10.3.2.1 The case of analytic singularities

def:POBanalsing

**Definition 10.3.2** When  $T$  is a Kähler current with analytic singularities, we take a modification  $\pi: Y \rightarrow X$  so that

(1)

$$\pi^*T = [D] + R, \tag{10.28}$$

{eq:resolveanalytic}

where  $D$  is an effective  $\mathbb{Q}$ -divisor on  $Y$  and  $R$  is a closed positive  $(1, 1)$ -current with bounded potential, and

(2) the lifting  $(Z_\bullet, g)$  of  $Y_\bullet$  to  $Y$  exists.

Define

$$\Delta_{Y_\bullet}(T) := \Delta_{Z_\bullet}([R])g^{-1} + \nu_{Z_\bullet}([D])g^{-1}.$$

The existence of  $\pi$  is guaranteed by [Theorem 1.6.1](#) and [Theorem 10.1.1](#).

**Lemma 10.3.1** The convex body  $\Delta_{Y_\bullet}(T)$  defined in [Definition 10.3.2](#) is independent of the choice of  $\pi$ .

**Proof** Take another map  $\pi': Y' \rightarrow X$  with the same properties. We want to show that  $\pi$  and  $\pi'$  defines the same  $\Delta_{Y_\bullet}(T)$ . We may assume that  $\pi'$  dominates  $\pi$  through  $p: Y' \rightarrow Y$ , so that we have a commutative diagram

$$\begin{array}{ccc}
 Y' & \xrightarrow{p} & Y \\
 \searrow \pi' & & \swarrow \pi \\
 & X &
 \end{array}$$

We take  $D$  and  $R$  as in (10.28). Then

$$\pi'^*T = [p^*D] + p^*R.$$

Write  $(Z_\bullet, g)$  and  $(Z'_\bullet, g')$  for the liftings of  $Y_\bullet$  to  $Y$  and  $Y'$  respectively. We need to prove that

$$\Delta_{Z_\bullet}([R])g^{-1} + \nu_{Z_\bullet}([D])g^{-1} = \Delta_{Z'_\bullet}([p^*R])g'^{-1} + \nu_{Z'_\bullet}([p^*D])g'^{-1}.$$

This follows [Theorem 10.3.1](#), [Proposition 10.1.4](#) and [Proposition 10.1.3](#).  $\square$

Note that from the above proof, we could describe the bimeromorphic behaviour of  $\Delta_{Y_\bullet}(T)$  as follows:

lma:liftOkounana

**Lemma 10.3.2** *Let  $T \in \mathcal{Z}_+(X, \alpha)$  be a Kähler current with analytic singularities. Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism and  $(W_\bullet, g)$  be the lifting of  $Y_\bullet$  to  $Y$ . Then*

$$\Delta_{W_\bullet}(\pi^*T) = \Delta_{Y_\bullet}(T)g.$$

lma:Okounkovanalycomp

**Lemma 10.3.3** *Assume that  $T, S \in \mathcal{Z}_+(X, \alpha)$  are two Kähler currents with analytic singularities and  $T \leq S$ , then*

$$\Delta_{Y_\bullet}(T) \subseteq \Delta_{Y_\bullet}(S) \subseteq \Delta_{Y_\bullet}(\alpha).$$

Moreover,

$$\text{vol } \Delta_{Y_\bullet}(T) = \frac{1}{n!} \int_X T^n. \quad (10.29)$$

{eq:volpobanaly}

**Proof** We first show that

$$\Delta_{Y_\bullet}(T) \subseteq \Delta_{Y_\bullet}(S).$$

Using [Lemma 10.3.2](#), we may assume that  $T$  and  $S$  have log singularities along effective  $\mathbb{Q}$ -divisors  $E$  and  $F$  respectively. By assumption,  $E \geq F$ . Replacing  $T$  and  $S$  by  $T - [F]$  and  $S - [F]$  respectively, we may assume that  $F = 0$ .

In this case, we need to show that

$$\Delta_{Y_\bullet}(\alpha) \supseteq \Delta_{Y_\bullet}(\alpha - [E]) + \nu_{Y_\bullet}([E]),$$

which is obvious.

Next we prove that

$$\Delta_{Y_\bullet}(T) \subseteq \Delta_{Y_\bullet}(\alpha).$$

By [Lemma 10.3.2](#) and [Theorem 10.3.1](#) again, we may assume that  $T$  has log singularities. We take  $D$  and  $\beta$  as in (10.28). We need to show that

$$\Delta_{Y_\bullet}(\alpha - [D]) + \nu_{Y_\bullet}([D]) \subseteq \Delta_{Y_\bullet}(\alpha),$$

which is again obvious.

Finally, (10.29) follows immediately from [Theorem 10.3.1](#).  $\square$

### 10.3.2.2 The case of Kähler currents

def:POBKahcurr

**Definition 10.3.3** Let  $T \in \mathcal{Z}_+(X, \alpha)$  be a Kähler current. Take a quasi-equisingular approximation  $(T_j)_j$  of  $T$  in  $\mathcal{Z}_+(X, \alpha)$ . Then we define

$$\Delta_{Y_\bullet}(T) := \bigcap_{j=1}^{\infty} \Delta_{Y_\bullet}(T_j).$$

**Lemma 10.3.4** *The convex body  $\Delta_{Y_\bullet}(T)$  in [Definition 10.3.3](#) is independent of the choices of the  $T_j$ 's.*

In particular, if  $T$  also has analytic singularities, then the  $\Delta_{Y_\bullet}(T)$ 's defined in [Definition 10.3.3](#) and in [Definition 10.3.2](#) coincide.

**Proof** Let  $(S_j)_j$  be another quasi-equisingular approximation of  $T$  in  $\mathcal{Z}_+(X, \alpha)$ . By [Proposition 1.6.3](#), for any small rational  $\epsilon > 0$ ,  $j > 0$ , we can find  $k > 0$  so that

$$S_k \leq (1 - \epsilon)T_j.$$

It is more convenient to use the language of  $\theta$ -psh functions at this point. Let  $\psi_k$  (resp.  $\varphi_k$ ) denote the potentials in  $\text{PSH}(X, \theta)$  corresponding to  $S_k$  (resp.  $T_k$ ) for each  $k \geq 1$ . Note that  $\psi_k$  and  $\varphi_k$  are unique up to additive constants.

By [Lemma 10.3.3](#),

$$\bigcap_{k=1}^{\infty} \Delta_{Y_\bullet}(\theta, \psi_k) \subseteq \Delta_{Y_\bullet}(\theta, (1 - \epsilon)\varphi_j).$$

On the other hand, observe that

$$\bigcap_{\epsilon \in \mathbb{Q}_{>0} \text{ small enough}} \Delta_{Y_\bullet}(\theta, (1 - \epsilon)\varphi_j) = \Delta_{Y_\bullet}(\theta, \varphi_j).$$

In fact, the  $\supseteq$  direction follows from [Lemma 10.3.3](#), so it suffices to show that the two sides have the same volume, which follows from (10.29).

It follows that

$$\bigcap_{k=1}^{\infty} \Delta_{Y_\bullet}(\theta, \psi_k) \subseteq \bigcap_{j=1}^{\infty} \Delta_{Y_\bullet}(\theta, \varphi_j).$$

The other inclusion follows by symmetry.  $\square$

The same argument shows that

cor:Kahlercurrentcase

**Corollary 10.3.1** Suppose that  $T, S \in \mathcal{Z}_+(X, \alpha)$  are two Kähler currents satisfying  $T \preceq_I S$ . Then

$$\Delta_{Y_\bullet}(T) \subseteq \Delta_{Y_\bullet}(S) \subseteq \Delta_{Y_\bullet}(\alpha).$$

**Proposition 10.3.1** Let  $T \in \mathcal{Z}_+(X, \alpha)$  be a Kähler current. Then

$$\text{vol } \Delta_{Y_\bullet}(T) = \frac{1}{n!} \text{vol } T. \quad (10.30)$$

{eq:vol0kocur}

*Proof* Take a quasi-equisingular approximation  $(T_j)_j$  of  $T$  in  $\mathcal{Z}_+(X, \alpha)$ . Note that  $\Delta_{Y_\bullet}(T_j)$  is decreasing in  $j$ , as follows from [Lemma 10.3.3](#). Our assertion follows from [\(10.29\)](#) and [Theorem 6.2.5](#).  $\square$

lma:Okomonotone

**Lemma 10.3.5** Let  $T \in \mathcal{Z}_+(X, \alpha)$  be a Kähler current and  $\omega$  be a Kähler form on  $X$ . Then

$$\Delta_{Y_\bullet}(T) \subseteq \Delta_{Y_\bullet}(T + \omega). \quad (10.31)$$

{eq:DeltaTincreaseomegatemp1}

Moreover,

$$\Delta_{Y_\bullet}(T) = \bigcap_{\epsilon > 0} \Delta_{Y_\bullet}(T + \epsilon\omega). \quad (10.32)$$

{eq:DeltaTincreaseomegatemp2}

*Proof* We first prove [\(10.31\)](#). Taking quasi-equisingular approximations, we reduce immediately to the case where  $T$  has analytic singularities. By [Lemma 10.3.2](#), we may assume that  $T$  has log singularities. Take  $D$  and  $R$  as in [\(10.28\)](#). By definition again, it suffices to show that

$$\Delta_{Y_\bullet}([\beta]) \subseteq \Delta_{Y_\bullet}([\beta + \omega]),$$

which is clear by definition.

Next we prove [\(10.32\)](#). Thanks to [\(10.31\)](#), it remains to prove that both sides have the same volume:

$$\lim_{\epsilon \rightarrow 0+} \text{vol}(T + \epsilon\omega) = \text{vol } T.$$

This is proved in [Proposition 7.2.3](#).  $\square$

### 10.3.2.3 The general case

def:generalPOB

**Definition 10.3.4** Let  $T \in \mathcal{Z}_+(X, \alpha)$ . Take a Kähler form  $\omega$  on  $X$ , we define

$$\Delta_{Y_\bullet}(T) = \bigcap_{j=1}^{\infty} \Delta_{Y_\bullet}(T + j^{-1}\omega). \quad (10.33)$$

{eq:DeltaTgeneral}

The same definition makes sense when  $\alpha$  is only pseudo-effective.

This definition is clearly independent of the choice of  $\omega$  by [Lemma 10.3.5](#). Moreover, it extends [Definition 10.3.3](#) and [Definition 10.3.2](#) as a result of [Lemma 10.3.5](#).

*Remark 10.3.1* When  $\alpha$  is pseudoeffective but not big and  $T$  has minimal singularities, **Definition 10.3.4** differs from all known definitions of  $\Delta_{Y_\bullet}(\alpha)$  in the literature. But in view of **Lemma 10.3.7**, our definition seems to be the most natural one.

The main properties of  $\Delta_{Y_\bullet}(T)$  are summarized as follows:

thm:pobmain

**Theorem 10.3.2** *The convex bodies  $\Delta_{Y_\bullet}(T)$ 's satisfies the following properties:*

(1) Suppose that  $T \in \mathcal{Z}_+(X, \alpha)_{>0}$ . We have

$$\text{vol } \Delta_{Y_\bullet}(T) = \frac{1}{n!} \text{vol } T. \quad (10.34)$$

{eq:volpobgeneral}

(2) For  $T, S \in \mathcal{Z}_+(X, \alpha)$  satisfying  $T \leq_I S$ , we have

$$\Delta_{Y_\bullet}(T) \subseteq \Delta_{Y_\bullet}(S) \subseteq \Delta_{Y_\bullet}(\alpha).$$

(3) For any current  $T \in \mathcal{Z}_+(X, \alpha)$  with minimal singularities, we have

$$\Delta_{Y_\bullet}(T) = \Delta_{Y_\bullet}(\alpha).$$

(4) The map  $\mathcal{Z}_+(X, \alpha)_{>0} \rightarrow \mathcal{K}_n$  given by  $T \mapsto \Delta_{Y_\bullet}(T)$  is continuous, where we endow the  $d_S$ -pseudometric on  $\mathcal{Z}_+(X, \alpha)_{>0}$  and the Hausdorff topology on  $\mathcal{K}_n$ .

(5) Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism with  $Y$  being a Kähler manifold. Assume that the lifting  $(W_\bullet, g)$  of  $Y_\bullet$  to  $Y$  exists, then for any  $T \in \mathcal{Z}_+(X, \alpha)_{>0}$ , we have

$$\Delta_{W_\bullet}(\pi^*T) = \Delta_{Y_\bullet}(T)g.$$

(6) For  $T, S \in \mathcal{Z}_+(X, \alpha)$ , we have

$$\Delta_{Y_\bullet}(T) + \Delta_{Y_\bullet}(S) \subseteq \Delta_{Y_\bullet}(T + S). \quad (10.35)$$

{eq:pobadditiv}

**Proof** (1) By (10.33) and (10.30), for any Kähler form  $\omega$  on  $X$ ,

$$\text{vol } \Delta_{Y_\bullet}(T) = \lim_{j \rightarrow \infty} \Delta_{Y_\bullet}(T + j^{-1}\omega) = \frac{1}{n!} \lim_{j \rightarrow \infty} \text{vol}(T + j^{-1}\omega).$$

The right-hand side is computed in **Proposition 7.2.3**. Hence, (10.34) follows.

(2) Fix a Kähler form  $\omega$  on  $X$ . By **Corollary 10.3.1**, for each  $j \geq 1$ ,

$$\Delta_{Y_\bullet}(T + j^{-1}\omega) \subseteq \Delta_{Y_\bullet}(S + j^{-1}\omega) \subseteq \Delta_{Y_\bullet}(\alpha + j^{-1}[\omega]).$$

It remains to show that

$$\Delta_{Y_\bullet}(\alpha) = \bigcap_{j=1}^{\infty} \Delta_{Y_\bullet}(\alpha + j^{-1}[\omega]).$$

The  $\subseteq$  direction is clear. Comparing the volumes using **Theorem 10.3.1**, we conclude that equality holds.

(3) This follows from (1) and (2).



(4) Let  $(T_j)_j$  be a sequence in  $\mathcal{Z}_+(X, \alpha)_{>0}$  converging to  $T \in \mathcal{Z}_+(X, \alpha)_{>0}$  with respect to  $d_S$ . We want to show that  $\Delta_{Y_\bullet}(T_j) \xrightarrow{d_{\text{Haus}}} \Delta_{Y_\bullet}(T)$ . By [Proposition 6.2.3](#) and (2), we may assume that the singularity type of  $T_j$  is either increasing or decreasing. In both cases, the continuity follows from (1).

(5) We may assume that  $T$  is  $\mathcal{I}$ -good. It follows from (4) and [Theorem 7.1.1](#) that we could reduce to the case where  $T$  has analytic singularities. Our assertion follows from [Lemma 10.3.2](#).

(6) By [\(10.33\)](#), in order to prove [\(10.35\)](#), we may assume that  $T$  and  $S$  are both Kähler currents. Take quasi-equisingular approximations  $(T_j)_j$  and  $(S_j)_j$  of  $T$  and  $S$  respectively. By [Theorem 6.2.2](#),  $T_j + S_j \xrightarrow{d_S} T + S$ . By (4), we may therefore assume that  $T$  and  $S$  have analytic singularities. Replacing  $X$  by a suitable modification, we may assume that  $T$  and  $S$  both have log singularities, say

$$T = [D] + R, \quad S = [D'] + R',$$

where  $D$  and  $D'$  are  $\mathbb{Q}$ -divisors on  $X$  and  $\beta$  and  $\beta'$  are closed positive  $(1, 1)$ -currents with bounded potentials. We need to show that

$$\Delta_{Y_\bullet}([R]) + \Delta_{Y_\bullet}([R']) + \nu_{Y_\bullet}([D]) + \nu_{Y_\bullet}([D']) \subseteq \Delta_{Y_\bullet}([R + R']) + \nu_{Y_\bullet}([D + D']).$$

By [Proposition 10.1.2](#), this is equivalent to

$$\Delta_{Y_\bullet}([R]) + \Delta_{Y_\bullet}([R']) \subseteq \Delta_{Y_\bullet}([R + R']),$$

which is already proved in [Theorem 10.3.1](#).  $\square$

**Corollary 10.3.2** *Assume that  $L$  is a big line bundle on  $X$  and  $h$  is a plurisubharmonic metric on  $L$  with positive volume. Then*

$$\Delta_{Y_\bullet}(\text{dd}^c h) = \Delta_{Y_\bullet}(L, h). \quad (10.36)$$

{eq: tranOkounandalgOkoun}

Similarly, the definition [\(10.19\)](#) is compatible with the definition in [Definition 10.3.4](#).

**Proof** We may assume that  $\text{dd}^c h$  has positive mass and is  $\mathcal{I}$ -good. By the  $d_S$ -continuity of both sides of [\(10.36\)](#) as proved in [Theorem 10.3.2](#) and [Theorem 10.2.2](#), together with [Theorem 7.1.1](#), we may assume that  $\text{dd}^c h$  has analytic singularities.

In this case, using the birational invariance of both sides of [\(10.36\)](#) as proved in [Proposition 10.2.9](#) and [Theorem 10.3.2](#), we may assume that  $\text{dd}^c h$  has log singularities. Finally, after all these reductions, the equality [\(10.36\)](#) holds by construction.  $\square$

### 10.3.3 The valuative characterization

In this section, we will characterize the partial Okounkov bodies using valuations of currents.

lma:Kahlerclassokounrest

**Lemma 10.3.6** *Let  $\beta$  be a nef class on  $X$ . Then*

$$\{y \in \mathbb{R}^{n-1} : (0, y) \in \Delta_{Y_\bullet}(\beta)\} = \Delta_{Y_1 \supseteq \dots \supseteq Y_n}(\beta|_{Y_1}). \quad (10.37)$$

{eq:Deltaresttox10}

**Proof Step 1.** We first reduce to the case where  $\beta$  is a Kähler class.

Take a Kähler class  $\alpha$  on  $X$ . It follows from the volume formula in [Theorem 10.3.1](#) that

$$\Delta_{Y_\bullet}(\beta) = \bigcap_{\epsilon > 0} \Delta_{Y_\bullet}(\beta + \epsilon\alpha), \quad \Delta_{Y_1 \supseteq \dots \supseteq Y_n}(\beta|_{Y_1}) = \bigcap_{\epsilon > 0} \Delta_{Y_1 \supseteq \dots \supseteq Y_n}(\beta|_{Y_1} + \epsilon\alpha|_{Y_1}).$$

So it suffices to prove (10.37) with  $\beta + \epsilon\alpha$  in place of  $\beta$ .

**Step 2.** Assume that  $\alpha$  is a Kähler class. The  $\supseteq$  direction in (10.37) follows from the extension theorem [Theorem 1.6.3](#). To prove the other direction, recall that by [Theorem 10.3.1](#), for  $t > 0$  small enough, we have

$$\{y \in \mathbb{R}^{n-1} : (t, y) \in \Delta_{Y_\bullet}(\beta)\} = \Delta_{Y_1 \supseteq \dots \supseteq Y_n}((\beta - t[Y_1])|_{Y_1}).$$

As  $t \rightarrow 0+$ , the right-hand side converges to  $\Delta_{Y_1 \supseteq \dots \supseteq Y_n}(\beta|_{Y_1})$  with respect to the Hausdorff metric as a consequence of [Theorem 10.3.1](#), while the left-hand side converges to

$$\{y \in \mathbb{R}^{n-1} : (0, y) \in \Delta_{Y_\bullet}(\beta)\}$$

by [Lemma C.1.2](#). We conclude our assertion.  $\square$

lma:slicebob

**Lemma 10.3.7** *Let  $T \in \mathcal{Z}_+(X, \alpha)$  be a Kähler current. Assume that  $\nu(T, Y_1) = 0$ , then*

$$\{y \in \mathbb{R}^{n-1} : (0, y) \in \Delta_{Y_\bullet}(T)\} = \Delta_{Y_1 \supseteq \dots \supseteq Y_n}(\text{Tr}_{Y_1}^{\alpha|_{Y_1}}(T)). \quad (10.38)$$

{eq:Deltaslice}

More generally, if  $T \in \mathcal{Z}_+(X, \alpha)$  and  $\nu(T, Y_1) = 0$ , suppose in addition that  $\text{Tr}_{Y_1}^{\alpha|_{Y_1}}(T)$  is defined, then (10.38) still holds.

See [Remark 8.1.1](#) for the definition of  $\text{Tr}_{Y_1}^{\alpha|_{Y_1}}(T)$ . Note that  $\Delta_{Y_1 \supseteq \dots \supseteq Y_n}(\text{Tr}_{Y_1}^{\alpha|_{Y_1}}(T))$  is independent of the choice of the representative  $\text{Tr}_{Y_1}^{\alpha|_{Y_1}}(T)$ .

**Remark 10.3.2** More generally, the same argument shows the following result: Let  $k = 0, \dots, n$  and  $T \in \mathcal{Z}_+(X, \alpha)$  such that  $\nu(T, Y_k) = 0$ . Assume that  $\text{Tr}_{Y_k}^{\alpha|_{Y_k}}(T)$  is defined, then

$$\{y \in \mathbb{R}^{n-k} : (0, \dots, 0, y) \in \Delta_{Y_\bullet}(T)\} = \Delta_{Y_k \supseteq \dots \supseteq Y_n}(\text{Tr}_{Y_k}^{\alpha|_{Y_k}}(T)). \quad (10.39)$$

This contradicts [\[CPW17, Theorem 1.1\]](#).<sup>2</sup> Also note that this result extends [\[Jow10, Theorem 3.4\]](#) and hence gives simpler proofs of [\[Jow10, Theorem A, Theorem B\]](#).

<sup>2</sup> The mistake with this paper happens in the proof of Theorem 3.6, the second paragraph, where the authors asserted that it is easy to check that  $\Delta_{Y_{n-k} \bullet}(D) \subseteq \Delta_{Y_\bullet}(D)$ .

**Proof** Let  $\omega$  be a Kähler form on  $X$ . The last assertion follows from the first by perturbing  $\theta$  to  $\theta + \epsilon\omega$ .

**Step 1.** We first handle the case where  $T$  has analytic singularities. Let  $\pi: Z \rightarrow X$  be a modification such that

- (1)  $Y_\bullet$  admits a lifting  $(W_\bullet, g)$ , and
- (2)  $\pi^*T = [D] + R$ , where  $D$  is an effective  $\mathbb{Q}$ -divisor on  $Z$  and  $R$  is closed positive  $(1, 1)$ -current with bounded potential.

This is possible by [Theorem 1.6.1](#) and [Theorem 10.1.1](#).

By [Lemma 8.2.1](#),

$$\Pi^* \text{Tr}_{Y_1}(T) \sim_P \text{Tr}_{W_1}(\pi^*T),$$

where  $\Pi: W_1 \rightarrow Y_1$  is the restriction of  $\pi$ . It follows from [Theorem 10.3.2](#) that

$$\begin{aligned} \Delta_{W_1 \supseteq \dots \supseteq W_n}(\text{Tr}_{W_1}(\pi^*T)) &= \Delta_{Y_1 \supseteq \dots \supseteq Y_n}(\text{Tr}_{Y_1}(T)) \text{cor}(Y_1 \supseteq \dots \supseteq Y_n, \Pi), \\ \Delta_{W_\bullet}(\pi^*T) &= \Delta_{Y_\bullet}(T)g. \end{aligned}$$

Taking (10.3) into account, we find that it suffices to show that

$$\{y \in \mathbb{R}^{n-1} : (0, y) \in \Delta_{W_\bullet}(\pi^*T)\} = \Delta_{W_1 \supseteq \dots \supseteq W_n}(\text{Tr}_{W_1}(\pi^*T)).$$

We may assume that  $\pi$  is the identity map. Then we have

$$T = [D] + R, \quad T|_{Y_1} = [D]|_{Y_1} + R|_{Y_1}.$$

Note that  $[D]|_{Y_1}$  is the current of integration along an effective  $\mathbb{Q}$ -divisor on  $Y_1$ .

In particular,

$$\begin{aligned} \Delta_{Y_\bullet}(T) &= \Delta_{Y_\bullet}([R]) + \nu_{Y_\bullet}([D]), \\ \Delta_{Y_1 \supseteq \dots \supseteq Y_n}(T|_{Y_1}) &= \Delta_{Y_1 \supseteq \dots \supseteq Y_n}([R]|_{Y_1}) + \nu_{Y_1 \supseteq \dots \supseteq Y_n}([D]|_{Y_1}). \end{aligned}$$

So it suffices to show that

$$\{y \in \mathbb{R}^{n-1} : (0, y) \in \Delta_{Y_\bullet}([R])\} = \Delta_{Y_1 \supseteq \dots \supseteq Y_n}([R]|_{Y_1}),$$

which is exactly [Lemma 10.3.6](#).

**Step 2.** Next we consider the case where  $T$  is a Kähler current. Take a quasi-equisingular approximation  $(T_j)_j$  of  $T$  in  $\mathcal{Z}_+(X, \alpha)$ . From Step 1, we know that for large  $j \geq 1$ ,

$$\{y \in \mathbb{R}^{n-1} : (0, y) \in \Delta_{Y_\bullet}(T_j)\} = \Delta_{Y_1 \supseteq \dots \supseteq Y_n}(\text{Tr}_{Y_1}(T_j)).$$

Letting  $j \rightarrow \infty$  and applying [Theorem 10.3.2](#) and [Proposition 8.2.2](#), we conclude (10.38).  $\square$

thm:KahcurrminOkoun

**Theorem 10.3.3** Assume that  $T \in \mathcal{Z}_+(X, \alpha)_{>0}$  is a Kähler current. We have

$$\min_{\text{lex}} \Delta_{Y_\bullet}(T) = \nu_{Y_\bullet}(T). \quad (10.40)$$

{eq:minOkounkov}

Here the minimum is with respect to the lexicographic order.

**Proof** We make induction on  $n \geq 0$ . The case  $n = 0$  is of course trivial. Let us assume that  $n > 0$  and the case  $n - 1$  has been proved.

We first observe that by [Theorem 10.3.2](#),

$$\Delta_{Y_\bullet}(T - \nu(T, Y_1)[Y_1]) + (\nu(T, Y_1), 0, \dots, 0) \subseteq \Delta_{Y_\bullet}(T).$$

Comparing the volumes of both sides using [Theorem 10.3.2](#) and [Proposition 7.2.3](#), we find that equality holds:

$$\Delta_{Y_\bullet}(T - \nu(T, Y_1)[Y_1]) + (\nu(T, Y_1), 0, \dots, 0) = \Delta_{Y_\bullet}(T).$$

Replacing  $T$  by  $T - \nu(T, Y_1)[Y_1]$ , we may therefore assume that  $\nu(T, Y_1) = 0$ . It suffices to apply [Lemma 10.3.7](#) and the inductive hypothesis.  $\square$

cor:valuationcurrentinPOB

**Corollary 10.3.3** For any  $T \in \mathcal{Z}_+(X, \alpha)$ ,

$$\nu_{Y_\bullet}(T) \in \Delta_{Y_\bullet}(T) \subseteq \Delta_{Y_\bullet}(\alpha).$$

**Proof** When  $T$  is a Kähler current, this follows from [Theorem 10.3.3](#).

In general, by definition,  $\nu_{Y_\bullet}(T) = \nu_{Y_\bullet}(T + \omega)$  for any Kähler form  $\omega$  on  $X$ . It follows that

$$\nu_{Y_\bullet}(T) \in \Delta_{Y_\bullet}(T + \omega)$$

for any Kähler form  $\omega$ . It follows that  $\nu_{Y_\bullet}(T) \in \Delta_{Y_\bullet}(T)$ .  $\square$

thm:Deltapartialint

**Theorem 10.3.4** For any  $T \in \mathcal{Z}_+(X, \alpha)_{>0}$ ,

$$\Delta_{Y_\bullet}(T) = \overline{\{\nu_{Y_\bullet}(S) : S \in \mathcal{Z}_+(X, \alpha), S \leq_I T\}}. \quad (10.41)$$

{eq:DeltaTequalallval}

In particular,

$$\Delta_{Y_\bullet}(\alpha) = \overline{\{\nu_{Y_\bullet}(T) : T \in \mathcal{Z}_+(X, \alpha)\}}.$$

*Remark 10.3.3* We expect that the closure operation in (10.41) is not necessary. This problem is closely related to the Dirichlet problem of the trace operator, see Page 239 for more details.

**Proof** The  $\supseteq$  direction in (10.41) follows from [Corollary 10.3.3](#) and [Theorem 10.3.2\(2\)](#).

Let us write

$$D_{Y_\bullet}(T) = \{\nu_{Y_\bullet}(S) : S \in \mathcal{Z}_+(X, \alpha), S \leq_I T\}$$

for the time being.

**Step 1.** Assume that  $T$  has analytic singularities. We have

$$\begin{aligned} \Delta_{Y_\bullet}(T) &\supseteq \overline{D_{Y_\bullet}(T)} \\ &\supseteq \overline{\{\nu_{Y_\bullet}(S) : \mathcal{Z}_+(X, \alpha) \ni S \text{ has gentle analytic singularities, } S \leq T\}}. \end{aligned}$$

It follows easily from [Theorem 10.3.1](#) that the volume of the right-hand side is equal to the volume of  $\Delta_{Y_\bullet}(T)$ , so (10.41) holds.

**Step 2.** Assume that  $T$  is a Kähler current. Take a quasi-equisingular approximation  $T_j \in \mathcal{Z}_+(X, \alpha)$  of  $T$ . Next we use the language of psh functions. Let  $\varphi_j, \varphi \in \text{PSH}(X, \theta)$  be the potentials corresponding to  $T_j, T$  for each  $j \geq 1$ .

Fix an integer  $N > 0$ . For large enough  $j \geq 1$ , we can find  $\psi \in \text{PSH}(X, \theta)_{>0}$  such that

$$P_\theta[\varphi]_T \geq (1 - N^{-1})\varphi_j + N^{-1}\psi_j.$$

The existence of  $\psi_j$  follows from [Lemma 2.3.1](#). It follows that

$$\begin{aligned} D_{Y_\bullet}(T) &\supseteq D_{Y_\bullet}\left(\theta + \text{dd}^c\left((1 - N^{-1})\varphi_j + N^{-1}\psi_j\right)\right) \\ &\supseteq (1 - N^{-1})D_{Y_\bullet}(T_j) + N^{-1}D_{Y_\bullet}(\theta + \text{dd}^c\psi_j). \end{aligned}$$

By [Theorem C.1.1](#), up to replacing  $T_j$  by a subsequence, we may guarantee that  $\overline{D_{Y_\bullet}(\theta + \text{dd}^c\psi_j)}$  admits a Hausdorff limit contained in  $\Delta_{Y_\bullet}(\alpha)$  as  $j \rightarrow \infty$ . Let  $j \rightarrow \infty$  and  $N \rightarrow \infty$  then it follows that

$$\overline{D_{Y_\bullet}(T)} \supseteq \bigcap_{j=1}^{\infty} D_{Y_\bullet}(T_j).$$

By [Lemma C.1.3](#),

$$\overline{D_{Y_\bullet}(T)} \supseteq \overline{\bigcap_{j=1}^{\infty} D_{Y_\bullet}(T_j)} = \bigcap_{j=1}^{\infty} \overline{D_{Y_\bullet}(T_j)}.$$

Therefore, by Step 1, we conclude that

$$\Delta_{Y_\bullet}(T) = \bigcap_{j=1}^{\infty} \overline{\Delta_{Y_\bullet}(T_j)} = \bigcap_{j=1}^{\infty} \overline{D_{Y_\bullet}(T_j)} \subseteq \overline{D_{Y_\bullet}(T)}.$$

The reverse direction is already known.

**Step 3.** Finally, consider the general case. Take a Kähler current  $T' \in \mathcal{Z}_+(X, \alpha)$  more singular than  $T$ . For each  $\epsilon \in (0, 1)$ . The existence of  $T'$  is proved in [Lemma 2.3.2](#). We know that

$$\Delta_{Y_\bullet}((1 - \epsilon)T + \epsilon T') = \overline{D_{Y_\bullet}((1 - \epsilon)T + \epsilon T')} \subseteq \overline{D_{Y_\bullet}(T)}.$$

Letting  $\epsilon \rightarrow 0+$  and using [Proposition 7.2.3](#), we find that

$$\Delta_{Y_\bullet}(T) \subseteq \overline{D_{Y_\bullet}(T)}.$$

As the other inclusion is already known, we conclude. □

cor:KahcurrminOkoun

**Corollary 10.3.4** Assume that  $T \in \mathcal{Z}_+(X, \alpha)_{>0}$ . We have

$$\min_{\text{lex}} \Delta_{Y_\bullet}(T) = \nu_{Y_\bullet}(T). \quad (10.42)$$

{eq:minOkounkov3}

**Proof** By [Theorem 10.3.4](#), it is clear that

$$\min_{\text{lex}} \Delta_{Y_\bullet}(T) \leq_{\text{lex}} \nu_{Y_\bullet}(T).$$

On the other hand, we clearly have

$$\Delta_{Y_\bullet}(T) \subseteq \Delta_{Y_\bullet}(T + \omega)$$

for any Kähler form  $\omega$  on  $X$ . It follows that

$$\min_{\text{lex}} \Delta_{Y_\bullet}(T) \geq_{\text{lex}} \min_{\text{lex}} \Delta_{Y_\bullet}(T + \omega).$$

By [Theorem 10.3.3](#), the right-hand side is just  $\nu_{Y_\bullet}(T + \omega) = \nu_{Y_\bullet}(T)$ . We conclude the proof.  $\square$

## 10.4 Okounkov test curves

Fix  $n \in \mathbb{N}$ . Let  $\Delta, \Delta' \subseteq \mathbb{R}^n$  be convex bodies with positive volume. The standard Lebesgue measure on  $\mathbb{R}^n$  is denoted by  $\text{vol}$ .

We refer to [Appendix C](#) for the notations  $\mathcal{K}_n$  and  $d_{\text{Haus}}$ .

def:Otc

**Definition 10.4.1** An *Okounkov test curve* relative to  $\Delta$  consists of

- (1) a number  $\Delta_{\max} \in \mathbb{R}$  and
- (2) an assignment  $(-\infty, \Delta_{\max}) \ni \tau \mapsto \Delta_\tau \in \mathcal{K}_n$  satisfying
  - a. the assignment  $\tau \mapsto \Delta_\tau$  is a decreasing and concave;
  - b. we have  $\Delta_\tau \xrightarrow{d_{\text{Haus}}} \Delta$  as  $\tau \rightarrow -\infty$ .

The set of Okounkov test curves relative to  $\Delta$  is denoted by  $\text{TC}(\Delta)$ .

An Okounkov test curve  $\Delta_\bullet$  is *bounded* if  $\Delta_\tau = \Delta$  when  $\tau$  is small enough. The subset of bounded Okounkov test curves is denoted by  $\text{TC}^\infty(\Delta)$ .

An Okounkov test curve  $\Delta_\bullet$  is said to have *finite energy* if

$$\mathbf{E}(\Delta_\bullet) := n! \Delta_{\max} \text{vol } \Delta + n! \int_{-\infty}^{\Delta_{\max}} (\text{vol } \Delta_\tau - \text{vol } \Delta) \, d\tau > -\infty. \quad (10.43)$$

{eq:0testcurveenergy}

The subset of Okounkov test curves with finite energy is denoted by  $\text{TC}^1(\Delta)$ .

Given  $\Delta_\bullet \in \text{TC}(\Delta)$  and  $\Delta'_\bullet \in \text{TC}(\Delta')$ , we say  $\Delta_\bullet \leq \Delta'_\bullet$  if  $\Delta_{\max} \leq \Delta'_{\max}$  and for any  $\tau < \Delta_{\max}$ , we have  $\Delta_\tau \subseteq \Delta'_\tau$ .

Here concavity in (2)b refers to the concavity with respect to the Minkowski sum. Sometimes it is convenient to introduce

$$\Delta_{\Delta_{\max}} = \bigcap_{\tau < \Delta_{\max}} \Delta_{\tau} \in \mathcal{K}_n. \quad (10.44)$$

{eq:DeltaDeltamax}

We shall always make this extension in the sequel when we talk about  $\Delta_{\Delta_{\max}}$ . Observe that  $(-\infty, \Delta_{\max}] \ni \tau \mapsto \Delta_{\tau}$  is still concave.

prop:Otccont

**Proposition 10.4.1** Any Okounkov test curve  $(\Delta_{\tau})_{\tau < \Delta_{\max}}$  relative to  $\Delta$  is continuous in  $\tau$ . Moreover,  $\text{vol } \Delta_{\tau} > 0$  for all  $\tau < \Delta_{\max}$ .

**Proof** We first claim that  $\text{vol } \Delta_{\tau'} > 0$  for all  $\tau' < \Delta_{\max}$ . By Condition (2)b in Definition 10.4.1 and Theorem C.1.2, we know that  $\text{vol } \Delta_{\tau''} > 0$  when  $\tau''$  is small enough. Fix one such  $\tau''$ . We may assume that  $\tau'' \leq \tau'$  since otherwise there is nothing to prove. Next take  $\tau''' \in (\tau', \Delta_{\max})$ . Take  $t \in (0, 1)$  such that  $\tau' = t\tau''' + (1-t)\tau''$ . It follows that

$$\text{vol } \Delta_{\tau'} \geq \text{vol } (t\Delta_{\tau'''} + (1-t)\Delta_{\tau''}) \geq (1-t)^n \text{vol } \Delta_{\tau''} > 0.$$

Next we claim that  $\text{vol } \Delta_{\tau}$  is continuous for  $\tau < \Delta_{\max}$ . In fact, it follows from Theorem C.1.4 that  $(-\infty, \Delta_{\max}) \ni \tau \mapsto \log \text{vol } \Delta_{\tau}$  is concave, the continuity follows.

Next we show that

$$\Delta_{\tau} = \bigcap_{\tau' < \tau} \Delta_{\tau'}.$$

The  $\supseteq$  direction is obvious. By the continuity of the volume, both sides have the same volume and the volume is positive, we therefore obtain the equality.

Similarly, we have

$$\Delta_{\tau} = \overline{\bigcup_{\tau' > \tau} \Delta_{\tau'}}.$$

The continuity of  $\Delta_{\tau}$  at  $\tau < \Delta_{\max}$  is proved.  $\square$

def:tf

**Definition 10.4.2** A test function on  $\Delta$  is a function  $F: \Delta \rightarrow [-\infty, \infty)$  such that

- (1)  $F$  is concave,
- (2)  $F$  is finite on  $\text{Int } \Delta$ , and
- (3)  $F$  is upper semicontinuous.

A test function  $F$  is *bounded* if  $F$  is bounded from below.

A test function  $F$  has *finite energy* if

$$\mathbf{E}(F) := n! \int_{\Delta} F \, d\lambda > -\infty. \quad (10.45)$$

{eq:EF}

def:LegOkoun

**Definition 10.4.3** Let  $\Delta_{\bullet} \in \text{TC}(\Delta)$ . We define its *Legendre transform* as

$$G[\Delta_{\bullet}]: \Delta \rightarrow [-\infty, \infty), \quad a \mapsto \sup \{ \tau < \Delta_{\max} : a \in \Delta_{\tau} \}.$$

Given a test function  $F: \Delta \rightarrow [-\infty, \infty)$ , we define its *inverse Legendre transform*  $\Delta[F]_{\bullet}$  as the Okounkov test curve relative to  $\Delta$  defined as follows:

- (1)  $\Delta[F]_{\max} = \sup_{\Delta} F$ , and

(2) for each  $\tau < \sup_{\Delta} F$ , we set

$$\Delta[F]_{\tau} = \{x \in \Delta : F \geq \tau\}.$$

We observe that

$$G[\Delta_{\bullet}](a) = \max \{\tau \leq \Delta_{\max} : a \in \Delta_{\tau}\}, \text{ if } G[\Delta_{\bullet}](a) > -\infty. \quad (10.46)$$

{eq:GDeltamax}

lma:convbodyLegendre

**Lemma 10.4.1** *Let  $\Delta_{\bullet} \in \text{TC}(\Delta)$ . Then  $G[\Delta_{\bullet}]$  defined in [Definition 10.4.3](#) is a test function.*

*Similar, if  $F: \Delta \rightarrow [-\infty, \infty)$  is a test function, then  $\Delta[F]_{\bullet}$  is an Okounkov test curve.*

**Proof** First suppose that  $\Delta_{\bullet} \in \text{TC}(\Delta)$ . We want to verify that  $G[\Delta_{\bullet}]$  satisfies the conditions in [Definition 10.4.2](#).

We first verify the concavity. Take  $a, b \in \Delta$ . We want to prove that for any  $t \in (0, 1)$ ,

$$G[\Delta_{\bullet}](ta + (1-t)b) \geq tG[\Delta_{\bullet}](a) + (1-t)G[\Delta_{\bullet}](b). \quad (10.47)$$

{eq:GDeltaconc}

There is nothing to prove if  $G[\Delta_{\bullet}](a)$  or  $G[\Delta_{\bullet}](b)$  is  $-\infty$ . So we assume that both are finite. In this case, by (10.46),

$$a \in \Delta_{G[\Delta_{\bullet}](a)}, \quad b \in \Delta_{G[\Delta_{\bullet}](b)}.$$

Thus,

$$ta + (1-t)b \in t\Delta_{G[\Delta_{\bullet}](a)} + (1-t)\Delta_{G[\Delta_{\bullet}](b)} \subseteq \Delta_{tG[\Delta_{\bullet}](a) + (1-t)G[\Delta_{\bullet}](b)}.$$

We deduce that

$$G[\Delta_{\bullet}](ta + (1-t)b) \geq tG[\Delta_{\bullet}](a) + (1-t)G[\Delta_{\bullet}](b).$$

Therefore, (10.47) follows.

It is clear that  $F$  is finite on the interior of  $\Delta$ . It remains to argue that  $F$  is upper semicontinuous.

Let  $(a_i)_{i \geq 1}$  be a sequence in  $\Delta$  with limit  $a \in \Delta$ . Define  $\tau_i = G[\Delta_{\bullet}](a_i)$ . Let  $\tau = \lim_i \tau_i$ . We need to show that

$$G[\Delta_{\bullet}](a) \geq \tau. \quad (10.48)$$

{eq:ainDelta1}

There is nothing to prove if  $\tau = -\infty$ . We assume that it is not this case. Up to subtracting a subsequence we may assume that  $\tau_i \rightarrow \tau$ . In particular, we can assume that  $\tau_i \neq -\infty$  for all  $i \geq 1$ . It follows from (10.46) that  $a_i \in \Delta_{\tau_i}$  for all  $i \geq 1$ . Since  $\Delta_{\tau_i} \xrightarrow{d_{\text{Haus}}} \Delta_{\tau}$ . By [Theorem C.1.3](#) it follows that  $a \in \Delta_{\tau}$ . Thus, (10.48) follows.

Conversely, suppose that  $F: \Delta \rightarrow [-\infty, \infty)$  is a test function. We argue that  $\Delta[F]_{\bullet}$  is an Okounkov test curve. We verify the conditions in [Definition 10.4.1](#).



Firstly, for each  $\tau < \sup_{\Delta} F$ , the set  $\Delta[F](\tau)$  is a convex body as  $F$  is concave and usc. Moreover,  $\Delta[F]_{\tau}$  is clearly decreasing in  $\tau$ .

Secondly, for each  $a \in \Delta$ , we can write  $a = \lim_i a_i$  with  $a_i \in \text{Int } \Delta$ . By assumption,  $F$  is finite at  $a_i$ . Thus,

$$a \in \overline{\{F > -\infty\}} = \overline{\bigcup_{\tau < \sup_{\Delta} F} \Delta[F]_{\tau}}.$$

By [Theorem C.1.3](#),  $\Delta[F]_{\tau} \xrightarrow{d_{\text{Haus}}} \Delta$  as  $\tau \rightarrow -\infty$ .

Thirdly,  $\Delta[F]$  is concave. To see, take  $\tau, \tau' < \Delta_{\max}$ , we need to prove that for any  $t \in (0, 1)$ ,

$$\Delta[F]_{t\tau + (1-t)\tau'} \supseteq t\Delta[F]_{\tau} + (1-t)\Delta[F]_{\tau'}. \quad (10.49)$$

{eq:Deconc}

Let  $a \in \Delta[F]_{\tau}$  and  $b \in \Delta[F]_{\tau'}$ . We have  $F(a) \geq \tau$  and  $F(b) \geq \tau'$ . As  $F$  is concave, we have  $F(ta + (1-t)b) \geq t\tau + (1-t)\tau'$ . Thus,

$$ta + (1-t)b \in \Delta[F]_{t\tau + (1-t)\tau'}$$

and (10.49) follows.  $\square$

thm:Okotestcurve

**Theorem 10.4.1** *The Legendre transform and inverse Legendre transform are inverse to each other, defining a bijection between  $\text{TC}(\Delta)$  and the set of test functions on  $\Delta$ .*

*Under this bijection,  $\text{TC}^1(\Delta)$  corresponds to test functions on  $\Delta$  with finite energy and  $\text{TC}^{\infty}(\Delta)$  corresponds to bounded test functions on  $\Delta$ .*

**Proof** Thanks to [Lemma 10.4.1](#), in order to prove the first assertion, it only remains to see that the Legendre transform and the inverse Legendre transform are inverse to each other, which is immediate by definition.

It is obvious that  $\text{TC}^{\infty}(\Delta)$  corresponds to bounded test curves. Moreover, a direct computation shows that if  $\Delta_{\bullet} \in \text{TC}(\Delta)$ , then

$$\mathbf{E}(\Delta_{\bullet}) = \mathbf{E}(G[\Delta_{\bullet}]),$$

concluding the  $\text{TC}^1(\Delta)$  case.  $\square$

prop:decnetLegend

**Proposition 10.4.2** *Let  $(\Delta^i)_{i \in I}$  be a decreasing net in  $\mathcal{K}_n$ . Consider a decreasing net  $(\Delta_{\bullet}^i)_{i \in I}$  with  $\Delta_{\bullet}^i \in \text{TC}(\Delta^i)$  for all  $i \in I$  such that there is  $\Delta_{\bullet} \in \text{TC}(\Delta)$  satisfying the following properties:*

- (1)  $\Delta_{\max} = \lim_{i \in I} \Delta_{\max}^i$ ;
- (2) for any  $\tau < \Delta_{\max}$ , we have  $\Delta_{\tau}^i \xrightarrow{d_{\text{Haus}}} \Delta_{\tau}$ .

*Then for any  $a \in \Delta$ , we have*

$$\lim_{i \in I} G[\Delta_{\bullet}^i](a) = G[\Delta_{\bullet}](a). \quad (10.50)$$

{eq:pwconvLegendre}

Note that in general,

$$\Delta \subsetneq \bigcap_{i \in I} \Delta^i.$$

**Proof** Fix  $a \in \Delta$ . It follows immediately from the definition of  $G$  that the net  $(G[\Delta_\bullet^i](a))_{i \in I}$  is decreasing and the  $\geq$  direction in (10.50) holds. Let us prove the reverse inequality. Let  $\tau$  denote the left-hand side of (10.50) for the moment. By definition, for any  $\epsilon > 0$  and any  $i \in I$ , we have  $a \in \Delta_{\tau-\epsilon}^i$ . It follows that

$$a \in \Delta_{\tau-\epsilon}^\infty.$$

Therefore,

$$\tau \leq G[\Delta_\bullet](a).$$

Similarly, for increasing nets, we have:

prop:incnetLegend

**Proposition 10.4.3** *Let  $(\Delta^i)_{i \in I}$  be an increasing net in  $\mathcal{K}_n$  with Hausdorff limit  $\Delta$  such that  $\text{vol } \Delta^i > 0$  for all  $i \in I$ . Consider an increasing net  $(\Delta_\bullet^i)_{i \in I}$  with  $\Delta_\bullet^i \in \text{TC}(\Delta^i)$  for all  $i \in I$ . Let  $\Delta_{\max} = \lim_{i \in I} \Delta_{\max}^i$ . For any  $\tau < \Delta_{\max}$ , let  $\Delta_\tau$  be the Hausdorff limit of  $\Delta_\tau^i$ . Then  $\Delta_\bullet \in \text{TC}(\Delta)$  and*

$$\lim_{i \in I} G[\Delta_\bullet^i](a) = G[\Delta_\bullet](a) \quad (10.51)$$

{eq:apwconvLegendre}

for any  $a \in \text{Int } \Delta$ .

**Proof** It is obvious that  $\Delta_\bullet \in \text{TC}(\Delta)$ .

Fix  $a \in \text{Int } \Delta$ . Then up to replacing  $I$  by a subnet, we may assume that  $a \in \Delta^i$  for all  $i \in I$ . By definition, the net  $(G[\Delta_\bullet^i](a))_{i \in I}$  is increasing and the  $\leq$  direction in (10.51) holds. Let us write  $\tau = G[\Delta_\bullet](a)$  for the time being. By definition of  $G$ , for any  $\epsilon > 0$ , we have

$$a \in \Delta_{\tau-\epsilon/2}.$$

The concavity of  $\Delta_\bullet$  guarantees that

$$a \in \text{Int } \Delta_{\tau-\epsilon}.$$

It follows that there is a subnet  $J$  in  $I$  such that for all  $j \in J$ ,

$$a \in \Delta_{\tau-\epsilon}^j.$$

Therefore,

$$\tau - \epsilon \leq G[\Delta_\bullet^j](a).$$

Taking the limit with respect to  $j$  and then with respect to  $\epsilon$ , we conclude the desired inequality.  $\square$

def:DHmeasureOTC

**Definition 10.4.4** Let  $\Delta_\bullet$  be an Okounkov test curve relative to  $\Delta$ . We define the *Duistermaat–Heckman measure*  $\text{DH}(\Delta_\bullet)$  as

$$\text{DH}(\Delta_\bullet) := G[\Delta_\bullet]_*(\text{vol}).$$

It is a Radon measure on  $\mathbb{R}$ .

In other words,  $\mathrm{DH}(\Delta_\bullet)$  is the distribution of the random variable  $G[\Delta_\bullet]$ .

prop:DHmoments

**Proposition 10.4.4** *Let  $\Delta_\bullet \in \mathrm{TC}(\Delta)$ . Let  $m \in \mathbb{Z}_{>0}$ . Then the  $m$ -th moment of the  $\mathrm{DH}(\Delta_\bullet)$  is given by*

$$\int_{\mathbb{R}} x^m \mathrm{DH}(\Delta_\bullet)(x) = \Delta_{\max}^m \mathrm{vol} \Delta + m \int_{-\infty}^{\Delta_{\max}} \tau^{m-1} (\mathrm{vol} \Delta_\tau - \mathrm{vol} \Delta) d\tau \quad (10.52)$$

{eq:momentcalc}

and

$$\int_{\mathbb{R}} \mathrm{DH}(\Delta_\bullet) = \mathrm{vol} \Delta. \quad (10.53)$$

{eq:massDHm1}

**Proof** In fact, (10.53) follows immediately from the definition, while (10.52) follows from a straightforward computation:

$$\begin{aligned} & \int_{\mathbb{R}} x^m \mathrm{DH}(\Delta_\bullet)(x) \\ &= \int_{\Delta} G[\Delta_\bullet](a)^m d \mathrm{vol}(a) \\ &= \int_{\Delta} \left( \Delta_{\max}^m - \int_{G[\Delta_\bullet](a)}^{\Delta_{\max}} m \tau^{m-1} d\tau \right) d \mathrm{vol}(a) \\ &= \Delta_{\max}^m \mathrm{vol} \Delta - m \int_{\mathbb{R}} \int_{\Delta} \mathbb{1}_{[G(\Delta_\bullet)(a), \Delta_{\max}]}(\tau) \tau^{m-1} d \mathrm{vol}(a) d\tau \\ &= \Delta_{\max}^m \mathrm{vol} \Delta - m \int_{-\infty}^{\Delta_{\max}} \int_{\Delta \setminus \Delta_\tau} \tau^{m-1} d \mathrm{vol}(a) d\tau \\ &= \Delta_{\max}^m \mathrm{vol} \Delta - m \int_{-\infty}^{\Delta_{\max}} \tau^{m-1} (\mathrm{vol} \Delta - \mathrm{vol} \Delta_\tau) d\tau. \end{aligned}$$

lma:DHmconv

**Lemma 10.4.2** *Let  $(\Delta^i)_{i \in I}$  be a decreasing net in  $\mathcal{K}_n$  with limit  $\Delta$ . Suppose that  $(\Delta_\bullet^i)_{i \in I}$  is a decreasing net with  $\Delta_\bullet^i \in \mathrm{TC}(\Delta^i)$ . Suppose that there is  $\Delta_\bullet \in \mathrm{TC}(\Delta)$  such that*

- (1)  $\Delta_{\max} = \lim_{i \in I} \Delta_{\max}^i$ ;
- (2) for any  $\tau < \Delta_{\max}$ , we have  $\Delta_\tau^i \xrightarrow{d_{\mathrm{Haus}}} \Delta_\tau$ .

Then  $\mathrm{DH}(\Delta_\bullet^i) \rightarrow \mathrm{DH}(\Delta_\bullet)$ .

**Proof** It follows from Proposition 10.4.2 that

$$G[\Delta_\bullet^i] \rightarrow G[\Delta_\bullet]$$

pointwisely on  $\Delta$ . Our assertion then follows from the dominated convergence theorem.  $\square$

Similarly, we have

lma:DHmconv2

**Lemma 10.4.3** *Let  $(\Delta^i)_{i \in I}$  be an increasing net in  $\mathcal{K}_n$  with Hausdorff limit  $\Delta$  such that  $\text{vol } \Delta^i > 0$  for all  $i \in I$ . Consider an increasing net  $(\Delta_\bullet^i)_{i \in I}$  with  $\Delta_\bullet^i \in \text{TC}(\Delta^i)$  for all  $i \in I$ . Let  $\Delta_\bullet \in \text{TC}(\Delta)$  be defined as*

- (1)  $\Delta_{\max} = \lim_{i \in I} \Delta_{\max}^i$ ;
- (2) for any  $\tau < \Delta_{\max}$ ,  $\Delta_\tau$  is the Hausdorff limit of  $\Delta_\tau^i$ .

Then we have

$$\text{DH}(\Delta_\bullet^i) \rightarrow \text{DH}(\Delta_\bullet).$$

**Proof** It follows from [Proposition 10.4.3](#) that

$$G[\Delta_\bullet^i] \rightarrow G[\Delta_\bullet]$$

almost everywhere on  $\Delta$ . Our assertion then follows from the dominated convergence theorem.  $\square$

The main source of Okounkov test curves is the following:

thm:Okountescurve

**Theorem 10.4.2** *Let  $X$  be a connected compact Kähler manifold and  $\theta$  be a closed smooth real  $(1, 1)$ -form on  $X$  representing a big cohomology class  $\alpha$ . Let  $Y_\bullet$  be a smooth flag on  $X$  and  $\Gamma \in \text{TC}(X, \theta)_{>0}$ . Then the map*

$$(-\infty, \Gamma_{\max}) \ni \tau \mapsto \Delta_{Y_\bullet}(\theta, \Gamma)_\tau := \Delta_{Y_\bullet}(\theta, \Gamma_\tau)$$

*defines an Okounkov test curve relative to  $\Delta_{Y_\bullet}(\theta, \Gamma_{-\infty})$ .*

*If furthermore  $\Gamma \in \text{TC}^1(X, \theta; \Gamma_{-\infty})$  (resp.  $\text{TC}^\infty(X, \theta; \Gamma_{-\infty})$ ), then we have  $\Delta_{Y_\bullet}(\theta, \Gamma) \in \text{TC}^1(\Delta_{Y_\bullet}(\theta, \Gamma_{-\infty}))$  (resp.  $\text{TC}^\infty(\Delta_{Y_\bullet}(\theta, \Gamma_{-\infty}))$ ).*

See [Definition 9.1.1](#) and [Definition 9.1.2](#) for the relevant definitions.

**Proof** Consider  $\Gamma \in \text{TC}(X, \theta)_{>0}$ . We need to verify that  $\Delta_{Y_\bullet}(\theta, \Gamma)$  is an Okounkov test curve relative to  $\Delta_{Y_\bullet}(\theta, \Gamma_{-\infty})$ .

First observe that  $\tau \mapsto \Delta_{Y_\bullet}(\theta, \Gamma_\tau)$  is concave and decreasing for  $\tau < \Gamma_{\max}$ . This is a direct consequence of [Theorem 10.3.4](#).

Next we show that as  $\tau \rightarrow -\infty$ , we have

$$\Delta_{Y_\bullet}(\theta, \Gamma_\tau) \xrightarrow{d_{\text{Haus}}} \Delta_{Y_\bullet}(\theta, \Gamma_{-\infty}).$$

It suffices to compute

$$\begin{aligned} \lim_{\tau \rightarrow -\infty} \text{vol } \Delta_{Y_\bullet}(\theta, \Gamma_\tau) &= \frac{1}{n!} \lim_{\tau \rightarrow -\infty} \text{vol}(\theta + \text{dd}^c \Gamma_\tau) = \frac{1}{n!} \text{vol}(\theta + \text{dd}^c \Gamma_{-\infty}) \\ &= \text{vol } \Delta_{Y_\bullet}(\theta, \Gamma_{-\infty}), \end{aligned}$$

where we applied [Theorem 10.3.2](#) and [Theorem 6.2.5](#).

When  $\Gamma \in \text{TC}^\infty(X, \theta; \Gamma_{-\infty})$ , it is clear that  $\Delta_{Y_\bullet}(\theta, \Gamma) \in \text{TC}^\infty(\Delta_{Y_\bullet}(\theta, \Gamma_{-\infty}))$ .

When  $\Gamma \in \text{TC}^1(X, \theta; \Gamma_{-\infty})$ , by [Theorem 10.3.2\(1\)](#), [\(9.3\)](#) and [\(10.43\)](#), we have

$$\mathbf{E}^{\Gamma_{-\infty}}(\Gamma) = \mathbf{E}(\Delta_{Y_\bullet}(\theta, \Gamma)).$$

So  $\Gamma \in \mathrm{TC}^1(\Delta_{Y_*}(\theta, \Gamma_{-\infty}))$ .

□



# Chapter 11

## The theory of b-divisors

chap:bdiv

In this chapter, we study the theory of b-divisors. In [Section 11.2](#), we prove a Chern–Weil type formula, which relates volumes of currents to intersection numbers.

In [Section 11.3](#), we prove that the algebraic partial Okounkov bodies constructed in [Chapter 10](#) have natural interpretations in terms of the b-divisors.

### 11.1 The intersection theory of b-divisors

In this section, we briefly recall the intersection theory of Dang–Favre [\[DF20, DF22\]](#).

Let  $X$  be a connected smooth projective variety of dimension  $n$ .

**Definition 11.1.1** A *birational model* of  $X$  is a projective birational morphism  $\pi : Y \rightarrow X$  from a *smooth* variety  $Y$ . A morphism between two birational models  $\pi : Y \rightarrow X$  and  $\pi' : Y' \rightarrow X$  is a morphism  $Y \rightarrow Y'$  over  $X$ .

We write  $\text{Bir}(X)$  for the isomorphism classes of birational models of  $X$ . It is a directed set under the partial ordering of domination.

We will usually be sloppy by omitting  $\pi$  and say  $Y$  is a birational model of  $X$ .

We write  $\text{NS}^1(X)$  for the Néron–Severi group of  $X$  and  $\text{NS}^1(X)_K$  for  $\text{NS}^1(X) \otimes_{\mathbb{Z}} K$  for any subfield  $K$  of  $\mathbb{R}$ . Given  $\alpha, \beta \in \text{NS}^1(X)_K$ , we write  $\alpha \leq \beta$  if  $\beta - \alpha$  is pseudo-effective.

**Definition 11.1.2** A *Weil b-divisor*  $\mathbb{D}$  on  $X$  is an assignment that associates with each  $(\pi : Y \rightarrow X) \in \text{Bir}(X)$  a class  $\mathbb{D}_Y = \mathbb{D}_\pi \in \text{NS}^1(Y)_{\mathbb{R}}$  such that when  $\pi' : Y' \rightarrow X$  dominates  $\pi$  through  $p : Y' \rightarrow Y$ , we have

$$p_* \mathbb{D}_{Y'} = \mathbb{D}_Y.$$

The set of Weil b-divisors on  $X$  is denoted by  $\text{bWeil}(X)$ .

A Weil b-divisor  $\mathbb{D}$  on  $X$  is *Cartier* if there is  $(\pi : Y \rightarrow X) \in \text{Bir}(X)$  such that for any  $(\pi' : Y' \rightarrow X) \in \text{Bir}(X)$  which dominates  $\pi$  through  $p : Y' \rightarrow Y$ , we have

$$\mathbb{D}_{Y'} = p^* \mathbb{D}_Y.$$

In this case we say  $\mathbb{D}$  is *determined* on  $Y$  or  $\mathbb{D}$  has an *incarnation*  $\mathbb{D}_Y$  on  $Y$  and write  $\mathbb{D} = \mathbb{D}(\mathbb{D}_Y)$ . We also say  $\mathbb{D}$  is a Cartier b-divisor. The linear space of Cartier b-divisors is denoted by  $\text{bCart}(X)$ .

Our definition simply means

$$\begin{aligned} \text{bWeil}(X) &= \varprojlim_{(\pi: Y \rightarrow X) \in \text{Bir}(X)} \text{NS}^1(Y)_{\mathbb{R}}, \\ \text{bCart}(X) &= \varinjlim_{(\pi: Y \rightarrow X) \in \text{Bir}(X)} \text{NS}^1(Y)_{\mathbb{R}}, \end{aligned} \quad (11.1) \quad \{\text{eq:bdivprojlim}\}$$

in the category of vector spaces.

We endow  $\text{bWeil}(X)$  with the projective limit topology, then the first equation in (11.1) becomes a projective limit in the category of locally convex linear spaces. Clearly,  $\text{bCart}(X)$  is dense in  $\text{bWeil}(X)$ .

def:nef

**Definition 11.1.3** A Cartier b-divisor  $\mathbb{D}$  on  $X$  is *nef* (resp. *big*) if some incarnation is (equivalently all incarnations are) nef (resp. *big*).

A Weil b-divisor  $\mathbb{D}$  on  $X$  is *nef* if it lies in the closure of the set of nef Cartier b-divisors.

Write  $\text{bWeil}_{\text{nef}}(X)$  for the set of nef Weil b-divisors on  $X$ .

A Weil b-divisor  $\mathbb{D}$  on  $X$  is *pseudo-effective* if for all  $(\pi: Y \rightarrow X) \in \text{Bir}(X)$ ,  $\mathbb{D}_Y \geq 0$ .

We introduce a partial ordering on  $\text{bWeil}(X)$ :

$$\mathbb{D} \leq \mathbb{D}' \text{ if and only if } \mathbb{D}_Y \leq \mathbb{D}'_Y \text{ for all } (\pi: Y \rightarrow X) \in \text{Bir}(X).$$

We summarise Dang–Favre’s results:

thm:DF1

**Theorem 11.1.1** (<sup>DF20</sup> [DF22, Theorem 2.1]) *Let  $\mathbb{D} \in \text{bWeil}(X)$  be a nef Weil b-divisor. Then there is a decreasing net  $(\mathbb{D}_i)_{i \in I}$  of nef Cartier b-divisors such that*

$$\mathbb{D} = \lim_{i \in I} \mathbb{D}_i.$$

def:nefint

**Definition 11.1.4** Let  $\mathbb{D}_i \in \text{bWeil}(X)$  ( $i = 1, \dots, n$ ) be nef Cartier b-divisors on  $X$ . We define  $(\mathbb{D}_1, \dots, \mathbb{D}_n) \in \mathbb{R}$  as follows: take  $(\pi: Y \rightarrow X) \in \text{Bir}(X)$  such that all  $\mathbb{D}_i$ ’s are determined on  $Y$ . Then define

$$(\mathbb{D}_1, \dots, \mathbb{D}_n) := (\mathbb{D}_{1,Y}, \dots, \mathbb{D}_{n,Y}). \quad (11.2)$$

The intersection number  $(\mathbb{D}_1, \dots, \mathbb{D}_n)$  does not depend on the choice of  $Y$ .

thm:DF2

**Theorem 11.1.2** (<sup>DF20</sup> [DF22, Proposition 3.1, Theorem 3.2]) *There is a unique pairing*

$$(\text{bWeil}_{\text{nef}}(X))^n \rightarrow \mathbb{R}_{\geq 0}$$

extending the pairing in Definition 11.1.4 such that



- (1) *The pairing is monotonically increasing in each variable.*
- (2) *The pairing is continuous along decreasing nets in each variable.*

Moreover, this pairing has the following properties:

- (1) *It is symmetric, multilinear.*
- (2) *It is usc in each variable.*

**Definition 11.1.5** We define the *volume* of  $\mathbb{D} \in \text{bWeil}_{\text{nef}}(X)$  by

$$\text{vol } \mathbb{D} = (\mathbb{D}, \dots, \mathbb{D}). \quad (11.3)$$

{eq:volbdivdef}

We say  $\mathbb{D} \in \text{bWeil}_{\text{nef}}(X)$  is *big* if  $\text{vol } \mathbb{D} > 0$ .

Note that the definition of bigness is compatible with the definition in [Definition 11.1.3](#) in the case of Cartier b-divisors.

lma:volbdivaslim

**Lemma 11.1.1** *Let  $\mathbb{D} \in \text{bWeil}_{\text{nef}}(X)$ , then*

$$\text{vol } \mathbb{D} = \inf_{(Y \rightarrow X) \in \text{Bir}(X)} \text{vol } \mathbb{D}_Y = \lim_{(Y \rightarrow X) \in \text{Bir}(X)} \text{vol } \mathbb{D}_Y.$$

**Proof** By [Theorem 11.1.1](#), we can find a decreasing net  $\mathbb{D}^\alpha$  of nef Cartier b-divisors on  $X$  converging to  $\mathbb{D}$ . Clearly,

$$\text{vol } \mathbb{D}^\alpha = \inf_{Y \rightarrow X} \text{vol } \mathbb{D}_Y^\alpha.$$

It follows from [Theorem 11.1.2](#) and the continuity of the volume functional [\[ELMNP05, Corollary 2.6\]](#) that

$$\text{vol } \mathbb{D} = \inf_{\alpha} \inf_{Y \rightarrow X} \text{vol } \mathbb{D}_Y^\alpha = \inf_{Y \rightarrow X} \text{vol } \mathbb{D}_Y.$$

On the other hand, as in general push-forward will increase the volume, we see that  $\text{vol } \mathbb{D}_Y$  is decreasing in  $Y$ , so we conclude.  $\square$

## 11.2 The singularity b-divisors

sec:bdiv1

Let  $X$  be a connected smooth projective variety over  $\mathbb{C}$  of dimension  $n$ . Let  $\alpha \in \text{NS}^1(X)_{\mathbb{R}}$  be a big class and  $T$  be a closed positive  $(1, 1)$ -current in  $\alpha$ .

Fix a closed real smooth  $(1, 1)$ -form  $\theta$  in  $c_1(L)$  and we can write  $T = \theta_\varphi$  for some  $\varphi \in \text{PSH}(X, \theta)$ .

**Definition 11.2.1** Define the *singularity divisor*  $\text{Sing}_X T$  of  $T$  as the formal sum

$$\text{Sing}_X T := \sum_E \nu(T, E) E, \quad (11.4)$$

{eq:singhatL}

where  $E$  runs over all prime divisors contained in  $X$ .

The singularity divisor is *not* a Weil divisor in general.

Note that this is a countable sum by Siu's semicontinuity theorem. Although  $\text{Sing}_X T$  is not a divisor in general, it does define a closed positive  $(1, 1)$ -current due to Siu's decomposition. Moreover, the numerical class  $[\text{Sing}_X T]$  in  $\text{NS}^1(X)_{\mathbb{R}}$  is also well-defined by treating the sum in (11.4) as a sum of numerical classes [BFJ09, Proposition 1.3].

def:singbdiv

**Definition 11.2.2** The *singularity b-divisor*  $\text{Sing } T$  of  $T$  is the b-divisor over  $X$  defined by

$$(\text{Sing } T)_Y := [\text{Sing}_Y \pi^* T],$$

where  $(\pi: Y \rightarrow X) \in \text{Bir}(X)$ .

Define

$$\mathbb{D}(T) := \mathbb{D}(\alpha) - \text{Sing } T.$$

Here  $\mathbb{D}(\alpha)$  is the Cartier b-divisor determined by  $\alpha$  on  $X$ .

We are ready to derive the first version of the Chern–Weil formula.

thm:nefbvolume

**Theorem 11.2.1** The b-divisor  $\mathbb{D}(T)$  is a nef b-divisor and if in addition  $\text{vol } T > 0$ ,

$$\text{vol } \mathbb{D}(T) = \text{vol } T. \quad (11.5)$$

{eq:volbandline}

**Proof Step 1.** We first handle the case where  $T$  has analytic singularities. After replacing  $X$  by a modification, we may assume that  $T$  has log singularities along an effective  $\mathbb{Q}$ -divisor  $D$  on  $X$ . Namely, we can write

$$T = [D] + R,$$

where  $R$  is a closed positive  $(1, 1)$ -current with bounded potential. In this case,  $\mathbb{D}(T) = \mathbb{D}(\alpha - D)$ , which is nef. In order to prove (11.5), it suffices to show that

$$\int_X T^n = ((\alpha - D)^n), \quad (11.6)$$

{eq:temp14}

which is obvious.

**Step 2.** Assume that  $T$  is a Kähler current. Take a quasi-equisingular approximation  $(T_j)_j$  of  $T$  in  $\mathcal{Z}_+(X, \theta)$ . By Theorem 6.2.5, we have

$$\lim_{j \rightarrow \infty} \text{vol } T_j = \text{vol } T.$$

In view of Step 1 and Theorem 11.1.2, it remains to show that  $\mathbb{D}(T_j) \rightarrow \mathbb{D}(T)$  as  $j \rightarrow \infty$ . In more concrete terms, this means that for any  $(\pi: Y \rightarrow X) \in \text{Bir}(X)$ ,

$$[\text{Sing}_Y(\pi^* T_j)] \rightarrow [\text{Sing}_Y(\pi^* T)]$$

in  $\text{NS}^1(Y)_{\mathbb{R}}$ . This obviously follows from Theorem 6.2.4 if  $\text{Sing}(\pi^* T)$  has only finitely many components. In general, fix an ample class  $\omega$  in  $\text{NS}^1(Y)$ . We want to show that

for any  $\epsilon > 0$ , we can find  $j_0 > 0$  so that when  $j \geq j_0$ ,

$$[\text{Sing}_Y(\pi^*T_j)] \geq [\text{Sing}_Y(\pi^*T)] - \epsilon\omega. \quad (11.7) \quad \{\text{eq:temp55}\}$$

Write

$$[\text{Sing}_Y(\pi^*T)] = \sum_{i=1}^{\infty} a_i E_i, \quad [\text{Sing}(\pi^*T_j)] = \sum_{i=1}^{\infty} a_i^j E_i.$$

Then  $a_i^j \leq a_i$ . We can find  $N > 0$  large enough, so that

$$[\text{Sing}_Y(\pi^*T)] \leq \sum_{i=1}^N a_i E_i + \frac{\epsilon}{2}\omega.$$

By [Theorem 6.2.4](#), we can take  $j_0$  large enough so that for  $j > j_0$ ,

$$(a_i - a_i^j)E_i \leq \frac{\epsilon}{2N}\omega, \quad i = 1, \dots, N.$$

Then (11.7) follows.

**Step 3.** Assume that  $\text{vol } T > 0$ .

By [Lemma 2.3.2](#), we can take a Kähler current  $S \in \alpha$  such that  $S \leq T$ . Consider  $\epsilon S + (1 - \epsilon)T$  for  $\epsilon \in (0, 1)$ . When  $\epsilon \rightarrow 0+$ , we have  $\epsilon S + (1 - \epsilon)T \xrightarrow{d_S} T$ . Using [Theorem 6.2.5](#), we reduce immediately to the situation of Step 2.

**Step 4.** We handle the general case.

Take a Kähler form  $\omega$  on  $X$ . From Step 3, we know that for any  $\epsilon > 0$ ,  $\mathbb{D}(T) + \epsilon\mathbb{D}(\omega)$  is a nef b-divisor. It follows immediately that  $\mathbb{D}(T)$  is nef.  $\square$

`cor:Imodcharbdiv`

**Corollary 11.2.1** Assume that  $\text{vol } T > 0$ , then  $T$  is  $\mathcal{I}$ -good if and only if

$$\text{vol } \mathbb{D}(T) = \int_X T^n.$$

**Proof** This follows from [Theorem 11.2.1](#) and [Theorem 7.3.1](#).  $\square$

`thm:pshbdivcont`

**Theorem 11.2.2** The map  $\mathbb{D}: \text{PSH}(X, \theta) \rightarrow \text{bWeil}(X)$  is continuous. Here on  $\text{PSH}(X, \theta)$  we take the  $d_S$ -pseudometric.

**Proof** Let  $\varphi_i \in \text{PSH}(X, \theta)$  be a sequence converging to  $\varphi \in \text{PSH}(X, \theta)$  with respect to  $d_S$ . We want to show that

$$\mathbb{D}(\theta + \text{dd}^c \varphi_i) \rightarrow \mathbb{D}(T).$$

As  $\varphi_i \xrightarrow{d_S} \varphi$  implies that  $\pi^* \varphi_i \xrightarrow{d_S} \pi^* \varphi$  for any  $(\pi: Y \rightarrow X) \in \text{Bir}(X)$ , it suffices to prove

$$[\text{Sing}_X \varphi_i] \rightarrow [\text{Sing}_X \varphi] \quad \text{in } \text{NS}^1(X)_{\mathbb{R}}. \quad (11.8) \quad \{\text{eq:temp7}\}$$

Write

$$\text{Sing}_X \varphi_i = \sum_E a_i^E E, \quad \text{Sing}_X \varphi = \sum_E a^E E,$$

where  $E$  runs over all prime divisors on  $X$ . By [Theorem 6.2.4](#),  $a_i^E \rightarrow a^E$  as  $i \rightarrow \infty$ . When the number of  $E$ 's is finite, (11.8) follows trivially. Otherwise, we write the prime divisors on  $X$  having positive coefficients in either  $\text{Sing}_X \varphi_i$  or  $\text{Sing}_X \varphi$  as  $E_1, E_2, \dots$ .

We fix a basis  $e_1, \dots, e_N$  of the finite-dimensional vector space  $\text{NS}^1(X)_{\mathbb{R}}$ , so that the pseudo-effective cone is contained in the cone  $\sum_d \mathbb{R}_{\geq 0} e_d$ . Write

$$E_i = \sum_{d=1}^N f_i^d e_d, \quad i = 1, 2, \dots$$

Then we need to show that for any  $d = 1, \dots, N$ ,

$$\lim_{i \rightarrow \infty} \sum_{j=1}^{\infty} a_i^{E_j} f_j^d = \sum_{j=1}^{\infty} a^{E_j} f_j^d.$$

This follows from the dominated convergence theorem, since

$$\sum_{j=1}^{\infty} a_i^{E_j} [E_j] \leq \alpha, \quad \sum_{j=1}^{\infty} a^{E_j} [E_j] \leq \alpha.$$

A mixed version of [Theorem 11.2.1](#) is also true:

thm:nefbvolume2

**Theorem 11.2.3** *Let  $T_1, \dots, T_n \in \mathcal{Z}_+(X)$  such that  $\text{vol } T_i > 0$  for each  $i = 1, \dots, n$ . Then*

$$\frac{1}{n!} (\mathbb{D}(T_1), \dots, \mathbb{D}(T_n)) \geq \frac{1}{n!} \int_X T_1 \wedge \dots \wedge T_n. \quad (11.9)$$

{eq:bdivmixint}

*If the  $T_i$ 's are  $\mathcal{I}$ -good, then equality holds.*

**Proof** This follows from [Theorem 11.2.1](#) and [Proposition 7.2.1](#).  $\square$

### 11.3 Okounkov bodies of b-divisors

sec:Okounkovbdiv

Let  $X$  be a connected projective manifold of dimension  $n$  and  $(L, h)$  be a Hermitian big line bundle on  $X$ .

Fix a smooth flag  $Y_{\bullet}$  on  $X$ . Let  $\nu = \nu_{Y_{\bullet}}: \mathbb{C}(X)^{\times} \rightarrow \mathbb{Z}^n$  be the valuation associated with  $Y_{\bullet}$ .

thm:pobbd

**Theorem 11.3.1** *The partial Okounkov body  $\Delta_{Y_{\bullet}}(L, h)$  admits the following expression:*

$$\Delta_{Y_{\bullet}}(L, h) = \nu_{Y_{\bullet}}(\text{dd}^c h) + \lim_{\pi: Z \rightarrow X} \Delta_{Y_{\bullet}}(c_1(\pi^* L) - [\text{Sing}_Z(\pi^* h)]), \quad (11.10)$$

{eq:DeltaasHlim}

*where  $\pi$  runs over the directed set of projective birational morphisms to  $X$  with  $Z$  normal.*

Here the limit is a Hausdorff limit.

This theorem suggests that we define

$$\Delta_{Y_\bullet}(\mathbb{D}(\mathrm{dd}^c h)) := \lim_{\pi: Z \rightarrow X} \Delta_{Y_\bullet}(c_1(\pi^* L) - [\mathrm{Sing}_Z(\pi^* h)]). \quad (11.11)$$

{eq:Okoubodbdiv}

Then one could rewrite (11.10) as

$$\Delta_{Y_\bullet}(L, h) = \Delta_{Y_\bullet}(\mathbb{D}(\mathrm{dd}^c h)) + \nu_{Y_\bullet}(\mathrm{dd}^c h).$$

*Remark 11.3.1* (11.11) shows that the partial Okounkov bodies are *algebraic* objects in nature.

One should be able to prove the existence of the limits like (11.11) over other base fields, at least after assuming the existence of resolution of singularities. If so, one would get an interesting extension of the theory of partial Okounkov bodies.

lma:valuationT

**Lemma 11.3.1** *Let  $T$  be a closed positive  $(1, 1)$ -current on  $X$ . Then we have*

$$\lim_{\pi: Z \rightarrow X} \nu(\mathrm{Sing}_Z(\pi^* T)) = \nu(T), \quad (11.12)$$

{eq:nuTaslimit}

where  $\pi$  runs over the directed set of projective birational morphisms to  $X$  with  $Z$  normal.

**Proof** Given  $\pi: Z \rightarrow X$ , we let  $W_1$  denote the strict transform of  $Y_1$  in  $Z$ . The restriction  $\pi_1: W_1 \rightarrow Y_1$  is necessarily birational. Let  $\widetilde{W}_1$  be the normalization of  $W_1$ . Let  $\widetilde{\pi}_1$  denote the normalization of  $\pi_1$  so that we have a commutative diagram

$$\begin{array}{ccccc} \widetilde{W}_1 & \longrightarrow & W_1 & \hookrightarrow & Z \\ \downarrow \widetilde{\pi}_1 & & \downarrow \pi_1 & & \downarrow \pi \\ Y_1 & \xlongequal{\quad} & Y_1 & \hookrightarrow & X. \end{array}$$

We will argue by induction. The case  $n = 0$  is trivial. Assume that  $n > 0$  and the case  $n - 1$  is known.

We may clearly assume that  $\nu(T, Y_1) = 0$ . By definition, we have

$$\nu(T) = (0, \mu(\mathrm{Tr}_{Y_1}(T))),$$

where  $\mu$  denotes the valuation induced by the flag  $Y_1 \supseteq Y_2 \supseteq \cdots \supseteq Y_n$ .

Observe that birational morphisms of the form  $\pi_1: \widetilde{W}_1 \rightarrow Y_1$  are cofinal in the directed set of projective birational morphisms of  $Y_1$ . This is obvious since the modifications given by compositions of blow-ups with smooth centers on  $Y_1$  are cofinal. It suffices to blow-up  $X$  with the same centers.

Therefore, by the inductive hypothesis applied to  $\mathrm{Tr}_{Y_1} T$ , it suffices to argue that

$$\nu(\mathrm{Sing}_Z(\pi^* T)) = \left(0, \mu\left(\mathrm{Sing}_{\widetilde{W}_1} \widetilde{\pi}_1^*(\mathrm{Tr}_{Y_1}(T))\right)\right). \quad (11.13)$$

{eq:indstep}

From [Lemma 8.2.1](#), we know that

$$\tilde{\pi}_1^* \operatorname{Tr}_{Y_1}(T) \sim_P \operatorname{Tr}_{W_1}(\pi^*T).$$

So we only need to prove

$$\nu(\operatorname{Sing}_Z(\pi^*T)) = \left(0, \mu(\operatorname{Sing}_{\widehat{W_1}}(\operatorname{Tr}_{W_1}(\pi^*T)))\right),$$

This is reduced to the following statement:

$$\operatorname{Tr}_{W_1} \operatorname{Sing}_Z(\pi^*T) \sim_P \operatorname{Sing}_{\widehat{W_1}}(\operatorname{Tr}_{W_1}(\pi^*T)). \quad (11.14)$$

{eq:nusingzpistarTtemp1}

In order to prove this, we may add a Kähler form to  $T$  and assume that  $T$  is a Kähler current. Take a quasi-equisingular approximation  $(T_j)_j$  of  $T$ . Then  $(\pi^*T_j)_j$  is a quasi-equisingular approximation of  $\pi^*T$ . Thanks to [Proposition 8.2.2](#), we have

$$\operatorname{Tr}_{W_1}(\pi^*T_j) \xrightarrow{d_S} \operatorname{Tr}_{W_1}(\pi^*T)$$

Therefore, as in the proof of [Theorem 11.2.2](#), we find that  $\operatorname{Sing}_Z$  and  $\operatorname{Sing}_{\widehat{W_1}}$  are both continuous along this sequence as well. So we finally reduce to the case where  $T$  has analytic singularities.

In this case, arguing as before, we may assume replace  $\pi$  by a modification dominating it so that  $\pi^*T \sim [D]$  for an effective  $\mathbb{Q}$ -divisor  $D$  on  $Z$ , in which case [\(11.14\)](#) is clear.  $\square$

**Proof (The proof of [Theorem 11.3.1](#))** It would be more convenient to use the language of currents. We shall write  $T = \operatorname{dd}^c h$ .

Instead of arguing [\(11.10\)](#), we shall argue a slightly more general version: for any  $\alpha \in \operatorname{NS}^1(X)_{\mathbb{R}}$ , we have

$$\Delta_{Y_\bullet}(T) = \nu(T) + \lim_{\pi: Z \rightarrow X} \Delta_{Y_\bullet}(\alpha - [\operatorname{Sing}_Z(\pi^*T)]). \quad (11.15)$$

{eq:mainvar}

We argue by induction on  $n$ . The case  $n = 0$  is of course trivial. Let us assume that  $n > 0$  and the result is known in dimension  $n - 1$ .

We may replace  $T$  by  $T - \nu(T, Y_1)[Y_1]$  and  $\alpha$  by  $\alpha - \nu(T, Y_1)[Y_1]$ , so that we may reduce to the case where  $\nu(T, Y_1) = 0$ .

For any projective birational morphism  $\pi: Z \rightarrow X$  with  $Z$  normal, it follows from [Theorem 10.3.4](#) (which also holds for a normal variety, as can be seen after passing to a resolution) that we have

$$\Delta_{Y_\bullet}(\pi^*\alpha - [\operatorname{Sing}_Z(\pi^*T)]) = \overline{\{\nu(S) : S \in \pi^*\alpha - [\operatorname{Sing}_Z(\pi^*T)]\}}.$$

Therefore,

$$\Delta_{Y_\bullet}(\pi^*\alpha - [\operatorname{Sing}_Z(\pi^*T)]) + \nu(\operatorname{Sing}_Z(\pi^*T)) \subseteq \overline{\{\nu(S) : S \in \alpha, \pi^*S \geq \operatorname{Sing}_Z(\pi^*T)\}}.$$

We observe that the right-hand side is decreasing with respect to  $\pi$ , which together with [Lemma 11.3.1](#) implies that the net of convex bodies  $\Delta_{Y_\bullet}(c_1(\pi^*L) - [\text{Sing}_Z(\pi^*T)])$  for various  $Z$  is uniformly bounded. Suppose that  $\Delta$  is the limit of a subnet. Then we have

$$\Delta + \nu(T) \subseteq \overline{\{\nu(S) : S \in c_1(L), S \leq_I T\}}.$$

As shown in [Theorem 10.3.4](#), the right-hand side is exactly  $\Delta_{Y_\bullet}(T)$ . So

$$\Delta + \nu(T) \subseteq \Delta_{Y_\bullet}(T).$$

But observe that both sides have the same volume, as computed in [Theorem 10.3.2](#) and [Theorem 11.2.1](#). So equality holds.

It follows from the Blaschke selection theorem [Theorem C.1.1](#) that the limit in [\(11.15\)](#) exists and [\(11.15\)](#) holds.  $\square$





## **Part III**

# **Applications**

In this part, we explain a few applications of the theory developed in this book.

In [Chapter 12](#), we develop the pluripotential theory on big line bundles on toric varieties. This theory depends crucially on the theory of partial Okounkov bodies developed in [Chapter 10](#).

In [Chapter 13](#), we develop the transcendental theory of non-Archimedean metrics based on the theory of test curves developed in [Chapter 9](#).

In [Chapter 14](#), we prove the convergence of partial Bergman measures.

## Chapter 12

# Toric pluripotential theory on big line bundles

chap:toricbig

In this chapter, we develop the toric pluripotential theory on big line bundles. Our development here is based on the theory of partial Okounkov bodies developed in [Chapter 10](#). We will deduce two non-trivial consequences from the general theory: [Corollary 12.2.2](#) and [Theorem 12.2.2](#). The author does not know how to prove either result without relying on partial Okounkov bodies.

### 12.1 Toric setup

Let  $T$  be a complex torus of dimension  $n$  with character lattice  $M$  and cocharacter lattice  $N$ . Consider a rational polyhedral fan  $\Sigma$  in  $N_{\mathbb{R}}$  corresponding to an  $n$ -dimensional smooth toric variety  $X$ .

Let  $D$  be a  $T$ -invariant big divisor on  $X$ . Then  $P_D \subseteq M_{\mathbb{R}}$  be the lattice polytope generated by  $u \in M$  such that

$$D + \operatorname{div} \chi^u \geq 0.$$

Let  $L = \mathcal{O}_X(D)$ . Note that replacing  $D$  by a linearly equivalent divisor amounts to replace  $D$  by an integral translation.

We shall fix a smooth  $T_c$ -invariant metric  $h_0$  on  $L$ . Let  $\theta = c_1(L, h_0)$ . Fix a smooth function  $F_{\theta}: N_{\mathbb{R}} \rightarrow \mathbb{R}$  such that

$$\theta = \operatorname{dd}^c \operatorname{Trop}^* F_{\theta}.$$

Note that  $F_{\theta}$  is well-defined up to a linear term.

We will consider a  $T$ -invariant subvariety  $Y \subseteq X$ . Since  $X$  is smooth, so is  $Y$ . Let  $\sigma$  be the cone in  $\Sigma$  corresponding to  $Y$  and  $Q$  be the face of  $P$  corresponding to  $Y$ .

Recall that the cocharacter lattice  $N(\sigma)$  of  $Y$  is given by  $N/N \cap \langle \sigma \rangle$ , where  $\langle \sigma \rangle$  is the linear span of  $\sigma$ . See [\[CLS11, \(3.2.6\)\]](#). In particular, the character lattice  $M(\sigma)$  of  $Y$  can be naturally identified with the linear span of  $Q$ . Let  $i_{\sigma}: M(\sigma) \rightarrow M$  be the corresponding inclusion.

Take  $m_\sigma \in M \cap P_D$  so that  $-\text{Supp}_{-P_D}$  coincides with  $m_\sigma$  on  $\sigma$ . Observe that  $m_\sigma$  is uniquely determined only when  $\sigma$  has full dimension.

## 12.2 Toric partial Okounkov bodies

### 12.2.1 Newton bodies

Let  $\text{PSH}_{\text{tor}}(X, \theta)$  be the set of  $T_c$ -invariant functions in  $\text{PSH}(X, \theta)$ .

**Definition 12.2.1** A function  $\varphi \in \text{PSH}_{\text{tor}}(X, \theta)$  can be written as

$$\varphi|_{T(\mathbb{C})} = \text{Trop}^* f$$

for some unique  $f: N_{\mathbb{R}} \rightarrow [-\infty, \infty)$ . Then we define

$$F_\varphi: N_{\mathbb{R}} \rightarrow \mathbb{R}$$

as follows:

$$F_\varphi = F_\theta + f. \quad (12.1)$$

Observe that  $F_\varphi$  is a convex function and takes finite values by [Lemma 5.2.1](#). It is well-defined up to a linear term.

**Definition 12.2.2** Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \theta)$ , we define its *Newton body* as

$$\Delta(\theta, \varphi) := \overline{\nabla F_\varphi(N_{\mathbb{R}})} \subseteq M_{\mathbb{R}}.$$

Observe that  $\Delta(\theta, \varphi)$  depends only on the current  $\theta_\varphi$ , not on the choices of  $\theta$  and  $F_\theta$ .

### 12.2.2 Partial Okounkov bodies

subsec:pobtorgeneral

There are some canonical choices of smooth flags in the toric setting.

Recall that for each  $\rho \in \Sigma(1)$ ,  $u_\rho$  denotes the ray generator of  $\rho$ . Since  $X$  is smooth and projective, we could choose a full-dimensional cone  $\sigma$  in  $\Sigma$  with rays  $\rho_1, \dots, \rho_n \in \Sigma(1)$  such that  $u_{\rho_1}, \dots, u_{\rho_n}$  form a basis of  $N$ . Define

$$Y_i = D_{\rho_1} \cap \dots \cap D_{\rho_i}, \quad i = 1, \dots, n.$$

Then  $Y_\bullet$  is a smooth flag on  $X$ . Let

$$\Phi: M \rightarrow \mathbb{Z}^n, \quad m \mapsto (\langle m - m_\sigma, u_{\rho_1} \rangle, \dots, \langle m - m_\sigma, u_{\rho_n} \rangle). \quad (12.2)$$

{eq:isoMZcanonical}

Then  $\Phi$  is an isomorphism of lattices. It induces an  $\mathbb{Z}$ -affine isomorphism

$$\Phi_{\mathbb{R}}: M_{\mathbb{R}} \rightarrow \mathbb{R}^n.$$

prop:toricusual0ko

**Proposition 12.2.1** *We have*

$$\nu_{Y_\bullet} \left( H^0(X, L^k)^\times \right) = \Phi((kP_D) \cap M) \quad (12.3)$$

{eq:DeltakLtoric}

for any  $k \in \mathbb{Z}_{>0}$ . In particular,

$$\Delta_{Y_\bullet}(L) = \Phi_{\mathbb{R}}(P_D). \quad (12.4)$$

**Proof** Up to replacing  $D$  by a linearly equivalent divisor, we may assume that  $D|_{U_\sigma} = 0$ , where  $U_\sigma$  is the affine subvariety of  $X$  corresponding to  $\sigma$ . Then  $m_\sigma = 0$ .

It suffices to prove (12.3) for  $k = 1$ . Let  $s \in H^0(X, L)$  be a non-zero section, say  $\chi^u$  for some  $u \in P_D \cap M$ . The zero-locus of  $s$  is given by

$$D + \sum_{i=1}^n \langle u, u_{\rho_i} \rangle D_{\rho_i}.$$

Therefore,

$$\nu_{Y_\bullet}(s) = (\langle u, u_{\rho_1} \rangle, \dots, \langle u, u_{\rho_n} \rangle) = \Phi(u).$$

So (12.3) follows.  $\square$

thm:toricpob

**Theorem 12.2.1** *Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \theta)_{>0}$ , then*

$$\Phi_{\mathbb{R}}(\Delta(\theta, \varphi)) = \Delta_{Y_\bullet}(\theta, \varphi). \quad (12.5)$$

{eq:toricOkounkovcomp}

**Proof** Up to replacing  $D$  by a linearly equivalent divisor, we may assume that  $D|_{U_\sigma} = 0$ , where  $U_\sigma$  is the affine subvariety of  $X$  corresponding to  $\sigma$ . Then  $m_\sigma = 0$ .

**Step 1.** We first reduce to the case where  $\theta_\varphi$  is a Kähler current.

By Lemma 2.3.2, we can find  $\psi \in \text{PSH}(X, \theta)$  such that  $\psi \leq \varphi$  and  $\theta_\psi$  is a Kähler current. Taking the average along  $T_c$ , we may assume that  $\psi$  is  $T_c$ -invariant.

For each  $t \in (0, 1)$ , we let

$$\varphi_t = (1 - t)\psi + t\varphi.$$

Suppose that Kähler current case is known. Then we get

$$\Phi_{\mathbb{R}}(\Delta(\theta, \varphi_t)) = \Delta_{Y_\bullet}(\theta, \varphi_t)$$

for any  $t \in (0, 1)$ . It follows from Theorem A.4.2 that

$$\Phi_{\mathbb{R}}(\Delta(\theta, \varphi)) \supseteq \Phi_{\mathbb{R}}(\Delta(\theta, \varphi_t)) \supseteq \Delta_{Y_\bullet}(\theta, \varphi_t)$$

for any  $t \in (0, 1)$ . Thanks to [Theorem 10.2.2](#), we have

$$\Phi_{\mathbb{R}}(\Delta(\theta, \varphi)) \supseteq \Delta_{Y_*}(\theta, \varphi).$$

Compare the volumes of both sides using [Proposition 12.2.2](#) and (10.11), we find that

$$n! \operatorname{vol} \Phi_{\mathbb{R}}(\Delta(\theta, \varphi)) = \int_X \theta_{\varphi}^n = \operatorname{vol} \theta_{\varphi} = n! \operatorname{vol} \Delta_{Y_*}(\theta, \varphi).$$

In particular, we conclude (12.5).

**Step 2.** We handle the case where  $\theta_{\varphi}$  is a Kähler current.

Let  $(\varphi_j)_j$  be a quasi-equisingular approximation of  $\varphi$  in  $\operatorname{PSH}(X, \theta)$ .

We may assume that  $\varphi_j$  is  $T_c$ -invariant for each  $j \geq 1$  from the construction of [Dem12](#), [Theorem 13.21](#).

Now assume that the result is known for each  $\varphi_j$ . Then

$$\Phi_{\mathbb{R}}(\Delta(\theta, \varphi_j)) = \Delta_{Y_*}(\theta, \varphi_j).$$

In particular, by [Proposition 12.2.2](#) again,

$$\Phi_{\mathbb{R}}(\Delta(\theta, \varphi)) \subseteq \Delta_{Y_*}(\theta, \varphi_j)$$

for each  $j \geq 1$ . It follows from [Theorem 10.2.2](#) that

$$\Phi_{\mathbb{R}}(\Delta(\theta, \varphi)) \subseteq \Delta_{Y_*}(\theta, \varphi).$$

Compare the volumes of both sides using [Proposition 12.2.2](#), (10.11) and [Theorem 5.3.1](#), we conclude (12.5).

**Step 3.** It remains to handle the case where  $\varphi$  has analytic singularities and  $\theta_{\varphi}$  is a Kähler current. In fact, we may assume that  $\varphi$  has the form

$$\varphi = \log \sum_{i=1}^a |s_i|_{h_0}^2 + O(1),$$

where  $s_1, \dots, s_a \in H^0(X, L)$ . This follows from the proof of Step 2 and the construction of [Dem12](#), [Theorem 13.21](#).

Let  $u_1, \dots, u_a \in P_D \cap M$  be the lattice points corresponding to  $s_1, \dots, s_a$ . Observe that  $\Delta(\theta, \varphi)$  is the convex envelope of  $u_1, \dots, u_a$  by [Lemma A.5.2](#).

Then for any  $m \in M$  and  $k \in \mathbb{Z}_{>0}$ ,  $m \in kP_D$  if and only if

$$|\chi^m|_{h_0}^2 e^{-k\varphi}$$

is bounded from above. It follows that

$$\Phi(k\Delta(\theta, \varphi) \cap M) \subseteq k\Delta_k(\theta, \varphi).$$

The notation  $\Delta_k$  is defined [Section 10.2](#). Letting  $k \rightarrow \infty$  and applying [Theorem 10.2.4](#), we find that

$$\Phi_{\mathbb{R}}(\Delta(\theta, \varphi)) \subseteq \Delta(\theta, \varphi).$$

Compare the volumes of both sides using [Proposition 12.2.2](#) and [\(10.11\)](#), we conclude that the equality holds and [\(12.5\)](#) follows.  $\square$

As another consequence we have

cor:toricLelong

**Corollary 12.2.1** *Let  $E$  be a  $T$ -invariant prime divisor on  $X$  corresponding to a ray with ray generator  $n \in N$ . Then for any  $\varphi \in \text{PSH}_{\text{tor}}(X, \theta)_{>0}$ , we have*

$$v(\varphi, E) = \inf \{ \langle m - m_{\sigma}, n \rangle : m \in \Delta(\theta, \varphi) \},$$

where  $\sigma$  is the ray in  $\Sigma$  corresponding to  $E$ .

**Proof** This follows immediately from [Theorem 12.2.1](#) and [Theorem 10.2.5](#). In fact, since  $X$  is projective and smooth, there is always a  $T$ -invariant smooth flag  $Y_{\bullet}$  with  $Y_1 = E$ .  $\square$

cor:toricLelong2

**Corollary 12.2.2** *For any  $T$ -invariant subvariety  $Y \subseteq X$  corresponding to a cone  $\sigma$  in  $\Sigma$  and any  $\varphi \in \text{PSH}_{\text{tor}}(X, \theta)_{>0}$ . Then the following are equivalent:*

- (1)  $v(\varphi, Y) = 0$ ;
- (2) *There is a point  $m \in \Delta(\theta, \varphi)$  such that  $(m - m_{\rho}) \cdot u_{\rho} = 0$  for any 1-dimensional face  $\rho$  of  $\sigma$ .*

**Proof** Let  $\rho_1, \dots, u_r$  be the rays of  $\sigma$ . Up to replacing  $D$  by a translation, we may assume that  $m_{\sigma} = 0$ .

Let  $\pi: Z \rightarrow X$  be the blow-up of  $X$  along  $Y$ . Observe that  $\Delta(\theta, \varphi) = \Delta(\pi^*\theta, \pi^*\varphi)$ . On the other hand, the ray corresponding to the exceptional divisor  $E$  is generated by  $u_{\rho_1} + \dots + u_{\rho_r}$ . Since  $X$  is smooth, this vector is primitive. Bou02

It follows from [Corollary 12.2.1](#) and [\[Bou02a, Corollaire 1.1.8\]](#) that

$$v(\varphi, Y) = v(\pi^*\varphi, E) = \inf \{ \langle m, u_{\rho_1} + \dots + u_{\rho_r} \rangle : m \in \Delta(\theta, \varphi) \}. \quad (12.6)$$

{eq:nuvarphiYtoric1}

Our assertion follows.  $\square$

It follows from [\(12.6\)](#) that

$$v(\varphi, Y) \geq \sum_{i=1}^a v(\varphi, E_i),$$

where the  $E_i$ 's are the prime divisors corresponding to the rays of  $\sigma$ . This inequality seems to be new as well.

thm:FVtheta

**Theorem 12.2.2** *We have*

$$F_{V_{\theta}} \in \mathcal{E}(N_{\mathbb{R}}, P_D).$$

**Proof** Take  $\varphi = V_\theta$  in [Theorem 12.2.1](#), we find

$$\Phi_{\mathbb{R}}(\Delta(\theta, V_\theta)) = \Delta_{Y_\bullet}(\theta, V_\theta) = \Phi_{\mathbb{R}}(P_D),$$

where we applied [Proposition 12.2.1](#) in the second equality. Therefore,

$$\Delta(\theta, V_\theta) = P_D.$$

prop:toricMAandrealMA2

**Proposition 12.2.2** *Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \theta)$ , then*

$$\text{Trop}_*(\theta|_{T(\mathbb{C})} + \text{dd}^c \varphi|_{T(\mathbb{C})})^n = \text{MA}_{\mathbb{R}}(F_\varphi). \quad (12.7) \quad \{\text{eq:toricMAmea2}\}$$

*In particular,*

$$\int_X \theta_\varphi^n = \int_{N_{\mathbb{R}}} \text{MA}_{\mathbb{R}}(F_\varphi) = n! \text{vol } \Delta(\theta, \varphi) \quad (12.8) \quad \{\text{eq:toricmass2}\}$$

*and*

$$\int_X \theta_{V_\theta}^n = n! \text{vol } P. \quad (12.9) \quad \{\text{eq:toricminsingmass}\}$$

**Proof** Take  $F_0$  as in (5.4) and  $\omega$  denotes the corresponding Kähler form.

Then for any large enough  $C > 0$ ,  $\theta + C\omega$  is a Kähler form. So we conclude from [Proposition 5.2.5](#) that

$$\text{Trop}_*((\theta + C\omega)|_{T(\mathbb{C})} + \text{dd}^c \varphi|_{T(\mathbb{C})})^n = \text{MA}_{\mathbb{R}}(F_\varphi + CF_0).$$

Since both sides are polynomials in  $C$ , we conclude that the same holds for  $C = 0$ . Therefore, (12.7) follows.

(12.8) is a direct consequence, while (12.9) follows from [Theorem 12.2.2](#).  $\square$

## 12.3 The pluripotential theory

thm:toricpshbig

**Theorem 12.3.1** *There is a canonical bijection between the following sets:*

- (1) *the set of  $\varphi \in \text{PSH}_{\text{tor}}(X, \theta)$ ;*
- (2) *the set of  $F \in \mathcal{P}(N_{\mathbb{R}}, P_D)$  satisfying  $F \leq F_{V_\theta}$ , and*
- (3) *the set of closed proper convex functions  $G \in \text{Conv}(M_{\mathbb{R}})$  satisfying*

$$G \geq F_{V_\theta}^*.$$

As before, we write  $F_\varphi, G_\varphi$  for the functions determined by this construction.

**Proof** The proof is similar to that of [Theorem 5.2.1](#), but due to its importance, we give the proof. Again, the correspondence between (2) and (3) is proved in [Proposition A.2.4](#).

Given  $\varphi$ , we can construct  $F_\varphi$  in (2) as explained earlier. Conversely, given  $F \in \mathcal{P}(N_{\mathbb{R}}, P_D)$  such that  $F \leq F_{V_\theta}$ . Then



$$\text{Trop}^*(F - F_\theta) \in \text{PSH}(T(\mathbb{C}), \theta|_{T(\mathbb{C})}).$$

Since  $F \leq F_{V_\theta}$ , we see that  $\text{Trop}^*(F - F_\theta)$  is bounded from above. It follows that Grauert–Remmert’s extension theorem [Theorem 1.2.1](#) is applicable, and this function extends to a unique  $\theta$ -psh function  $\varphi$ . The uniqueness of the extension guarantees that  $\varphi \in \text{PSH}_{\text{tor}}(X, \theta)$ .

The two maps are clearly inverse to each other.  $\square$

We fix a model potential  $\phi \in \text{PSH}_{\text{tor}}(X, \theta)_{>0}$  with Newton body  $\Delta(\theta, \phi)$ .

A similar argument guarantees the following:

**Corollary 12.3.1** *There is a canonical bijection between the following sets:*

- (1) the set of  $\varphi \in \text{PSH}_{\text{tor}}(X, \theta; \phi)$ ,
- (2) the set of  $F \in \mathcal{P}(N_{\mathbb{R}}, \Delta(\theta, \phi))$  satisfying  $F \leq F_{V_\theta}$ , and
- (3) the set of closed proper convex functions  $G \in \text{Conv}(M_{\mathbb{R}})$  satisfying

$$G \geq F_{V_\theta}^*, \quad G|_{M_{\mathbb{R}} \setminus \Delta(\theta, \phi)} = \infty.$$

Moreover, under these correspondences, we have the following bijections:

- (1) the set  $\mathcal{E}_{\text{tor}}(X, \theta; \phi)$ ,
- (2) the set of  $F \in \mathcal{E}(N_{\mathbb{R}}, \Delta(\theta, \phi))$  satisfying  $F \leq F_{V_\theta}$ , and
- (3) the set of closed proper convex functions  $G \in \text{Conv}(M_{\mathbb{R}})$  satisfying

$$G \geq F_{V_\theta}^*, \quad G|_{\text{Int } P} < \infty.$$

Here the notation  $\mathcal{E}_{\text{tor}}(X, \theta; \phi)$  means  $\mathcal{E}(X, \theta; \phi) \cap \text{PSH}_{\text{tor}}(X, \theta)$ .

With an almost identical argument, we arrive at

prop:toricsubgeod

**Proposition 12.3.1** *Let  $\varphi_0, \varphi_1 \in \text{PSH}_{\text{tor}}(X, \theta)$ . There is a canonical bijection between the following sets:*

- (1) the set of  $T_c$ -invariant subgeodesics from  $\varphi_0$  to  $\varphi_1$ ,
- (2) the set of convex functions  $F: N_{\mathbb{R}} \times (0, 1) \rightarrow \mathbb{R}$  such that for each  $r \in (0, 1)$ , the function

$$F_r: N_{\mathbb{R}} \rightarrow \mathbb{R}, \quad n \mapsto F(n, r)$$

satisfies  $F_r \rightarrow F_{\varphi_1}$  (resp.  $F_r \rightarrow F_{\varphi_0}$ ) everywhere as  $r \rightarrow 1-$  (resp.  $r \rightarrow 0+$ ), and

- (3) the set of convex functions  $\Psi$  on  $M_{\mathbb{R}} \times \mathbb{R}$  such that

$$\Psi(m, s) \geq G_{\varphi_0}(m) \vee (G_{\varphi_1}(m) + s).$$

Note that  $\Psi$  in (3) is nothing but the Legendre transform of  $F$ .

As an immediate corollary,

cor:toricgeodgeneral

**Corollary 12.3.2** *Let  $\varphi_0, \varphi_1 \in \mathcal{E}_{\text{tor}}(X, \theta)$ . Then the geodesic  $(\varphi_t)_{t \in (0, 1)}$  from  $\varphi_0$  to  $\varphi_1$  corresponds to the lower convex envelope [Definition A.1.4](#) of the function*

$$N_{\mathbb{R}} \times [0, 1] \rightarrow \mathbb{R}, \quad (n, t) \mapsto tF_{\varphi_1}(n) + (1 - t)F_{\varphi_0}(n).$$

Moreover, we have

$$G_{\varphi_t} = (1-t)G_{\varphi_1} + tG_{\varphi_0}. \quad (12.10)$$

{eq:Glinear}

**Proof** The first assertion follows immediately from [Proposition 12.3.1](#). It remains to argue [\(12.10\)](#).

Let  $F: N_{\mathbb{R}} \times [0, 1] \rightarrow \mathbb{R}$  be the map  $(n, t) \mapsto F_{\varphi_t}(n)$ .

It follows from the correspondence in [Proposition 12.3.1](#) that the Legendre transform of  $F$  is given by  $G_{\varphi_0} \vee (G_{\varphi_1} + s)$ . From this we conclude that

$$G_{\varphi_t}(m) = -\sup_{s \in \mathbb{R}} (st - G_{\varphi_0}(m) \vee (G_{\varphi_1}(m) + s)) = (1-t)G_{\varphi_1}(m) + tG_{\varphi_0}(m).$$

The proofs of the following results are similar to the ample case studied in [Chapter 5](#). We omit the details.

prop:toricpluscstbig

**Proposition 12.3.2** *Given  $\varphi \in \text{PSH}_{\text{tor}}(X, \theta)$  and  $C \in \mathbb{R}$ . We have*

$$F_{\varphi+C} = F_{\varphi} + C, \quad G_{\varphi+C} = G_{\varphi} - C.$$

prop:toricrooftopbig

**Proposition 12.3.3** *Given  $\varphi, \psi \in \text{PSH}_{\text{tor}}(X, \theta)$ , then  $\varphi \wedge \psi \in \text{PSH}_{\text{tor}}(X, \theta)$  and*

$$F_{\varphi \wedge \psi} = F_{\varphi} \wedge F_{\psi}, \quad G_{\varphi \wedge \psi} = G_{\varphi} \vee G_{\psi}.$$

prop:toricseqbig

**Proposition 12.3.4** *Let  $\{\varphi_i\}_{i \in I}$  be a family in  $\text{PSH}_{\text{tor}}(X, \theta)$  uniformly bounded from above. Then  $\sup_{i \in I} \varphi_i \in \text{PSH}_{\text{tor}}(X, \theta)$  and*

$$F_{\sup_{i \in I} \varphi_i} = \sup_{i \in I} F_{\varphi_i}, \quad G_{\sup_{i \in I} \varphi_i} = \text{cl} \bigwedge_{i \in I} G_{\varphi_i}.$$

Moreover, if  $I$  is finite, then

$$G_{\max_{i \in I} \varphi_i} = \bigwedge_{i \in I} G_{\varphi_i}.$$

Similarly, if  $\{\varphi_i\}_{i \in I}$  is a decreasing net in  $\text{PSH}_{\text{tor}}(X, \theta)$  such that  $\inf_{i \in I} \varphi_i \not\equiv -\infty$ , then  $\inf_{i \in I} \varphi_i \in \text{PSH}_{\text{tor}}(X, \theta)$  and

$$F_{\inf_{i \in I} \varphi_i} = \inf_{i \in I} F_{\varphi_i}, \quad G_{\inf_{i \in I} \varphi_i} = \sup_{i \in I} G_{\varphi_i}.$$

prop:GPenvelopebig

**Proposition 12.3.5** *Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \theta)$ . Then  $P_{\theta}[\varphi] \in \text{PSH}_{\text{tor}}(X, \theta)$  and*

$$G_{P_{\theta}[\varphi]}(x) = \begin{cases} G_{V_{\theta}}(x), & \text{if } x \in \overline{\{G_{\varphi}(x) < \infty\}}; \\ \infty, & \text{otherwise.} \end{cases} \quad (12.11)$$

{eq:toricPenvbig}

As a consequence, we have

**Corollary 12.3.3** *Let  $\varphi, \psi \in \text{PSH}_{\text{tor}}(X, \theta)_{>0}$ . Then the following are equivalent:*

- (1)  $\varphi \sim_P \psi$ ;

$$(2) \Delta(\theta, \varphi) = \Delta(\theta, \psi).$$

Next we consider the trace operator. For this purpose, we will need to fix a  $T$ -invariant subvariety  $Y \subseteq X$ . Since  $X$  is smooth, so is  $Y$ . Let  $\sigma$  be the cone in  $\Sigma$  corresponding to  $Y$  and  $Q$  be the face of  $P$  corresponding to  $Y$ .

prop:traceoptic

**Proposition 12.3.6** *Let  $\varphi \in \text{PSH}_{\text{tor}}(X, \theta)_{>0}$ . Consider a  $T$ -invariant subvariety  $Y$  corresponding to a face  $Q$  of  $P$ . Suppose that  $\nu(\varphi, Y) = 0$  and  $\text{vol}(\theta|_Y, \text{Tr}_Y^\theta(\varphi)) > 0$ . Then*

$$\Delta(\theta|_Y, \text{Tr}_Y^\theta(\varphi)) = (i_\sigma + m_\sigma)_\mathbb{R}^* (\Delta(\theta, \varphi) \cap Q). \quad (12.12)$$

{eq:traceticNewton}

In particular,  $\text{Tr}_Y(\varphi) \sim_P \varphi|_Y$  if  $\varphi|_Y \not\equiv -\infty$ .

Observe that the condition  $\nu(\varphi, Y) = 0$  means exactly that  $\Delta(\theta, \varphi) \cap Q \neq \emptyset$  by [Corollary 12.2.2](#).

**Proof** Perturbing  $\theta$  slightly, we may assume that  $\theta_\varphi$  is a Kähler current. Let  $(\varphi_j)_j$  be a quasi-equisingular approximation of  $\varphi$  in  $\text{PSH}_{\text{tor}}(X, \theta)$ . It follows from the continuity of the partial Okounkov bodies [Theorem 10.2.2](#) and the continuity of the trace operator [Proposition 8.2.2](#) that it suffices to handle the case where  $\varphi$  has analytic singularities. We need to show that

$$\Delta(\theta|_Y, \varphi|_Y) = (i_\sigma + m_\sigma)_\mathbb{R}^* (\Delta(\theta, \varphi) \cap Q).$$

It is enough to observe that

$$G_{\varphi|_Y} = (i_\sigma + m_\sigma)_\mathbb{R}^* G_\varphi|_Q.$$

The argument is contained in [\[BGPS14, Proof of Proposition 4.8.9\]](#).

Finally, observe that if  $\varphi|_Y \not\equiv -\infty$ , the right-hand side of [\(12.12\)](#) is nothing but  $\Delta(\theta|_Y, \varphi|_Y)$  using [\[BGPS14, Proof of Proposition 4.8.9\]](#). So we conclude that  $\varphi|_Y \sim_P \text{Tr}_Y(\varphi)$ .  $\square$



## Chapter 13

# Non-Archimedean pluripotential theory

chap:NAapp

In this chapter, we will establish the non-Archimedean pluripotential theory using the theory of  $\mathcal{I}$ -good singularities.

We also construct the Duistermaat–Heckman measure of a non-Archimedean metric in [Section 13.3](#).

### 13.1 The definition of non-Archimedean metrics

Let  $X$  be a connected compact Kähler manifold of dimension  $n$ . Let  $\text{Käh}(X)$  be the set of Kähler forms on  $X$  with the partial order given as follows: we say  $\omega \leq \omega'$  if  $\omega \geq \omega'$ . Note that the ordered set  $\text{Käh}(X)$  is a directed set.

Let  $\theta$  be a closed smooth real  $(1, 1)$ -form.

**Definition 13.1.1** We define

$$\text{PSH}^{\text{NA}}(X, \theta) = \varprojlim_{\omega \in \text{Käh}(X)} \text{PSH}^{\text{NA}}(X, \theta + \omega)_{>0}$$

in the category of sets, where the transition maps are given as follows: suppose that  $\omega, \omega' \in \text{Käh}$  and  $\omega \geq \omega'$ , then the transition map is defined in [Proposition 9.3.4](#):

$$P_{\theta+\omega'}[\bullet]_{\mathcal{I}}: \text{PSH}^{\text{NA}}(X, \theta + \omega')_{>0} \rightarrow \text{PSH}^{\text{NA}}(X, \theta + \omega)_{>0}. \quad (13.1)$$

{eq:PItransPSHNApositive}

In general, we denote the components of  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$  in  $\text{PSH}^{\text{NA}}(X, \theta + \omega)$  by  $P_{\theta+\omega}[\Gamma]_{\mathcal{I}}$ .

*Remark 13.1.1* Thanks to [Proposition 9.3.2](#), for any other  $\theta'$  representing  $[\theta]$ , we have a canonical bijection

$$\text{PSH}^{\text{NA}}(X, \theta) \xrightarrow{\sim} \text{PSH}^{\text{NA}}(X, \theta').$$

Moreover, these bijections satisfy the cocycle condition. If we view the set of closed real smooth  $(1, 1)$ -forms representing  $[\theta]$  as a category with a unique morphism between any two objects, then we can define

$$\mathrm{PSH}^{\mathrm{NA}}(X, [\theta]) = \varprojlim_{\theta} \mathrm{PSH}^{\mathrm{NA}}(X, \theta).$$

This definition is independent of the choice of the explicit representative of the cohomology class  $[\theta]$ .

However, given the fact that our notations are already quite heavy, we decide to stick to the set  $\mathrm{PSH}^{\mathrm{NA}}(X, \theta)$ . The readers should verify that all constructions below are independent of the choice of  $\theta$  within its cohomology class.

prop:testcminftyPrela

**Proposition 13.1.1** *Let  $\Gamma \in \mathrm{PSH}^{\mathrm{NA}}(X, \theta)$ . Then given  $\omega, \omega' \in \mathrm{K\ddot{a}h}(X)$  with  $\omega \leq \omega'$ , we have*

$$P_{\theta+\omega} [P_{\theta+\omega'} [\Gamma]_{I, -\infty}] = P_{\theta+\omega} [\Gamma]_{I, -\infty}.$$

**Proof** Since  $P_{\theta+\omega'} [\Gamma]_{I, -\infty}$  is  $I$ -good by [Example 7.1.2](#), it follows that

$$P_{\theta+\omega} [P_{\theta+\omega'} [\Gamma]_{I, -\infty}] = P_{\theta+\omega} [P_{\theta+\omega'} [\Gamma]_{I, -\infty}]_I.$$

Our assertion follows from [Proposition 3.2.12](#).  $\square$

prop:NAposNAemb

**Proposition 13.1.2** *There is a natural injective map*

$$\mathrm{PSH}^{\mathrm{NA}}(X, \theta)_{>0} \hookrightarrow \mathrm{PSH}^{\mathrm{NA}}(X, \theta), \quad \Gamma \mapsto (P_{\theta+\omega} [\Gamma]_I)_{\omega \in \mathrm{K\ddot{a}h}(X)}.$$

In the sequel, we will not distinguish an element in  $\mathrm{PSH}^{\mathrm{NA}}(X, \theta)_{>0}$  with its image in  $\mathrm{PSH}^{\mathrm{NA}}(X, \theta)$ .

**Proof** It is obvious that this map is well-defined. It suffices to argue its injectivity. Suppose that  $\Gamma, \Gamma' \in \mathrm{PSH}^{\mathrm{NA}}(X, \theta)_{>0}$  and

$$P_{\theta+\omega} [\Gamma]_I = P_{\theta+\omega} [\Gamma']_I$$

for some Kähler form  $\omega$  on  $X$ . Then for any  $\tau < \Gamma_{\max}$ , we have

$$\Gamma_{\tau} \sim_I \Gamma'_{\tau}$$

by [Proposition 6.1.3](#). It follows again from [Proposition 6.1.3](#) that

$$\Gamma_{\tau} = \Gamma'_{\tau}.$$

**Definition 13.1.2** Let  $\Gamma \in \mathrm{PSH}^{\mathrm{NA}}(X, \theta)$ . We define  $\Gamma_{\max}$  as  $P_{\theta+\omega} [\Gamma]_{I, \max}$  for any Kähler form  $\omega$  on  $X$ .

Note that under the identification of [Proposition 13.1.2](#), for any  $\Gamma \in \mathrm{PSH}^{\mathrm{NA}}(X, \theta)_{>0}$ , this definition is compatible with the notion of  $\Gamma_{\max}$  in [Definition 9.1.1](#).

**Definition 13.1.3** Let  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ , we define its *volume* as follows:

$$\text{vol } \Gamma := \lim_{\omega \in \text{K\"ah}(X)} \int_X (\theta + \omega + \text{dd}^c P_{\theta+\omega'}[\Gamma]_{I, -\infty})^n \in [0, \infty).$$

Observe that the net is decreasing, so the limit exists.

**Proposition 13.1.3** Let  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ . Then

$$\text{vol } \Gamma = \int_X (\theta + \text{dd}^c \Gamma_{-\infty})^n.$$

*Proof* This follows from [Proposition 3.1.8](#), [Corollary 3.1.3](#) and [Proposition 13.1.1](#).  $\square$

def:PSHNAtarangeneral

**Definition 13.1.4** Let  $\omega$  be a closed real smooth positive  $(1, 1)$ -form on  $X$ . We define the map

$$P_{\theta+\omega}[\bullet]_I : \text{PSH}^{\text{NA}}(X, \theta) \rightarrow \text{PSH}^{\text{NA}}(X, \theta + \omega)$$

as follows: given  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ , we define  $P_{\theta+\omega}[\Gamma]_I$  as the element such that for any  $\omega' \in \text{K\"ah}(X)$ , we have

$$P_{\theta+\omega+\omega'}[P_{\theta+\omega}[\Gamma]_I]_I = P_{\theta+\omega+\omega'}[\Gamma]_I.$$

It is straightforward to check that under the identification of [Proposition 13.1.2](#), the map  $P_{\theta+\omega}[\bullet]_I$  extends the map [\(13.1\)](#).

**Proposition 13.1.4** The maps  $P_{\theta+\omega}[\bullet]_I$  in [Definition 13.1.4](#) together induce a bijection

$$\text{PSH}^{\text{NA}}(X, \theta) \xrightarrow{\sim} \varprojlim_{\omega \in \text{K\"ah}(X)} \text{PSH}^{\text{NA}}(X, \theta + \omega). \quad (13.2)$$

{eq:PSHNAprojlimigeneral2}

*Proof* It is a tautology that the maps  $P_{\theta+\omega}[\bullet]_I$  in [Definition 13.1.4](#) are compatible with the transition maps. So the map [\(13.2\)](#) is well-defined. It is injective by the same argument as [Proposition 13.1.2](#). We argue the surjectivity.

By unfolding the definitions, an object in the target of [\(13.2\)](#) is an assignment: with each  $\omega \in \text{K\"ah}(X)$ , we associate a family  $(\Gamma^{\omega, \omega'})_{\omega' \in \text{K\"ah}(X)}$  satisfying:

- (1)  $\Gamma^{\omega, \omega'} \in \text{PSH}^{\text{NA}}(X, \theta + \omega + \omega')_{>0}$  for each  $\omega, \omega' \in \text{K\"ah}(X)$ ;
- (2) for each  $\omega, \omega', \omega'' \in \text{K\"ah}(X)$  satisfying  $\omega'' \geq \omega'$ , we have

$$P_{\theta+\omega+\omega''}[\Gamma^{\omega, \omega'}]_I = \Gamma^{\omega, \omega''};$$

- (3) for each  $\omega, \omega', \omega'' \in \text{K\"ah}(X)$  satisfying  $\omega \leq \omega'$ , we have

$$P_{\theta+\omega'+\omega''}[\Gamma^{\omega, \omega''}]_I = \Gamma^{\omega', \omega''}.$$

The preimage of such an object is given by the family  $(\Gamma^{\omega})_{\omega \in \text{K\"ah}(X)}$  given by

$$\Gamma^\omega = \Gamma^{\omega/2, \omega/2}.$$

The fact that the image of  $\Gamma$  is as expected is a tautology, which we leave to the readers.  $\square$

With an almost identical argument involving [Proposition 3.1.8](#), we get

prop:PSHNAreform1

**Proposition 13.1.5** *The maps  $P_{\theta+\omega}[\bullet]_I$  in [Definition 13.1.4](#) and the injective maps [Proposition 13.1.2](#) together induce bijections*

$$\text{PSH}^{\text{NA}}(X, \theta) \xrightarrow{\sim} \varprojlim_{\omega} \text{PSH}^{\text{NA}}(X, \theta + \omega)_{>0} \xrightarrow{\sim} \varprojlim_{\omega} \text{PSH}^{\text{NA}}(X, \theta + \omega), \quad (13.3)$$

{eq:PSHNAprojlimigeneral}

where  $\omega$  runs over either the partially ordered set of all smooth closed real positive  $(1, 1)$ -forms with positive volume on  $X$  or  $\text{K\"ah}(X)$ .

cor:PSHNAbimero

**Corollary 13.1.1** *Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism from a compact K\"ahler manifold  $Y$ . Then  $\pi^*$  induces a bijection*

$$\text{PSH}^{\text{NA}}(X, \theta) \xrightarrow{\sim} \text{PSH}^{\text{NA}}(Y, \pi^*\theta).$$

**Proof** This follows immediately from [Proposition 13.1.5](#).  $\square$

It is immediate to verify that  $\pi^*$  in [Corollary 13.1.1](#) extends the map [Proposition 9.3.3](#).

## 13.2 Operations on non-Archimedean metrics

Let  $X$  be a connected compact K\"ahler manifold of dimension  $n$  and  $\theta, \theta', \theta''$  be closed real smooth  $(1, 1)$ -forms on  $X$  representing big cohomology classes.

**Definition 13.2.1** Let  $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)$ . We say  $\Gamma \leq \Gamma'$  if  $\Gamma_{\max} \leq \Gamma'_{\max}$  and for some  $\omega \in \text{K\"ah}(X)$ , we have

$$P_{\theta+\omega}[\Gamma]_I \geq P_{\theta+\omega}[\Gamma']_I.$$

This notion is independent of the choice of  $\omega$  thanks to [\(9.13\)](#).

Moreover, we have the following:

**Proposition 13.2.1** *Let  $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)$  and  $\omega$  be a closed smooth positive  $(1, 1)$ -form on  $X$ , then the following are equivalent:*

- (1)  $\Gamma \leq \Gamma'$ ;
- (2)  $P_{\theta+\omega}[\Gamma]_I \leq P_{\theta+\omega}[\Gamma']_I$ .

**Proof** This follows immediately from [\(9.13\)](#).  $\square$

Observe that this definition coincides with the corresponding definition in [Definition 9.4.1](#) when  $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)_{>0}$ .



def:sumNAmetrics

**Definition 13.2.2** Let  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$  and  $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')$ . Then we define  $\Gamma + \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta + \theta')$  as the unique element such that for any  $\omega \in \text{K\"ah}(X)$ , we have

$$P_{\theta+\omega}[\Gamma + \Gamma']_I = P_{\theta+\omega}[\Gamma]_I + P_{\theta+\omega}[\Gamma']_I.$$

This definition yields an element in  $\text{PSH}^{\text{NA}}(X, \theta + \theta')$  by [Lemma 9.4.3](#).

**Proposition 13.2.2** Let  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$  and  $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')$ . Suppose that  $\omega, \omega'$  are two smooth closed positive  $(1, 1)$ -forms on  $X$ . Then

$$P_{\theta+\omega+\theta'+\omega'}[\Gamma + \Gamma']_I = P_{\theta+\omega}[\Gamma]_I + P_{\theta'+\omega'}[\Gamma']_I.$$

*Proof* This is a direct consequence of [Lemma 9.4.3](#).  $\square$

**Proposition 13.2.3** The operation  $+$  is commutative and associative: for any  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ ,  $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')$  and  $\Gamma'' \in \text{PSH}^{\text{NA}}(X, \theta'')$ , we have

$$\Gamma + \Gamma' = \Gamma' + \Gamma, \quad (\Gamma + \Gamma') + \Gamma'' = \Gamma + (\Gamma' + \Gamma'').$$

*Proof* This is a direct consequence of [Proposition 9.4.1](#).  $\square$

**Definition 13.2.3** Let  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$  and  $C \in \mathbb{R}$ . We define  $\Gamma + C \in \text{PSH}^{\text{NA}}(X, \theta)$  as the unique element such that for any  $\omega \in \text{K\"ah}(X)$ , we have

$$P_{\theta+\omega}[\Gamma + C] = P_{\theta+\omega}[\Gamma] + C.$$

It is obvious from [Definition 9.4.3](#) that  $\Gamma + C \in \text{PSH}^{\text{NA}}(X, \theta)$ . It is also obvious that this definition extends [Definition 9.4.3](#).

**Proposition 13.2.4** Let  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$  and  $C \in \mathbb{R}$ . Suppose that  $\omega$  is a smooth closed positive  $(1, 1)$ -form on  $X$ . Then

$$P_{\theta+\omega}[\Gamma]_I + C = P_{\theta+\omega}[\Gamma + C]_I.$$

*Proof* This is clear by definition.  $\square$

prop:NAmetricplusC

**Proposition 13.2.5** Let  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ ,  $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')$  and  $C, C' \in \mathbb{R}$ , then

- (1)  $(\Gamma + \Gamma') + C = \Gamma + (\Gamma' + C) = (\Gamma + C) + \Gamma'$ ;
- (2)  $\Gamma + (C + C') = (\Gamma + C) + C'$ .

*Proof* This is a direct consequence of [Proposition 9.4.2](#).  $\square$

def:PSHNAlor

**Definition 13.2.4** Let  $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)$ , we define  $\Gamma \vee \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)$  as the unique element such that for any  $\omega \in \text{K\"ah}(X)$ , we have

$$P_{\theta+\omega}[\Gamma \vee \Gamma']_I = P_{\theta+\omega}[\Gamma]_I \vee P_{\theta+\omega}[\Gamma']_I.$$

It follows from [Lemma 9.4.5](#) that  $\Gamma \vee \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)$  and this definition extends the corresponding definition in [Definition 9.4.4](#).

**Proposition 13.2.6** *Let  $\Gamma, \Gamma' \in \text{PSH}^{\text{NA}}(X, \theta)$  and  $\omega$  be a closed smooth positive  $(1, 1)$ -form on  $X$ . Then*

$$P_{\theta+\omega}[\Gamma \vee \Gamma']_I = P_{\theta+\omega}[\Gamma]_I \vee P_{\theta+\omega}[\Gamma']_I.$$

**Proof** This is a direct consequence of [Lemma 9.4.5](#).  $\square$

**Proposition 13.2.7** *The operation  $\vee$  is commutative and associative.*

In particular, given a finite non-empty family  $(\Gamma^i)_{i \in I}$  in  $\text{PSH}^{\text{NA}}(X, \theta)$ , we then define  $\bigvee_{i \in I} \Gamma^i$  in the obvious way.

**Proof** This is a direct consequence of [Corollary 9.4.1](#).  $\square$

**Definition 13.2.5** Let  $(\Gamma^i)_{i \in I}$  be a non-empty family in  $\text{PSH}^{\text{NA}}(X, \theta)$ . Assume that

$$\sup_{i \in I} \Gamma^i_{\max} < \infty. \quad (13.4)$$

`{eq: supPSHNAmaxfinite}`

Then we define  $\sup_{i \in I}^* \Gamma^i \in \text{PSH}^{\text{NA}}(X, \theta)$  as the unique element such that for any  $\omega \in \text{K\"ah}(X)$ , we have

$$P_{\theta+\omega} \left[ \sup_{i \in I} \Gamma^i \right] = \sup_{i \in I}^* P_{\theta+\omega} [\Gamma^i].$$

It follows immediately from [Lemma 9.4.7](#) that  $\sup_{i \in I}^* \Gamma^i \in \text{PSH}^{\text{NA}}(X, \theta)$  and this definition extends [Definition 9.4.6](#). Moreover, this definition clearly extends [Definition 13.2.4](#) as well.

**Proposition 13.2.8** *Let  $(\Gamma^i)_{i \in I}$  be a non-empty in  $\text{PSH}^{\text{NA}}(X, \theta)$  satisfying (13.4). Assume that  $\omega$  is a closed smooth positive  $(1, 1)$ -form on  $X$ . Then*

$$P_{\theta+\omega} \left[ \sup_{i \in I}^* \Gamma^i \right] = \sup_{i \in I}^* P_{\theta+\omega} [\Gamma^i].$$

**Proof** This is a direct consequence of [Lemma 9.4.7](#).  $\square$

`prop: NAChoquet`

**Proposition 13.2.9** *Let  $(\Gamma^i)_{i \in I}$  be a non-empty in  $\text{PSH}^{\text{NA}}(X, \theta)$  satisfying (13.4). Then there exists a countable subfamily  $I' \subseteq I$  such that*

$$\sup_{i \in I}^* \Gamma^i = \sup_{i \in I'}^* \Gamma^i.$$

**Proof** For any fixed  $\omega \in \text{K\"ah}(X)$ , thanks to [Proposition 9.4.5](#), we could find a countable subfamily  $I' \subseteq I$  such that

$$\sup_{i \in I}^* P_{\theta+\omega} [\Gamma^i]_I = \sup_{i \in I'}^* P_{\theta+\omega} [\Gamma^i]_I.$$

It suffices to show that for any other  $\omega' \in \text{K\"ah}(X)$ , we have

$$\sup_{i \in I}^* P_{\theta+\omega'}[\Gamma^i]_I = \sup_{i \in I'}^* P_{\theta+\omega'}[\Gamma^i]_I.$$

This is an immediate consequence of [Proposition 6.1.6](#).  $\square$

prop:supGammiotherprop2

**Proposition 13.2.10** *Let  $(\Gamma^i)_{i \in I}$  be a non-empty family in  $\text{PSH}^{\text{NA}}(X, \theta)$  satisfying (13.4). Let  $C \in \mathbb{R}$ . Then*

$$\sup_{i \in I}^*(\Gamma^i + C) = \sup_{i \in I}^* \Gamma^i + C.$$

*Suppose that  $(\Gamma'^i)_{i \in I}$  is another family in  $\text{PSH}^{\text{NA}}(X, \theta)$  satisfying (13.4). Suppose that  $\Gamma^i \leq \Gamma'^i$  for all  $i \in I$ , then*

$$\sup_{i \in I}^* \Gamma^i \leq \sup_{i \in I}^* \Gamma'^i.$$

**Proof** This is an immediate consequence of [Proposition 9.4.6](#).  $\square$

**Definition 13.2.6** Let  $(\Gamma_i)_{i \in I}$  be a decreasing net in  $\text{PSH}^{\text{NA}}(X, \theta)$ . Assume that

$$\inf_{i \in I} \Gamma_{i, \max} > -\infty, \quad (13.5)$$

{eq:decretcontition}

then we define  $\inf_{i \in I} \Gamma_i \in \text{PSH}^{\text{NA}}(X, \theta)$  as the unique element such that for each  $\omega \in \text{K\"ah}(X)$ , the component

$$P_{\theta+\omega} \left[ \inf_{i \in I} \Gamma_i \right]_I \in \text{PSH}^{\text{NA}}(X, \theta + \omega)_{>0}$$

is defined as follows:

(1) We set

$$\left( P_{\theta+\omega} \left[ \inf_{i \in I} \Gamma_i \right]_I \right)_{\max} = \inf_{i \in I} \Gamma_{i, \max};$$

(2) for any  $\tau < \inf_{i \in I} \Gamma_{i, \max}$ , we define

$$\left( P_{\theta+\omega} \left[ \inf_{i \in I} \Gamma_i \right]_I \right)_{\tau} = \inf_{i \in I} P_{\theta+\omega}[\Gamma_i, \tau]_I. \quad (13.6)$$

{eq:decrettestcurdef}

We observe that

$$P_{\theta+\omega} \left[ \inf_{i \in I} \Gamma_i \right]_I \in \text{PSH}^{\text{NA}}(X, \theta + \omega)_{>0}.$$

This follows from [Proposition 3.2.11](#). Now it is clear that  $\inf_{i \in I} \Gamma_i \in \text{PSH}^{\text{NA}}(X, \theta)$ .

prop:infGammiotherprop2

**Proposition 13.2.11** *Let  $(\Gamma^i)_{i \in I}$  be a decreasing net in  $\text{PSH}^{\text{NA}}(X, \theta)$  satisfying (13.5). Let  $C \in \mathbb{R}$ . Then*

$$\inf_{i \in I}(\Gamma^i + C) = \inf_{i \in I} \Gamma^i + C.$$

Suppose that  $(\Gamma^i)_{i \in I}$  is another decreasing net in  $\text{PSH}^{\text{NA}}(X, \theta)$  satisfying (13.5). Suppose that  $\Gamma^i \leq \Gamma'^i$  for all  $i \in I$ , then

$$\inf_{i \in I} \Gamma^i \leq \inf_{i \in I} \Gamma'^i.$$

**Proof** This is clear by definition.  $\square$

**Definition 13.2.7** Let  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$  and  $\lambda \in \mathbb{R}_{>0}$ , then we define  $\lambda\Gamma \in \text{PSH}^{\text{NA}}(X, \lambda\theta)$  as the unique element such that for any  $\omega \in \text{K\"ah}(X)$ , we have

$$P_{\lambda\theta+\omega}[\lambda\Gamma]_I = \lambda P_{\theta+\lambda^{-1}\omega}[\Gamma]_I.$$

It follows immediately from Lemma 9.4.8 that  $\lambda\Gamma \in \text{PSH}^{\text{NA}}(X, \lambda\theta)$  and this definition extends Definition 9.4.7.

**Proposition 13.2.12** Let  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$  and  $\lambda \in \mathbb{R}_{>0}$ . Then for any closed smooth positive  $(1, 1)$ -form  $\omega$  on  $X$ , we have

$$P_{\lambda\theta+\omega}[\lambda\Gamma]_I = \lambda P_{\theta+\lambda^{-1}\omega}[\Gamma]_I.$$

**Proof** This follows immediately from Lemma 9.4.8.  $\square$

prop:resclacomp2

**Proposition 13.2.13** Let  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ ,  $\Gamma' \in \text{PSH}^{\text{NA}}(X, \theta')$ ,  $C \in \mathbb{R}$  and  $\lambda, \lambda' > 0$ , we have

$$\begin{aligned} \lambda(\Gamma + \Gamma') &= \lambda\Gamma + \lambda\Gamma', \\ (\lambda\lambda')\Gamma &= \lambda(\lambda'\Gamma), \\ \lambda(\Gamma + C) &= \lambda\Gamma + \lambda C. \end{aligned}$$

Suppose that  $(\Gamma^i)_{i \in I}$  is a non-empty family in  $\text{PSH}^{\text{NA}}(X, \theta)$  satisfying (13.4), then

$$\lambda \left( \sup_{i \in I}^* \Gamma^i \right) = \sup_{i \in I}^* (\lambda \Gamma^i).$$

If  $(\Gamma^i)_{i \in I}$  is a decreasing net in  $\text{PSH}^{\text{NA}}(X, \theta)$  satisfying (13.5), then

$$\lambda \left( \inf_{i \in I} \Gamma^i \right) = \inf_{i \in I} (\lambda \Gamma^i).$$

**Proof** Everything except the last assertion follows from Proposition 9.4.8. The last assertion is obvious by definition.  $\square$

**Definition 13.2.8** Let  $\Gamma \in \text{PSH}^{\text{NA}}(X, \theta)$ . Let  $Y \subseteq X$  be an irreducible analytic subset. We say that the trace operator of  $\Gamma$  along  $Y$  is *well-defined* if

$$\nu(P_{\theta+\omega''}[\Gamma_\tau]_I, Y) = 0$$

for small enough  $\tau$  and any  $\omega'' \in \text{K\"ah}(X)$ . We define

$$(\text{Tr}_Y(\Gamma))_{\max} := \sup \{ \tau < \Gamma_{\max} : \nu(P_{\theta+\omega''}[\Gamma_\tau]_I, Y) = 0 \}.$$

In this case, we define  $\mathrm{Tr}_Y(\Gamma) \in \mathrm{PSH}^{\mathrm{NA}}(\tilde{Y}, \theta|_{\tilde{Y}})$  as the unique element such that for any  $\omega \in \mathrm{K\ddot{a}h}(\tilde{Y})$ , the component

$$P_{\theta|_{\tilde{Y}}+\omega} [\mathrm{Tr}_Y(\Gamma)]_I \in \mathrm{PSH}^{\mathrm{NA}}(Y, \theta|_{\tilde{Y}} + \omega)_{>0}$$

is defined as follows:

(1) We let

$$\left( P_{\theta|_{\tilde{Y}}+\omega} [\mathrm{Tr}_Y(\Gamma)]_I \right)_{\max} = (\mathrm{Tr}_Y(\Gamma))_{\max}; \quad (13.7) \quad \boxed{\{\text{eq: tracemax}\}}$$

(2) for each  $\tau \in \mathbb{R}$  less than the common value (13.7), we define

$$P_{\theta|_{\tilde{Y}}+\omega} [\mathrm{Tr}_Y(\Gamma)]_{I,\tau} := P_{\theta|_{\tilde{Y}}+\omega} \left[ \mathrm{Tr}_Y^{\theta+\tilde{\omega}} (P_{\theta+\tilde{\omega}}[\Gamma]_{I,\tau}) \right],$$

where  $\tilde{\omega}$  is an arbitrary Kähler form on  $X$  such that  $\omega \geq \tilde{\omega}|_{\tilde{Y}}$ .

It follows from [GK20, Proposition 3.5] that  $\tilde{Y}$  is a normal Kähler space. We observe that the choice of the trace operator  $\mathrm{Tr}_Y^{\theta+\tilde{\omega}} (P_{\theta+\tilde{\omega}}[\Gamma]_{I,\tau})$  is irrelevant since two different choice are  $I$ -equivalent. Moreover,

$$\left( P_{\theta|_{\tilde{Y}}+\omega} [\mathrm{Tr}_Y(\Gamma)]_I \right)_{\tau}$$

is  $I$ -model by Proposition 8.1.2.

Furthermore,

$$P_{\theta|_{\tilde{Y}}+\omega} [\mathrm{Tr}_Y(\Gamma)]_I \in \mathrm{PSH}^{\mathrm{NA}}(Y, \theta|_{\tilde{Y}} + \omega)_{>0}$$

is a consequence of Proposition 8.2.1. It is therefore clear that  $\mathrm{Tr}_Y(\Gamma) \in \mathrm{PSH}^{\mathrm{NA}}(X, \theta)$ .

**Proposition 13.2.14** *Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism from a compact Kähler manifold  $Y$ . Then all definitions in this section are invariant under pulling-back to  $Y$ .*

The meaning is clear in most cases. In the case of the trace operator, this means the following: suppose that  $Z \subseteq X$  is an analytic subset and  $\Gamma \in \mathrm{PSH}^{\mathrm{NA}}(X, \theta)$  has non-trivial restriction to  $Z$ . Suppose that  $Z$  is not contained in the non-isomorphism locus of  $\pi$  so that the strict transform  $W$  of  $Z$  is defined. If we write  $\Pi: W \rightarrow Z$  for the restriction of  $\pi$  and  $\tilde{\Pi}: \tilde{W} \rightarrow \tilde{Z}$  the strict transform of  $\Pi$ , then we have

$$\tilde{\Pi}^* \mathrm{Tr}_Z(\Gamma) = \mathrm{Tr}_W(\pi^* \Gamma).$$

**Proof** We only prove the assertion for the trace operator, as the other proofs are similar.

We shall use the notations above. Observe that for any closed positive smooth  $(1, 1)$ -form on  $X$  with positive mass, we have

$$(\tilde{\Pi}^* \mathrm{Tr}_Z(\Gamma))_{\max} = (\mathrm{Tr}_Z(\Gamma))_{\max} = \sup \{ \tau < \Gamma_{\max} : \nu(P_{\theta+\omega}[\Gamma]_{I,\tau}, Z) = 0 \}$$

and

$$\begin{aligned} (\mathrm{Tr}_W(\pi^*\Gamma))_{\max} &= \sup \{ \tau < \Gamma_{\max} : \nu(P_{\pi^*\theta+\pi^*\omega}[\pi^*\Gamma_\tau]_I, W) = 0 \} \\ &= \sup \{ \tau < \Gamma_{\max} : \nu(\pi^*P_{\theta+\omega}[\Gamma_\tau]_I, W) = 0 \} \\ &= \sup \{ \tau < \Gamma_{\max} : \nu(P_{\theta+\omega}[\Gamma_\tau]_I, Z) = 0 \}. \end{aligned}$$

Here we applied implicitly [Proposition 13.1.5](#). Therefore,

$$(\tilde{\Pi}^* \mathrm{Tr}_Z(\Gamma))_{\max} = (\mathrm{Tr}_W(\pi^*\Gamma))_{\max}.$$

Let  $\tau \in \mathbb{R}$  be less than this common value. Take a closed smooth Kähler form  $\omega$  (resp.  $\omega'$ ) on  $\tilde{Z}$  (resp.  $\tilde{W}$ ) with positive mass. We may assume that  $\omega' \geq \tilde{\Pi}^*\omega$ . Take a Kähler form  $\tilde{\omega}$  on  $Y$  (resp.  $\tilde{\omega}'$  on  $X$ ) such that

$$\omega' \geq \tilde{\omega}'|_{\tilde{W}}, \quad \omega \geq \tilde{\omega}|_{\tilde{Z}}.$$

Without loss of generality, we may assume that

$$\tilde{\omega}' \geq \pi^*\tilde{\omega}.$$

It suffices to show that

$$\mathrm{Tr}_W^{\pi^*\theta+\tilde{\omega}'}(P_{\pi^*\theta+\tilde{\omega}'}[\pi^*\Gamma]_{I,\tau}) \sim_P \tilde{\Pi}^* \mathrm{Tr}_Z^{\theta+\tilde{\omega}}(P_{\theta+\tilde{\omega}}[\Gamma]_{I,\tau}).$$

Using [Proposition 8.2.1](#), this is equivalent to

$$\mathrm{Tr}_W(P_{\pi^*\theta+\pi^*\omega}[\pi^*\Gamma]_{I,\tau}) \sim_P \tilde{\Pi}^* \mathrm{Tr}_Z(P_{\theta+\tilde{\omega}}[\Gamma]_{I,\tau}).$$

This is a consequence of [Lemma 8.2.1](#). □

### 13.3 Duistermaat–Heckman measures

sec:DHmeasure

Let  $X$  be a connected compact Kähler manifold of dimension  $n$  and  $\theta$  be a closed real smooth  $(1, 1)$ -form on  $X$  representing a big cohomology class.

def:DHm

**Definition 13.3.1** Assume that  $X$  admits a smooth flag  $Y_\bullet$ . Let  $\Gamma \in \mathrm{PSH}^{\mathrm{NA}}(X, \theta)_{>0}$ . The *Duistermaat–Heckman measure*  $\mathrm{DH}(\Gamma)$  of an element  $\Gamma \in \mathrm{PSH}^{\mathrm{NA}}(X, \theta)_{>0}$  is defined as

$$\mathrm{DH}(\Gamma) := n! \cdot \mathrm{DH}(\Delta_{Y_\bullet}(\theta, \Gamma)).$$

Recall that  $\Delta_{Y_\bullet}(\theta, \Gamma) \in \mathrm{TC}(\Delta_{Y_\bullet}(\theta, \Gamma_{-\infty}))$  is defined in [Theorem 10.4.2](#). See [Definition 10.4.4](#) for the definition of the Duistermaat–Heckman measure of an Okounkov test curve..

thm:DHindep

**Theorem 13.3.1** The *Duistermaat–Heckman measure*  $\mathrm{DH}(\Gamma)$  of  $\Gamma \in \mathrm{PSH}^{\mathrm{NA}}(X, \theta)_{>0}$  in [Definition 13.3.1](#) is independent of the choice of the smooth flag  $Y_\bullet$ . Furthermore, for any  $m \in \mathbb{Z}_{>0}$ , the  $m$ -th moment of  $\mathrm{DH}(\Gamma)$  is given by

$$\int_{\mathbb{R}} x^m \mathrm{DH}(\Gamma)(x) = \Gamma_{\max}^m \mathrm{vol} \Gamma + m \int_{-\infty}^{\Gamma_{\max}} \tau^{m-1} (\mathrm{vol}(\theta + \mathrm{dd}^c \Gamma_{\tau}) - \mathrm{vol} \Gamma) \, d\tau \quad (13.8)$$

{eq:momentDHmtc1}

if  $m > 0$  and

$$\int_{\mathbb{R}} \mathrm{DH}(\Gamma) = \mathrm{vol} \Gamma. \quad (13.9)$$

{eq:momentDHmtc2}

**Proof** Assume furthermore that  $\Gamma$  is bounded, we observe that the moments of the random variable  $G[\Delta_{Y_{\bullet}}(\theta, \Gamma)]$  as computed in [Proposition 10.4.4](#) are independent of the choice of the flag: In fact, they are given by (13.8) and (13.9) thanks to [Theorem 10.3.2\(1\)](#). Since the Duistermaat–Heckman measure has bounded support in this case (c.f. [Theorem 10.4.1](#)), we conclude that  $\mathrm{DH}(\Gamma)$  is uniquely determined.

In general,  $\Gamma$  is the decreasing limit of the sequence  $\Gamma \vee \Gamma^k$  as  $k \rightarrow \infty$ , where  $\Gamma^k: (-\infty, -k) \rightarrow \mathrm{PSH}(X, \theta)$  takes the constant value  $\Gamma_{-\infty}$ . It follows from the argument of [Theorem 9.2.1](#) that  $\Delta_{Y_{\bullet}}(\Gamma)_{\tau}$  is the decreasing limit of  $\Delta_{Y_{\bullet}}(\Gamma \vee \Gamma^k)_{\tau}$  for any  $\tau < \Gamma_{\max}$ . So  $\mathrm{DH}(\Gamma \vee \Gamma^k) \rightarrow \mathrm{DH}(\Gamma)$  by [Lemma 10.4.2](#). It follows that  $\mathrm{DH}(\Gamma)$  is independent of the choice of the flag.  $\square$

More generally, when  $X$  does not admit a smooth flag, we could make a modification  $\pi: Y \rightarrow X$  so that  $Y$  admits a flag. We define

$$\mathrm{DH}(\Gamma) := \mathrm{DH}(\pi^* \Gamma). \quad (13.10)$$

{eq:DHmgeneral}

It follows from [Theorem 10.3.2\(5\)](#) that this measure is independent of the choice of  $\pi$ .

prop:contDH

**Proposition 13.3.1** *Let  $(\Gamma^i)_{i \in I}$  be a net in  $\mathrm{PSH}^{\mathrm{NA}}(X, \theta)_{>0}$  and  $\Gamma \in \mathrm{PSH}^{\mathrm{NA}}(X, \theta)_{>0}$ . Assume one of the following conditions holds:*

(1) *The net  $(\Gamma^i)_{i \in I}$  is decreasing and  $\Gamma = \inf_{i \in I} \Gamma^i$ . Assume that*

$$\mathrm{vol} \Gamma = \lim_{i \in I} \mathrm{vol} \Gamma^i.$$

(2) *The net  $(\Gamma^i)_{i \in I}$  is increasing and  $\Gamma = \sup_{i \in I}^* \Gamma^i$ .*

Then

$$\mathrm{DH}(\Gamma^i) \rightarrow \mathrm{DH}(\Gamma). \quad (13.11)$$

{eq:contDHm}

**Proof** We may assume that  $X$  admits a smooth flag  $Y_{\bullet}$ .

Assume (1). We want to derive (13.11) from [Proposition 10.4.2](#). It boils down to prove the following: for any  $\tau < \Gamma_{\max}$ , we have

$$\Delta_{Y_{\bullet}}(\theta, \Gamma_{\tau}^i) \xrightarrow{d_{\mathrm{Haus}}} \Delta_{Y_{\bullet}}(\theta, \Gamma_{\tau}).$$

This follows immediately from [Theorem 10.3.2\(1\)](#).

The proof under the assumption (2) is similar. We only need to apply [Proposition 10.4.3](#) instead of [Proposition 10.4.2](#).  $\square$





## Chapter 14

### Partial Bergman kernels

chap:Berg

In this chapter, we prove the convergence of the partial Bergman kernels.

#### 14.1 Partial envelopes

sec:envrel

In this section, let  $X$  be a connected compact Kähler manifold of dimension  $n$  and  $K \subseteq X$  be a closed non-pluripolar set. Let  $\theta$  be a smooth closed real  $(1, 1)$ -form on  $X$  representing a pseudoeffective cohomology class. Fix  $\varphi \in \text{PSH}(X, \theta)$ .

**Definition 14.1.1** Given a function  $v: K \rightarrow [-\infty, \infty)$ , we introduce the *relative  $P$ -envelope* of  $\varphi$  (with respect to  $K, v, \theta$ ) as

$$P_{\theta, K}[\varphi](v) := \sup^* \{ \eta \in \text{PSH}(X, \theta) : \eta|_K \leq v \text{ and } \eta \leq \varphi \}. \quad (14.1)$$

Similarly, we define the *relative  $I$ -envelope* of  $\varphi$  (with respect to  $K, v, \theta$ ) as

$$P_{\theta, K}[\varphi]_I(v) := \sup^* \{ \eta \in \text{PSH}(X, \theta) : \eta|_K \leq v \text{ and } \eta \leq_I \varphi \}. \quad (14.2)$$

Observe that when  $v$  is bounded, the neither envelope is identically  $-\infty$ . When  $K = X$  and  $v = 0$ , these definitions reduce to the usual  $P$ -envelope and  $I$ -envelope of  $\varphi$ .

It would be helpful to consider the following auxiliary functions:

$$\begin{aligned} P'_{\theta, K}[\varphi](v) &:= \sup \{ \eta \in \text{PSH}(X, \theta) : \eta|_K \leq v \text{ and } \eta \leq \varphi \}, \\ P'_{\theta, K}[\varphi]_I(v) &:= \sup \{ \eta \in \text{PSH}(X, \theta) : \eta|_K \leq v \text{ and } \eta \leq_I \varphi \}. \end{aligned}$$

We note the following maximum principles, that follow from the above definitions:

lem: max\_princ

**Lemma 14.1.1** Let  $v \in C^0(K)$ . Let  $\eta \in \text{PSH}(X, \theta)$ . Assume that  $\eta \leq \varphi$ , then

$$\sup_K (\eta - v) = \sup_{\{\eta \neq -\infty\}} (\eta - P'_{\theta,K}[\varphi](v)) = \sup_{\{P'_{\theta,K}[\varphi](v) \neq -\infty\}} (\eta - P'_{\theta,K}[\varphi](v)). \quad (14.3)$$

{eq: max\_princ}

**Proof** We prove the first equality at first. We write  $S = \{\eta = -\infty\}$ .

By definition,  $P'_{\theta,K}[\varphi](v)|_K \leq v$ , so

$$\left( h - P'_{\theta,K}[\varphi](v) \right) \Big|_{K \setminus S} \geq \eta|_{K \setminus S} - v|_{K \setminus S}.$$

This implies that

$$\sup_K (\eta - v) \leq \sup_{X \setminus S} (\eta - P'_{\theta,K}[\varphi](v)).$$

Conversely, observe that  $\sup_K (\eta - v) > -\infty$  as  $K$  is non-pluripolar. Let  $\eta' := \eta - \sup_K (\eta - v)$ , then  $\eta'$  is a candidate in the definition of  $P'_{\theta,K}[\varphi](v)$ , hence  $\eta' \leq P'_{\theta,K}[\varphi](v)$ , namely,

$$\eta - \sup_K (\eta - v) \leq P'_{\theta,K}[\varphi](v),$$

the latter implies that

$$\sup_K (\eta - v) \geq \sup_{X \setminus S} (\eta - P'_{\theta,K}[\varphi](v)),$$

finishing the proof of the first identity.

We have  $\{P'_{\theta,K}[\varphi](v) = -\infty\} \subseteq S$ , and we notice that points in  $S \setminus \{P'_{\theta,K}[\varphi](v) = -\infty\}$  do not contribute to the supremum of  $\eta - P'_{\theta,K}[\varphi](v)$  on  $X \setminus \{P'_{\theta,K}[\varphi](v) = -\infty\}$ , hence the last equality of (14.3) also follows.  $\square$

Next, we make the following observations about the singularity types of our envelopes:

lma:same\_sing\_type

**Lemma 14.1.2** For any  $v \in C^0(K)$  we have

$$P_{\theta,K}[\varphi](v) \sim P_{\theta}[\varphi], \quad P_{\theta,K}[\varphi]_I(v) \sim P_{\theta}[\varphi]_I.$$

If  $\varphi$  has analytic singularities, we have

$$P_{\theta,K}[\varphi](v) = P_{\theta,K}[\varphi]_I(v). \quad (14.4)$$

{eq:relativePandPIana}

**Proof** Let  $C > 0$  such that  $-C \leq v \leq C$ . Then

$$P_{\theta}[\varphi] - C \leq P_{\theta,K}[\varphi](v).$$

Since  $K$  is non-pluripolar, for  $\eta \in \text{PSH}(X, \theta)$  the condition  $\eta|_K \leq v \leq C$  implies that  $\eta \leq \tilde{C}$  on  $X$  for some  $\tilde{C} := \tilde{C}(C, K) > 0$  [GZ07, Corollary 4.3]. This implies that

$$P_{\theta,K}[\varphi](v) \leq P_{\theta}[\varphi] + \tilde{C},$$

giving

$$P_{\theta,K}[\varphi](v) \sim P_{\theta}[\varphi].$$

The exact same argument applies in case of the relative  $\mathcal{I}$ -envelope.

Next assume that  $\varphi$  has analytic singularities, then we have that

$$\varphi \sim P_{\theta}[\varphi]_{\mathcal{I}}$$

by **Proposition 3.2.9**. In particular, for  $\eta \in \text{PSH}(X, \theta)$ ,  $\eta \leq \varphi$  if and only if  $\eta \leq P_{\theta}[\varphi]_{\mathcal{I}}$ . So (14.4) follows.  $\square$

cor:projectivity

**Corollary 14.1.1** *Let  $v \in C^0(X)$ . Then*

$$P_{\theta,K}[\varphi]_{\mathcal{I}}(v) = P_{\theta,X}[P_{\theta,K}[\varphi]_{\mathcal{I}}(v)]_{\mathcal{I}}(v).$$

**Proof** By definition, we have

$$\begin{aligned} & P_{\theta,X}[P_{\theta,K}[\varphi]_{\mathcal{I}}(v)]_{\mathcal{I}}(v) \\ &= \sup^* \{ \eta \in \text{PSH}(X, \theta) : \eta|_K \leq v, \eta \leq_{\mathcal{I}} P_{\theta,K}[\varphi]_{\mathcal{I}}(v) \} \\ &= \sup^* \{ \eta \in \text{PSH}(X, \theta) : \eta|_K \leq v, \eta \leq_{\mathcal{I}} \varphi \} \\ &= P_{\theta,K}[\varphi]_{\mathcal{I}}(v), \end{aligned}$$

where we applied **Lemma 14.1.2** on the third line.  $\square$

lma:PKoutsidepps

**Lemma 14.1.3** *Assume that  $\varphi \in \text{PSH}(X, \theta)_{>0}$ . Let  $v \in C^0(K)$ . Let  $S \subseteq X$  be a pluripolar set and  $\eta \in \text{PSH}(X, \theta)_{>0}$  with  $\eta \leq \varphi$ . Assume that  $\eta|_{K \setminus S} \leq v|_{K \setminus S}$ , then  $\eta \leq P_{\theta,K}[\varphi](v)$ .*

**Proof** By **Theorem 1.1.5**, there is  $\chi \in \text{PSH}(X, \theta)$ , such that  $\chi|_S \equiv -\infty$ . We claim that we can choose  $\chi$  so that

$$\chi \leq \eta.$$

In fact, since  $\int_X \theta_{\eta}^n > 0$ , fixing some  $\chi$  and  $\epsilon \in (0, 1)$  small enough, we have

$$\int_X \theta_{\epsilon\chi + (1-\epsilon)V_{\theta}}^n + \int_X \theta_{\eta}^n > \int_X \theta_{V_{\theta}}^n.$$

Thus, by **Proposition 3.1.3**, we have

$$(\epsilon\chi + (1-\epsilon)V_{\theta}) \wedge \eta \in \text{PSH}(X, \theta).$$

It suffices to replace  $\chi$  by  $(\epsilon\chi + (1-\epsilon)V_{\theta}) \wedge \eta$ .

Fix  $\chi \leq \eta$  as above. For any  $\delta \in (0, 1)$ , we have

$$(1-\delta)\eta|_K + \delta\chi|_K \leq v, \quad (1-\delta)\eta + \delta\chi \leq \varphi.$$

Hence,

$$(1-\delta)\eta + \delta\chi \leq P_{\theta,K}[\varphi](v).$$

Letting  $\delta \rightarrow 0+$ , we conclude that  $\eta \leq P_{\theta,K}[\varphi](v)$ .  $\square$

cor:PKtoPX

**Corollary 14.1.2** Assume that  $\varphi \in \text{PSH}(X, \theta)_{>0}$ . Let  $v \in C^0(K)$ . Then

$$P_{\theta,K}[\varphi](v) = P_{\theta,X}[\varphi](P_{\theta,K}[V_\theta](v)).$$

**Proof** It is clear that

$$P_{\theta,K}[\varphi](v) \leq P_{\theta,X}[\varphi](P_{\theta,K}[V_\theta](v)).$$

For the reverse direction, it suffices to prove that any  $\eta \in \text{PSH}(X, \theta)$  such that

$$\eta \leq \varphi, \quad \eta \leq P_{\theta,K}[V_\theta](v),$$

we have

$$\eta \leq P_{\theta,K}[\varphi](v). \quad (14.5)$$

{eq:etaleqPthetaKtempl}

As  $\varphi$  has positive mass, we can assume that  $\eta$  has positive mass as well. Let

$$S = \{P_{\theta,K}[V_\theta](v) > P'_{\theta,K}[V_\theta](v)\}.$$

By [Proposition 1.2.3](#),  $S$  is a pluripolar set. Observe that

$$\eta|_{K \setminus S} \leq v|_{K \setminus S}.$$

Hence, (14.5) follows from [Lemma 14.1.3](#).  $\square$

The next result motivates our terminology to call the measures  $\theta_{P_{\theta,K}[\varphi](v)}^n$  the *partial equilibrium measures* of our context:

lma:balayage

**Lemma 14.1.4** Let  $v \in C^0(K)$ . Then

$$\int_{X \setminus K} \theta_{P_{\theta,K}[\varphi](v)}^n = 0.$$

Moreover,  $P_{\theta,K}[\varphi](v)|_K = v$  almost everywhere with respect to  $\theta_{P_{\theta,K}[\varphi](v)}^n$ . More precisely, we have

$$\theta_{P_{\theta,K}[\varphi](v)}^n \leq \mathbb{1}_{K \cap \{P_{\theta,K}[\varphi](v) = P_{\theta,K}[V_\theta](v) = v\}} \theta_{P_{\theta,K}[V_\theta](v)}^n. \quad (14.6)$$

{eq:thetaPKuv}

**Proof Step 1.** We address the case where  $\varphi = V_\theta$ .

Let  $S \subseteq X$  be a closed pluripolar set, such that  $V_\theta$  is locally bounded on  $X \setminus S$ . This is possible because we can always find a Kähler current with analytic singularities in the cohomology class  $[\theta]$ , as a consequence of [Theorem 1.6.2](#).

For the first assertion, it suffices to show that  $\theta_{P_{\theta,K}[V_\theta](v)}^n$  does not charge any open ball  $B \Subset X \setminus (S \cup K)$ .

By [Proposition 1.2.2](#), we can take an increasing sequence  $(\eta_j)_j$  in  $\text{PSH}(X, \theta)$  such that

$$\eta_j \rightarrow P_{\theta,K}[V_\theta](v) \text{ almost everywhere, } \eta_j|_K \leq v \text{ for all } j \geq 1.$$

By [BT82](#), Proposition 9.1], for each  $j \geq 1$ , we can find  $\gamma_j \in \text{PSH}(X, \theta)$ , such that  $(\theta + \text{dd}^c \gamma_j|_B)^n = 0$  and  $w_j$  agrees with  $\eta_j$  outside  $B$ . Note that  $(\gamma_j)_j$  is clearly increasing and

$$\gamma_j \geq \eta_j, \quad \gamma_j|_K \leq v.$$

for all  $j \geq 1$ .

It follows that  $\gamma_j$  converges to  $P_{\theta,K}[V_\theta](v)$  almost everywhere as well. By [Theorem 2.3.1](#), we find that  $\theta_{P_{\theta,K}[V_\theta](v)}^n$  does not charge  $B$ , as desired.

For the second assertion, let  $x \in (X \setminus S) \cap K$  be a point such that  $P_{\theta,K}[V_\theta](v)(x) < v(x) - \epsilon$  for some  $\epsilon > 0$ . Let  $B$  be a ball centered at  $x$ , small enough so that  $\theta$  has a local potential on  $B$ , allowing us to identify  $\theta$ -psh functions with psh functions (on  $B$ ). By shrinking  $B$ , we can further guarantee

- (1)  $\overline{B} \subseteq X \setminus S$ .
- (2)  $P_{\theta,K}[V_\theta](v)|_{\overline{B}} < v(x) - \epsilon$ .
- (3)  $v|_{\overline{B} \cap K} > v(x) - \epsilon$ .

Construct the sequences  $\eta_j, \gamma_j$  as above. On  $B$ , by choosing a local potential of  $\theta$ , we may identify  $\eta_j, \gamma_j$  with the corresponding psh functions in a neighborhood of  $\overline{B}$ . By (2), we have  $\gamma_j \leq v(x) - \epsilon$  on  $\partial B$ , hence by the comparison principle,  $\gamma_j|_B \leq v(x) - \epsilon$ . By (3), we have  $\gamma_j|_{B \cap K} \leq v|_{B \cap K}$ . Thus, we conclude that  $\theta_{P_{\theta,K}[V_\theta](v)}^n$  does not charge  $B$ , as in the previous paragraph.

**Step 2.** We handle the general case. We can assume  $\varphi \in \text{PSH}(X, \theta)_{>0}$ . Indeed, due to [Lemma 14.1.2](#) and [Theorem 2.3.2](#), we have that

$$\int_X \theta_{P_{\theta,K}[\varphi](v)}^n = \int_X \theta_\varphi^n.$$

Hence, there is nothing to prove if  $\int_X \theta_\varphi^n = 0$ .

By [Corollary 14.1.2](#),

$$P_{\theta,K}[\varphi](v) = P_{\theta,X}[\varphi](P_{\theta,K}[V_\theta](v)).$$

Now [DDNL18mono](#), Theorem 3.8] gives

$$\begin{aligned} \theta_{P_{\theta,K}[\varphi](v)}^n &\leq \mathbb{1}_{\{P_{\theta,K}[\varphi](v) = P_{\theta,K}[V_\theta](v)\}} \theta_{P_{\theta,K}[V_\theta](v)}^n \\ &\leq \mathbb{1}_{\{P_{\theta,K}[\varphi](v) = v\}} \theta_{P_{\theta,K}[V_\theta](v)}^n, \end{aligned}$$

where in the second inequality we have used Step 1. □

cor:supptheta

**Corollary 14.1.3** Let  $v \in C^0(K)$ .

$$\begin{aligned} \int_{(X \setminus K) \cup \{P_{\theta,K}[\varphi](v) < v\}} \theta_{P_{\theta,K}[\varphi](v)}^n &= 0, \\ \int_{(X \setminus K) \cup \{P_{\theta,K}[\varphi]_I(v) < v\}} \theta_{P_{\theta,K}[\varphi]_I(v)}^n &= 0. \end{aligned} \tag{14.7}$$

{eq:theta does not charge 1}

**Proof** The first equation in (14.7) follows from Lemma 14.1.4. For the second, we can assume that

$$\int_X \theta_{P_{\theta,K}[\varphi]_I}^n > 0, \quad (14.8)$$

otherwise there is nothing to prove. By definition, we have

$$P_{\theta,K}[\varphi]_I(v) = P_{\theta,K}[P_{\theta}[\varphi]_I]_I(v).$$

Next we show that

$$P_{\theta,K}[P_{\theta}[\varphi]_I]_I(v) = P_{\theta,K}[P_{\theta}[\varphi]_I](v).$$

The  $\geq$  direction is trivial. It remains to prove the reverse inequality. By Lemma 14.1.2, we get that

$$P_{\theta,K}[P_{\theta}[\varphi]_I]_I(v) \sim P_{\theta}[\varphi]_I.$$

Due to Proposition 1.2.3, we get that

$$P_{\theta,K}[P_{\theta}[\varphi]_I]_I(v) \leq v$$

on  $K \setminus S$ , where  $S \subseteq X$  is a pluripolar set. As a result, due to (14.8), Lemma 14.1.3 allows to conclude that

$$P_{\theta,K}[P_{\theta}[\varphi]_I]_I(v) \leq P_{\theta,K}[P_{\theta}[\varphi]_I](v).$$

Since

$$P_{\theta,K}[P_{\theta}[\varphi]_I]_I(v) = P_{\theta,K}[\varphi]_I(v),$$

we get that the second equation in (14.7), using the first.  $\square$

prop:PKdependsonmodeltype

**Proposition 14.1.1** Assume that  $\varphi \in \text{PSH}(X, \theta)_{>0}$ . Let  $v \in C^0(K)$ . Then

$$P_{\theta,K}[\varphi](v) = P_{\theta,K}[P_{\theta}[\varphi]](v). \quad (14.9)$$

{eq: interm\_eq}

In particular,

$$P_{\theta,K}[\varphi](v) = P_{\theta,K}[P_{\theta,K}[\varphi](v)](v).$$

**Proof** The  $\leq$  direction in (14.9) is obvious. We to prove the reverse inequality. As  $P_{\theta,K}[\varphi](v)$  and  $P_{\theta,K}[P_{\theta}[\varphi]](v)$  have the same singularity types by Lemma 14.1.2, by the domination principle [DDNL18, Corollary 3.10], it suffices to show that

$$P_{\theta,K}[\varphi](v) \geq P_{\theta,K}[P_{\theta}[\varphi]](v) \text{ almost everywhere with respect to } \theta_{P_{\theta,K}[\varphi]}^n. \quad (14.10)$$

{eq:PthetaKtemp1}

By (14.6),

$$P_{\theta,K}[\varphi](v) = P_{\theta,K}[V_{\theta}](v) = v$$

almost everywhere with respect to  $\theta_{P_{\theta,K}[\varphi]}^n$ . Hence,

$$P_{\theta,K}[P_{\theta}[\varphi]](v) = v$$

almost everywhere with respect to  $\theta_{P_{\theta,K}[\varphi]}^n(v)$ . We conclude that

$$P_{\theta,K}[\varphi](v) = P_{\theta,K}[P_{\theta}[\varphi]](v).$$

Finally, (14.10) follows from Lemma 14.1.2 and (14.9).  $\square$

def:partialequienergy

**Definition 14.1.2** Given  $\varphi \in \text{PSH}(X, \theta)_{>0}$ , the *partial equilibrium energy functional*  $\mathcal{E}_{[\varphi],K}^{\theta} : C^0(K) \rightarrow \mathbb{R}$  of  $v \in C^0(K)$  as follows

$$\mathcal{E}_{\theta,K}^{\varphi}(v) := E_{\theta}^{P_{\theta}[\varphi]_I}(P_{\theta,K}[\varphi]_I(v)). \quad (14.11)$$

Recall that the energy  $E_{\theta}^{P_{\theta}[\varphi]_I}$  functional is defined in Definition 3.1.5.

Note that by Lemma 14.1.2, we have

$$P_{\theta,K}[\varphi]_I(v) \in \mathcal{E}^{\infty}(X, \theta; P_{\theta}[\varphi]_I),$$

so  $\mathcal{E}_{\theta,K}^{\varphi}(v) \in \mathbb{R}$ .

prop: differential\_P

**Proposition 14.1.2** Let  $K \subseteq X$  be a closed non-pluripolar set,  $v, f \in C^0(K)$  and  $\varphi \in \text{PSH}(X, \theta)_{>0}$ . Then  $\mathbb{R} \ni t \mapsto \mathcal{E}_{\theta,K}^{\varphi}(v + tf)$  is differentiable and

$$\frac{d}{dt} \mathcal{E}_{\theta,K}^{\varphi}(v + tf) = \int_K f \theta_{P_{\theta,K}[\varphi]_I}^n(v + tf) \quad (14.12)$$

{eq: ddtI}

for all  $t \in \mathbb{R}$ .

**Proof** We may assume that  $\varphi$  is  $I$ -model by replacing  $\varphi$  by  $P_{\theta}[\varphi]_I$ .

Note that it suffices to prove (14.12) at  $t = 0$ , which is equivalent to

$$\lim_{t \rightarrow 0} \frac{E_{\theta}^{\varphi}(P_{\theta,K}[\varphi]_I(v + tf)) - E_{\theta}^{\varphi}(P_{\theta,K}[\varphi]_I(v))}{t} = \int_K f \theta_{P_{\theta,K}[\varphi]_I}^n(v). \quad (14.13)$$

{eq: to\_prove\_1}

By switching  $f$  to  $-f$ , we may assume that  $t > 0$  in the above limit.

By the comparison principle [DDNL18, Proposition 3.5] and Proposition 3.1.11, we find

$$\begin{aligned} & E_{\theta}^{\varphi}(P_{\theta,K}[\varphi]_I(v + tf)) - E_{\theta}^{\varphi}(P_{\theta,K}[\varphi]_I(v)) \\ &= \frac{1}{n+1} \sum_{i=0}^n \int_X (P_{\theta,K}[\varphi]_I(v + tf) - P_{\theta,K}[\varphi]_I(v)) \theta_{P_{\theta,K}[\varphi]_I}^i(v + tf) \wedge \theta_{P_{\theta,K}[\varphi]_I}^{n-i}(v) \\ &\leq \int_X (P_{\theta,K}[\varphi]_I(v + tf) - P_{\theta,K}[\varphi]_I(v)) \theta_{P_{\theta,K}[\varphi]_I}^n(v). \end{aligned}$$

By Lemma 14.1.4,

$$\int_X (P_{\theta,K}[\varphi]_I(v + tf) - P_{\theta,K}[\varphi]_I(v)) \theta_{P_{\theta,K}[\varphi]_I}^n(v) \leq t \int_K f \theta_{P_{\theta,K}[\varphi]_I}^n(v).$$

Thus, we get the inequality,

$$\lim_{t \rightarrow 0+} \frac{E_{\theta}^{\varphi}(P_{\theta,K}[\varphi]_I(v+tf)) - E_{\theta}^{\varphi}(P_{\theta,K}[\varphi]_I(v))}{t} \leq \int_K f \theta_{P_{\theta,K}[\varphi]_I(v)}^n.$$

Similarly, we have

$$\begin{aligned} & E_{\theta}^{\varphi}(P_{\theta,K}[\varphi]_I(v+tf)) - E_{\theta}^{\varphi}(P_{\theta,K}[\varphi]_I(v)) \\ & \geq \int_X (P_{\theta,K}[\varphi]_I(v+tf) - P_{\theta,K}[\varphi]_I(v)) \theta_{P_{\theta,K}[\varphi]_I(v+tf)}^n \\ & \geq t \int_K f \theta_{P_{\theta,K}[\varphi]_I(v+tf)}^n. \end{aligned}$$

Together with the above, this implies (14.13).  $\square$

lem: global\_env\_approx

**Lemma 14.1.5** Fix a Kähler form  $\omega$  on  $X$ . For  $v \in C^0(K)$  there exists an increasing bounded sequence  $(v_j^-)_j$  in  $C^\infty(X)$  and a decreasing bounded sequence  $(v_j^+)_j$  in  $C^\infty(X)$ , such that for all  $\varphi \in \text{PSH}(X, \theta)_{>0}$  and  $\delta \in [0, 1]$  we have

- (1)  $P_{\theta+\delta\omega,X}[\varphi](v_j^+) \searrow P_{\theta+\delta\omega,K}[\varphi](v)$ ,
- (2)  $P_{\theta+\delta\omega,X}[\varphi](v_j^-) \nearrow P_{\theta+\delta\omega,K}[\varphi](v)$  almost everywhere,
- (3)  $\sup_X |v_j^-| \leq C$ ,  $\sup_X |v_j^+| \leq C$  for some constant  $C$  depending only on  $\|v\|_{C^0(K)}$ ,  $K$  and  $\theta + \omega$ , and
- (4)

$$\lim_{j \rightarrow \infty} \mathcal{E}_{\theta,K}^{\varphi}(v_j^-) = \mathcal{E}_{\theta,K}^{\varphi}(v), \quad \lim_{j \rightarrow \infty} \mathcal{E}_{\theta,K}^{\varphi}(v_j^+) = \mathcal{E}_{\theta,K}^{\varphi}(v).$$

**Proof** We fix  $\delta \in [0, 1]$ . First we prove the existence of  $(v_j^-)_j$ . Let

$$C_{K,v} := \sup \left\{ \sup_X \eta : \eta \in \text{PSH}(X, \theta + \omega), \eta|_K \leq v \right\}.$$

Since  $K$  is non-pluripolar, we have that  $C_{K,v} \in \mathbb{R}$ . Now define  $\tilde{v}: X \rightarrow \mathbb{R}$  as

$$\tilde{v}(x) = \begin{cases} v(x), & x \in K; \\ C_{K,v} + 1, & x \in X \setminus K. \end{cases}$$

Since  $\tilde{v}$  is lower semicontinuous, there exists an increasing and uniformly bounded sequence  $(v_j^-)_j$  in  $C^\infty(X)$ , such that  $v_j^- \nearrow \tilde{v}$ .

Observe that  $P_{\theta+\delta\omega,X}[\varphi](v_j^-)$  is increasing in  $j \geq 1$ , and

$$P_{\theta+\delta\omega,X}[\varphi](v_j^-) \leq P_{\theta+\delta\omega,K}[\varphi](v).$$

To prove that

$$P_{\theta+\delta\omega,X}[\varphi](v_j^-) \nearrow P_{\theta+\delta\omega,K}[\varphi](v)$$

almost everywhere, let  $\eta$  be a candidate for  $P_{\theta+\delta\omega,K}[\varphi](v)$  such that  $\sup_K (\eta - v) < 0$ . Then, since  $\eta$  is upper semicontinuous and  $\eta < \tilde{v}$ , by Dini's lemma there exists  $j_0 > 0$



such that  $\eta < v_j^-$  for  $j \geq j_0$ , i.e.

$$\eta \leq P_{\theta+\delta\omega, X}[\varphi](v_j^-),$$

proving existence of  $(v_j^-)_j$ .

Next, we prove the existence of  $(v_j^+)_j$ . Since

$$h := P_{\theta+\omega, K}[V_{\theta+\omega}](v) \vee (\inf_K v - 1)$$

is usc, there exists a decreasing and uniformly bounded sequence  $(v_j^+)_j$  in  $C^\infty(X)$ , such that  $v_j^+ \searrow h$ . Trivially,

$$\chi := \lim_{j \rightarrow \infty} P_{\theta+\delta\omega, X}[\varphi](v_j^+) \geq P_{\theta+\delta\omega, K}[\varphi](v).$$

In particular,  $\chi$  has positive mass, since it has the same singularity types as  $P_{\theta+\delta\omega, K}[\varphi](v)$  by [Lemma 14.1.2](#). We introduce

$$S := \{P'_{\theta+\omega, K}[V_{\theta+\omega}](v) < P_{\theta+\omega, K}[V_{\theta+\omega}](v)\}.$$

By [Proposition 1.2.3](#),  $S$  is a pluripolar set. Observe that

$$P_{\theta+\delta\omega, X}[\varphi](v_j^+) \leq v_j^+$$

for all  $j \geq 1$ . Thus,  $\chi \leq h$ . On the other hand,  $h \leq v$  on  $K \setminus S$ . So in particular,  $\chi|_{K \setminus S} \leq v|_{K \setminus S}$ . By [Lemma 14.1.2](#) we also have that  $\chi \sim P_{\theta+\delta\omega, K}[\varphi](v)$ . Hence, by [Lemma 14.1.3](#),

$$\chi \leq P_{\theta+\delta\omega, K}[P_{\theta+\delta\omega, K}[\varphi](v)](v) = P_{\theta+\delta\omega, K}[\varphi](v),$$

where we also used the last statement of [Proposition 14.1.1](#).

Finally observe that (4) follows from [Lemma 14.1.2](#), [Lemma 14.1.5](#) and [Theorem 2.3.1](#).  $\square$

prop: conv\_of\_K\_env

**Proposition 14.1.3** *Let  $K \subseteq X$  be a compact and non-pluripolar subset. Let  $v \in C^0(K)$ . Let  $\varphi_j, \varphi \in \text{PSH}(X, \theta)_{>0}$  ( $j \geq 1$ ) with  $\varphi_j \xrightarrow{d_S} \varphi$ . Then the following hold:*

- (1) *If  $\varphi_j \searrow \varphi$ , then  $P_{\theta, K}[\varphi_j]_I(v) \searrow P_{\theta, K}[\varphi]_I(v)$  and  $P_{\theta, K}[\varphi_j](v) \searrow P_{\theta, K}[\varphi](v)$ .*
- (2) *If  $\varphi_j \nearrow \varphi$  almost everywhere then  $P_{\theta, K}[\varphi_j]_I(v) \nearrow P_{\theta, K}[\varphi]_I(v)$  almost everywhere, and  $P_{\theta, K}[\varphi_j](v) \nearrow P_{\theta, K}[\varphi](v)$  almost everywhere.*

**Proof** (1) By [Theorem 6.2.1](#), we have

$$\lim_{j \rightarrow \infty} \int_X \theta_{\varphi_j}^n = \int_X \theta_{\varphi}^n.$$

It follows from [Lemma 2.3.1](#) that there is a decreasing sequence  $\epsilon_j \searrow 0$  with  $\epsilon_j \in (0, 1)$  and  $\eta_j \in \text{PSH}(X, \theta)$  such that

$$(1 - \epsilon_j)\varphi_j + \epsilon_j\eta_j \leq \varphi.$$

By the concavity similar to [Proposition 3.2.10](#), we get

$$\begin{aligned} (1 - \epsilon_j)P_{\theta,K}[\varphi_j]_I(v) + \epsilon_j P_{\theta,K}[\eta_j]_I(v) &\leq P_{\theta,K}[(1 - \epsilon_j)\varphi_j + \epsilon_j\eta_j]_I(v) \\ &\leq P_{\theta,K}[\varphi]_I(v). \end{aligned}$$

Since  $(\varphi_j)_j$  is decreasing, so is  $(P_{\theta,K}[\varphi_j]_I(v))_j$ , hence

$$\psi := \lim_{j \rightarrow \infty} P_{\theta,K}[\varphi_j]_I(v) \geq P_{\theta,K}[\varphi]_I(v)$$

exists. Since  $\epsilon_j \rightarrow 0$  and  $\sup_X P_{\theta,K}[\eta_j]_I(v)$  is bounded, we can let  $j \rightarrow \infty$  in the above estimate to conclude that

$$\psi = P_{\theta,K}[\varphi]_I(v).$$

The same ideas yield that

$$P_{\theta,K}[\varphi_j](v) \searrow P_{\theta,K}[\varphi](v).$$

The proof of (2) is similar and is left to the readers.  $\square$

## 14.2 Quantization of partial equilibrium measures

sec:quant

Let  $X$  be a connected compact Kähler manifold of dimension  $n$  and  $L$  be a pseudoeffective line bundle on  $X$ . Let  $h$  be a Hermitian metric on  $L$  and set  $\theta = c_1(L, h)$ . Let  $(T, h_T)$  be a Hermitian line bundle on  $X$ . Take a Kähler form  $\omega$  on  $X$  so that

$$\int_X \omega^n = 1.$$

### 14.2.1 Bernstein–Markov measures

Let  $K \subseteq X$  be a closed non-pluripolar subset. Let  $v$  be a measurable function on  $K$  and let  $\mu$  be a positive Borel probability measure on  $K$ . We introduce the following functions on  $H^0(X, L^k \otimes T)$  ( $k \geq 1$ ), with values possibly equaling  $\infty$ :

$$\begin{aligned} N_{v,v}^k(s) &:= \left( \int_K h^k \otimes h_T(s, s) e^{-kv} d\mu \right)^{1/2}, \\ N_{v,K}^k(s) &:= \sup_{K \setminus \{v=-\infty\}} \left( h^k \otimes h_T(s, s) e^{-kv} \right)^{1/2}. \end{aligned}$$

We start with the following elementary observation:

lma:mononorm

**Lemma 14.2.1** *Let  $v_1 \leq v_2$  be two measurable functions on  $X$ . Assume that  $\{v_1 = -\infty\} = \{v_2 = -\infty\}$ . Then for any  $s \in H^0(X, L^k \otimes T)$  ( $k \geq 1$ ), we have*

$$N_{v_1, K}^k(s) \geq N_{v_2, K}^k(s).$$

If  $v$  puts no mass on  $\{v = -\infty\}$  then we always have

$$N_{v, v}^k(s) \leq N_{v, K}^k(s). \quad (14.14)$$

{eq:Nkinfcomp}

def:weightedss

**Definition 14.2.1** A *weighted subset* of  $X$  is a pair  $(K, v)$  consisting of a closed non-pluripolar subset  $K \subseteq X$  and a function  $v \in C^0(K)$ .

def:BMmeasure

**Definition 14.2.2** Let  $(K, v)$  be a weighted subset of  $X$ . A positive Borel probability measure  $\nu$  on  $K$  is *Bernstein–Markov* with respect to  $(K, v)$  if for each  $\epsilon > 0$ , there is a constant  $C_\epsilon > 0$  such that

$$N_{\nu, K}^k(s) \leq C_\epsilon e^{\epsilon k} N_{v, \nu}^k(s) \quad (14.15)$$

{eq:BM}

for any  $s \in H^0(X, L^k \otimes T)$  and any  $k \in \mathbb{N}$ . We write  $\text{BM}(K, v)$  for the set of Bernstein–Markov measures with respect to  $(K, v)$ .

As pointed out in [BBWN11], any volume form on  $X$  is Bernstein–Markov with respect to  $(X, v)$ , with  $v \in C^\infty(X)$ .

prop:BMimplynorm

**Proposition 14.2.1** *Assume that  $(K, v)$  is a weighted subset of  $X$ , then*

- (1)  $N_{v, K}^k$  is a norm on  $H^0(X, L^k \otimes T)$ .
- (2) For any  $\nu \in \text{BM}(K, v)$ ,  $N_{\nu, \nu}^k$  is a norm on  $H^0(X, L^k \otimes T)$ .

**Proof** (1) As  $v$  is bounded,  $N_{v, K}^k$  is clearly finite on  $H^0(X, L^k \otimes T)$ . In order to show that it is a norm, it suffices to show that for any  $s \in H^0(X, L^k \otimes T)$ ,  $N_{v, K}^k(s) = 0$  implies that  $s = 0$ . In fact, we have  $s|_K = 0$ , hence  $s = 0$  by the connectedness of  $X$ .

(2) As  $v$  is bounded, clearly  $N_{\nu, \nu}^k$  is finite and satisfies the triangle inequality. Non-degeneracy follows from the fact that  $N_{v, K}^k$  is a norm and (14.15).  $\square$

## 14.2.2 Partial Bergman kernels

In this section, we fix a weighted subset  $(K, v)$  of  $X$  and  $\nu \in \text{BM}(K, v)$ .

**Definition 14.2.3** For any  $\varphi \in \text{PSH}(X, \theta)$ , we introduce the *partial Bergman kernels* of  $\varphi$  (with respect to  $(K, v)$ ) as follows: For any  $k \geq 0$ , we introduce

$$B_{v, \varphi, \nu}^k(x) := \sup \left\{ h^k \otimes h_T(s, s) e^{-k\varphi}(x) : N_{v, \nu}^k(s, s) \leq 1, \right. \\ \left. s \in H^0(X, L^k \otimes T \otimes I(k\varphi)) \right\}, \quad x \in K. \quad (14.16)$$

We extend  $B_{v,\varphi,v}^k$  to the whole  $X$  by setting it to be 0 outside  $K$ .

The *partial Bergman measures* of  $\varphi$  (with respect to  $(K, v)$ ) are defined as

$$\beta_{v,\varphi,v}^k := \frac{n!}{k^n} B_{v,\varphi,v}^k dv \quad (14.17)$$

for each  $k \geq 0$ .

Observe that

$$\int_K \beta_{v,\varphi,v}^k = \frac{n!}{k^n} h^0(X, T \otimes L^k \otimes \mathcal{I}(k\varphi)). \quad (14.18) \quad \{\text{eq:intbeta}\}$$

The goal of this section is to prove the following theorem:

thm: pBMconvergence

**Theorem 14.2.1** *Suppose that  $\varphi \in \text{PSH}(X, \theta)_{>0}$ . Let  $(K, v)$  be a weighed subset of  $X$ , let  $v \in \text{BM}(K, v)$ . Then*

$$\beta_{v,\varphi,v}^k \rightarrow \theta_{P_{\theta,K}[\varphi]_I(v)}^n \quad (14.19) \quad \{\text{eq:pbkconvgeneral}\}$$

as  $k \rightarrow \infty$ .

prop: smooth\_weak\_conv

**Proposition 14.2.2** *Let  $\varphi \in \text{PSH}(X, \theta)$  be a potential with analytic singularities such that  $\theta_\varphi$  is a Kähler current. If  $v \in C^\infty(X)$ , then*

$$\beta_{v,\varphi,\omega^n}^k \rightarrow \theta_{P_{\theta,X}[\varphi]_I(v)}^n = \theta_{P_{\theta,X}[\varphi](v)}^n \quad (14.20) \quad \{\text{eq:pbmconvanaly}\}$$

as  $k \rightarrow \infty$ .

**Proof** The equality part in (14.20) follows from Lemma 14.1.2. We start with noticing that as  $k \rightarrow \infty$ ,

$$\beta_{v,\varphi,\omega^n}^k \leq \beta_{v,V_\theta,\omega^n}^k \rightarrow \theta_{P_{\theta,X}[V_\theta](v)}^n = \mathbb{1}_{\{v=P_{\theta,X}[V_\theta](v)\}} \theta_v^n,$$

where the convergence follows from [Ber11, Theorem 1.2], and the last identity is due to [DNT21, Corollary 3.4]. Let  $\mu$  be the weak limit of a subsequence of  $\beta_{v,\varphi,\omega^n}^k$ , then we obtain that

$$\mu \leq \mathbb{1}_{\{v=P_{\theta,X}[V_\theta](v)\}} \theta_v^n. \quad (14.21) \quad \{\text{eq: Bergmanmeasure}\}$$

Let  $k > 0$ ,  $s \in H^0(X, L^k \otimes T \otimes \mathcal{I}(k\varphi))$  be a section such that  $N_{v,\omega^n}^k(s, s) \leq 1$ . Then by [Ber11, Lemma 4.1], there exists  $C > 0$  such that

$$h^k \otimes h_T(s, s) e^{-kv} \leq B_{v,\varphi,\omega^n}^k \leq B_{v,V_\theta,\omega^n}^k \leq k^n C.$$

This implies that

$$\frac{1}{k} \log h^k \otimes h_T(s, s) \leq v + \frac{\log C}{k} + n \frac{\log k}{k}.$$

We define  $\varphi_k$  as in Proposition 1.8.2. Take  $\alpha_k \nearrow 1$  as in Proposition 1.8.2. Then

$$\frac{1}{k} \log h^k \otimes h_T(s, s) \leq \varphi_k \leq \alpha_k \varphi.$$

Let  $\epsilon > 0$ . We notice that  $\frac{1}{k} \log h^k \otimes h_T(s, s) \in \text{PSH}(X, \theta + \epsilon\omega)$  for all  $k \geq k_0(\epsilon)$ . In particular,

$$\frac{1}{k} \log h^k \otimes h_T(s, s) - \frac{\log C}{k} - n \frac{\log k}{k} \leq P_{\theta+\epsilon\omega, X}[\alpha_k \varphi](v).$$

Now taking supremum over all candidates  $s$ , we obtain that

$$B_{v, \varphi, \omega^n}^k \leq C k^n e^{k(P_{\theta+\epsilon\omega, X}[\alpha_k \varphi](v) - v)}, \quad k \geq k_0. \quad (14.22)$$

{eq: smooth\_Berg\_est}

We claim that  $\mu$  does not put mass on  $\{P_{\theta+\epsilon\omega, X}[\varphi](v) < v\}$  for any  $\epsilon > 0$ . Since

$$P_{\theta+\epsilon\omega, X}[\alpha_k \varphi](v) \searrow P_{\theta+\epsilon\omega, X}[\varphi](v)$$

by [Proposition 14.1.3](#), we get that

$$\{P_{\theta+\epsilon\omega, X}[\alpha_k \varphi](v) < v\} \nearrow \{P_{\theta+\epsilon\omega, X}[\varphi](v) < v\}.$$

As a result, to argue the claim, it suffices to show that  $\mu$  does not put mass on the set  $\{P_{\theta+\epsilon\omega, X}[\alpha_k \varphi](v) < v\}$  for any  $k$ . Note that the latter set is open, hence [\(14.22\)](#) implies our claim.

Since  $\varphi$  has analytic singularities, we have that

$$P_{\theta+\epsilon\omega, X}[\varphi](v) \sim \varphi$$

for all  $\epsilon \geq 0$  by [Lemma 14.1.2](#) and [Proposition 3.2.9](#). As a result,

$$P_{\theta+\epsilon\omega, X}[\varphi](v) \searrow P_{\theta, X}[\varphi](v),$$

and we can let  $\epsilon \searrow 0$  to conclude that  $\mu$  does not put mass on  $\{P_{\theta, X}[\varphi](v) < v\} = \bigcup_{\epsilon > 0} \{P_{\theta+\epsilon\omega, X}[\varphi](v) < v\}$ . Putting this together with [\(14.21\)](#), we obtain that

$$\mu \leq \mathbb{1}_{\{P_{\theta, X}[\varphi](v) = v\}} \theta_v^n = \theta_{P_{\theta, X}[\varphi](v)}^n,$$

where the last equality is due to [\[DNT19, Corollary 3.4\]](#). Comparing total masses via [\(14.18\)](#) and [Theorem 7.3.1](#), we conclude that  $\mu = \theta_{P_{\theta, X}[\varphi](v)}^n$ . As  $\mu$  is an arbitrary cluster point of  $\beta_{v, \varphi, \omega^n}^k$ , we conclude that  $\beta_{v, \varphi, \omega^n}^k$  converges weakly to  $\theta_{P_{\theta, X}[\varphi](v)}^n$ , as  $k \rightarrow \infty$ .  $\square$

**Definition 14.2.4** Take  $k \geq 0$  and  $\varphi \in \text{PSH}(X, \theta)$ , let  $\text{Norm}(\text{H}^0(X, L^k \otimes T \otimes I(k\varphi)))$  be the space of Hermitian norms on the vector space  $\text{H}^0(X, L^k \otimes T \otimes I(k\varphi))$ .

Let  $\mathcal{L}_{k, \varphi} : \text{Norm}(\text{H}^0(X, L^k \otimes T \otimes I(k\varphi))) \rightarrow \mathbb{R}$  be the *partial Donaldson functional*:

$$\mathcal{L}_{k, \varphi}(H) = \frac{n!}{k^{n+1}} \log \frac{\text{vol}\{s : H(s) \leq 1\}}{\text{vol}\{s : N_{0, \omega^n}^k(s) \leq 1\}}, \quad (14.23)$$

where  $\text{vol}$  is simply the Euclidean volume.

prop: quant\_I\_smooth

**Proposition 14.2.3** *Let  $w, w' \in C^0(X)$  and  $\varphi \in \text{PSH}(X, \theta)$  be a potential with analytic singularities such that  $\theta_\varphi$  is a Kähler current, then*

$$\lim_{k \rightarrow \infty} \left( \mathcal{L}_{k, \varphi}(N_{w, \omega^n}^k) - \mathcal{L}_{k, \varphi}(N_{w', \omega^n}^k) \right) = \mathcal{E}_{\theta, X}^\varphi(w) - \mathcal{E}_{\theta, X}^\varphi(w'). \quad (14.24)$$

{eq: LdiffonXsmoothmeasure}

In particular,

$$\lim_{k \rightarrow \infty} \mathcal{L}_{k, \varphi}(N_{w, \omega^n}^k) = \mathcal{E}_{\theta, X}^\varphi(w). \quad (14.25)$$

{eq: LdiffonXsmoothmeasure2}

**Proof** First observe that by [Proposition 14.2.1](#), for any  $k \geq 0$ ,  $N_{w, \omega^n}^k$  and  $N_{w', \omega^n}^k$  are both norms, hence the expressions inside the limit in [\(14.24\)](#) make sense.

To start, we make the following observation:

$$\begin{aligned} \mathcal{L}_{k, \varphi}(N_{w, \omega^n}^k) - \mathcal{L}_{k, \varphi}(N_{w', \omega^n}^k) &= \int_0^1 \frac{d}{dt} \mathcal{L}_{k, \varphi}(N_{w+t(w'-w), \omega^n}^k) dt \\ &= \int_0^1 \int_X (w' - w) \beta_{w+t(w'-w), \varphi, \omega^n}^k dt. \end{aligned}$$

By [Proposition 14.2.2](#), we have

$$\lim_{k \rightarrow \infty} \int_X (w' - w) \beta_{w+t(w'-w), \varphi, \omega^n}^k = \int_X (w' - w) \theta_{P_{\theta, X}[\varphi](w+t(w'-w))}^n.$$

By [Theorem 7.3.1](#), we have  $|\int_X (w' - w) \beta_{w+t(w'-w), u, \omega^n}^k| \leq C \sup_X |w - w'|$ . Hence, by the dominated convergence theorem we obtain that

$$\begin{aligned} \lim_{k \rightarrow \infty} \left( \mathcal{L}_{k, \varphi}(N_{w, \omega^n}^k) - \mathcal{L}_{k, \varphi}(N_{w', \omega^n}^k) \right) &= \int_0^1 \int_X (w' - w) \theta_{P_{\theta, X}[\varphi](w+t(w'-w))}^n dt \\ &= \mathcal{E}_{\theta, X}^\varphi(w) - \mathcal{E}_{\theta, X}^\varphi(w'), \end{aligned}$$

where in the last line we have used [Proposition 14.1.2](#).

Finally, [\(14.25\)](#) is just a special case of [\(14.24\)](#) with  $w' = 0$ .  $\square$

lem:BML

**Lemma 14.2.2** *Let  $\varphi \in \text{PSH}(X, \theta)$ . Let  $(K, \nu)$  be a weighted subset of  $X$ . Let  $\nu \in \text{BM}(K, \nu)$ . Then*

$$\lim_{k \rightarrow \infty} \left( \mathcal{L}_{k, \varphi}(N_{\nu, K}^k) - \mathcal{L}_{k, \varphi}(N_{\nu, \nu}^k) \right) = 0. \quad (14.26)$$

{eq: Bern\_Mark\_implies}

**Proof** This is a direct consequence of the definition of Bernstein–Markov measures [\(14.15\)](#).  $\square$

cor:Ninfdifflim

**Corollary 14.2.1** *Let  $w \in C^0(X)$ ,  $\varphi \in \text{PSH}(X, \theta)$  be a potential with analytic singularities such that  $\theta_\varphi$  is a Kähler current. Then*

$$\lim_{k \rightarrow \infty} \mathcal{L}_{k, \varphi}(N_{w, X}^k) = \mathcal{E}_{\theta, X}^\varphi(w).$$

**Proof** This follows from [Lemma 14.2.2](#) and [Proposition 14.2.3](#) and the fact that  $\omega^n \in \text{BM}(X, 0)$ .  $\square$

**Proposition 14.2.4** *Let  $\varphi \in \text{PSH}(X, \theta)$  be a potential with analytic singularities such that  $\theta_\varphi$  is a Kähler current. Let  $(K, v)$ ,  $(K', v')$  be two weighted subsets of  $X$ . Then*

$$\lim_{k \rightarrow \infty} (\mathcal{L}_{k, \varphi}(N_{v, K}^k) - \mathcal{L}_{k, \varphi}(N_{v', K'}^k)) = \mathcal{E}_{\theta, K}^\varphi(v) - \mathcal{E}_{\theta, K'}^\varphi(v'). \quad (14.27)$$

{eq:LkdiffconvtoI}

In particular,

$$\lim_{k \rightarrow \infty} \mathcal{L}_{k, \varphi}(N_{v, K}^k) = \mathcal{E}_{\theta, K}^\varphi(v). \quad (14.28)$$

{eq:Lkconv}

**Proof** First observe that by [Proposition 14.2.1](#), for any  $k > 0$ ,  $N_{v, K}^k$  and  $N_{v', K'}^k$  are both norms, hence the expressions inside the limit in (14.27) make sense. Moreover, (14.28) is just a special case of (14.27) for  $K' = X$  and  $v' = 0$ .

To prove (14.27) it is enough to show that for any fixed  $w \in C^\infty(X)$  we have

$$\lim_{k \rightarrow \infty} (\mathcal{L}_{k, \varphi}(N_{v, K}^k) - \mathcal{L}_{k, \varphi}(N_{w, \omega^n}^k)) = \mathcal{E}_{\theta, K}^\varphi(v) - \mathcal{E}_{\theta, X}^\varphi(w). \quad (14.29)$$

{eq:inproofLdiffsmw}

For  $\epsilon \in (0, 1)$  small enough we have that  $\theta_{(1-\epsilon)\varphi}$  is still a Kähler current. Let us fix such  $\epsilon$ , along with an arbitrary  $\epsilon' \in (0, 1)$ .

Let  $(v_j^-)_j, (v_j^+)_j$  be the sequences of smooth functions constructed in [Lemma 14.1.5](#) for the data  $(K, v)$ .

By [Proposition 1.8.2](#) there exists  $k_0(\epsilon, \epsilon') \in \mathbb{N}$  such that

$$\frac{1}{k} \log h^k \otimes h_T(s, s) \leq (1 - \epsilon)u,$$

and  $\frac{1}{k} \log h^k \otimes h_T(s, s) \in \text{PSH}(X, \theta + \epsilon'\omega)$  for any  $s \in H^0(X, T \otimes L^k \otimes I(k\varphi))$ , as long as  $k \geq k_0(\epsilon, \epsilon')$ .

In particular, [Lemma 14.1.1](#) gives that

$$\begin{aligned} N_{P'_{\theta+\epsilon'\omega, K}[(1-\epsilon)\varphi](v), X}^k(s) &= N_{v, K}^k(s), \\ N_{P'_{\theta+\epsilon'\omega, X}[(1-\epsilon)\varphi](v_j^-), X}^k(s) &= N_{v_j^-, X}^k(s), \\ N_{P'_{\theta+\epsilon'\omega, X}[(1-\epsilon)\varphi](v_j^+), X}^k(s) &= N_{v_j^+, X}^k(s). \end{aligned}$$

As

$$P'_{\theta+\epsilon'\omega, X}[(1-\epsilon)\varphi](v_j^-) \leq P'_{\theta+\epsilon'\omega, K}[(1-\epsilon)\varphi](v) \leq P'_{\theta+\epsilon'\omega, X}[(1-\epsilon)\varphi](v_j^+),$$

by [Lemma 14.2.1](#) we have

$$N_{v_j^+, X}^k(s) \leq N_{v, K}^k(s) \leq N_{v_j^-, X}^k(s), \quad s \in H^0(X, T \otimes L^k \otimes I(k\varphi)), k \geq k_0(\epsilon, \epsilon').$$

Composing with  $\mathcal{L}_{k, \varphi}$  we arrive at

$$\mathcal{L}_{k,\varphi}(N_{v_j^-,X}^k) \leq \mathcal{L}_{k,\varphi}(N_{v,K}^k) \leq \mathcal{L}_{k,\varphi}(N_{v_j^+,X}^k), \quad k \geq k_0(\epsilon, \epsilon').$$

For any  $j > 0$ , by [Corollary 14.2.1](#) we get

$$\begin{aligned} \mathcal{E}_{\theta,X}^\varphi(v_j^-) - \mathcal{E}_{\theta,X}^\varphi(w) &= \lim_{k \rightarrow \infty} \left( \mathcal{L}_{k,\varphi}(N_{v_j^+,X}^k) - \mathcal{L}_{k,\varphi}(N_{w,X}^k) \right) \\ &\leq \varliminf_{k \rightarrow \infty} \left( \mathcal{L}_{k,\varphi}(N_{v,K}^k) - \mathcal{L}_{k,\varphi}(N_{w,X}^k) \right) \\ &\leq \overline{\lim}_{k \rightarrow \infty} \left( \mathcal{L}_{k,\varphi}(N_{v,K}^k) - \mathcal{L}_{k,\varphi}(N_{w,X}^k) \right) \\ &\leq \lim_{k \rightarrow \infty} \left( \mathcal{L}_{k,\varphi}(N_{v_j^-,X}^k) - \mathcal{L}_{k,\varphi}(N_{w,X}^k) \right) \\ &= \mathcal{E}_{\theta,X}^\varphi(v_j^+) - \mathcal{E}_{\theta,X}^\varphi(w). \end{aligned}$$

Using [Lemma 14.1.5](#), we can let  $j \rightarrow \infty$  to arrive at

$$\begin{aligned} \mathcal{E}_{\theta,K}^\varphi(v) - \mathcal{E}_{\theta,K}^\varphi(w) &\leq \varliminf_{k \rightarrow \infty} \left( \mathcal{L}_{k,\varphi}(N_{v,K}^k) - \mathcal{L}_{k,\varphi}(N_{w,X}^k) \right) \\ &\leq \overline{\lim}_{k \rightarrow \infty} \left( \mathcal{L}_{k,\varphi}(N_{v,K}^k) - \mathcal{L}_{k,\varphi}(N_{w,X}^k) \right) \\ &\leq \mathcal{E}_{\theta,K}^\varphi(v) - \mathcal{E}_{\theta,K}^\varphi(w). \end{aligned}$$

Hence, [\(14.29\)](#) follows.  $\square$

cor:LktoI

**Corollary 14.2.2** *Let  $\varphi \in \text{PSH}(X, \theta)$  be a potential with analytic singularities such that  $\theta_\varphi$  is a Kähler current. Let  $(K, v)$  be a weighted subset of  $X$ . Assume that  $v \in \text{BM}(K, v)$ . Then*

$$\lim_{k \rightarrow \infty} \mathcal{L}_{k,\varphi}(N_{v,v}^k) = \mathcal{E}_{\theta,K}^\varphi(v).$$

**Proof** Our claim follows from [Proposition 14.2.4](#) and [Lemma 14.2.2](#).  $\square$

prop:weakconvana

**Proposition 14.2.5** *Suppose that  $\varphi \in \text{PSH}(X, \theta)$  be a potential with analytic singularities such that  $\theta_\varphi$  is a Kähler current. Let  $(K, v)$  be a weighted subset of  $X$ . Let  $v \in \text{BM}(K, v)$ . Then*

$$\beta_{v,\varphi,v}^k \rightharpoonup \theta_{P_{\theta,K}[\varphi]}^n(v) = \theta_{P_{\theta,K}[\varphi]}^n(v)$$

weakly as  $k \rightarrow \infty$ .

**Proof** For  $w \in C^0(X)$ , let

$$f_k(t) := \mathcal{L}_{k,\varphi}(N_{v+tw,v}^k), \quad g(t) := \mathcal{E}_{\theta,K}^\varphi(v + tw).$$

By [Corollary 14.2.2](#)  $\varliminf_{k \rightarrow \infty} f_k(t) = g(t)$ . Note that  $f_k$  is concave by Hölder's inequality (see [\[BBW11, Proposition 2.4\]](#)), so by [\[BB10, Lemma 7.6\]](#),  $\lim_{k \rightarrow \infty} f'_k(0) = g'(0)$ , which is equivalent to  $\beta_{v,\varphi,v}^k \rightharpoonup \theta_{P_{\theta,K}[\varphi]}^n(v)$ , by [Proposition 14.1.2](#).  $\square$



prop:mainKahcurr

**Proposition 14.2.6** Suppose that  $\varphi \in \text{PSH}(X, \theta)$  such that  $\theta_\varphi$  is a Kähler current. Let  $(K, \nu)$  be a weighted subset of  $X$  and  $\nu \in \text{BM}(K, \nu)$ . Then

$$\beta_{\nu, \varphi, \nu}^k \rightharpoonup \theta_{P_{\theta, K}[\varphi]_I(\nu)}^n \quad (14.30)$$

{eq:pbkconv1}

as  $k \rightarrow \infty$ .

**Proof** Let  $\mu$  be the weak limit of a subsequence of  $\beta_{\nu, \varphi, \nu}^k$ . We claim that

$$\mu \leq \theta_{P_{\theta, K}[\varphi]_I(\nu)}^n. \quad (14.31)$$

{eq:inproofmuleq}

Observe that this claim implies the conclusion. In fact, by [Theorem 7.3.1](#), we have equality of the total masses, so equality holds in (14.31). As  $\mu$  is an arbitrary cluster point of the sequence  $(\beta_{\nu, \varphi, \nu}^k)_k$ , we get (14.30).

It remains to prove (14.31). Let  $(\varphi_j)$  be a quasi-equisingular approximation of  $\varphi$  in  $\text{PSH}(X, \theta)$ . We may assume that  $\theta_{\varphi_j}$  is a Kähler current for all  $j \geq 1$ . By [Lemma 14.1.2](#), [Corollary 7.1.2](#), we know that

$$\varphi_j \xrightarrow{d_S} P_{\theta, K}[\varphi]_I(\nu).$$

In particular,

$$\lim_{j \rightarrow \infty} \int_X \theta_{P_{\theta, K}[\varphi_j]_I(\nu)}^n = \int_X \theta_{P_{\theta, K}[\varphi]_I(\nu)}^n. \quad (14.32)$$

{eq:inproofeqmass}

Observe that

$$\beta_{\nu, \varphi, \nu}^k \leq \beta_{\nu, \varphi_j, \nu}^k$$

for any  $k \geq 1$ . As  $\nu \in \text{BM}(K, \nu)$ , by [Proposition 14.2.5](#),

$$\mu \leq \theta_{P_{\theta, K}[\varphi_j]_I(\nu)}^n,$$

for any  $j \geq 1$  fixed. By [Proposition 14.1.3](#),

$$P_{\theta, K}[\varphi_j]_I(\nu) \searrow P_{\theta, K}[\varphi]_I(\nu)$$

as  $j \rightarrow \infty$ . Hence, by (14.32) and [Theorem 2.3.1](#), (14.31) follows.  $\square$

**Proof (Proof of [Theorem 14.2.1](#))** By [Lemma 14.1.2](#), we have that

$$\begin{aligned} H^0(X, L^k \otimes T \otimes I(k\varphi)) &= H^0(X, L^k \otimes T \otimes I(kP_\theta[\varphi]_I)) \\ &= H^0(X, L^k \otimes T \otimes I(kP_{\theta, K}[\varphi]_I(\nu))). \end{aligned}$$

This allows us to replace  $\varphi$  with  $P_{\theta, K}[\varphi]_I(\nu)$ .

By [Lemma 2.3.2](#), there exists  $\varphi_j \in \text{PSH}(X, \theta)$ , such that  $\varphi_j \nearrow \varphi$  a.e. and  $\theta_{\varphi_j}$  is a Kähler current for each  $j \geq 1$ . This gives

$$\beta_{\nu, \varphi_j, \nu}^k \leq \beta_{\nu, \varphi, \nu}^k.$$

Let  $\mu$  be the weak limit of a subsequence of  $(\beta_{v,\varphi,v}^k)_k$ . Then by [Proposition 14.2.6](#),

$$\theta_{P_{\theta,K}[\varphi_j]_I(v)}^n \leq \mu.$$

By [Proposition 14.1.3](#) and [Theorem 2.3.1](#) we have that

$$\theta_{P_{\theta,K}[\varphi_j]_I(v)}^n \nearrow \theta_{P_{\theta,K}[\varphi]_I(v)}^n.$$

Hence,

$$\theta_{P_{\theta,K}[\varphi]_I(v)}^n \leq \mu. \tag{14.33}$$

{eq:inproofmulower}

A comparison of total masses using [\(14.18\)](#) and [Theorem 7.3.1](#) gives that equality holds in [\(14.33\)](#). As  $\mu$  is an arbitrary cluster limit of the weak compact sequence  $(\beta_{v,\varphi,\mu}^k)_k$ , we obtain [\(14.19\)](#).  $\square$

*Remark 14.2.1* The results in this chapter could also be reformulated as the large deviation principle of a determinantal point process on  $X$  using the Gärtner–Ellis theorem exactly as in [\[Ber14\]](#). We leave the details to the readers.

# Comments

chap:history

## A brief history

Here we recall the origin of various results.

### Chapter 1.

The global Josefson theorem [Theorem 1.1.5](#) was due to Vu [\[Vu19\]](#). In the projective setting, it was due to Dinh–Sibony [\[DS06\]](#) and in the Kähler setting, it was established by Guedj–Zeriahi [\[GZ05\]](#).

The extension theorem [Theorem 1.2.1](#) was proved in [\[GR56\]](#). In fact, they proved a more general version for complex spaces, see [Theorem B.2.2](#). We reproduced their arguments almost word by word for the convenience of the readers.

The plurifine topology was introduced by Bedford–Taylor [\[BT87\]](#) based on Cartan’s works on the fine topology. This area lacks a rigorous foundation until the appearance of [\[EMW06\]](#), which gave the first proof of [Theorem 1.3.2](#).

The strong openness was first established by Guan–Zhou [\[GZ15\]](#). A more elegant proof was due to Hiep [\[Hie14\]](#).

The idea of [Theorem 1.4.3](#) first appeared in the ground-breaking work of Boucksom–Favre–Jonsson [\[BFJ08\]](#).

[Proposition 1.2.6](#) was due to Kiselman [\[Kis78\]](#).

The semicontinuity theorem was due to Siu [\[Siu74\]](#).

### Chapter 2

The Monge–Ampère operators for bound plurisubharmonic functions were introduced by Bedford–Taylor [\[BT76, BT82\]](#). The non-pluripolar product is due to Bedford–Taylor [\[BT87\]](#), Guedj–Zeriahi [\[GZ07\]](#) and Boucksom–Eyssidieux–Guedj–Zeriahi [\[BEGZ10\]](#).

### Chapter 3

The notion of the  $P$ -envelope is due to Ross–Witt Nyström [\[RWN14\]](#) based on the ideas of Rashkovskii–Sigurdsson [\[RS05\]](#).

The  $\mathcal{I}$ -envelope was introduced by Darvas–Xia [\[DX22\]](#), inspired by the works of Dano Kim [\[Kim15\]](#) and Boucksom–Favre–Jonsson [\[BFJ08\]](#). The notion of  $\mathcal{I}$ -model singularities was first formulated in the explicit way in [\[DX22\]](#) in 2020, although it

was already essentially known in Boucksom–Jonsson’s work. In fact, they correspond exactly to the homogeneous non-Archimedean potentials assuming that the relevant masses do not vanish. A less explicit equivalent formulation of  $\mathcal{I}$ -model potentials also appeared in [Dem15]. A few months later, the same notion was rediscovered by Trusiani [Tru22].

Proposition 3.1.3 was first proved in [DDNLmetric, DDNL21b].

#### Chapter 4

The notion of weak geodesics was studied in detail by Darvas [Dar17] in the Kähler case.

The case of general big classes was partly handled in [DDNL18fullma, DDNL18big, DDNL18c], [DDNL18a]. However, the key fact that the geodesics between two full mass potentials have the correct limit at the end points does not seem to have been proved in any references. We give a proof in Proposition 4.2.1. We also extend the relevant results to the relative setting.

Previously, Proposition 4.2.2 and Proposition 4.2.4 were only known in the Kähler case.

#### Chapter 5

The toric framework was first written down by Coman–Guedj–Sahin–Zeriahi in [CGSZ19].

The beautiful theorem Theorem 5.3.1 was first proved by Yi Yao, who did not publish the result. Later on, a new proof was found by Botero–Burgos Gil–Holmes–de Jong [BBGHdJ21]. We chose to present the approach of Yao, which integrates naturally with our framework.

#### Chapter 6

The notion of  $P$ -partial order is new, as well as most results in Section 6.1.

The  $d_S$ -pseudometric was introduced in [DDNLmetric, DDNL21b]. The basic properties are proved in [DDNLmetric, DDNL21b] and [Xia21].

Theorem 6.2.4 is proved in [Xia22b]. Theorem 6.2.6 and Theorem 6.2.5 appear to be new. These results appeared previously in the form of lecture notes.

#### Chapter 7

The notion of  $\mathcal{I}$ -good singularities was due to [DX21]. The name  $\mathcal{I}$ -good was chosen in [Xia22b].

Example 7.1.3 was due to Berman–Boucksom–Jonsson [BBJ21].

Theorem 7.1.1 and Theorem 7.3.1 are due to [DX21, DX22].

There are some further examples of  $\mathcal{I}$ -good singularities provided by [BBGHdJ21] with applications in the theory of modular forms in [BBGHdJ22].

#### Chapter 8

The trace operator was introduced in [DX24]. Here we present a different point of view. Theorem 8.3.1 was proved in [DX24].

The analytic Bertini theorem Theorem 8.4.1 was proved in [Xia22a], based on the works of Matsumura–Fujino [FM21] and [Fuj23]. A weaker result was established by Meng–Zhou [MZ23].

#### Chapter 9

The technique of test curves originates from [RWN14]. It was generalized by Darvas–Di Nezza–Lu [DDNL18b], [DX21], [DZ22] and [DXZ23]. We give the full details of the proofs.

Test curves in Definition 9.1.1 are called *maximal test curves* in the literature, a terminology which I do not like. I prefer to call the usual notion of test curves in the literature *sub-test curves*.

Results in Section 9.4 are easy generalizations of the results proved in [Xia23Operations].

#### Chapter 10

The algebraic theory of partial Okounkov bodies was developed in [Xia21]. The transcendental Okounkov body was first defined by Deng [Deng17] as suggested by Demailly. The volume identity was proved in [DRWN+23]. The transcendental theory of partial Okounkov bodies is new. Results in Section 11.3 are also new.

#### Chapter 11

The applications of b-divisors in pluripotential theory began with [BF109]. The intersection theory of nef b-divisors was introduced by Dang–Favre [DF20]. The technique of singularity b-divisors was introduced in [Xia23c] in 2020. The general form first appeared in [Xia22b]. One year later, a special case was rediscovered in [BBGHJ21]. In 2023, another special case was rediscovered by Trusiani [Tru23].

#### Chapter 12

The whole chapter appears to be new. The study of toric pluripotential theory on big line bundles was made possible by the development of partial Okounkov bodies. The key result is Theorem 12.2.2.

Most results in this chapter resulted from discussions with Yi Yao.

#### Chapter 13

Most results from this chapter are from [Xia23Operations]. Results from Section 13.3 are new, although the main idea was already contained in [Xia21].

We deliberately avoid talking about the non-Archimedean point of view, which is explained in [DX22] and [Xia23b]. The reason is that the Berkovich analytification has not been constructed in written literature yet. This theory will be studied in the forthcoming thesis of Pietro Piccione.

Special cases of the results in this section have been applied to study K-stability, see [Xia23c], [DZ22], [DXZ23] and [DR22]. In [DX22], we established the bijective correspondence between a class of  $\mathcal{I}$ -model test curves with the maximal geodesic rays in the sense of [BBJ21].

#### Chapter 14

The special case of Theorem 14.2.1 without the prescribed singularity  $\varphi$  was due to Berman–Boucksom–Witt Nyström, see [BB10], [BBWN11]. The general case is due to [DX21].

## Open problems

We give a list of important open problem in this theory.

conj:exttracegeneral

*Conjecture 14.2.1* Let  $X$  be a connected compact Kähler manifold and  $Y$  be a submanifold. Fix a Kähler class  $\alpha$  on  $X$ . For each Kähler current  $S \in \alpha|_Y$ , we can find a Kähler current  $T \in \alpha$  such that

$$\mathrm{Tr}_Y(T) \sim_I S.$$

If we formally view  $\mathrm{Tr}_Y$  as an analogue of the trace operator in the theory of Sobolev spaces, then this conjecture corresponds exactly to the Dirichlet problem.

Using [Proposition 8.2.2](#), one could also reduce this conjecture to a strong version of the extension theorem [Theorem 1.6.3](#).

*Conjecture 14.2.2* Let  $X$  be a connected compact Kähler manifold and  $Y$  be a submanifold. Fix a Kähler class  $\alpha$  on  $X$ . Consider Kähler currents  $R \in \alpha$ ,  $S \in \alpha|_Y$  with analytic singularities such that  $S \leq R|_Y$ . Assume in addition that  $S$  has gentle analytic singularities. Then there is a Kähler current  $T \in \alpha$  with analytic singularities such that

$$\mathrm{Tr}_Y(T) \sim_I S, \quad T \leq R.$$

This conjecture was proposed by Darvas for different purposes.

*Conjecture 14.2.3* Let  $X$  be a connected smooth projective variety of dimension  $n$ . Assume that  $(L_i, h_i)$  is a Hermitian big line bundle on  $X$  for each  $i = 1, \dots, n$  with the  $h_i$ 's being  $\mathcal{I}$ -good. Then

$$\int_X c_1(L_1, h_1) \wedge \cdots \wedge c_1(L_n, h_n) = \sup_v \mathrm{vol}(\Delta_v(L_1, h_1), \dots, \Delta_v(L_n, h_n)),$$

where  $v: \mathbb{C}(X)^\times \rightarrow \mathbb{Z}^n$  runs over all (surjective) valuation of rank  $n$ .

See [\[Sch14\]](#) or [\[Sch93, Section 5.1\]](#) for the notion of mixed volumes.

This conjecture seems reasonable in view of [Corollary 10.2.3](#) and [Corollary 10.2.2](#).

Even when  $h_1, \dots, h_n$  have minimal singularities, this conjecture remains open:

*Conjecture 14.2.4* Let  $X$  be a connected smooth projective variety of dimension  $n$ . Assume that  $L_1, \dots, L_n$  are big line bundles on  $X$ . Then

$$\langle L_1, \dots, L_n \rangle = \sup_v \mathrm{vol}(\Delta_v(L_1), \dots, \Delta_v(L_n)),$$

where  $v: \mathbb{C}(X)^\times \rightarrow \mathbb{Z}^n$  runs over all (surjective) valuation of rank  $n$ .

Here on the left-hand side, we are using the movable intersection theory [\[BDPP13\]](#).

**Problem 14.2.1** Is it possible to extend the definition of the trace operator  $\mathrm{Tr}_Y$  to the case where the ambient variety is only unibranch?

The difficulty lies in the lack of Demailly type regularization theorems.

**Problem 14.2.2** What is the relation between the Duistermaat–Heckman measure in [Section 13.3](#) and the definition in [\[Ino22\]](#)?

**Problem 14.2.3** Is there a natural definition of the transcendental Okounkov body of a closed positive  $(1, 1)$ -current  $T$  with 0-mass so that its dimension is equal to the numerical dimension of  $T$ ?

See [\[Cao14\]](#) for the definition of the numerical dimension of a current.





## Appendix A

### Convex functions and convex bodies

chap:convex

We recall some basic facts about convex functions in this section. Our basic reference is [Roc70]. The results in this appendix can be applied to concave functions after considering their negatives.

#### A.1 The notion of convex functions

Let  $N$  be a real vector space of finite dimension.

**Definition A.1.1** Let  $F: N \rightarrow [-\infty, \infty]$  be a function. The *epigraph* of  $F$  is defined as the following set

$$\text{epi } F := \{(n, r) \in N \times \mathbb{R} : r \geq F(n)\}.$$

**Definition A.1.2** A *convex function* on  $N$  is a function  $F: N \rightarrow [-\infty, \infty]$  such that the epigraph  $\text{epi } F$  is a convex subset of  $N \times \mathbb{R}$ .

The *effective domain* of  $F$  is the set

$$\text{Dom } F := \{n \in N : F(n) < \infty\}.$$

A convex function  $F$  on  $N$  such that  $\text{Dom } F \neq \emptyset$  and  $F(n) \neq -\infty$  for all  $n \in N$  is said to be *proper*.

The set of convex functions on  $N$  is denoted by  $\text{Conv}(N)$ . The subset set of proper convex functions is denoted by  $\text{Conv}^{\text{prop}}(N)$ .

The following characterization of convex functions is well-known.

lma:charconvex

**Lemma A.1.1** Let  $F: N \rightarrow [-\infty, \infty]$ . Then  $F$  is convex if and only if the following condition holds: suppose that  $n, r \in N$  and  $a, b \in \mathbb{R}$  such that  $a > F(n)$ ,  $b > F(r)$ , then for any  $t \in (0, 1)$ , we have

$$F(tn + (1-t)r) < ta + (1-t)b.$$

See [Roc70, Theorem 4.2] for the proof.

*Example A.1.1* Let  $A \subseteq N$  be a convex subset. Then the *characteristic function*  $\chi_A: N \rightarrow \{0, \infty\}$  of  $A$  is defined by

$$\chi_A(n) := \begin{cases} 0, & n \in A; \\ \infty, & n \notin A. \end{cases}$$

The function  $\chi_A$  lies in  $\text{Conv}(N)$ .

ex:supfun

*Example A.1.2* Let  $M$  be the dual vector space of  $N$  and  $P \subseteq M$  be a convex subset. The *support function*  $\text{Supp}_P \in \text{Conv}(N)$  of  $P$  is defined as follows:

$$\text{Supp}_P(n) := \sup\{\langle m, n \rangle : m \in P\}.$$

It is well-known that convexity is preserved by a number of natural operations. We recall a few to fix the notation.

**Definition A.1.3** Let  $F_1, \dots, F_m \in \text{Conv}^{\text{prop}}(N)$  ( $m \in \mathbb{Z}_{>0}$ ). We define their *infimal convolution*  $F_1 \square \dots \square F_m \in \text{Conv}(N)$  as follows:

$$F_1 \square \dots \square F_m(n) := \inf \left\{ \sum_{i=1}^m F_i(n_i) : n_i \in N, \sum_{i=1}^m n_i = n \right\}.$$

The fact  $F_1 \square \dots \square F_m \in \text{Conv}(N)$  is proved in [Roc70, Theorem 5.4]. One should note that  $F_1 \square \dots \square F_m$  is not always proper.

prop:supconv

**Proposition A.1.1** Let  $\{F_i\}_{i \in I}$  be a non-empty family in  $\text{Conv}(N)$ . Then  $\sup_{i \in I} F_i \in \text{Conv}(N)$ .

This follows from [Roc70, Theorem 5.5]. In particular, this allows us to introduce

def:LCE

**Definition A.1.4** Let  $f: N \rightarrow [-\infty, \infty]$ . The *lower convex envelope* of  $f$  is defined as

$$\text{CE } f := \sup\{F \in \text{Conv}(N) : F \leq f\}.$$

It follows from Proposition A.1.1 that  $\text{CE } f \in \text{Conv}(N)$ .

def:convwedge

**Definition A.1.5** Given a non-empty family  $\{F_i\}_{i \in I}$  in  $\text{Conv}(N)$ , we define

$$\bigwedge_{i \in I} F_i := \text{CE} \left( \inf_{i \in I} F_i \right).$$

When the family  $I$  is finite, say  $I = \{1, \dots, m\}$ , we also write

$$F_1 \wedge \dots \wedge F_m = \bigwedge_{i \in I} F_i.$$

prop:concavhull

**Proposition A.1.2** Let  $F_1, \dots, F_m \in \text{Conv}^{\text{prop}}(N)$ , then

$$F_1 \wedge \dots \wedge F_m(x) = \inf \left\{ \sum_{i=1}^m \lambda_i F_i(x_i) : x_i \in \text{Dom}(F_i), \right. \\ \left. \lambda_i \in [0, 1], \sum_{i=1}^m \lambda_i = 1, \sum_{i=1}^m \lambda_i x_i = x \right\}.$$

See [Roc70, Theorem 5.6] for the more general result.

lma:convdecnet

**Lemma A.1.2** Let  $\{F_i\}_{i \in I}$  be a decreasing net in  $\text{Conv}(N)$ . Then  $\inf_{i \in I} F_i \in \text{Conv}(N)$ .

**Proof** Write  $F = \inf_{i \in I} F_i$ . We shall apply the characterization in Lemma A.1.1. Take  $n, r \in N$ ,  $a, b \in \mathbb{R}$  such that  $a > F(n)$ ,  $b > F(r)$  and  $t \in (0, 1)$ . We need to show that

$$F(tn + (1-t)r) < ta + (1-t)b. \quad (\text{A.1})$$

{eq:convtempl}

By definition, there exists  $j \in I$  such that for any  $i \geq I$  with  $i \geq j$ , we have

$$a > F_i(n), \quad b > F_i(r).$$

It follows from Lemma A.1.1 that

$$F_i(tn + (1-t)r) < ta + (1-t)b$$

for any  $i \geq j$ . Since  $F_i$  is decreasing in  $i$ , we conclude (A.1).  $\square$

def:convexclosure

**Definition A.1.6** Let  $F \in \text{Conv}(N)$ . The *closure*  $\text{cl } F \in \text{Conv}(N)$  of  $F$  is defined as follows: if  $F(n) = -\infty$  for some  $n \in N$ , then  $\text{cl } F := -\infty$ . Otherwise, we define  $\text{cl } F$  as the lower semicontinuity regularization of  $F$ .

A convex function  $F \in \text{Conv}(N)$  is *closed* if  $F = \text{cl } F$ . In other words,  $F \in \text{Conv}(N)$  if one of the following conditions hold:

- (1)  $F \equiv -\infty$ ;
- (2)  $F \equiv \infty$ ;
- (3)  $F$  is proper and lower semi-continuous.

**Proposition A.1.3** Let  $F \in \text{Conv}(N)$  be a closed convex function. Then  $F$  is the supremum of all affine functions lying below  $F$ .

See [Roc70, Theorem 12.1].

**Theorem A.1.1** Let  $F \in \text{Conv}^{\text{prop}}(N)$ . Then  $\text{cl } F$  is a closed proper convex function. Moreover,  $\text{cl } F$  agrees with  $F$  except possibly on the relative boundary of  $\text{Dom } F$ .

See [Roc70, Theorem 7.4].

def:partialorderconv

**Definition A.1.7** Given  $F, F' \in \text{Conv}(N)$ , we write  $F \leq F'$  if there is  $C \in \mathbb{R}$  such that

$$F \leq F' + C.$$

We say  $F \sim F'$  if  $F \leq F'$  and  $F' \leq F$  both hold.

## A.2 Legendre transform

Let  $N$  be a real vector space of finite dimension and  $M$  be the dual vector space. The pairing  $M \times N \rightarrow \mathbb{R}$  will be denoted by  $\langle \bullet, \bullet \rangle$ .

def:Legendregeneral

**Definition A.2.1** Let  $F \in \text{Conv}(N)$  be a convex function. We define the *Legendre transform* of  $F$  as the function  $F^* \in \text{Conv}(M)$ :

$$F^*(m) := \sup_{n \in N} (\langle m, n \rangle - F(n)) = \sup_{n \in \text{RelInt Dom } F} (\langle m, n \rangle - F(n)).$$

The latter equality follows from [\[Roc70, Corollary 12.2.2\]](#).

Recall the well-known Legendre–Fenchel duality [\[Roc70, Theorem 12.2\]](#).

thm:Legendredual

**Theorem A.2.1** Let  $F \in \text{Conv}(N)$ . Then  $F^*$  is a closed convex function. The function  $F^*$  is proper if and only if  $F$  is.

Moreover, we have  $(\text{cl } F)^* = F^*$  and

$$F^{**} = \text{cl } F.$$

ex:suppfundual

**Example A.2.1** Let  $P \subseteq M$  be a closed convex subset. Then

$$\text{Supp}_P^* = \chi_P, \quad \chi_P^* = \text{Supp}_P.$$

See [\[Roc70, Theorem 13.2\]](#).

**Definition A.2.2** Let  $F \in \text{Conv}(N)$  and  $n \in N$ . An element  $m \in M$  is a *subgradient* of  $F$  at  $n$  if

$$F(n') \geq F(n) + \langle n' - n, m \rangle, \quad \forall n' \in N. \quad (\text{A.2})$$

{eq:subgrad}

The set of subgradients of  $F$  at  $n$  is denoted by  $\nabla F(n)$ .

More generally, for any subset  $E \subseteq N$ , we write

$$\nabla F(E) = \bigcup_{n \in E} \nabla F(n).$$

def:convexPorder

**Definition A.2.3** Given  $F, F' \in \text{Conv}(N)$ , we write  $F \leq_P F'$  if

$$\overline{\nabla F(N)} \subseteq \overline{\nabla F'(N)}.$$

We write  $F \sim_P F'$  if  $F \leq_P F'$  and  $F' \leq_P F$ .

**Theorem A.2.2** Suppose that  $F \in \text{Conv}^{\text{prop}}(N)$ . Then the following hold:

- (1) For any  $n \notin \text{Dom } F$ ,  $\nabla F(n) = \emptyset$ ;
- (2) for any  $n \in \text{RelInt Dom } F$ ,  $\nabla F(n) \neq \emptyset$ ; Moreover, for any  $n' \in N$ , we have

$$\partial_{n'} F(n) = \sup \{ \langle n', m \rangle : m \in \nabla F(n) \};$$

- (3) for  $n \in N$ , the set  $\nabla F(n)$  is bounded if and only if  $n \in \text{Int Dom } F$ .

For the proof, we refer to [Roc70, Theorem 23.4].

prop:gradDomFstar

**Proposition A.2.1** *Let  $F \in \text{Conv}^{\text{prop}}(N)$ . Then*

$$\nabla F(N) \subseteq \text{Dom } F^*.$$

*If moreover  $F$  is closed, we have*

$$\text{RelInt Dom } F^* \subseteq \nabla F(N). \quad (\text{A.3})$$

{eq:relintdomFstar}

*In particular, if  $F$  is a proper closed convex function on  $N$ , then*

$$\overline{\nabla F(N)} = \overline{\text{Dom } F^*}.$$

**Proof** Suppose that  $m \in \nabla F(n)$  for some  $n \in N$ , it follows that (A.2) holds. In particular,

$$\langle m, n' \rangle - F(n') \leq \langle m, n \rangle - F(n).$$

It follows that

$$F^*(m) \leq \langle m, n \rangle - F(n) < \infty.$$

(A.3) is proved in [Roc70, Corollary 23.5.1]. For the last assertion, it suffices to observe that  $\overline{\text{RelInt Dom } F^*} = \overline{\text{Dom } F^*}$ .  $\square$

prop:Legendretranssup

**Proposition A.2.2** *Let  $\{F_i\}_{i \in I}$  be a non-empty family in  $\text{Conv}^{\text{prop}}(N)$ . Then*

$$\left( \bigwedge_{i \in I} F_i \right)^* = \sup_{i \in I} F_i^*, \quad \left( \sup_{i \in I} \text{cl } F_i \right)^* = \text{cl } \bigwedge_{i \in I} F_i^*.$$

*If  $I$  is finite and  $\overline{\text{Dom } F_i}$  is independent of the choice of  $i \in I$ , then*

$$\left( \sup_{i \in I} F_i \right)^* = \bigwedge_{i \in I} F_i^*.$$

Recall that  $\wedge$  is defined in Definition A.1.5. See [Roc70, Theorem 16.5] for the proof.

prop:sumLegendre

**Proposition A.2.3** *Let  $F_1, \dots, F_r \in \text{Conv}^{\text{prop}}(N)$  ( $r \in \mathbb{Z}_{>0}$ ). Assume that*

$$\bigcap_{i=1}^r \text{RelInt Dom}(F_i) \neq \emptyset,$$

*then*

$$\left( \sum_{i=1}^r F_i \right)^*(m) = \inf \left\{ \sum_{i=1}^r F_i^*(m_i) : m_1, \dots, m_r \in M, \sum_{i=1}^r m_i = m \right\}.$$

prop:Fsuppchar

**Proposition A.2.4** Let  $P \subseteq M$  be a convex body<sup>1</sup> and  $F \in \text{Conv}^{\text{prop}}(N)$ . The following are equivalent:

- (1)  $F \leq \text{Supp}_P$ ;
- (2)  $\text{Dom } F = N$  and  $F^*|_{M \setminus P} \equiv \infty$ ;
- (3)  $\text{Dom } F = N$  and  $\nabla F(N) \subseteq P$ .

Moreover, under these conditions,

$$F(n) - \text{Supp}_P(n) \leq F(0), \quad \forall n \in N. \quad (\text{A.4})$$

{eq:Fsupequal}

**Proof** (1)  $\implies$  (2). It is clear that  $\text{Dom } F = N$  since  $\text{Dom } \text{Supp}_P = N$ . From  $F \leq \text{Supp}_P$  and [Example A.2.1](#), we know that

$$\chi_P = \text{Supp}_P^* \leq F^*.$$

So it follows.

(2)  $\implies$  (3). This follows from [Proposition A.2.1](#).

(3)  $\implies$  (1). Taken  $n \in N$ , we know that  $F$  is locally Lipschitz [[Roc70](#), [Theorem 10.4](#)], so we can compute

$$\begin{aligned} F(n) - F(0) &= \int_0^1 \frac{d}{dt} \Big|_{t=0} F(tn) dt = \int_0^1 \langle \nabla F(tn), n \rangle dt \\ &\leq \int_0^1 \text{Supp}_P(n) dt = \text{Supp}_P(n). \end{aligned}$$

In particular, (A.4) also follows.  $\square$

### A.3 Classes of convex functions

Let  $N$  be a real vector space of finite dimension and  $M$  be the dual vector space.

We shall fix a convex body  $P \subseteq M$ .

The following classes are introduced in [[BB13](#)].

def:convexPfunctions

**Definition A.3.1** We define the set  $\mathcal{P}(N, P)$  as the set of proper convex functions  $F \in \text{Conv}(N)$  such that  $F \leq \text{Supp}_P$ .

We define the set  $\mathcal{E}^\infty(N, P)$  as the set of closed convex functions  $F \in \text{Conv}(N)$  such that  $F \sim \text{Supp}_P$ .

We define the set  $\mathcal{E}(N, P)$  as follows: suppose that  $\text{Int } P = \emptyset$ , then  $\mathcal{E}(N, P) := \mathcal{P}(N, P)$ ; otherwise, let

$$\mathcal{E}(N, P) = \left\{ F \in \mathcal{P}(N, P) : P = \overline{\nabla F(N)} \right\}.$$

<sup>1</sup> Here a convex body refers to a non-empty closed convex subset, not necessarily having non-empty interior.

Observe that for any  $F \in \mathcal{P}(N, P)$ , we have  $\text{Dom } F = N$  and  $F$  is necessarily closed.

**Proposition A.3.1** *We have*

$$\mathcal{E}^\infty(N, P) \subseteq \mathcal{E}(N, P) \subseteq \mathcal{P}(N, P).$$

**Proof** When  $\text{Int } P = \emptyset$ , the assertion is clear. We assume that  $\text{Int } P \neq \emptyset$ . The second inclusion follows from definition. We only hand the first inequality. Take  $F \in \mathcal{E}^\infty(N, P)$ . By definition,  $F \sim \text{Supp}_P$  and hence  $F^* \sim \chi_P$ . It follows that  $P = \text{Dom } F^*$ .

By [Proposition A.2.4](#), we already know that

$$\nabla F(N) \subseteq P = \text{Dom } F^*.$$

On the other hand, by [Proposition A.2.1](#), we have

$$\text{Int } P \subseteq \nabla F(N).$$

So it follows that

$$P = \overline{\nabla F(N)}.$$

**Proposition A.3.2** *For any  $F \in \mathcal{E}^\infty(N, P)$ , we have  $F^*|_{M \setminus P} \equiv \infty$  and  $F^*$  is bounded on  $P$ .*

**Proof** From  $F \sim \text{Supp}_P$ , we take the Legendre transform to get  $F^* \sim \text{Supp}_P^* = \chi_P$ , where we applied [Example A.2.1](#).  $\square$

**Definition A.3.2** We endow the topology of pointwise convergence on  $\mathcal{P}(N, P)$ . Note that this topology coincides with the compact-open topology.

**Proposition A.3.3** *Let  $F \in \mathcal{P}(N, P)$ . Then there is a decreasing sequence  $F_j \in \mathcal{E}^\infty(N, P) \cap C^\infty(N)$  converging to  $F$ .*

See [\[BB13, Lemma 2.2\]](#).

We observe that the point  $0 \in N$  plays a special role since it does in the definition of the support function.

**Proposition A.3.4** *For any  $F \in \text{Conv}(N, P)$ , we have*

$$\max_N (F - \text{Supp}_P) = F(0).$$

**Proof** It follows from [\(A.4\)](#) that

$$\sup_N (F - \text{Supp}_P) \leq F(0).$$

The equality is clearly obtained at  $0 \in N$ .  $\square$

## A.4 Monge–Ampère measures

Let  $N$  be a free Abelian group of finite rank (i.e. a lattice) and  $M$  be its dual lattice. There is a canonical Lebesgue type measure on  $M_{\mathbb{R}}$ , denoted by  $\mathrm{d vol}$ , normalized so that the smallest cubes in  $M$  have volume 1. Similarly, the canonical measure on  $N_{\mathbb{R}}$  is normalized in the same way and is denoted by  $\mathrm{d vol}$  as well.

We will write

$$N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}, \quad M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}.$$

**Definition A.4.1** Let  $F \in \mathrm{Conv}(N_{\mathbb{R}})$ , we define  $\mathrm{MA}_{\mathbb{R}} F$  as the Borel measure on  $N_{\mathbb{R}}$  given as follows: for each Borel measurable set  $E \subseteq N_{\mathbb{R}}$ , define

$$\mathrm{MA}_{\mathbb{R}} F(E) := n! \int_{\nabla F(E)} \mathrm{d vol}.$$

**Proposition A.4.1** Let  $P \in M_{\mathbb{R}}$  be a convex body and  $F \in \mathcal{P}(N_{\mathbb{R}}, P)$ . Then  $F \in \mathcal{E}(N_{\mathbb{R}}, P)$  if and only if

$$\int_{M_{\mathbb{R}}} \mathrm{MA}_{\mathbb{R}} F = n! \mathrm{vol} P. \quad (\text{A.5}) \quad \boxed{\text{\{eq:cvxfullmass\}}}$$

*Proof* By definition of  $\mathrm{MA}_{\mathbb{R}}$ , (A.5) is equivalent to

$$\mathrm{vol} \overline{\nabla F(N_{\mathbb{R}})} = \mathrm{vol} P.$$

We first handle the case where  $\mathrm{Int} P \neq \emptyset$ . By [Proposition A.2.4](#), the latter is equivalent to

$$\overline{\nabla F(N_{\mathbb{R}})} = P.$$

Now assume that  $\mathrm{Int} P = \emptyset$ , then  $\mathrm{vol} \overline{\nabla F(N_{\mathbb{R}})} = \mathrm{vol} P = 0$  by [Proposition A.2.4](#). The assertion is clear.  $\square$

thm:realMAcont

**Theorem A.4.1** Let  $F, F_j \in \mathcal{P}(N_{\mathbb{R}}, P)$  ( $j \in \mathbb{Z}_{>0}$ ). Assume that  $F_j \rightarrow F$ , then  $\mathrm{MA}_{\mathbb{R}}(F_j)$  converges to  $\mathrm{MA}_{\mathbb{R}}(F)$  weakly.

See [Fig17](#), Proposition 2.6].

There is a well-known comparison principle.

thm:convcomp

**Theorem A.4.2** Let  $F, F' \in \mathcal{P}(N_{\mathbb{R}}, P)$ . Assume that  $F \leq F'$ , then

$$\overline{\nabla F(N_{\mathbb{R}})} \subseteq \overline{\nabla F'(N_{\mathbb{R}})}.$$

$$\int_{N_{\mathbb{R}}} \mathrm{MA}_{\mathbb{R}}(F) \leq \int_{N_{\mathbb{R}}} \mathrm{MA}_{\mathbb{R}}(F').$$

See [BB13](#), Lemma 2.5].



## A.5 Separation lemmata

lma:polybdd

**Lemma A.5.1** Let  $\alpha, \beta_1, \dots, \beta_m \in \mathbb{Z}^n$ . Let  $\Delta$  be the polytope generated by  $\beta_1, \dots, \beta_m$ . Then the following are equivalent:

(1)

$$|z^\alpha|^2 \left( \sum_{i=1}^m |z^{\beta_i}|^2 \right)^{-1} \quad (\text{A.6})$$

{eq:zalpha}

is a bounded function on  $\mathbb{C}^{*n}$ .

(2)  $\alpha \in \Delta$ .

**Proof** (2)  $\implies$  (1). Write  $\alpha = \sum_i t_i \beta_i$ , where  $t_i \in [0, 1]$ ,  $\sum_i t_i = 1$ . Then

$$\begin{aligned} |z^\alpha|^2 \left( \sum_{i=1}^m |z^{\beta_i}|^2 \right)^{-1} &= \prod_i |z^{\beta_i}|^{2t_i} \left( \sum_{i=1}^m |z^{\beta_i}|^2 \right)^{-1} \\ &\leq \prod_i \sum_j |z^{\beta_j}|^{2t_i} \left( \sum_{i=1}^m |z^{\beta_i}|^2 \right)^{-1} \leq 1. \end{aligned}$$

(1)  $\implies$  (2). Assume that  $\alpha \notin \Delta$ . Let  $H$  be a hyperplane that separates  $\alpha$  and  $\Delta$ . Say  $H$  is defined by  $a_1 x_1 + \dots + a_n x_n = C$ . Set

$$z(t) := (t^{a_1}, \dots, t^{a_n}).$$

Then clearly (A.6) evaluated at  $z(t)$  is not bounded.  $\square$

lma:polybdd2

**Lemma A.5.2** Let  $\beta_1, \dots, \beta_m \in \mathbb{N}^n$  and  $\beta \in \mathbb{R}^n$ . Then the following are equivalent

(1)  $\log \sum_{i=1}^m e^{x \cdot \beta_i} - (x, \beta)$  is bounded from below.(2)  $\beta$  is in the convex hull of the  $\beta_i$ 's.

**Proof** The proof follows the same pattern as Lemma A.5.1.  $\square$



## Appendix B

### Pluripotential theory on unibranch spaces

chap:unib

In this appendix, we extend the theory in the book to compact unibranch Kähler spaces.

#### B.1 Complex spaces

A complex space is assumed to be reduced, Hausdorff and paracompact in the whole book.

def:primdiv

**Definition B.1.1** A *prime divisor* over an irreducible complex space  $Z$  is a connected smooth hypersurface  $E \subseteq X'$ , where  $X' \rightarrow Z$  is a proper bimeromorphic morphism with  $X'$  smooth. Such a morphism  $X' \rightarrow Z$  is also called a *resolution* of  $Z$ .

Two prime divisors  $E_1 \subseteq X'_1$  and  $E_2 \subseteq X'_2$  over  $Z$  are *equivalent* if there is a common resolution  $X'' \rightarrow X$  dominating both  $X'_1$  and  $X'_2$  such that the strict transforms of  $E_1$  and  $E_2$  coincide.

The set  $Z^{\text{div}}$  is the set of pairs  $(c, E)$ , where  $c \in \mathbb{Q}_{>0}$  and  $E$  is an equivalence class of a prime divisor over  $Z$ . For simplicity, we will denote the pair  $(c, E)$  by  $c \text{ ord}_E$ , although one should not really think of this object as a valuation unless  $Z$  is projective and irreducible.

Note that a prime divisor on  $Z$  does not always define a prime divisor over  $Z$  if  $Z$  is singular.

**Definition B.1.2** A complex space  $X$  is *unibranch* if for all  $x \in X$ , the local ring  $\mathcal{O}_{X,x}$  is unibranch.

It is shown in the arXiv version of [\[Xia23Mabuchi\]](#), Remark 2.7] that when  $X$  is a projective variety, this notion coincides with the corresponding algebraic notion of unibranchness.

thm:Zariskimain

**Theorem B.1.1 (Zariski's main theorem)** Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism between complex spaces. Assume that  $X$  is unibranch, then  $\pi$  has connected fibers.

We refer to [\[Dem85\]](#), Proof of Théorème 1.7].

def:modif

**Definition B.1.3** A *modification* of a compact complex space  $X$  is a finite composition of blow-ups with smooth centers.

thm:HironakaChow

**Theorem B.1.2 (Hironaka's Chow lemma)** *Suppose that  $X$  is a compact complex space. Then every proper bimeromorphic morphism to  $X$  can be dominated by a modification.*

This follows from the proof of [\[Hir75\]](#), Corollary 2].

thm:res

**Theorem B.1.3** *Let  $X$  be a compact complex space. Then there is a modification  $\pi: Y \rightarrow X$  such that  $Y$  is smooth.*

See [\[BM97, Wlo09\]](#).

cor:primerealization

**Corollary B.1.1** *Let  $X$  be a compact complex space and  $E$  be a prime divisor over  $X$ . Then there is a modification  $\pi: Y \rightarrow X$  such that  $Y$  is smooth and  $E$  can be realized as a prime divisor on  $Y$ .*

## B.2 Plurisubharmonic functions

Let  $X$  be a complex space.

**Definition B.2.1** A function  $\varphi: X \rightarrow [-\infty, \infty)$  is *plurisubharmonic* if

- (1)  $\varphi$  is not identically  $-\infty$  on any irreducible component of  $X$ , and
- (2) for any  $x \in X$ , there is an open neighbourhood  $V$  of  $x$  in  $X$ , a domain  $\Omega \subseteq \mathbb{C}^N$ , a closed immersion  $V \hookrightarrow \Omega$  and a plurisubharmonic function  $\tilde{\varphi} \in \text{PSH}(\Omega)$  such that  $\varphi|_{\Omega \cap V} = \tilde{\varphi}|_{\Omega \cap V}$ .

The set of plurisubharmonic functions on  $X$  is denoted by  $\text{PSH}(X)$ .

Similarly, if  $\theta$  is a smooth closed<sup>1</sup> real  $(1, 1)$ -form on  $X$ , then a function  $\varphi: X \rightarrow [-\infty, \infty)$  is  *$\theta$ -plurisubharmonic* if for any  $x \in X$ , there is an open neighbourhood  $V$  of  $x$  in  $X$ , a domain  $\Omega \subseteq \mathbb{C}^N$ , a closed immersion  $V \hookrightarrow \Omega$  and a smooth function  $g$  on  $\Omega$  such that  $\theta = (\text{dd}^c g)|_{V \cap \Omega}$  and  $g + \varphi|_V \in \text{PSH}(V)$ .

thm:FN

**Theorem B.2.1 (Fornaess–Narasimhan)** *Let  $\varphi: X \rightarrow [-\infty, \infty)$  be a function. Assume that  $\varphi$  is not identically  $-\infty$  on any irreducible component of  $X$ , then the following are equivalent:*

- (1)  $\varphi$  is psh;
- (2)  $\varphi$  is usc and for any morphism  $f: \Delta \rightarrow X$  from the open unit disk  $\Delta$  in  $\mathbb{C}$  to  $X$  such that  $f^* \varphi$  is not identically  $-\infty$ , the pull-back  $f^* \varphi$  is psh.

*If further more  $X$  is unibranch, then these conditions are equivalent to*

<sup>1</sup> Here *closed* means that locally  $\theta$  is defined by a closed form under a local embedding.

(3)  $\varphi \in \text{PSH}(X^{\text{Reg}})$ , locally bounded from above near  $X^{\text{Sing}}$  and  $\varphi = \varphi^*$ .

See [\[FN80\]](#) and [\[Dem85, Section 1.8\]](#).

cor:PSH

**Corollary B.2.1** *Let  $\pi: Y \rightarrow X$  be a proper bimeromorphic morphism between compact Kähler spaces. Let  $\theta$  be a smooth closed real  $(1, 1)$ -form on  $X$ . Assume that  $X$  is unibranch, then the pull-back induces a bijection*

$$\pi^*: \text{PSH}(X, \theta) \xrightarrow{\sim} \text{PSH}(Y, \pi^*\theta).$$

See [\[Dem85, Théorème 1.7\]](#) for the details.

thm:GRekten2

**Theorem B.2.2 (Grauert–Remmert)** *Let  $X$  be a unibranch complex space and  $Z$  be an analytic subset in  $X$  and  $\varphi \in \text{PSH}(X \setminus Z)$ . Then the function  $\varphi$  admits an extension to  $\text{PSH}(X)$  in the following two cases:*

- (1) *The set  $Z$  has codimension at least 2 everywhere.*
- (2) *The set  $Z$  has codimension at least 1 everywhere and is locally bounded from above on an open neighbourhood of  $Z$ .*

*In both cases, the extension is unique and is given by*

$$\varphi(x) = \overline{\lim_{X \setminus Z \ni y \rightarrow x}} \varphi(y), \quad x \in X. \quad (\text{B.1})$$

{eq:GRextvarphi}

**Proof** The problem is local in natural. By the local description of complex spaces [\[GR84, Section 3.4\]](#), we may assume that there is a domain  $\Omega \subseteq \mathbb{C}^n$ , a finite  $s$ -sheet branched covering  $\Phi: X \rightarrow \Omega$  with branched locus contained in a proper analytic subset  $V \subseteq \Omega$ . We may assume that  $X$  is connected,  $n \geq 1$  and  $Z \subseteq \Phi^{-1}(V)$ .

We first prove the uniqueness in both cases. For this purpose, we may assume that  $Z = \Phi^{-1}(V)$ . Fix  $z \in Z$ , we can find a complex line  $L$  passing through  $\Phi(z)$  such that  $L \cap V \cap B = \{\Phi(z)\}$ , where  $B$  is a small open ball centered at  $\Phi(z)$ . After shrinking  $\Omega$ , we may choose one isomorphic copy  $L'$  of  $L \cap B \setminus \{z\}$  in a neighbourhood of  $z$ . Since  $\varphi$  restricts to a subharmonic function on  $L' \cap \{z\}$ , it follows that the value of  $\varphi(z)$  is uniquely determined.

(2) Let  $\psi$  be the function defined in (B.1). We claim that  $\psi \in \text{PSH}(X)$ . Since  $\psi$  clearly extends  $\varphi$ , so our assertion is proved.

Let  $f: \Delta \rightarrow X$  be a morphism. Due to [Theorem B.2.1](#), we only need to show that  $f^*\psi$  is subharmonic. We may assume that  $f$  is non-constant, so that  $\Phi \circ f$  has full rank outside a discrete subset  $S' \subseteq \Delta$ .

**Step 1.** We show that after enlarging  $S'$  to a larger discrete subset,  $f^*\psi$  is subharmonic outside  $S'$ . We may assume that  $0 \notin S'$  and it suffices to show that  $f^*\psi$  is subharmonic near 0 outside a discrete subset.

For this purpose, after shrinking  $\Delta$ , we may assume that  $\Phi \circ f$  has full rank everywhere. After shrinking  $\Omega$  and  $\Delta$ , we may furthermore assume that

- (1)  $A = \Phi \circ f(\Delta)$  is an analytic subset of  $\Omega$  of dimension 1, and
- (2)  $f(0)$  is the only preimage of  $\Phi(f(0))$  with respect to  $\Phi$ .

Thanks to the first condition, we may then find a discrete subset  $S'' \subseteq A$  such that  $\Phi$  restricts to an unbranched covering on  $A \setminus S''$ .

Now it would suffice to show that

$$\psi \in \text{PSH}(\Phi^{-1}(A \setminus S'')). \quad (\text{B.2})$$

{eq:psipshtemp1}

Let  $x \in A \setminus S''$ . After further shrinking  $\Omega$  around  $x$  (and replacing  $X$  by the corresponding connected component), we may assume that each point in  $A \setminus S''$  has exactly one preimage in  $X$ . By an elementary argument (see [GR56, Hilfssatz 6]), the fibral integration  $\Phi_*\psi \in \text{PSH}(\Omega)$  and (B.2) follows.

**Step 2.** We show that  $f^*\psi$  is subharmonic near  $S'$ . Let  $z \in S'$ , it suffices to show that  $f^*\psi$  is subharmonic in an open neighbourhood of  $z$ .

After shrinking  $\Phi$  along  $\Phi \circ f(z)$ , we may assume that  $X$  is connected and  $\Phi^{-1}(\Phi \circ f(z))$  consists only of  $f(z)$ . Let  $\eta \in \text{PSH}(\Omega)$  be the fibral integration of  $\psi$  along  $\Phi$ . Then  $f^*\Phi^*\eta \in \text{SH}(\Delta)$  and

$$\overline{\lim}_{w \rightarrow z} \frac{1}{s} f^*\Phi^*\eta(w) = f^*\psi(z).$$

Assume that

$$\overline{\lim}_{w \rightarrow z} f^*\varphi(w) < f^*\psi(z),$$

then

$$\overline{\lim}_{w \rightarrow z} \frac{1}{s} f^*\Phi^*\eta(w) \leq \frac{1}{s} \overline{\lim}_{w \rightarrow z} f^*\varphi(w) + \frac{s-1}{s} f^*\psi(z) < f^*\psi(z),$$

which is a contradiction. It follows that

$$f^*\psi = (f^*\psi)^* \in \text{SH}(\Delta).$$

(1) It suffices to show that  $\varphi$  is locally bounded near  $Z$ . Suppose that this fails. Then by (2) we can find  $z \in Z$  and  $x_i \in X \setminus (Z \cup V)$  ( $i \geq 1$ ) such that

$$\lim_{i \rightarrow \infty} \varphi(x_i) = \infty.$$

Let  $L$  be a complex line passing through  $\Phi(z)$  intersecting  $(\Phi(Z) \cup V) \cap B$  only at  $\Phi(z)$ , where  $B \Subset B'$  are two small open balls centered at  $\Phi(z)$ . We can find a sequence of lines  $L_i$  passing through  $\Phi(x_i)$  converging to  $L$  such that  $L_i \cap (B' \cap \Phi(Z)) = \emptyset$  while  $L_i \cap (B' \cap V)$  is discrete. The  $\Phi$  restricts to a branched covering over  $B' \cap L_i$  for all  $i \geq 1$ . Adding a constant to  $\varphi$ , we may assume that  $\varphi|_{\Phi^{-1}(L \cap \partial B)} < 0$ . We can then find an open neighbourhood  $U$  of  $\Phi^{-1}(L \cap \partial B)$  so that  $\varphi|_U < 0$ . For large  $i$  we have  $\Phi^{-1}(L_i \cap \partial B) \subseteq U$ , it follows from the maximum principle that  $\varphi(x_i) \leq 0$ , which is a contradiction.  $\square$

### B.3 Extensions of the results in the smooth setting

Let  $X$  be an irreducible unibranch compact Kähler space of dimension  $n$ . Let  $\theta$  be a closed real smooth  $(1, 1)$ -form on  $X$ . We say *the cohomology class*  $[\theta]$  is big if for any proper bimeromorphic morphism  $\pi: Y \rightarrow X$  from a compact Kähler manifold  $Y$ ,  $[\pi^*\theta]$  is big.

The non-pluripolar products can be defined exactly as in [Chapter 2](#) and the results in that chapter holds *mutadis mutandis*.

The results in [Chapter 3](#) can be also be easily extended. The definition of the  $P$ -envelope remains unchanged. As for the  $\mathcal{I}$ -envelope, we define

**Definition B.3.1** Given  $\varphi \in \text{PSH}(X, \theta)$ , we define  $P_\theta[\varphi]_{\mathcal{I}} \in \text{PSH}(X, \theta)$  as the unique element with the following property: if  $\pi: Y \rightarrow X$  is a proper bimeromorphic morphism from a compact Kähler manifold  $Y$ , then

$$\pi^*P_\theta[\varphi]_{\mathcal{I}} = P_{\pi^*\theta}[\pi^*\varphi]_{\mathcal{I}}.$$

It follows from [Corollary B.2.1](#) and [Proposition 3.2.5](#) that  $P_\theta[\varphi]_{\mathcal{I}}$  is independent of the choice of  $\pi$  and is well-defined. The other results can be easily extended.

[Chapter 4](#) and [Chapter 6](#) can be extended without big changes. The only exception is [Theorem 6.2.6](#), where we do not have the notion of multiplier ideal sheaves. So we do not know how to extend this theorem.

[Chapter 7](#) can be extended except for [Section 7.3](#) for the same reason as above.

The trace operator defined in [Chapter 8](#) can be extended as long as  $Y$  is not contained in  $X^{\text{Sing}}$  using the embedded resolution. In general, due to the lack of Demailly regularization, we do not know how to define the trace operator.

[Chapter 9](#) can be extended easily.

[Chapter 10](#) is easy to extend since the partial Okounkov bodies are bimeromorphically invariant in the sense of [Theorem 10.3.2](#).

[Chapter 11](#) is unchanged, since we always take projective limits with respect to all models in that section.

[Chapter 13](#) can be extended except for the parts involving the trace operator.

[Chapter 14](#) can be easily extended by considering a resolution.

I do not know how to extend the results in [Chapter 5](#) and [Chapter 12](#) to the singular setting.





## Appendix C

### Almost semigroups

chap:almostsg

We introduce and study almost semigroups. In particular, we will define the Okounkov bodies of almost semigroups.

#### C.1 Convex bodies

Fix  $n \in \mathbb{N}$ .

def:convbodies

**Definition C.1.1** A *convex body* in  $\mathbb{R}^n$  is a non-empty compact convex set.

We allow a convex body to have empty interior.

We write  $\mathcal{K}_n$  for the set of convex bodies in  $\mathbb{R}^n$ .

def:Hausdorffmetric

**Definition C.1.2** The *Hausdorff metric* between  $K_1, K_2 \in \mathcal{K}_n$  is given by

$$d_{\text{Haus}}(K_1, K_2) := \max \left\{ \sup_{x_1 \in K_1} \inf_{x_2 \in K_2} |x_1 - x_2|, \sup_{x_2 \in K_2} \inf_{x_1 \in K_1} |x_1 - x_2| \right\}.$$

It is well-known that the metric space  $(\mathcal{K}_n, d_{\text{Haus}})$  is complete. We will need the following fundamental theorem:

thm:Blaschke

**Theorem C.1.1 (Blaschke selection theorem)** *The metric space  $(\mathcal{K}_n, d_{\text{Haus}})$  is locally compact.*

We refer to [Sch14, Theorem 1.8.7] for details.

thm:contvol

**Theorem C.1.2** *The Lebesgue volume  $\text{vol}: \mathcal{K}_n \rightarrow \mathbb{R}_{\geq 0}$  is continuous.*

See [Sch14, Theorem 1.8.20].

thm:Hausconvcond

**Theorem C.1.3** *Let  $K_i, K \in \mathcal{K}_n$  ( $i \in \mathbb{N}$ ). Then  $K_i \xrightarrow{d_{\text{Haus}}} K$  if and only if the following conditions hold:*

- (1) *each point  $x \in K$  is the limit of a sequence  $x_i \in K_i$ , and*

(2) the limit of any convergent sequence  $(x_{i_j})_{j \in \mathbb{N}}$  with  $x_{i_j} \in K_{i_j}$  lies in  $K$ , where  $i_j$  is a strictly increasing sequence in  $\mathbb{Z}_{>0}$ .

See [Sch14, Theorem 1.8.8].

lma:latcvb

**Lemma C.1.1** Let  $K \in \mathcal{K}_n$  be a convex body with positive volume and  $K' \in \mathcal{K}_n$ . Assume that for some large enough  $k \in \mathbb{Z}_{>0}$ ,  $K'$  contains  $K \cap (k^{-1}\mathbb{Z})^n$ , then  $K' \supseteq K^{n^{1/2}k^{-1}}$ .

**Proof** Let  $x \in K^{n^{1/2}k^{-1}}$ , by assumption, the closed ball  $B$  with center  $x$  and radius  $n^{1/2}k^{-1}$  is contained in  $K$ . Observe that  $x$  can be written as a convex combination of points in  $B \cap (k^{-1}\mathbb{Z})^n$ , which are contained in  $K'$  by assumption. It follows that  $x \in K'$ .  $\square$

Given a sequence of convex bodies  $K_i$  ( $i \in \mathbb{N}$ ), we set

$$\varliminf_{i \rightarrow \infty} K_i = \bigcup_{i=0}^{\infty} \bigcap_{j \geq i} K_j.$$

Suppose  $K$  is the limit of a subsequence of  $K_i$ , we have

$$\varliminf_{i \rightarrow \infty} K_i \subseteq K. \quad (\text{C.1}) \quad \{\text{eq:liminflimsup}\}$$

This is a simple consequence of [Theorem C.1.3](#).

lma:Hausdorffconvslice

**Lemma C.1.2** Let  $K \subseteq \mathbb{R}^n$  be a convex body. Let

$$t_{\min} := \min\{t \in \mathbb{R} : \{x_1 = t\} \cap K \neq \emptyset\}, \quad t_{\max} := \max\{t \in \mathbb{R} : \{x_1 = t\} \cap K \neq \emptyset\}.$$

Then for  $t \in [t_{\min}, t_{\max}]$ , the map

$$t \mapsto \{x_1 = t\} \cap K$$

is continuous with respect to the Hausdorff metric.

Here  $x_1$  denotes the first coordinate in  $\mathbb{R}^n$ .

**Proof** We may assume that  $t_{\min} < t_{\max}$  as otherwise there is nothing to prove.

For each  $t \in [t_{\min}, t_{\max}]$ , we write  $K_t = \{x_1 = t\} \cap K$ . Let  $t_j \rightarrow t$  be a convergent sequence in  $[t_{\min}, t_{\max}]$ , we want to show that  $K_{t_j}$  converges to  $K_t$  with respect to the Hausdorff metric. Recall that this amounts to the following two assertions:

- (1) For each convergent sequence  $x_j \in K_{t_j}$  with limit  $x$ , we have  $x \in K_t$ ;
- (2) Given any  $x \in K_t$ , up to replacing  $t_j$  by a subsequence, we can find  $x_j \in K_{t_j}$  converging to  $x$ .  $\square$

The first assertion is obvious. Let us prove the second. Take  $x = (t, x') \in K_t$ . Up to replacing  $t_j$  by a subsequence and taking the symmetry into account, we may assume that  $t_j > t$  for all  $t$ . In particular,  $t < t_{\max}$ .

We can find a point  $y = (y^1, y') \in K$  such that  $y^1 > t$  (for example, there is always such a point with  $y^1 = t_{\max}$ ). Replacing  $t_j$  by a subsequence, we may assume that  $t_j \in (t, y^1)$  for all  $j$ . Then it suffices to take

$$x_j = \frac{y^1 - t_j}{y^1 - t} x + \frac{t_j - t}{y^1 - t} y.$$

lma:intconvexset

**Lemma C.1.3** *Let  $D_j \subseteq \mathbb{R}^n$  ( $j \geq 1$ ) be a decreasing sequence of convex sets. Assume that  $\text{vol} \bigcap_j D_j > 0$ , then*

$$\overline{\bigcap_{j=1}^{\infty} D_j} = \bigcap_{j=1}^{\infty} \overline{D_j}.$$

**Proof** The  $\subseteq$  direction is clear. By convexity, it suffices to show that both sides have the same positive volume. As the boundary of convex sets has zero Lebesgue measure, it follows that the volumes of both sides are equal to  $\lim_{j \rightarrow \infty} \text{vol } D_j$ .  $\square$

**Definition C.1.3** Let  $K, K' \in \mathcal{K}_n$ , their *Minkowski sum* is given by

$$K + K' := \{x + x' : x \in K, x' \in K'\}.$$

**Proposition C.1.1** *The Minkowski sum  $\mathcal{K}_n \times \mathcal{K}_n \rightarrow \mathcal{K}_n$  is continuous.*

See [Sch14, Sch93, Page 139].

thm:BrunnMin

**Theorem C.1.4 (Brunn–Minkowski)** *Let  $K, K' \in \mathcal{K}_n$ , then for any  $t \in (0, 1)$ , we have*

$$\text{vol}((1-t)K' + tK) \geq (\text{vol } K')^{(1-t)} (\text{vol } K)^t.$$

In other words, the volume is log concave. See [Sch14, Sch93, Page 372].

## C.2 The Okounkov bodies of almost semigroups

sec:clo

Fix an integer  $n \geq 0$ . Fix a closed convex cone  $C \subseteq \mathbb{R}^n \times \mathbb{R}_{\geq 0}$  such that  $C \cap \{x_{n+1} = 0\} = \{0\}$ . Here  $x_{n+1}$  is the last coordinate of  $\mathbb{R}^{n+1}$ .

### C.2.1 Generalities on semigroups

Write  $\hat{\mathcal{S}}(C)$  for the set of subsets of  $C \cap \mathbb{Z}^{n+1}$  and  $\mathcal{S}(C)$  for the set of sub-semigroups  $S \subseteq C \cap \mathbb{Z}^{n+1}$ . For each  $k \in \mathbb{N}$  and  $S \in \hat{\mathcal{S}}(C)$ , we write

$$S_k := \{x \in \mathbb{Z}^n : (x, k) \in S\}.$$

Note that  $S_k$  is a finite set by our assumption on  $C$ .

We introduce a pseudometric on  $\hat{\mathcal{S}}(C)$  as follows:

$$d_{\text{sg}}(S, S') := \overline{\lim}_{k \rightarrow \infty} k^{-n} (|S_k| + |S'_k| - 2|(S \cap S')_k|).$$

Here  $|\bullet|$  denotes the cardinality of a finite set.

lma:dps

**Lemma C.2.1** *The above defined  $d_{\text{sg}}$  is a pseudometric on  $\hat{\mathcal{S}}(C)$ .*

**Proof** Only the triangle inequality needs to be argued. Take  $S, S', S'' \in \hat{\mathcal{S}}(C)$ . We claim that for any  $k \in \mathbb{N}$ ,

$$|S_k| + |S'_k| - 2|S_k \cap S'_k| + |S''_k| + |S'_k| - 2|S''_k \cap S'_k| \geq |S_k| + |S''_k| - 2|S_k \cap S''_k|.$$

From this the triangle inequality follows. To argue the claim, we rearrange it to the following form:

$$|S'_k| - |S_k \cap S'_k| \geq |S'_k \cap S''_k| - |S_k \cap S''_k|,$$

which is obvious.  $\square$

Given  $S, S' \in \hat{\mathcal{S}}(C)$ , we say  $S$  is equivalent to  $S'$  and write  $S \sim S'$  if  $d_{\text{sg}}(S, S') = 0$ . This is an equivalence relation by [Lemma C.2.1](#).

lma:dBi1

**Lemma C.2.2** *Given  $S, S', S'' \in \hat{\mathcal{S}}(C)$ , we have*

$$d_{\text{sg}}(S \cap S'', S' \cap S'') \leq d_{\text{sg}}(S, S').$$

*In particular, if  $S^i, S'^i \in \hat{\mathcal{S}}(C)$  ( $i \in \mathbb{N}$ ) and  $S^i \rightarrow S, S'^i \rightarrow S'$ , then*

$$S^i \cap S'^i \rightarrow S \cap S'.$$

**Proof** Observe that for any  $k \in \mathbb{N}$ ,

$$|S_k \cap S''_k| - |S_k \cap S'_k \cap S''_k| \leq |S_k| - |S_k \cap S'_k|.$$

The same holds if we interchange  $S$  with  $S'$ . It follows that

$$|S_k \cap S''_k| + |S'_k \cap S''_k| - 2|S_k \cap S'_k \cap S''_k| \leq |S_k| + |S'_k| - 2|S_k \cap S'_k|.$$

The first assertion follows.

Next we compute

$$\begin{aligned} d_{\text{sg}}(S^i \cap S'^i, S \cap S') &\leq d_{\text{sg}}(S^i \cap S'^i, S^i \cap S') + d_{\text{sg}}(S^i \cap S', S \cap S') \\ &\leq d_{\text{sg}}(S'^i, S') + d_{\text{sg}}(S^i, S) \end{aligned}$$

and the second assertion follows.  $\square$

The volume of  $S \in \mathcal{S}(C)$  is defined as

$$\text{vol } S := \lim_{k \rightarrow \infty} (ka)^{-n} |S_{ka}| = \overline{\lim}_{k \rightarrow \infty} k^{-n} |S_k|,$$

where  $a$  is a sufficiently divisible positive integer. The existence of the limit and its independence from  $a$  both follow from the more precise result [KK12, Theorem 2].

lma:volliip

**Lemma C.2.3** *Let  $S, S' \in \mathcal{S}(C)$ , then*

$$|\operatorname{vol} S - \operatorname{vol} S'| \leq d_{\text{sg}}(S, S').$$

**Proof** By definition, we have

$$d_{\text{sg}}(S, S') \geq \operatorname{vol} S + \operatorname{vol} S' - 2 \operatorname{vol}(S \cap S').$$

It follows that  $\operatorname{vol} S - \operatorname{vol} S' \leq d_{\text{sg}}(S, S')$  and  $\operatorname{vol} S' - \operatorname{vol} S \leq d_{\text{sg}}(S, S')$ .  $\square$

We define  $\overline{\mathcal{S}}(C)$  as the closure of  $\mathcal{S}(C)$  in  $\hat{\mathcal{S}}(C)$  with respect to the topology defined by the pseudometric  $d$ . By Lemma C.2.3,  $\operatorname{vol}: \mathcal{S}(C) \rightarrow \mathbb{R}$  admits a unique 1-Lipschitz extension to

$$\operatorname{vol}: \overline{\mathcal{S}}(C) \rightarrow \mathbb{R}. \quad (\text{C.2})$$

{eq:volex}

lma:volcompa

**Lemma C.2.4** *Suppose that  $S, S' \in \overline{\mathcal{S}}(C)$  and  $S \subseteq S'$ . Then*

$$\operatorname{vol} S \leq \operatorname{vol} S'.$$

**Proof** Take sequences  $S^j, S'^j$  in  $\mathcal{S}(C)$  such that  $S^j \rightarrow S, S'^j \rightarrow S'$ . By Lemma C.2.2, after replacing  $S^j$  by  $S^j \cap S'^j$ , we may assume that  $S^j \subseteq S'^j$  for each  $j$ . Then our assertion follows easily.  $\square$

## C.2.2 Okounkov bodies of semigroups

Given  $S \in \hat{\mathcal{S}}(C)$ , we will write  $C(S) \subseteq C$  for the closed convex cone generated by  $S \cup \{0\}$ . Moreover, for each  $k \in \mathbb{Z}_{>0}$ , we define

$$\Delta_k(S) := \operatorname{Conv} \{k^{-1}x \in \mathbb{R}^n : x \in S_k\} \subseteq \mathbb{R}^n.$$

Here  $\operatorname{Conv}$  denotes the convex hull.

**Definition C.2.1** Let  $\mathcal{S}'(C)$  be the subset of  $\mathcal{S}(C)$  consisting of semigroups  $S$  such that  $S$  generates  $\mathbb{Z}^{n+1}$  (as an Abelian group).

Note that for any  $S \in \mathcal{S}'(C)$ , the cone  $C(S)$  has full dimension (i.e. the topological interior is non-empty). Given a full-dimensional subcone  $C' \subseteq C$ , it is clear that  $C' \cap \mathbb{Z}^{n+1} \in \mathcal{S}'(C)$ .

This class behaves well under intersections:

lma:intersecS'

**Lemma C.2.5** *Let  $S, S' \in \mathcal{S}'(C)$ . Assume that  $\operatorname{vol}(S \cap S') > 0$ , then  $S \cap S' \in \mathcal{S}'(C)$ .*

The lemma obviously fails if  $\operatorname{vol}(S \cap S') = 0$ .

**Proof** We first observe that the cone  $C(S) \cap C(S')$  has full dimension since otherwise  $\text{vol}(S \cap S') = 0$ . Take a full-dimensional subcone  $C'$  in  $C(S) \cap C(S')$  such that  $C'$  intersects the boundary of  $C(S) \cap C(S')$  only at 0. It follows from [KK12, Theorem 1] that there is an integer  $N > 0$  such that for any  $x \in \mathbb{Z}^{n+1} \cap C'$  with Euclidean norm no less than  $N$  lies in  $S \cap S'$ . Therefore,  $S \cap S' \in \mathcal{S}'(C)$ .  $\square$

We recall the following definition from [KK12].

def:Okokk

**Definition C.2.2** Given  $S \in \mathcal{S}'(C)$ , its *Okounkov body* is defined as follows

$$\Delta(S) := \{x \in \mathbb{R}^n : (x, 1) \in C(S)\}.$$

thm:HausOkoun

**Theorem C.2.1** For each  $S \in \mathcal{S}'(C)$ , we have

$$\text{vol } S = \lim_{k \rightarrow \infty} k^{-n} |S_k| = \text{vol } \Delta(S) > 0. \quad (\text{C.3})$$

{eq:volWvolDelta}

Moreover, as  $k \rightarrow \infty$ ,

$$\Delta_k(S) \xrightarrow{d_{\text{Haus}}} \Delta(S). \quad (\text{C.4})$$

{eq:HausconvDeltaGLS}

This is essentially proved in [WN14, Lemma 4.8], which itself follows from a theorem of Khovanskii [Kho92]. We remind the readers that (C.3) fails for a general  $W \in \mathcal{S}(C)$ , see [KK12, Theorem 2].

**Proof** The equalities (C.3) follow from the general theorem [KK12, Theorem 2].

It remains to prove (C.4). By the argument of [WN14, Lemma 4.8], for any compact set  $K \subseteq \text{Int } \Delta(S)$ , there is  $k_0 > 0$  such that for any  $k \geq k_0$ ,  $\alpha \in K \cap (k^{-1}\mathbb{Z})^n$  implies that  $\alpha \in \Delta_k(S)$ .

In particular, taking  $K = \Delta(S)^\delta$  for any  $\delta > 0$  and applying Lemma C.1.1, we find

$$d_{\text{Haus}}(\Delta(S), \Delta_k(S)) \leq n^{1/2} k^{-1} + \delta$$

when  $k$  is large enough. This implies (C.4).  $\square$

cor:dist

**Corollary C.2.1** Let  $S, S' \in \mathcal{S}'(C)$ . Assume that  $\text{vol}(S \cap S') > 0$ , then we have

$$d_{\text{sg}}(S, S') = \text{vol}(S) + \text{vol}(S') - 2 \text{vol}(S \cap S').$$

**Proof** This is a direct consequence of Lemma C.2.5 and (C.3).  $\square$

lma:regularizat

**Lemma C.2.6** Given  $S \in \mathcal{S}'(C)$ , we have  $S \sim \text{Reg}(S)$ .

Recall that the regularization  $\text{Reg}(S)$  of  $S$  is defined as  $C(S) \cap \mathbb{Z}^{n+1}$ .

**Proof** Since  $S$  and  $\text{Reg}(S)$  have the same Okounkov body, we have  $\text{vol } S = \text{vol } \text{Reg}(S)$  by Theorem C.2.1. By Corollary C.2.1 again,

$$d_{\text{sg}}(\text{Reg}(S), S) = \text{vol } \text{Reg}(S) - \text{vol } S = 0.$$

lma:Deltaindclass

**Lemma C.2.7** Let  $S, S' \in \mathcal{S}'(C)$ . Assume that  $d_{\text{sg}}(S, S') = 0$ , then  $\Delta(S) = \Delta(S')$ .

**Proof** Observe that  $\text{vol}(S \cap S') > 0$ , as otherwise

$$d_{\text{sg}}(S, S') \geq \text{vol } S + \text{vol } S' > 0,$$

which is a contradiction.

It follows from [Lemma C.2.5](#) that  $S \cap S' \in \mathcal{S}'(C)$ . It suffices to show that  $\Delta(S) = \Delta(S \cap S')$ . In fact, suppose that this holds, since  $\text{vol } \Delta(S') = \text{vol } S' = \text{vol } S = \text{vol } \Delta(S)$ , the inclusion  $\Delta(S') \supseteq \Delta(S \cap S') = \Delta(S)$  is an equality.

By [Lemma C.2.2](#), we can therefore replace  $S'$  by  $S \cap S'$  and assume that  $S \supseteq S'$ . Then clearly  $\Delta(S) \supseteq \Delta(S')$ . By [\(C.3\)](#),

$$\text{vol } \Delta(S) = \text{vol } \Delta(S') > 0.$$

Thus,  $\Delta(S) = \Delta(S')$ . □

lma:Sprimeint

**Lemma C.2.8** Suppose that  $S^i \in \mathcal{S}'(C)$  is a decreasing sequence such that

$$\lim_{i \rightarrow \infty} \text{vol } S^i > 0.$$

Then there is  $S \in \mathcal{S}'(C)$  such that  $S^i \rightarrow S$ .

In general, one cannot simply take  $S = \bigcap_i S^i$ . For example, consider the sequence  $S^i = S^1 \cap \{x_{n+1} \geq i\}$ .

**Proof** By [Lemma C.2.6](#), we may replace  $S^i$  by its regularization and assume that  $S^i = C(S^i) \cap \mathbb{Z}^{n+1}$ . We define

$$S = \left( \bigcap_{i=1}^{\infty} C(S^i) \right) \cap \mathbb{Z}^{n+1}.$$

Since  $\bigcap_{i=1}^{\infty} C(S^i)$  is a full-dimensional cone by assumption, we have  $S \in \mathcal{S}'(C)$ . By [Corollary C.2.1](#) and [Theorem C.2.1](#), we can compute the distance

$$d_{\text{sg}}(S, S^i) = \text{vol } S^i - \text{vol } S = \text{vol } \Delta(S^i) - \text{vol } \Delta(S),$$

which tends to 0 by construction. □

### C.2.3 Okounkov bodies of almost semigroups

subsec:Okobalmosg

**Definition C.2.3** We define  $\overline{\mathcal{S}'(C)}_{>0}$  as elements in the closure of  $\mathcal{S}'(C)$  in  $\hat{\mathcal{S}}(C)$  with positive volume. An element in  $\overline{\mathcal{S}'(C)}_{>0}$  is called an *almost semigroup* in  $C$ .

Recall that the volume here is defined in [\(C.2\)](#).

Our goal is to prove the following theorem:

thm:Okocont

**Theorem C.2.2** *The Okounkov body map  $\Delta: \mathcal{S}'(C) \rightarrow \mathcal{K}_n$  as defined in Definition C.2.2 admits a unique continuous extension*

$$\Delta: \overline{\mathcal{S}'(C)}_{>0} \rightarrow \mathcal{K}_n. \quad (\text{C.5}) \quad \{\text{eq:Deltagensg}\}$$

Moreover, for any  $S \in \overline{\mathcal{S}'(C)}_{>0}$ , we have

$$\text{vol } S = \text{vol } \Delta(S). \quad (\text{C.6}) \quad \{\text{eq:volWfinal}\}$$

**Proof** The uniqueness of the extension is clear as long as it exists. Moreover, (C.6) follows easily from Theorem C.2.1 and Theorem C.1.2 by continuity. It remains to argue the existence of the continuous extension. We first construct an extension and prove its continuity.

**Step 1.** We construct the desired map (C.5). Let  $S \in \overline{\mathcal{S}'(C)}_{>0}$ . We wish to construct a convex body  $\Delta(S) \in \mathcal{K}_n$ .

Let  $S^i \in \mathcal{S}'(C)$  be a sequence that converges to  $S$  such that

$$d_{\text{sg}}(S^i, S^{i+1}) \leq 2^{-i}.$$

For each  $i, j \geq 0$ , we introduce

$$S^{i,j} = S^i \cap S^{i+1} \cdots \cap S^{i+j}.$$

Then by Lemma C.2.2,

$$d_{\text{sg}}(S^{i,j}, S^{i,j+1}) \leq 2^{-i-j}.$$

Take  $i_0 > 0$  large enough so that for  $i \geq i_0$ ,  $\text{vol } S^i > 2^{-1} \text{vol } S$  and  $2^{2-i} < \text{vol } S$  and hence

$$\text{vol } S^i - \text{vol } S^{i,j} \leq d_{\text{sg}}(S^{i,0}, S^{i,1}) + d_{\text{sg}}(S^{i,1}, S^{i,2}) + \cdots + d_{\text{sg}}(S^{i,j-1}, S^{i,j}) \leq 2^{1-i}.$$

It follows that  $\text{vol } S^{i,j} > 2^{-1} \text{vol } S - 2^{1-i} > 0$  whenever  $i \geq i_0$ . In particular, by Lemma C.2.5,  $S^{i,j} \in \mathcal{S}'(C)$  for  $i \geq i_0$ .

By Lemma C.2.8, for  $i \geq i_0$ , there exists  $T^i \in \mathcal{S}'(C)$  such that  $S^{i,j} \rightarrow T^i$  as  $j \rightarrow \infty$ . Moreover,

$$d_{\text{sg}}(T^i, S) = \lim_{j \rightarrow \infty} d_{\text{sg}}(S^{i,j}, S) \leq \lim_{j \rightarrow \infty} d_{\text{sg}}(S^{i,j}, S^i) + d_{\text{sg}}(S^i, S) \leq 2^{1-i} + d_{\text{sg}}(S^i, S).$$

Therefore,  $T^i \rightarrow S$ . We then define

$$\Delta(S) := \overline{\bigcup_{i=i_0}^{\infty} \Delta(T^i)}.$$

In other words, we have defined

$$\Delta(S) := \varliminf_{i \rightarrow \infty} \Delta(S^i).$$



This is an honest limit: if  $\Delta$  is the limit of a subsequence of  $\Delta(S^i)$ , then  $\Delta(S) \subseteq \Delta$  by (C.1). Comparing the volumes, we find that equality holds. So by Theorem C.1.1,

$$\Delta(S) = \lim_{i \rightarrow \infty} \Delta(S^i). \quad (\text{C.7})$$

{eq:deltawtemp}

Next we claim that  $\Delta(S)$  as defined above does not depend on the choice of the sequence  $S^i$ . In fact, suppose that  $S'^i \in S'(C)$  is another sequence satisfying the same conditions as  $S^i$ . The same holds for  $R^i := S^{i+1} \cap S'^{i+1}$ . It follows that

$$\lim_{i \rightarrow \infty} \Delta(R^i) \subseteq \lim_{i \rightarrow \infty} \Delta(S^i).$$

Comparing the volumes, we find that equality holds. The same is true with  $S'^i$  in place of  $S^i$ . So we conclude that  $\Delta(S)$  as in (C.7) does not depend on the choices we made.

**Step 2.** It remains to prove the continuity of  $\Delta$  defined in Step 1. Suppose that  $S^i \in \overline{S'(C)}_{>0}$  is a sequence with limit  $S \in \overline{S'(C)}_{>0}$ . We want to show that

$$\Delta(S^i) \xrightarrow{d_{\text{Haus}}} \Delta(S). \quad (\text{C.8})$$

{eq:temp5}

We first reduce to the case where  $S^i \in S'(C)$ . By (C.7), for each  $i$ , we can choose  $T^i \in S'(C)$  such that  $d_{\text{sg}}(S^i, T^i) < 2^{-i}$  and  $d_{\text{Haus}}(\Delta(S^i), \Delta(T^i)) < 2^{-i}$ . If we have shown  $\Delta(T^i) \xrightarrow{d_{\text{Haus}}} \Delta(S)$ , then (C.8) follows immediately.

Next we reduce to the case where  $d_{\text{sg}}(S^i, S^{i+1}) \leq 2^{-i}$ . In fact, thanks to Theorem C.1.1, in order to prove (C.8), it suffices to show that each subsequence of  $\Delta(S^i)$  admits a subsequence that converges to  $\Delta(S)$ . Hence, we easily reduce to the required case.

After these reductions, (C.8) is nothing but (C.7).  $\square$

*Remark C.2.1* As the readers can easily verify from the proof, for any  $S \in \overline{S'(C)}_{>0}$ , there is  $S' \in S'(C)$  such that  $S \sim S'$ .

cor:Okocomp

**Corollary C.2.2** Suppose that  $S, S' \in \overline{S'(C)}_{>0}$  with  $S \subseteq S'$ , then

$$\Delta(S) \subseteq \Delta(S'). \quad (\text{C.9})$$

{eq:Deltacontain}

**Proof** Let  $S^j, S'^j \in S'(C)$  be elements such that  $S^j \rightarrow S$ ,  $S'^j \rightarrow S'$ . Then it follows from Lemma C.2.2 that  $S^j \cap S'^j \rightarrow S$ . Since  $\text{vol}$  is continuous, for large  $j$ ,  $S^j \cap S'^j$  has positive volume and hence lies in  $S'(C)$  by Lemma C.2.5. We may therefore replace  $S^j$  by  $S^j \cap S'^j$  and assume that  $S^j \subseteq S'^j$ . Hence, (C.9) follows from the continuity of  $\Delta$  proved in Theorem C.2.2.  $\square$

*Remark C.2.2* As the readers can easily verify, the construction of  $\Delta$  is independent of the choice of  $C$  in the following sense: Suppose that  $C'$  is another cone satisfying the same assumptions as  $C$  and  $C' \supseteq C$ , then the Okounkov body map  $\Delta: \overline{S'(C')}_{>0} \rightarrow \mathcal{K}_n$  is an extension of the corresponding map (C.5). We will constantly use this fact without further explanations.



# Index

## Symbols

$B_{v,\varphi,v}^k$	230	$\mathrm{PSH}^{\mathrm{NA}}(X, \theta)_{>0}$	136
$E_\theta^\phi$	44	$\mathrm{PSH}^{\mathrm{NA}}(X, \theta; \phi)$	136
$E_{\theta,K}[\varphi](v)$	219	$\mathrm{PSH}_{\mathrm{tor}}(X, \omega)$	65
$E_{\theta,K}[\varphi]_I(v)$	219	$\mathrm{QPSH}(X)$	19
$E_\theta$	44	$\mathrm{Res}_Y I$	19
$F_\varphi$	198	$\mathrm{SH}(\Omega)$	4
$P_{\theta+\omega'}[\bullet]_I$	207	$\mathrm{Sing} T$	188
$P_{\theta+\omega}[\bullet]_I$	209	$\mathrm{Sing}_X T$	188
$P_{\theta,K}[\varphi](v)$	219	$\mathrm{TC}(X, \theta)_{>0}$	127
$P_\theta[\Gamma]_I$	136	$\mathrm{TC}(X, \theta; \phi)$	127
$P_\theta[\varphi]$	36	$\mathrm{TC}(\Delta)$	176
$P_\theta[\varphi]_I$	47	$\mathrm{TC}^1(X, \theta; \phi)$	129
$V_\theta$	31	$\mathrm{TC}^1(\Delta)$	176
$Y_\bullet$	147	$\mathrm{TC}^\infty(X, \theta; \phi)$	129
$\mathrm{BM}(K, v)$	229	$\mathrm{TC}^\infty(\Delta)$	176
$\mathrm{Bir}(X)$	185	$\mathrm{Tr}_Y(T)$	110
$\mathrm{DH}(\Delta_\bullet)$	180	$\mathrm{Tr}_Y(\varphi)$	110
$\mathrm{DH}(\Gamma)$	216	$\mathrm{Trop}$	64
$\Delta(\omega, \varphi)$	67	$\beta_{v,\varphi,v}^k$	230
$\Delta(\theta, \varphi)$	198	$\mathrm{cor}(Y_\bullet, \pi)$	150
$\Delta_k(\theta, \varphi)$	152	$\mathrm{dd}^c \varphi_1 \wedge \cdots \wedge \mathrm{dd}^c \varphi_p$	28
$\Delta_{Y_\bullet}(T)$	166	$\ell \vee \ell'$	60
$\Delta_{Y_\bullet}(\alpha)$	165	$\mathbb{D}(T)$	188
$\Delta_v(L)$	154	$\mathbf{E}$	59
$\Delta_v(L, h)$	155	$\mathbf{E}(\Delta_\bullet)$	176
$\Delta_{k,T}(L)$	153	$\mathbf{E}^\phi$	58, 129
$\Delta_{k,T}(\theta, \varphi)$	153	$\mathcal{E}(X, \theta; \phi)$	43
$\Gamma(\theta, \varphi)$	152	$\mathcal{E}^1(X, \theta; \phi)$	43
$\Gamma^\infty(\theta, \varphi)$	153	$\mathcal{E}^\infty(X, \theta; \phi)$	43
$\Gamma_{\max}$	208	$\mathcal{E}_{[\varphi],K}^\theta$	225
$\mathrm{PSH}(X)$	6	$I(\varphi)$	17
$\mathrm{PSH}(X, \theta)$	19	$I_\infty(\varphi)$	22
$\mathrm{PSH}(X, \theta; \phi)$	43	$\mathcal{L}_{k,\varphi}$	231
$\mathrm{PSH}(\Omega)$	4	$\mathcal{R}(X, \theta)$	58
$\mathrm{PSH}^{\mathrm{NA}}(X, \theta)$	207	$\mathcal{R}(X, \theta; \phi)$	58
		$\mathcal{R}^1(X, \theta)$	58
		$\mathcal{R}^1(X, \theta; \phi)$	58

$\mathcal{R}^\infty(X, \theta)$  58  
 $\mathcal{R}^\infty(X, \theta; \phi)$  58  
 $\mathcal{Z}_+(X)$  24  
 $\mathcal{Z}_+(X, \alpha)$  24  
 $\nu(\varphi, E)$  17  
 $\nu(\varphi, F)$  17  
 $\nu(\varphi, x)$  15  
 $\nu_{Y_\bullet}$  147  
 $\nu_{Y_\bullet}(T)$  148  
 $\sim_I$  46  
 $\varphi \wedge \psi$  35  
 $\varphi \leq \psi$  19  
 $\varphi \leq_P \psi$  75  
 $\varphi \leq_I \psi$  78  
 $\varphi \sim \psi$  19  
 $\text{vol } \Gamma$  209  
 $\text{vol } \mathbb{D}$  187  
 $\text{vol } \theta_\varphi$  48  
 $\text{vol}(\theta, \varphi)$  48  
 $d_S$  83

**A**

admissible flag 147  
 analytic singularities 20

**B**

Bedford–Taylor product 28  
 Bernstein–Markov 229  
 big Cartier b-divisor 186  
 big class 24  
 birational model 185  
 bounded geodesic ray 58  
 bounded Okounkov test curve 176  
 bounded test curve 129  
 bounded test function 177

**C**

Cartier b-divisor 185  
 co-pluripolar set 6  
 complexification 53

**D**

Duistermaat–Heckman measure 180, 216

**F**

finite energy 43  
 full mass 43

**G**

generic Lelong number 17  
 gentle analytic singularities 22  
 geodesic 55  
 geodesic ray 58  
 geodesic ray with finite energy 58

**H**

Hermitian big line bundle 105  
 Hermitian form 24  
 Hermitian line bundle 25  
 Hermitian metric 25  
 Hermitian pseudoeffective line bundle 105

**I**

I-envelope 47, 136  
 I-equivalent 46  
 I-good 100  
 I-model potential 47  
 I-model test curve 136  
 I-more singular 78  
 inverse Legendre transform 177

**K**

Kähler currents 24

**L**

Legendre transform 130, 177  
 Lelong number 15  
 log singularities 21

**M**

minimal singularities 43  
 Minkowski sum 261  
 model potential 37  
 Monge–Ampère energy 44  
 multiplier ideal sheaf 17

**N**

neat analytic singularities 20  
 nef Cartier b-divisor 186  
 nef Weil b-divisor 186  
 Newton body 67, 198  
 non-pluripolar 6  
 non-pluripolar product 28, 30

**O**

Okounkov body 154, 165  
 Okounkov test curve 176  
 Okounkov test curve with finite energy 176

**P**

P-envelope 36  
 P-more singular 75  
 P-singularity type 75  
 partial Bergman kernels 229  
 partial Bergman measures 230  
 partial Donaldson functional 231  
 partial equilibrium energy 225  
 partial Okounkov body 155, 166  
 plurifine topology 11, 14  
 pluripolar set 6  
 plurisubharmonic function 4, 6  
 plurisubharmonic metric 25  
 polar locus 24  
 positive currents 23  
 pseudo-effective class 24  
 pseudo-effective Weil b-divisor 186

**Q**

quasi-equisingular approximation 22  
 quasi-plurisubharmonic function 19

**R**

radial Monge–Ampère energy 58  
 relative finite energy 43  
 relative full mass 43  
 relative I-envelope 219  
 relative minimal singularities 43

relative P-envelope 219  
 restriction ideal 19  
 rooftop operator 35

**S**

singular Hermitian form 25  
 singularity b-divisor 188  
 singularity divisor 187  
 smooth flag 148  
 subgeodesic 53  
 subharmonic function 3

**T**

test curve 127  
 test curve with finite energy 129  
 test function 177  
 test function with finite energy 177  
 thick subset 11  
 trace operator 110

**V**

valuation 148  
 volume 48, 187, 209

**W**

weighted subset 229  
 Weil b-divisor 185



## References

- BB10. Robert Berman and Sébastien Boucksom. Growth of balls of holomorphic sections and energy at equilibrium. *Invent. Math.*, 181(2):337–394, 2010.
- BB13. Robert J. Berman and Bo Berndtsson. Real Monge-Ampère equations and Kähler-Ricci solitons on toric log Fano varieties. *Ann. Fac. Sci. Toulouse Math. (6)*, 22(4):649–711, 2013.
- BBGHdJ21. A. Botero, J. I. Burgos Gil, D. Holmes, and R. de Jong. Chern-Weil and Hilbert-Samuel formulae for singular hermitian line bundles, 2021.
- BBGHdJ22. A. Botero, J. I. Burgos Gil, D. Holmes, and R. de Jong. Rings of Siegel-Jacobi forms of bounded relative index are not finitely generated, 2022.
- BBJ21. Robert J. Berman, Sébastien Boucksom, and Mattias Jonsson. A variational approach to the Yau-Tian-Donaldson conjecture. *J. Amer. Math. Soc.*, 34(3):605–652, 2021.
- BBWN11. Robert Berman, Sébastien Boucksom, and David Witt Nyström. Fekete points and convergence towards equilibrium measures on complex manifolds. *Acta Math.*, 207(1):1–27, 2011.
- BDPP13. Sébastien Boucksom, Jean-Pierre Demailly, Mihai Păun, and Thomas Peternell. The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension. *J. Algebraic Geom.*, 22(2):201–248, 2013.
- BEGZ10. Sébastien Boucksom, Philippe Eyssidieux, Vincent Guedj, and Ahmed Zeriahi. Monge-Ampère equations in big cohomology classes. *Acta Math.*, 205(2):199–262, 2010.
- Ber11. Robert J. Berman. Bergman kernels and equilibrium measures for line bundles over projective manifolds. *Amer. J. Math.*, 131(5):1485–1524, 2009.
- Ber14. Robert J. Berman. Determinantal point processes and fermions on complex manifolds: large deviations and bosonization. *Comm. Math. Phys.*, 327(1):1–47, 2014.
- BFJ08. Sébastien Boucksom, Charles Favre, and Mattias Jonsson. Valuations and plurisubharmonic singularities. *Publ. Res. Inst. Math. Sci.*, 44(2):449–494, 2008.
- BFJ09. Sébastien Boucksom, Charles Favre, and Mattias Jonsson. Differentiability of volumes of divisors and a problem of Teissier. *J. Algebraic Geom.*, 18(2):279–308, 2009.
- BGPS14. José Ignacio Burgos Gil, Patrice Philippon, and Martín Sombra. Arithmetic geometry of toric varieties. Metrics, measures and heights. *Astérisque*, pages vi+222, 2014.
- BM97. Edward Bierstone and Pierre D. Milman. Canonical desingularization in characteristic zero by blowing up the maximum strata of a local invariant. *Invent. Math.*, 128(2):207–302, 1997.
- Bon98. Laurent Bonavero. Inégalités de morse holomorphes singulières. *J. Geom. Anal.*, 8(3):409–425, 1998.
- Bou02. S. Boucksom. *Cônes positifs des variétés complexes compactes*. PhD thesis, Université Joseph-Fourier-Grenoble I, 2002.
- Bou02b. Sébastien Boucksom. On the volume of a line bundle. *Internat. J. Math.*, 13(10):1043–1063, 2002.

- Bou17. Sébastien Boucksom. Singularities of plurisubharmonic functions and multiplier ideals. <http://sebastien.boucksom.perso.math.cnrs.fr/notes/L2.pdf>, 2017.
- BT76. Eric Bedford and B. A. Taylor. The Dirichlet problem for a complex Monge-Ampère equation. *Invent. Math.*, 37(1):1–44, 1976.
- BT82. Eric Bedford and B. A. Taylor. A new capacity for plurisubharmonic functions. *Acta Math.*, 149(1-2):1–40, 1982.
- BT87. Eric Bedford and B. A. Taylor. Fine topology, Šilov boundary, and  $(dd^c)^n$ . *J. Funct. Anal.*, 72(2):225–251, 1987.
- Cao14. Junyan Cao. Numerical dimension and a Kawamata-Viehweg-Nadel-type vanishing theorem on compact Kähler manifolds. *Compos. Math.*, 150(11):1869–1902, 2014.
- Car83. Lennart Carleson. Selected problems on exceptional sets. In *Selected reprints*, Wadsworth Math. Ser., pages iv+100. Wadsworth, Belmont, CA, 1983.
- CDG03. David M. J. Calderbank, Liana David, and Paul Gauduchon. The Guillemin formula and Kähler metrics on toric symplectic manifolds. *J. Symplectic Geom.*, 1(4):767–784, 2003.
- CDM17. JunYan Cao, Jean-Pierre Demailly, and Shin-ichi Matsumura. A general extension theorem for cohomology classes on non reduced analytic subspaces. *Sci. China Math.*, 60(6):949–962, 2017.
- CFKLRS17. C<sup>+</sup>FK17. Ciro Ciliberto, Michal Farnik, Alex Küronya, Victor Lozovanu, Joaquim Roé, and Constantin Shramov. Newton-Okounkov bodies sprouting on the valuative tree. *Rend. Circ. Mat. Palermo (2)*, 66(2):161–194, 2017.
- CGSZ19. CGSZ19. Dan Coman, Vincent Guedj, Sibel Sahin, and Ahmed Zeriahi. Toric pluripotential theory. *Ann. Polon. Math.*, 123(1):215–242, 2019.
- CLS11. CLS11. David A. Cox, John B. Little, and Henry K. Schenck. *Toric varieties*, volume 124 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2011.
- CPW17. CPW17. Sung Rak Choi, Jinhyung Park, and Joonyeong Won. Okounkov bodies associated to pseudoeffective divisors II. *Taiwanese J. Math.*, 21(3):601–620, 2017.
- Da17. Dar17. Tamás Darvas. Weak geodesic rays in the space of Kähler potentials and the class  $\mathcal{E}(X, \omega)$ . *J. Inst. Math. Jussieu*, 16(4):837–858, 2017.
- DDNL18big. DDNL18a. Tamás Darvas, Eleonora Di Nezza, and Chinh H. Lu.  $L^1$  metric geometry of big cohomology classes. *Ann. Inst. Fourier (Grenoble)*, 68(7):3053–3086, 2018.
- DDNL18mono. DDNL18b. Tamás Darvas, Eleonora Di Nezza, and Chinh H. Lu. Monotonicity of nonpluripolar products and complex Monge-Ampère equations with prescribed singularity. *Anal. PDE*, 11(8):2049–2087, 2018.
- DDNL18fullmass. DDNL18c. Tamás Darvas, Eleonora Di Nezza, and Chinh H. Lu. On the singularity type of full mass currents in big cohomology classes. *Compos. Math.*, 154(2):380–409, 2018.
- DDNL19log. DDNL21a. Tamás Darvas, Eleonora Di Nezza, and Chinh H. Lu. Log-concavity of volume and complex Monge-Ampère equations with prescribed singularity. *Math. Ann.*, 379(1-2):95–132, 2021.
- DDNLmetric. DDNL21b. Tamás Darvas, Eleonora Di Nezza, and Hoang-Chinh Lu. The metric geometry of singularity types. *J. Reine Angew. Math.*, 771:137–170, 2021.
- DDNLsurv. DDNL23. Tamás Darvas, Eleonora Di Nezza, and Chinh H. Lu. Relative pluripotential theory on compact kähler manifolds, 2023.
- Dem85. Dem85. Jean-Pierre Demailly. Mesures de Monge-Ampère et caractérisation géométrique des variétés algébriques affines. *Mém. Soc. Math. France (N.S.)*, page 124, 1985.
- Dem12. Dem12a. Jean-Pierre Demailly. *Analytic methods in algebraic geometry*, volume 1 of *Surveys of Modern Mathematics*. International Press, Somerville, MA; Higher Education Press, Beijing, 2012.
- DemBook. Dem12b. Jean-Pierre Demailly. *Complex analytic and differential geometry*, 2012. Available on personal website, [link](#).
- Dem15. Dem15. Jean-Pierre Demailly. On the cohomology of pseudoeffective line bundles. In *Complex geometry and dynamics*, volume 10 of *Abel Symp.*, pages 51–99. Springer, Cham, 2015.



- Deng17 Den17. Ya Deng. Transcendental Morse inequality and generalized Okounkov bodies. *Algebr. Geom.*, 4(2):177–202, 2017.
- DF20 DF22. Nguyen-Bac Dang and Charles Favre. Intersection theory of nef  $b$ -divisor classes. *Compos. Math.*, 158(7):1563–1594, 2022.
- EGAIV-2 DG65. Jean Dieudonné and Alexandre Grothendieck. *Éléments de géométrie algébrique: IV. Étude locale des schémas et des morphismes de schémas, Seconde partie*, volume 24. Institut des hautes études scientifiques, 1965.
- DNT19 DNT21. Eleonora Di Nezza and Stefano Trapani. Monge-Ampère measures on contact sets. *Math. Res. Lett.*, 28(5):1337–1352, 2021.
- DPS01 DPS01. Jean-Pierre Demailly, Thomas Peternell, and Michael Schneider. Pseudo-effective line bundles on compact Kähler manifolds. *Internat. J. Math.*, 12(6):689–741, 2001.
- DR22 DR22. Ruadhai Dervan and Rémi Reboulet. Ding stability and Kähler-Einstein metrics on manifolds with big anticanonical class, 2022.
- DRWNXZ DRWN<sup>+</sup>23. Tamás Darvas, Rémi Reboulet, David Witt Nyström, Mingchen Xia, and Kewei Zhang. Transcendental okounkov bodies, 2023.
- DS06 DS06. Tien-Cuong Dinh and Nessim Sibony. Distribution des valeurs de transformations méromorphes et applications. *Comment. Math. Helv.*, 81(1):221–258, 2006.
- DX21 DX21. Tamás Darvas and Mingchen Xia. The volume of pseudoeffective line bundles and partial equilibrium. *Geometry & Topology (to appear)*, 2021.
- DX22 DX22. Tamás Darvas and Mingchen Xia. The closures of test configurations and algebraic singularity types. *Adv. Math.*, 397:Paper No. 108198, 56, 2022.
- DX24 DX24. Tamás Darvas and Mingchen Xia. The trace operator of quasi-plurisubharmonic functions on compact Kähler manifolds, 2024.
- DXZ23 DXZ23. Tamás Darvas, Mingchen Xia, and Kewei Zhang. A transcendental approach to non-Archimedean metrics of pseudoeffective classes. *Commentarii Mathematici Helvetici(to appear)*, 2023.
- DZ22 DZ22. T. Darvas and K. Zhang. Twisted kähler-einstein metrics in big classes, 2022.
- ELMNP05 ELM<sup>+</sup>05. Lawrence Ein, Robert Lazarsfeld, Mircea Mustață, Michael Nakamaye, and Mihnea Popa. Asymptotic invariants of line bundles. *Pure Appl. Math. Q.*, 1(2):379–403, 2005.
- EMSW06 EMW06. Said El Marzguioui and Jan Wiegerinck. The pluri-fine topology is locally connected. *Potential Anal.*, 25(3):283–288, 2006.
- Fig17 Fig17. Alessio Figalli. *The Monge-Ampère equation and its applications*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2017.
- Fin22 Fin22. Siarhei Finski. On the metric structure of section ring, 2022.
- FK18 FK18. Kazuhiro Fujiwara and Fumiharu Kato. *Foundations of rigid geometry. I*. EMS Monographs in Mathematics. European Mathematical Society (EMS), Zürich, 2018.
- FM21 FM21. Osamu Fujino and Shin-ichi Matsumura. Injectivity theorem for pseudo-effective line bundles and its applications. *Trans. Amer. Math. Soc. Ser. B*, 8:849–884, 2021.
- FN80 FsN80. John Erik Fornæss and Raghavan Narasimhan. The Levi problem on complex spaces with singularities. *Math. Ann.*, 248(1):47–72, 1980.
- Fuj23 Fuj23. Osamu Fujino. Relative Bertini type theorem for multiplier ideal sheaves. *Osaka J. Math.*, 60(1):207–226, 2023.
- GH14 GH14. P. Griffiths and J. Harris. *Principles of algebraic geometry*. John Wiley & Sons, 2014.
- GK20 GK20. Patrick Graf and Tim Kirschner. Finite quotients of three-dimensional complex tori. *Ann. Inst. Fourier (Grenoble)*, 70(2):881–914, 2020.
- GR56 GR56. Hans Grauert and Reinhold Remmert. Plurisubharmonische Funktionen in komplexen Räumen. *Math. Z.*, 65:175–194, 1956.
- CAS GR84. Hans Grauert and Reinhold Remmert. *Coherent analytic sheaves*, volume 265 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1984.
- SHC6 Gro60. Alexander Grothendieck. Techniques de construction en géométrie analytique. VI. étude locale des morphismes: germes d’espaces analytiques, platitude, morphismes simples. *Séminaire Henri Cartan*, 13(1):1–13, 1960.

- Gui94. Victor Guillemin. Kaehler structures on toric varieties. *J. Differential Geom.*, 40(2):285–309, 1994.
- GZ05. Vincent Guedj and Ahmed Zeriahi. Intrinsic capacities on compact Kähler manifolds. *J. Geom. Anal.*, 15(4):607–639, 2005.
- GZ07. Vincent Guedj and Ahmed Zeriahi. The weighted Monge-Ampère energy of quasi-plurisubharmonic functions. *J. Funct. Anal.*, 250(2):442–482, 2007.
- GZ15. Qi'an Guan and Xiangyu Zhou. Effectiveness of Demailly's strong openness conjecture and related problems. *Invent. Math.*, 202(2):635–676, 2015.
- GZ17. Vincent Guedj and Ahmed Zeriahi. *Degenerate complex Monge-Ampère equations*, volume 26 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2017.
- Har. Robin Hartshorne. *Algebraic geometry*, volume No. 52 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Heidelberg, 1977.
- Hiep14. Pham Hoang Hiep. The weighted log canonical threshold. *C. R. Math. Acad. Sci. Paris*, 352(4):283–288, 2014.
- Hir75. Heisuke Hironaka. Flattening theorem in complex-analytic geometry. *Amer. J. Math.*, 97:503–547, 1975.
- His12. Tomoyuki Hisamoto. Restricted Bergman kernel asymptotics. *Trans. Amer. Math. Soc.*, 364(7):3585–3607, 2012.
- HK76. W. K. Hayman and P. B. Kennedy. *Subharmonic functions. Vol. I*, volume No. 9 of *London Mathematical Society Monographs*. Academic Press [Harcourt Brace Jovanovich, Publishers], London-New York, 1976.
- HPS18. C. Hacon, M. Popa, and C. Schnell. Algebraic fiber spaces over abelian varieties: around a recent theorem by Cao and Păun. In *Local and global methods in algebraic geometry*, volume 712 of *Contemp. Math.*, pages 143–195. Amer. Math. Soc., [Providence], RI, 2018.
- Ino22. Eiji Inoue. Entropies in  $\mu$ -framework of canonical metrics and K-stability, II – Non-archimedean aspect: non-archimedean  $\mu$ -entropy and  $\mu$ K-semistability, 2022.
- Jow10. Shin-Yao Jow. Okounkov bodies and restricted volumes along very general curves. *Adv. Math.*, 223(4):1356–1371, 2010.
- Kho92. A. G. Khovanskii. The Newton polytope, the Hilbert polynomial and sums of finite sets. *Funktsional. Anal. i Prilozhen.*, 26(4):57–63, 96, 1992.
- Kim15. Dano Kim. Equivalence of plurisubharmonic singularities and Siu-type metrics. *Monatsh. Math.*, 178(1):85–95, 2015.
- Kis78. Christer O. Kiselman. The partial Legendre transformation for plurisubharmonic functions. *Invent. Math.*, 49(2):137–148, 1978.
- KK12. Kiumars Kaveh and A. G. Khovanskii. Newton-Okounkov bodies, semigroups of integral points, graded algebras and intersection theory. *Ann. of Math. (2)*, 176(2):925–978, 2012.
- LM09. Robert Lazarsfeld and Mircea Mustață. Convex bodies associated to linear series. *Ann. Sci. Éc. Norm. Supér. (4)*, 42(5):783–835, 2009.
- Mat89. Hideyuki Matsumura. *Commutative ring theory*, volume 8 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1989. Translated from the Japanese by M. Reid.
- MM07. Xiaonan Ma and George Marinescu. *Holomorphic Morse inequalities and Bergman kernels*, volume 254 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2007.
- MZ23. Xiankui Meng and Xiangyu Zhou. On the restriction formula. *J. Geom. Anal.*, 33(12):Paper No. 369, 30, 2023.
- PT18. Mihai Păun and Shigeharu Takayama. Positivity of twisted relative pluricanonical bundles and their direct images. *J. Algebraic Geom.*, 27(2):211–272, 2018.
- Rau15. Hossein Raufi. Singular hermitian metrics on holomorphic vector bundles. *Ark. Mat.*, 53(2):359–382, 2015.
- Roc70. R. Tyrrell Rockafellar. *Convex analysis*. Princeton Mathematical Series, No. 28. Princeton University Press, Princeton, N.J., 1970.

- RS05. Alexander Rashkovskii and Ragnar Sigurdsson. Green functions with singularities along complex spaces. *Internat. J. Math.*, 16(4):333–355, 2005.
- RWN14. Julius Ross and David Witt Nyström. Analytic test configurations and geodesic rays. *J. Symplectic Geom.*, 12(1):125–169, 2014.
- Sch14. Sch93. Rolf Schneider. *Convex bodies: the Brunn-Minkowski theory*, volume 44 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1993.
- Siu74. Yum Tong Siu. Analyticity of sets associated to Lelong numbers and the extension of closed positive currents. *Invent. Math.*, 27:53–156, 1974.
- Tru22. Antonio Trusiani. Kähler-Einstein metrics with prescribed singularities on Fano manifolds. *J. Reine Angew. Math.*, 793:1–57, 2022.
- Tru23. Antonio Trusiani. A relative Yau–Tian–Donaldson conjecture and stability thresholds, 2023.
- Vu19. Duc-Viet Vu. Locally pluripolar sets are pluripolar. *Internat. J. Math.*, 30(13):1950029, 13, 2019.
- WN14. David Witt Nyström. Transforming metrics on a line bundle to the Okounkov body. *Ann. Sci. Éc. Norm. Supér. (4)*, 47(6):1111–1161, 2014.
- Wlo09. Jarosław Włodarczyk. Resolution of singularities of analytic spaces. In *Proceedings of Gökova Geometry-Topology Conference 2008*, pages 31–63. Gökova Geometry/Topology Conference (GGT), Gökova, 2009.
- Xia21. Mingchen Xia. Partial Okounkov bodies and Duistermaat–Heckman measures of non-Archimedean metrics. *Geometry & Topology (to appear)*, 2021.
- XiaBer. Xia22a. Mingchen Xia. Analytic Bertini theorem. *Math. Z.*, 302(2):1171–1176, 2022.
- Xia22. Xia22b. Mingchen Xia. Non-pluripolar products on vector bundles and Chern–Weil formulae. *Math. Ann.*, 2022.
- Xia23Mabuchi. Xia23a. Mingchen Xia. Mabuchi geometry of big cohomology classes. *J. Reine Angew. Math.*, 798:261–292, 2023.
- Xia23Operations. Xia23b. Mingchen Xia. Operations on transcendental non-Archimedean metrics, 2023.
- XiaPPT. Xia23c. Mingchen Xia. Pluripotential-theoretic stability thresholds. *Int. Math. Res. Not. IMRN*, pages 12324–12382, 2023.