ProTherm and ProNIT: thermodynamic databases for proteins and protein–nucleic acid interactions

M. D. Shaji Kumar¹, K. Abdulla Bava¹, M. Michael Gromiha², Ponraj Prabakaran³, Koji Kitajima¹, Hatsuho Uedaira⁴ and Akinori Sarai*

Department of Bioscience and Bioinformatics, Kyushu Institute of Technology (KIT) 680-4 Kawazu, Iizuka, 820-8502 Japan, ¹Advanced Technology Institute, Inc. (ATI), 2-3-13-103 Tate, Shiki, Saitama 353-0006, Japan, ²Computational Biology Research Center (CBRC), National Institute of Advanced Industrial Science and Technology (AIST), AIST Tokyo Waterfront Bio-IT Research Building, 2-42 Aomi, Koto-ku, Tokyo 135-0064, Japan, ³Laboratory of Experimental and Computational Biology, NCI, NIH, Frederick, MD 21702, USA and ⁴Tsukuba Materials Information Laboratory, 3-23-4, Ninomiya, Tsukuba, 305-0051 Japan

Received September 15, 2005; Revised and Accepted October 14, 2005

ABSTRACT

ProTherm and ProNIT are two thermodynamic databases that contain experimentally determined thermodynamic parameters of protein stability and protein-nucleic acid interactions, respectively. The current versions of both the databases have considerably increased the total number of entries and enhanced search interface with added new fields, improved search, display and sorting options. As on September 2005, ProTherm release 5.0 contains 17113 entries from 771 proteins, retrieved from 1497 scientific articles (~20% increase in data from the previous version). ProNIT release 2.0 contains 4900 entries from 273 research articles, representing 158 proteins. Both databases can be gueried using WWW interfaces. Both quick search and advanced search are provided on this web page to facilitate easy retrieval and display of the data from these databases. ProTherm is freely available online at http://gibk26.bse.kyutech.ac.jp/jouhou/Protherm/ protherm.html and ProNIT at http://gibk26.bse. kyutech.ac.jp/jouhou/pronit/pronit.html.

INTRODUCTION

Thermodynamic database for proteins and mutants (Pro-Therm) and thermodynamic database for protein–nucleic acid interactions (ProNIT) are two comprehensive, integrated databases that document experimentally determined thermodynamic parameters published in the literature. Both ProTherm (1–4) and ProNIT (5) include several thermodynamic parameters along with sequence and structural information, experimental methods and conditions, and literature information. Recent years have seen tremendous progress in studies on proteins owing to the development of various experimental methods to analyze proteins at the genome scale. The correlation between structure and thermodynamics of these key molecules provides valuable insights into the way in which they function. Even though the information is available in scientific journals, books (6,7) and literature databases, retrieving useful, specific data from these resources is time consuming and laborious. Our major goal in developing these databases is to provide the scientific community a single, comprehensive data repository, where all the thermodynamic data related to protein stability and protein-nucleic acid interactions are available. The availability of such thermodynamic databases would be a valuable resource for understanding the protein folding mechanism, protein stability, molecular recognition and gene expressions. This can lead to a wide spectrum of applications such as developing algorithms/ methods for prediction systems, protein engineering and quantitative simulation of gene regulatory networks. The thermodynamic data available in ProTherm and ProNIT are widely used by researchers to study the underlying mechanisms of protein stability upon mutations and protein-nucleic acid interactions (see the reference sections on both the websites). This paper describes the major updates and enhancements to these databases for the last few years.

CONTENT, ORGANIZATION AND DATA COLLECTION

Both the databases contain information on protein, mutational information, experimental methods and conditions, several thermodynamic parameters and literature information.

The online version of this article has been published under an open access model. Users are entitled to use, reproduce, disseminate, or display the open access version of this article for non-commercial purposes provided that: the original authorship is properly and fully attributed; the Journal and Oxford University Press are attributed as the original place of publication with the correct citation details given; if an article is subsequently reproduced or disseminated not in its entirety but only in part or as a derivative work this must be clearly indicated. For commercial re-use, please contact journals.permissions@oxfordjournals.org

^{*}To whom correspondence should be addressed. Tel: +81 948 29 7811; Fax: +81 948 29 7841; Email: sarai@bse.kyutech.ac.jp

[©] The Author 2006. Published by Oxford University Press. All rights reserved.

Table 1. Contents of ProTherm and ProNIT

Contents of the databases ProTherm	ProNIT
Protein information Name, Source PIR, SWISSPROT PDB code EC, PMD number Mutation details Secondary structure Accessible surface area (ASA) Experimental condition: Temperature pH Buffer, Ion Protein concentration Measure (DSC, CD and so on) Method of denaturation Thermodynamic data: Denaturant denaturation: Free energy of unfolding: $\Delta G_{\text{H}_2\text{O}}$ Difference in $\Delta G_{\text{H}_2\text{O}}$: $\Delta \Delta G_{\text{H}_2\text{O}}$ Denaturation concentration: C_{m} Slope of denaturation curve: m Temperature: T Thermal denaturation: Free energy of unfolding: ΔG Difference in ΔG : $\Delta \Delta G$ Transition temperature: T_{m} Change in T_{m} : ΔT_{m} Enthalpy change: ΔH_{cal} , ΔH_{vH} Heat capacity change: ΔC_{p} Literature: Reference, Author Keywords, Remarks Related entries	Protein information Name, Synonyms Source, Sequence EC, PIR, SWISSPROT PDB code Biological unit Mutation details Secondary structure ASA Nucleic acid information: Name Source Type (DNA or RNA) Sequence (wild and mutant) Mutation details GenBank Number Complex information: PDB code, NDB code Conformation of protein Conformation of Nucleic Acid ASA Experimental condition: T, pH, Buffer, Ion, Additives Experimental method Binding data: Dissociation constant: K _d Association constant: K _a Free energy change: ΔG Enthalpy change: ΔH Heat capacity change: ΔC _p Literature: Reference, Author Keywords, Remarks

Previous publications (1–5) explain in detail the content and organization of the databases. Table 1 summarizes the contents of ProTherm and ProNIT. ProTherm and ProNIT are implemented in 3DinSight (8), a relational database system for structure, function and property of biomolecules. This facilitates more efficient search and retrieval of data by flexible queries, and enables users to gain insight into the relationship among structure, thermodynamics and function of proteins. We have been collecting the thermodynamic data from published original articles, by searching the PubMed literature database with a combination of specific terms, as well as by searching online journals probably containing thermodynamic data. The database does not contain any predicted or computational interaction data. Researchers then extract the relevant data from the selected articles. The input data are checked automatically by checking programs and also manually to avoid errors. Then, we upload the data first to a test site, where expert curators check and verify the data. After this checking, we upload the data to the public site for users. Furthermore, an email notification for each input entry is sent to the corresponding author, which enables the authors to check their own data and thereby improve the data validation.

DATABASE STATISTICS

We update both the databases frequently. The current release, ProTherm 5.0 contains 17113 entries from 771 proteins, retrieved from 1497 scientific articles, which is \sim 20% increase in data from the last version (4). Currently, the numbers of data for wild-type proteins, single, double and multiple mutants are 7014, 8202, 1277 and 620, respectively. Based on the solvent accessibility of mutants, 4426 mutations are buried, 2687 partially buried and 2751 exposed. In terms of secondary structures, 3993 mutations are in helix, 2622 in strand, 1227 in turn and 2467 in coil regions. Majority of data are obtained from CD (6825) and DSC (5294) experiments followed by fluorescence (3628). Further, 10154 data are obtained by thermal denaturation, 3890 and 2796 data from GdnHCl and urea denaturation, respectively.

Currently, ProNIT 2.0 contains 4900 entries from 273 research papers. There are 158 different DNA-binding proteins with 3489 wild-type entries and 1411 mutant entries. Majority of data are obtained by gel shift (1316), fluorescence (1143) and filter binding (1053), followed by calorimetry (727), surface plasmon resonance (185) and footprinting (168). Although proteins from a variety of organisms are present in ProNIT, majority of interaction data are from Escherichia coli proteins (1625) followed by Mus musculus (637) and *Homo sapiens* (569).

NEW FEATURES

There is a growing interest in the relationship between structure and thermodynamics of proteins. Thus, we try to provide link from thermodynamic data in ProTherm to structural information. So far, ProTherm data are connected to sequence and structural information of proteins through 3DinSight. We have added a new cross-link between ProTherm and STING (9), a comprehensive analysis tool for proteins with many structural descriptors. For given protein mutations searched within STING, each entry in STING report is connected to available thermodynamic data in ProTherm. Conversely, all the ProTherm data with available protein structure have pointers to the corresponding STING entry with detailed structural information. This cross-link will greatly facilitate the analysis of structure-thermodynamic relationship of proteins. The ProTherm page also provides cross-reference tables necessary for creating cross-links with PDB (10), PIR (11) and Swiss-Prot (12) databases. We have also added several new features in the search interface to make the search more efficient and convenient.

In the current release of ProNIT 2.0, we have included \sim 200 protein-RNA interaction data. To facilitate the retrieval of data based on DNA and RNA separately, we have added a new field called TYPE_NUC, where we provide the information about whether the nucleic acid sequence is DNA (singlestranded DNA, ssDNA, or double-stranded DNA, dsDNA) or RNA. Furthermore, a search option is added to retrieve data based on ssDNA, dsDNA or RNA. The protein nomenclature in the literature is not necessarily uniform. Hence, we have added a new field, SYNONYMS, in order to address this problem. Other additions of field are the SwissProt ID of the protein and 'RELATED_ENTRIES', which provides the list of entries that contain data from the same paper (the original paper usually contains multiple data and they are entered in different entries). We also provide a link to all homologous PDB codes with sequence identity of >95%. Also, display

options and sorting options are significantly improved. We have supplied the lists of ProNIT entries, protein names, protein sources, PDB code, NDB codes (13), authors and references in the advanced search page, along with a new query help page to help users for easy retrieval of the data. Several entries are deleted because of duplication, co-operative binding and so on, and the database entries are now renumbered. A mapping table, which relates the old and new entry numbers, is provided to help old users of ProNIT.

CITATION AND AVAILABILITY

The URLs for ProTherm and ProNIT are http://gibk26.bse. kyutech.ac.jp/jouhou/Protherm/protherm.html and http:// gibk26.bse.kyutech.ac.jp/jouhou/pronit/pronit.html, respectively. The users of ProTherm and ProNIT are requested to cite the references (4) and (5), respectively, in their publication including the above URLs. Users who use both the databases for their work may cite this article in their publications. Suggestions and other materials for inclusion in the databases are welcome and should be sent to either protherm@rtcmain.bse. kyutech.ac.jp or pronit@rtcmain.bse.kyutech.ac.jp.

ACKNOWLEDGEMENTS

The database development is partially supported by a Grant-in-Aid for Publication Scientific Research Results from the Japan Society for the Promotion of Sciences (JSPS). We also thank Advanced Technology Institute Inc. (ATI) for support. Funding to pay the Open Access publication charges for this article was provided by JSPS.

Conflict of interest statement. None declared.

REFERENCES

- 1. Gromiha, M.M., An, J., Kono, H., Oobatake, M., Uedaira, H. and Sarai, A. (1999) ProTherm: Thermodynamic database for proteins and mutants. Nucleic Acids Res., 27, 286-288.
- 2. Gromiha, M.M., Uedaira, H., An, J., Selvaraj, S., Prabakaran, P. and Sarai, A. (2002) ProTherm: thermodynamic database for proteins and mutants: developments in version 3.0. Nucleic Acids Res., 30, 301–302.
- 3. Sarai, A., Gromiha, M.M., An, J., Prabakaran, P., Selvaraj, S., Kono, H., Oobatake, M. and Uedaira, H. (2002) Thermodynamic databases for proteins and protein–nucleic acid interactions. *Biopolymers*, **61**, 121–126.
- 4. Abdulla Bava, K., Gromiha, M.M., Uedaira, H., Kitajima, K. and Sarai, A. (2004) ProTherm, version 4.0: thermodynamic database for proteins and mutants. Nucleic Acids Res., 32, D120-D121.
- 5. Prabakaran, P., An, J., Gromiha, M., Selvaraj, S., Uedaira, H., Kono, H. and Sarai, A. (2001) Thermodynamic database for protein-nucleic acid interactions (ProNIT). Bioinformatics, 17, 1027-1034.
- 6. Pfeil, W. (1998) Protein Stability and Folding: A Collection of Thermodynamic Data. Springer, NY.
- 7. Pfeil, W. (2001) Protein Stability and Folding, Supplement 1: A Collection of Thermodynamic Data. Springer, NY.
- 8. An,J., Nakama,T., Kubota,Y. and Sarai,A. (1998) 3DinSight: an integrated relational database and search tool for structure, function and property of biomolecules. Bioinformatics, 14, 188-195.
- 9. Neshich, G., Borro, L.C., Higa, R.H., Kuser, P.R., Yamagishi, M.E., Franco, E.H., Krauchenco, J.N., Fileto, R., Ribeiro, A.A., Bezerra, G.B. et al. (2005) The Diamond STING server. Nucleic Acids Res., 33, W29-W35
- 10. Berman, H.M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T.N., Weissig, H., Shindyalov, I.N. and Bourne, P.E. (2000) The Protein Data bank. Nucleic Acids Res., 28, 235-242.
- 11. Cathy, H.W., Yeh, L.L., Huang, H., Arminski, L., Jorge, C.A., Chen, Y., Hu,Z.Z., Ledley,R.S., Kourtesis,P., Suzek,B.E. et al. (2003) The Protein Information Resource. Nucleic Acids Res., 31, 345-347.
- 12. Boeckmann, B., Bairoch, A., Apweiler, R., Blatter, M.-C., Estreicher, A., Gasteiger, E., Martin, M.J., Michoud, K., O'Donovan, C., Phan, I. et al. (2003) The SWISS-PROT protein knowledgebase and its supplement TrEMBL in 2003. Nucleic Acids Res., 31, 365-370.
- 13. Berman, H.M., Zardecki, C. and Westbrook, J. (1998) The Nucleic Acid Database: a resource for nucleic acid science. Acta Crystallogr. D Biol. Crystallogr., **D54**, 1095–1104.