

This accepted version of the article may differ from the final published version.
This is an Accepted Manuscript for *Disaster Medicine and Public Health Preparedness* as
part of the Cambridge Coronavirus Collection
DOI: 10.1017/dmp.2020.348

Assessing Socioeconomic Vulnerabilities related to COVID-19 Risk in India: A State-level

Analysis

Dr. Praveen Kumar Pathak, Associate Professor, Department of Geography, Faculty of
Natural Sciences, Jamia Millia Islamia (Central University), New Delhi-110025, India

Mr. Yadawendra Singh, Assistant Professor, Department of Economics, Chandradhari Mithil
College, Lalit Narayan Mithila University, Darbhanga-846004, Bihar, India

Dr. Sandhya R. Mahapatro, Assistant Professor, A.N. Sinha Institute of Social Studies, Patna-
800001, Bihar, India

Dr. Niharika Tripathi, Research Associate, ICMR-National Institute of Medical Statistics, New
Delhi-110029, India

Ms. Jyoti Jee, Assistant Professor, Post-Graduate Department of Geography, Veer Kunwar
Singh University, Arrah-802301, Bihar, India

Corresponding Author:

Dr. Praveen Kumar Pathak, Associate Professor, Department of Geography, Faculty of
Natural Sciences, Jamia Millia Islamia (Central University), New Delhi-110025, India

Email: pathakprave@gmail.com

Abstract

Objective: There is a paucity of scientific analysis that has examined spatial heterogeneities in the socioeconomic vulnerabilities related to COVID-19 risk and potential mitigation strategies at the sub-national level in India. The present study examined the demographic, socioeconomic, and health system-related vulnerabilities shaping COVID-19 risk across 36 states and union territories in India.

Study Design: Using secondary data from the Ministry of Health and Family Welfare (MoHFW), Government of India; Census of India, 2011; National Family Health Survey, 2015-16; and various rounds of the National Sample Survey, we examined socioeconomic vulnerabilities associated with COVID-19 risk at the sub-national level in India from March 16, 2020 to May 03, 2020.

Methods: Descriptive statistics, principal component analysis (PCA), and the negative binomial regression model were used to examine the predictors of COVID-19 risk in India.

Results: There persist substantial heterogeneities in the COVID-19 risk across states and union territories in India. The underlying demographic, socioeconomic and health infrastructure characteristics drive the vulnerabilities related to COVID-19 in India.

Conclusion: This study emphasizes that concerted socially inclusive policy action and sustained livelihood/economic support for the most vulnerable population groups is critical to mitigate the impact of the COVID-19 pandemic in India.

INTRODUCTION

The novel Coronavirus disease, also referred to as COVID-19, is a global health emergency that has triggered an unprecedented catastrophe with respect to human lives and livelihood, disrupted economic systems cutting across sectors, halted public transportation networks, and restricted social interactions across the globe. On March 11, 2020, the World Health Organization (WHO) announced COVID-19 as a pandemic and called for decisive action including the need to: i) prepare and be ready; ii) detect, protect, and treat; iii) reduce transmission; iv) innovate and learn ¹. Since late December 2019, when early clusters of COVID-19 cases were reported from Wuhan City, Hubei Province of the People's Republic of China, more than 9.84 million confirmed COVID-19 cases of infection and 495,760 COVID-19 related deaths have been recorded across 216 countries and territories ². The impact of COVID-19 has been devastating across the globe, though its repercussions may be more serious for the developing countries characterized by a relatively large population base, great strain on the available resources (energy, food, water, land), inadequate public health infrastructure, fragile economic systems, and weak social safety programs ³.

Several national governments and global agencies have been struggling to tackle the COVID-19 pandemic⁴, seeking to contain the burgeoning risk of infection, fatality rate, improve access to affordable health care, ensure food security, and the protection of livelihood opportunities among the most vulnerable population groups (e.g. children, elderly, pregnant women, people with co-morbidities, people with disabilities, migrants, slum dwellers,)^{5–10}. Some of these vulnerable population groups are also the least likely to be able to practice the preventive measures related to physical distancing and self-isolation

given their health and livelihood constraints¹¹. Moreover, governments in many of these countries do not have enough resources for quarantine facilities^{12–14}.

It is important to note that any disaster, natural or man-made, spreads indiscriminately, although it disproportionately impacts the human population along the contours of socioeconomic and gendered inequities in the affected region^{15,16}. For instance, recent evidence has highlighted the disproportionate burden of COVID-19 risk among those aged 60 years and above; those afflicted with co-morbidities; disabilities, and poor nutritional status, as well as other marginalized groups^{17,18,19}. Therefore, regular assessment of underlying demographic, socioeconomic, epidemiological, and environmental exposure to COVID-19 risk is critical for an effective management and mitigation strategy, particularly among the low-and-middle-income countries^{20,21,22}.

India is among the top 10 countries most impacted by COVID-19 and has been fighting the pandemic since the first confirmed case was reported in January 2020 in Thrissur, one of the districts of the southern state of Kerala.²³ After the WHO declared COVID-19 a pandemic, the Union Government of India invoked the Disaster Management Act 2005 to implement a 21-day complete nationwide lockdown from March 25, 2020 to April 14, 2020 (Lockdown 1) to contain the spread of COVID-19²⁴. This was probably done to “flatten the epidemic curve” and to manage the contagion of the pandemic in such a way that it allows the public health system to meet the substantial surge of patients²⁵. Subsequently, the nationwide lockdown was extended from April 15, 2020 to May 03, 2020 (Lockdown 2); from May 04, 2020 to May 17, 2020 (Lockdown 3); and from May 18, 2020 to May 31, 2020 (Lockdown 4) with gradual relaxation of lockdown conditions in a phased

manner after continuous review of the emergency situation. The gradual slackening of lockdown, particularly in the low burden states and union territories, was provided to facilitate the revival of economic activities and routine human lives²⁶.

Given the large and relatively dense population base, a rising share of aging population with a poor healthcare system, regional inequalities in economic development and higher population mobility^{27–29}, India appears to be at greater risk of an increasing number of COVID-19 infected people. Identification of risk factors that are responsible for the rise/decline in the rate of disease spread is critical to provide direction for effective and long-term planning. It is opined that socioeconomic and environmental factors play a significant role in influencing the prevalence of any disaster including COVID-19³⁰. Although demographic pressures and geographical vulnerabilities enhance exposure to the disease, many pre-existing vulnerabilities such as age structure, morbidity, and poor environmental conditions also contribute to the spike in the spread of the pandemic. On the other hand, the resilience capacity of the state and communities, such as improved health infrastructure, better working conditions, and economic status may enable them to contain the disease spread³¹. Indian states are positioned at different stages of socioeconomic and demographic transition, hence, the vulnerability to COVID-19 risk may vary between and within states. One can expect that the vulnerability to COVID-19 would be high in the relatively laggard states of the country given their poorer socioeconomic, and health infrastructure status³².

The present study attempts to examine the demographic, socioeconomic, and health system-related vulnerabilities against COVID-19 risk across 36 states and union territories in India. Furthermore, a composite measure of socioeconomic vulnerabilities has been

generated by incorporating a diverse set of indicators ranging from demographic and socioeconomic characteristics to others such as health infrastructure, which may allow a comprehensive understanding of risk factors that augment the vulnerability to COVID-19 pandemic at the sub-national level in India³³. Findings of the present study may facilitate a long-term plan of action for containing the transmission of this pandemic. To the best of our knowledge, this is the first study to investigate the sub-national level vulnerabilities shaping COVID-19 risk across states and union territories in India.

METHODS

Data

Data employed for the study was compiled from multiple sources such as the Ministry of Health and Family Welfare, Government of India- 2020, COVID19India.org, Census of India- 2011, National Family Health Survey- 2015-2016, National Sample Survey 68th round- 2011-12, National Sample Survey 75th round- 2017-18, and National Sample Survey 76th round- 2018, Periodic Labour Force Survey- 2017-2018, and National Health Accounts- 2018³⁴⁻⁴³.

Study Design

The confirmed positive COVID-19 cases have been used as the main outcome variable in the study. The unit of analysis for the study is all 36 states and union territories of India. Table1 provides a detailed list of demographic, socioeconomic and health system-related indicators, data sources, and description of the indicators. We reviewed different

frameworks, ranging from the infectious disease vulnerability index, socioeconomic vulnerability indicator framework, and recent COVID-19 vulnerability assessment tools (focussed on socioeconomic inequalities, population characteristics, access to services, and epidemiological factors), to investigate the sub-national vulnerability assessment of COVID-19 risk in India^{33,44,45}. The infectious disease vulnerability framework identified seven domains (demographic, health care, public health, disease dynamics, political-domestic, political-international, and economic), which show a nation's/region's ability to prevent or contain a disease outbreak. . We adopted the modified infectious disease vulnerability framework according to its suitability and feasibility as per available data in the Indian context. The COVID-19 vulnerability assessment is performed using the selected domains related to demographic composition, socioeconomic structure, disease dynamics, health care, and public health system characteristics across 36 states and union territories in India.

Methods

We have applied bivariate and multivariate techniques to understand the sub-national level vulnerabilities associated with COVID-19 risk in India. The main outcome variable of the study is the number of COVID-19 positive cases for the period March 16, 2020 to May 03, 2020. We restricted the analysis to this date range because the national lockdown was implemented across the country with a comprehensive restriction on population movement between states and union territories. For the descriptive analysis, we classified the states and union territories into three broad categories (high, medium, and low burden states) based on the number of COVID-19 positive cases seen in each case. For instance, states and union territories with more than 3,000 positive cases until May 03, 2020 were categorized

into a high burden cluster, those with 1,000 to 3,000 COVID-19 positive cases were categorized as a medium burden cluster, and others with less than 1,000 COVID-19 positive cases were categorized as a low burden cluster (Appendix 1). We used COVID-19 positive cases, prevalence rate, positivity rate, recovery rate, and case-fatality rate to monitor the sub-national trends related to the COVID-19 pandemic in India during the study period. The exponential growth method was used to project the state and union territories total population using data from the Census of India, 2011, to estimate the COVID-19 prevalence rate and recovery rate per million population⁴⁶. A complete description of variables, measurements, and data sources is presented in Table 1. We used principal component analysis to compute the 'health infrastructure index' based on an array of health systems performance variables such as average population served per hospital, community health center, primary health center, sub-center, doctors, auxiliary nursing midwife, and hospital bed ratio^{47,48}. The negative value of the index suggests 'high vulnerability' whereas the positive value of the index indicates 'low vulnerability'. The reliability of the index was assessed using Cronbach's alpha test⁴⁹.

We used a negative binomial regression model to identify the association of various demographic, socioeconomic, and health system-related factors with the COVID-19 pandemic. A Poisson regression model is a commonly used model for count data. The assumption of the Poisson model is that variance is equal to mean. However, due to unobserved heterogeneity and clustering, data often presents an over dispersion. The negative Binomial regression model however, relaxes the equi-dispersion restriction of the Poisson model⁵⁰. Negative Binomial regression model typically uses the log link to relate the

mean of the data to the set of covariates. The regression model with log odds link function is given by:

$$\eta = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \dots + \beta_k X_k$$

where 'η' is log of the outcome variable that is the number of COVID-19 positive cases.

To summarize, from the sub-national heterogeneities related to the socioeconomic vulnerability to the COVID-19 pandemic in India, we generated four different sets of vulnerabilities indices (demographic susceptibility index, socioeconomic and disease exposure index, public health resilience index, and composite vulnerability index) using the principal component analysis. We mapped COVID-19 risk related vulnerability indices, prevalence rate, and recovery rate across states and union territories using ArcGIS software. Line graphs were plotted to present the trends of positive cases, positivity rate, recovery rate, deaths and case-fatality rate related to COVID-19. We performed all statistical analyses using STATA 13.0 software.

RESULTS

Sub-national Trends of COVID-19 in India

From March 16, 2020 to May 03, 2020, the prevalence rate of COVID-19 positive cases was 30 per million population in India with substantial inter-state differentials. For instance, 10 states and union territories had higher prevalence rates of COVID-19 positive

cases than the national average, which included among others, Delhi (228), Maharashtra (101), Gujarat (77), Tamil Nadu (37), Rajasthan (35) and Madhya Pradesh (33) (Figure1). The prevalence of recovered cases of COVID-19 in India was 8 per million population, marked by substantial inter-state variability during the period of March 16, 2020 to May 03, 2020.

The sub-national average weekly COVID-19 positive cases, and the corresponding positivity rate suggest that the average number of weekly new confirmed COVID-19 positive cases have increased continuously across states and union territories of India (Figure 2). On an all India basis, the cases increased exponentially from 41 on March 16, 2020 to 2227 on April 27, 2020. We calculated the positivity rate, which is the percent of confirmed COVID-19 positive cases out of the total tested case. The positivity rate has been presented from March 30, 2020 when data of COVID-19 testing became available (Figure 2). The positivity rate declined from 3.9 per cent to 2.9 per cent during the first week (March 16, 2020 to March 23, 2020) in India. After April 06, 2020, there was not much variation in the all India positivity rate; it was 2.6 per cent for the last week of our analysis (April 27, 2020 to May 03, 2020).

Subsequently, the average weekly recovered cases and recovery rate from COVID-19 has been analyzed (Figure 3). It is observed that the average weekly recovered cases have increased continuously for all the categories in states across India. The all India average weekly recovered cases increased from one in the first week to 792 in the last week. Data suggests that recovery rate, (recovered cases as a percentage of total COVID-19 positive cases), has improved for India and all its states.

While examining the COVID-19 related deaths and the case-fatality rate across states, it emerged that the average weekly deaths due to COVID-19 increased continuously over time (Figure 4). The average weekly deaths increased from none during the first week (March 16, 2020) to 76 during the week of April 27, 2020 in India. It was observed that of these 76 COVID-19 deaths, 66 occurred in the high category states, while the remaining casualties occurred in the low category states.

Sub-national Analysis of Different Dimension of the Socioeconomic Vulnerabilities related to COVID-19 positive cases

We analyzed the patterning of COVID-19 positive cases across a set of selected factors such as demographic susceptibility, socioeconomic and disease exposure, and public health resilience capacity characteristics at the sub-national level in India (Table 2, Table 3, and Table 4). The sub-national analysis of demographic susceptibility patterns related to COVID-19 shows that states such as Maharashtra, Gujarat, Delhi and Tamil Nadu which recorded the highest burden of COVID-19 positive cases, also happened to have relatively higher employment-related in-migration (Table 2). On the other hand, the north and eastern states including Bihar and Jharkhand, belonging to the low COVID-19 category, reported a smaller volume of migrants. A similar pattern was noticed concerning international migration as well. The association between level of economic development and prevalence of the COVID-19 pandemic shows that states having a higher burden of infections seemed to have a low and moderate level of poverty, whereas around 43 percent of laggard states reported a low level of infection and had a high level of poverty.

The relationship between education and COVID-19 infection indicates that states with a higher burden of COVID-19 infections have a moderate and higher literacy rate. Among demographic factors, age and co-morbidity are the pre-existing vulnerabilities which increases susceptibility to the disease. Among states with a moderate level of COVID-19 positive cases, a moderate and high percent of elderly population was observed. Half of the states with relatively high burden of COVID-19 positive cases also reported higher percent of the ageing population. Furthermore, with regard to the prevalence of morbidity, majority of the states reported a moderate burden of co-morbidities.

Sub-national analysis of socioeconomic and disease exposure patterns related to COVID-19 pandemic shows that states with a relatively higher burden of the disease reported moderate and higher population density (Table 3). Around three-fourth of the states reporting a higher number of infected cases such as Maharashtra, Delhi and Tamil Nadu also had a relatively higher slum population. States with a moderate burden of infection had a moderate and higher concentration of slum population. The association of level of urbanization with the burden of COVID-19 shows that high disease burden states recorded a moderate and higher level of urbanization. With regard to the demographic structure of the family, it was found that half the high and moderate level infected states recorded a higher proportion of multi-generational families, whereas only 28 percent of low COVID-19 states recorded a higher number of a multi-generational families.

The sub-national analysis of public health resilience patterns related to COVID-19 pandemic shows that among the low COVID-19 burden states, about 24 percent recorded low scores in the health infrastructure index, while, 38 percent scored high (Table 4).

Average healthcare expenditure in low disease burden states was substantially higher than in states with a higher burden of disease. Almost 38 percent of low COVID-19 burden states spent more on public health, whereas, none of the moderate and high COVID-19 burden states came into this category.

Multivariate Results

The descriptive analysis demonstrates substantial heterogeneities in the socioeconomic vulnerabilities associated with the burden of COVID-19 across states and union territories in India. We have fitted a negative binomial regression model to predict the risk of COVID-19 adjusting for demographic, socioeconomic, and health infrastructural characteristics across states and union territories in India (Table 5). The estimated regression model indicates that the proportion of elderly, the prevalence of any ailment, interstate migration, international migration, drinking of water outside household premises, health infrastructure, population density, proportion of urban population, proportion of non-agricultural casual laborers, and joint/extended families, had a statistically significant and positive association with the risk of COVID-19 across states and union territories in India. For example, for one unit increase in the proportion of elderly population, and prevalence of any ailment were associated with 0.701 unit and 0.005 unit increase respectively in the COVID-19 risk, after adjusting for other socioeconomic, demographic, and health infrastructure characteristics. Likewise, the proportion of literates, hand washing before the meal and mass media exposure also had a statistically significant and negative association with COVID-19 risk. For instance, a one unit increase in literacy rate, hand washing before the meal and mass media exposure were

associated with 0.306, 0.041, and 1.236 unit decreases respectively in the COVID-19 risk across states and union territories in India.

Assessment of Sub-national Level Vulnerabilities

The composite vulnerability index at the sub-national level considering demographic susceptibility, socioeconomic and disease exposure, and public health resilience indicators suggest that the eastern states and parts of the northern states (accounting for 62 percent of India's total population) were highly vulnerable to COVID-19 risk, and faced disproportionate challenges related to mitigation of this pandemic (Figure 5).

DISCUSSION AND CONCLUSION

The present study makes a novel attempt to examine sub-national heterogeneities in the underlying socioeconomic vulnerabilities related to COVID-19 risk across states and union territories in India. Drawing upon the infectious disease vulnerability framework, the study attempts to quantify the COVID-19 risk and examined the potential policy solutions to mitigate the menace of this pandemic across states and union territories in India. Major findings of the study suggest that there persist substantial heterogeneities in the burden of COVID-19 positive cases, recovery rates, case-fatality rates and testing performance across Indian states. Overall, the COVID-19 risk has been increasing at an exponential rate, particularly in the high COVID-19 burden states of Maharashtra, Delhi, and Gujarat (an almost 90-fold increase during the study period). On the other hand, the medium COVID-19

burden states observed close to a 28-fold increase in the disease burden during the same period, followed by a 12-fold increase among the low COVID-19 burden states across India.

The unequal progression of COVID-19 across different states is an outcome of the complex interplay of the underlying demographic, socioeconomic, and health infrastructure heterogeneities. Examining the patterns of COVID-19 risk by the selected susceptibility factors across Indian states through the descriptive analysis, it was found that the high burden COVID-19 states were characterized by relatively higher inter-state migration, international migration, a large proportion of elderly population and an increased share of Scheduled Castes/Scheduled Tribes. In addition, analysis of the patterns of COVID-19 risk by the selected exposure factors across states suggest that the high burden COVID-19 states had relatively higher population density, a disproportionately high share of urban and slum population, and a relatively higher average number of persons living per room. Furthermore, the patterns of COVID-19 risk by the selected resilience factors highlights that the high burden COVID-19 states were marked by relatively poor health infrastructure, lower public health expenditure and high mass media exposure. This finding highlights that spatial heterogeneities associated with the resilience vulnerabilities might determine the course of mitigation and recovery prospects in the fight against COVID-19 risk across states in India. The composite index of vulnerabilities to COVID-19 risk indicates that the northern, central, and eastern states suffered from multidimensional vulnerabilities, and also exhibited limited resilience capabilities to tackle the COVID-19 risk. However, vulnerable population groups/states are dynamic, and those falling in these clubs may change over time owing to socially inclusive policy actions⁶.

The multivariate regression analysis confirmed that a disproportionately large share of elderly population, prevalence of co-morbidities, predominant inter-state and international migration, proportion of non-agricultural casual laborers, joint/extended families, multipurpose/business-related use of households, and level of urbanization had a statistically significant and positive association with COVID-19 risk across states and union territories in India. On the other hand, educational status, hand washing before meals and exposure to mass media had a statistically significant and negative association with the COVID-19 risk across states and union territories in India. These findings have important implication for the mitigation of COVID-19 risk in India. The regression results revealed that older adults were more vulnerable to COVID-19 as compared to the rest of the population. Our finding is corroborated by various studies, which suggest that not only is the probability of the elderly getting infected with COVID-19 higher, but also the case fatality rate is found to be higher among them^{51–54}. It is obvious that states and union territories having a higher proportion of older adults have a higher burden of COVID-19 risk in comparison with other states. Therefore, there is need for policy action to encourage physical distancing among the elderly population. Apart from these social distancing policy measures, these states are also expected to divert more of its resources towards the care of older adults.

Our findings showed that the risk of COVID-19 was very high for those who were already suffering from pre-existing diseases such as diabetes, heart disease, or respiratory problems. Our finding is in line with studies, which have suggested that co-morbidity is an important risk factor for COVID-19^{55–57}. These results are critical for those states and union territories with a higher proportion of older adults since co-morbidity is often found to be

significantly higher among them. Therefore, it is imperative that the public health system is such that states are prepared to attend non-COVID related ailments with equal sensitivity.

The level of urbanization, population density and slum population emerged as important risk factors for COVID-19 across states and union territories in India. Most of the highly urbanized, densely populated and large slum cluster states and union territories have emerged as the hotspot of COVID-19 risk in India including Maharashtra, Gujarat, Delhi and Tamil Nadu. This scenario is further aggravated in states with a higher volume of labor migration working in the informal sector. The large-scale inter-state migration mainly for informal casual work as observed from the analysis has emerged as an important risk factor of COVID-19 transmission across states. The migrants, who were invariably engaged in low profile jobs owing to their limited education, were less aware of the seriousness of the disease, and lived under suboptimal conditions and hence, were at higher risk of being infected with COVID-19. Crowded living environments and congested living place limit the effective implementation of preventive measures such as social distancing, and exposure to the disease was disproportionately higher for urban poor and labor migrants living in slums^{7,8,58}. Therefore, it is crucial to generate public awareness about the risk of COVID-19 and precautions to avoid the same. This is further supported by the findings related to safe sanitation practices and exposure to mass media that seem to be associated with significantly lower risk of COVID-19 in India. Efforts must be made to promote the use of safe sanitation practices and dissemination of accurate and verified information through television, radio and newspaper related to restricting the spread of COVID-19.

Health infrastructure had a statistically significant and positive association with the COVID-19 risk across states and union territories in India. States with relatively robust health care systems are better able to perform a large number of testing, treatments and recording of COVID-19 cases. In states with poor health infrastructure and limited testing facilities, the risk of COVID-19 tends to remain under-reported. This underscores the significance of a strong public healthcare system to effectively implement preventive and curative measures and lower the risk of COVID-19 spread. It is essential to revitalize the public health system with the deployment of medical professionals, supply of personal protective equipment (PPE) kits, intensive care unit (ICU) and ventilator facilities in order to meet the demand of COVID-19 patients as required. Public health system preparedness can go a long way towards mitigating the burden of COVID-19 risk in India⁵⁹.

AUTHOR STATEMENTS

Funding. No funding was received for this research.

Conflicts of Interest. The authors have no conflicts of interest to declare.

Ethical Approval. The study was based on a secondary dataset with no identifiable information on the survey participants. This dataset is available in the public domain for research use and, hence, no approval was required from any institutional review board.

REFERENCES

1. World Health Organization. WHO Director-General's opening remarks at the Mission briefing on COVID-19 - 12 March 2020. World Health Organisation. Published 2020. Accessed May 17, 2020. <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-mission-briefing-on-covid-19---12-march-2020#:~:text=WHO Director-General's opening remarks at the Mission briefing,COVID-19 - 12 March 2020&text=Excellencies%2C dear col>
2. World Health Organization. *Coronavirus Disease (COVID-19) Situation Report-160*.; 2020. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200628-covid-19-sitrep-160.pdf?sfvrsn=2fe1c658_2
3. Ahmed F, Ahmed N, Pissarides C, Stiglitz J. Why inequality could spread COVID-19. *Lancet Public Heal*. Published online 2020. doi:10.1016/S2468-2667(20)30085-2
4. Hopman J, Allegranzi B, Mehtar S. Managing COVID-19 in Low- and Middle-Income Countries. *JAMA*. Published online 2020. doi:10.1001/jama.2020.4169
5. Smith JA, Judd J. COVID-19: Vulnerability and the power of privilege in a pandemic. *Heal Promot J Aust*. Published online 2020. doi:10.1002/hpja.333
6. The Lancet. Redefining vulnerability in the era of COVID-19. *Lancet*. Published online 2020. doi:10.1016/S0140-6736(20)30757-1
7. Corburn J, Vlahov D, Mberu B, et al. Slum Health: Arresting COVID-19 and Improving Well-Being in Urban Informal Settlements. *J Urban Heal*. Published online 2020. doi:10.1007/s11524-020-00438-6
8. International Organization for Migration. *COVID-19 and the Risk of Exacerbating Existing Vulnerabilities*.; 2019. https://www.iom.int/sites/default/files/our_work/ICP/MPR/migration_factsheet_6_covid-19_and_migrants.pdf
9. Wisner B, Adams J. *Environmental Health in Emergencies and Disasters- a Practical Guide*.; 2002.
10. Plough A, Bristow B, Fielding J, Caldwell S, Khan S. Pandemics and health equity: Lessons learned from the H1N1 response in Los Angeles County. *J Public Heal Manag Pract*. 2011;17(1):20-27. doi:10.1097/PHH.0b013e3181ff2ad7
11. Brown C, Ravallion M, van de Walle D. Can the World's Poor Protect Themselves from the New Coronavirus? *Natl Bur Econ Res*. Published online 2020. doi:10.3386/w27200
12. Thacker T. Covid-19 spread: Poor conditions of quarantine facilities come into focus in India - The Economic Times. *The Economic Times*. March 2020.
13. Pillai S. Cramped quarantine facilities for health care staff give rise to fear of Covid-19 spread.

Hindustan Times. May 2020.

14. Hussain Y, Muhammad K, Umar MF, Omerkhail A, Khan Z. COVID-19 in five neighbouring limited resources countries: A financial and health threats. *Value Heal Reg Issues*. Published online 2020. doi:10.1016/j.vhri.2020.06.001
15. Donner W, Rodriguez H. Population Composition, Migration and Inequality: The Influence of Demographic Changes on Disaster Risk and Vulnerability. *Soc Forces*. Published online 2008. doi:10.1353/sof.0.0141
16. Phelan JC, Link BG, Tehranifar P. Social Conditions as Fundamental Causes of Health Inequalities: Theory, Evidence, and Policy Implications. *J Health Soc Behav*. Published online 2010. doi:10.1177/0022146510383498
17. UNDRR, HelpAge International, UN Women. *UNDRR Asia-Pacific COVID-19 Brief: Leave No One Behind in COVID-19 Prevention, Response and Recovery.*; 2020.
<https://www.undrr.org/publication/undrr-asia-pacific-covid-19-brief-leave-no-one-behind-covid-19-prevention-response-and>
18. IRFC, OCHA, WHO. *COVID-19: How to Include Marginalized and Vulnerable People in Risk Communication and Community Engagement.*; 2020.
19. Jensen L, Molina GG. *COVID 19 and Health System Vulnerabilities in the Poorest Developing Countries.*; 2020.
20. Fakhruddin B (SHM), Blanchard K, Ragupathy D. Are we there yet? The transition from response to recovery for the COVID-19 pandemic. *Prog Disaster Sci*. 2020;7:100102. doi:10.1016/j.pdisas.2020.100102
21. Walker PGT, Walker PGT, Whittaker C, et al. The impact of COVID-19 and strategies for mitigation and suppression in low- and middle-income countries. *Science (80-)*. 2020;369(June):413-422. doi:10.1126/science.abc0035
22. Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will country-based mitigation measures influence the course of the COVID-19 epidemic ? *Lancet*. 2020;395(10228):931-934. doi:10.1016/S0140-6736(20)30567-5
23. Varghese G, John R. COVID-19 in India: Moving from containment to mitigation. *Indian J Med Res*. Published online 2020. doi:10.4103/ijmr.ijmr_860_20
24. Ministry of Home Affairs. *Disaster Management Act-2005*. Government of India; 2005:1-29.
25. The Lancet. India under COVID-19 lockdown. *Lancet*. 2020;395(10233):1315. doi:10.1016/S0140-6736(20)30938-7
26. Prusty N, Khanna S. India extends lockdown to May 31, to relax rules in some areas. *Reuters*. May 2020.

27. Balarajan Y, Selvaraj S, Subramanian S. Health care and equity in India. *Lancet*. Published online 2011. doi:10.1016/S0140-6736(10)61894-6
28. James KS. India's Demographic Change: Opportunities and Challenges. *Science* (80-). 2011;333(July):576-580.
29. Agarwal A, Lubet A, Mitgang E, Mohanty S, Bloom DE. *Population Aging in India: Facts, Issues, and Options.*; 2016. doi:10.1007/978-981-10-0230-4_13
30. Group WB. *Disease Control Priorities Improving Health and Reducing Poverty.*; 2018. doi:10.1596/978-1-4648-0527-1_ch8
31. Phibbs S, Kenney C, Severinsen C, Mitchell J, Hughes R. Synergising public health concepts with the sendai framework for disaster risk reduction: A conceptual glossary. *Int J Environ Res Public Health*. Published online 2016. doi:10.3390/ijerph13121241
32. Kumar N, Rani R. Regional Disparities in Social Development: Evidence from States and Union Territories of India. *South Asian Surv*. 2019;26(1):1-27. doi:10.1177/0971523118825388
33. Jhan HT, Ballinger R, Jaleel A, Ting KH. Development and application of a socioeconomic vulnerability indicator framework (SVIF) for local climate change adaptation in Taiwan. *Sustain*. 2020;12(4). doi:10.3390/su12041585
34. Ministry of Health and Family Welfare. COVID-19 State-wise status. Government of India. Published 2020. Accessed May 3, 2020. <https://www.mohfw.gov.in/#state-data>
35. [Coronavirus Outbreak in India - covid19india.org. https://www.covid19india.org/](https://www.covid19india.org/). Accessed May 04, 202036. International Institute for Population Sciences. *National Family Health Survey (NFHS-4) 2015-16 India.*; 2017. doi:kwm120 [pii]10.1093/aje/kwm120
37. Ministry of Statistics and Programme Implementation. *Key Indicators of Household Consumption Expenditure in India 2011-2012.*; 2013. <http://www.icssrdataservice.in/datarepository/index.php/catalog/91>
38. Ministry of Statistics and Programme Implementation. *Key Indicators of Social Consumption in India: Health (2017-2018).*; 2019. http://www.mospi.gov.in/sites/default/files/publication_reports/KI_Health_75th_Final.pdf
39. Ministry of Statistics and Programme Implementation. *Drinking Water, Sanitation, Hygiene and Housing Condition in India (2018).*; 2019. http://www.mospi.gov.in/sites/default/files/publication_reports/Report_584_final_0.pdf
40. Ministry of Statistics & Programme Implementation. *Periodic Labor Force Survey (2017-2018).*; 2019. http://www.mospi.gov.in/sites/default/files/publication_reports/Annual Report%2C PLFS 2017-18_31052019.pdf
41. Ministry of Health and Family Welfare. *National Health Accounts Estimates of India.*; 2018.

- https://main.mohfw.gov.in/sites/default/files/NHA_Estimates_Report_2015-16_0.pdf
42. Director General of Health Services. *National Health Profile 2015*.; 2015.
<https://www.cbhidghs.nic.in/WriteReadData/l892s/NHP-2015.pdf>
43. Director General of Health Services. *Nation Health Profile 2018 13th Issue*.; 2018.
www.who.int/classifications
44. Moore M, Gelfeld B, Okunogbe A, Paul C. *Identifying Future Disease Hot Spots: Infectious Disease Vulnerability Index*.; 2016. doi:10.7249/rr1605
45. Macharia PM, Joseph NK, Okiro EA. A vulnerability index for COVID-19: spatial analysis to inform equitable response in Kenya. *medRxiv*. Published online 2020:2020.05.27.20113803. doi:10.1101/2020.05.27.20113803
46. Office of the Registrar and Census Commissioner. Census of India 2011. Ministry of Home Affairs, Government of India. Published 2011. Accessed May 10, 2020.
<https://censusindia.gov.in/>
47. Reckien D. What is in an index? Construction method, data metric, and weighting scheme determine the outcome of composite social vulnerability indices in New York City. *Reg Environ Chang*. Published online 2018. doi:10.1007/s10113-017-1273-7
48. Opiyo FEO, Wasonga O V., Nyangito MM. Measuring household vulnerability to climate-induced stresses in pastoral rangelands of Kenya: Implications for resilience programming. *Pastoralism*. Published online 2014. doi:10.1186/s13570-014-0010-9
49. Tavakol M, Dennick R. Making sense of Cronbach's alpha. *Int J Med Educ*. 2011;2:53-55. doi:10.5116/ijme.4dfb.8dfd
50. An Q, Wu J, Fan X, Pan L, Sun W. Using a negative binomial regression model for early warning at the start of a hand foot mouth disease epidemic in Dalian, Liaoning Province, China. *PLoS One*. Published online 2016. doi:10.1371/journal.pone.0157815
51. Wang L, He W, Yu X, et al. Coronavirus disease 2019 in elderly patients: Characteristics and prognostic factors based on 4-week follow-up. *J Infect*. Published online 2020. doi:10.1016/j.jinf.2020.03.019
52. Liu K, Chen Y, Lin R, Han K. Clinical features of COVID-19 in elderly patients: A comparison with young and middle-aged patients. *J Infect*. Published online 2020. doi:10.1016/j.jinf.2020.03.005
53. Wu JT, Leung K, Bushman M, et al. Estimating clinical severity of COVID-19 from the transmission dynamics in Wuhan, China. *Nat Med*. 2020;26(4):506-510. doi:10.1038/s41591-020-0822-7
54. Lloyd-Sherlock P, Ebrahim S, Geffen L, McKee M. Bearing the brunt of covid-19: Older people

- in low and middle income countries. *BMJ*. 2020;368(March):1-2. doi:10.1136/bmj.m1052
55. McMichael TM, Currie DW, Clark S, et al. Epidemiology of Covid-19 in a Long-Term Care Facility in King County, Washington. *N Engl J Med*. Published online 2020:2005-2011. doi:10.1056/NEJMoa2005412
56. Jordan RE, Adab P, Cheng KK. Covid-19: Risk factors for severe disease and death. *BMJ*. 2020;368(March):1-2. doi:10.1136/bmj.m1198
57. Yang J, Zheng Y, Gou X, et al. Prevalence of comorbidities and its effects in coronavirus disease 2019 patients: A systematic review and meta-analysis. *Int J Infect Dis*. 2020;94:91-95. doi:10.1016/j.ijid.2020.03.017
58. Khatua S. Density, Distancing, Informal Settlements and the Pandemic. *Econ Polit Wkly*. 2020;55(20). <https://www.epw.in/journal/2020/20/commentary/density-distancing-informal-settlements-and.html>
59. Isaac TMT, Sadanandan R. COVID-19, Public Health System and Local Governance in Kerala. *Econ Polit Wkly*. 2020;55(21). <https://www.epw.in/journal/2020/21/perspectives/covid-19-public-health-system-and-local-governance.html>

Table 1: Description of the study variables

Risk Factors	Description of variables	Data Source
Outcome variable	Number of COVID-19 cases	Ministry of Health and Family Welfare, Government of India; www.covid19india.org
Demographic composition	<ul style="list-style-type: none"> ▪ Percent of Scheduled Caste/Scheduled Tribe population ▪ Percent of Muslim population ▪ Percent of elderly population (60 years and above) ▪ Percent of literate population ▪ Number of employment related inter-state migrants during last five year ▪ Number of employment related international migrants during last five year ▪ Percent of slum population ▪ Percent of urban population ▪ Population density (Number of persons per square kms) 	Census of India, 2011
Disease dynamics	<ul style="list-style-type: none"> ▪ Co-morbidity (At least one ailment per 1000 population). The diseases considered in the analysis are HIV AIDS, Cancer, Diabetes, Other endocrine, metabolic, nutritional diseases including obesity, Hypertension, Heart disease, and Bronchial asthma. 	National Sample Survey, 75 th round, 2017-18
Health care and public health	<ul style="list-style-type: none"> ▪ Percent washing hand before meal ▪ Percent washing hand after defecation ▪ Percent household used for purpose other than residential ▪ Percent of households bringing water from outside household premise ▪ Average people living per room ▪ Percent of joint/extended family 	National Sample Survey, 76 th round, 2018

	<ul style="list-style-type: none"> Health infrastructure Index constructed using average population covered per primary health center (PHC), Community Health center (CHC), Sub-Centre, District Hospital, Auxiliary Nursing Midwife (ANM), Doctors, Hospital-bed ratio Per capita health expenditure 	<p>National Health Profile, 2018</p> <p>National Health Profile, 2015</p>
Socio-economic structure	<ul style="list-style-type: none"> Mean score of mass media exposure based on the relative frequency of watching television/listening radio/reading newspaper 	National Family Health Survey, 2015-16
	<ul style="list-style-type: none"> Percent living below the poverty line 	National Sample Survey, 68 th round, 2011-12
	<ul style="list-style-type: none"> Percent casual laborers in agriculture Percent casual laborers in non-agriculture 	Periodic Labour Force survey, 2018

Table 2: Sub-national analysis of COVID-19 related demographic susceptibility patterns in India

Characteristics	Low Burden COVID-19 Cluster	Moderate Burden COVID-19 Cluster	High Burden COVID-19 Cluster
Inter-state migration			
Low	BR, JK, AN, PY, AN, ML, MN, AR, MZ (42.9)	(0.0)	(0.0)
Middle	WB, KL, OR, JH, CH, UT, CT, AS, HP, GA (47.6)	(0.0)	(0.0)
High	KR, HR (9.5)	RJ, MP, UP, AP, PB, TG (100.0)	MH, DL, GJ, TN (100.0)
International Migration			
Low	JK, AN, PY, NE (38.1)	(0.0)	(0.0)
Middle	BR,OR, JH,CH,CTGA,MZ(33.3)	RJ, MP, AP, TG (66.7)	(0.0)
High	WB, KR,KL,HR,UT, HP ((28.6)	UP, PB (33.3)	MH, DL, GJ, TN (100.0)
Casual labour in non-agriculture			
Low	CH, UT, GA, MN, AR, MZ (28.6)	(0.0)	MAH, DEL, GJ (75.0)
Middle	JK, KR, BR, HR, AS, AN, TR (33.3)	RJ, UP, PB, TG (75.0)	(0.0)
High	WB, KL, OR, JH, CT, HP, ML, PY (38.1)	MP, AP (25.0)	TN(25.0)
Poverty			
Low	KL, HP, AN, PY, GA (23.8)	AP, TG, PB (50.0)	DL (25.0)
Middle	WB, JK, HR, UT, TR, MG, MZ (33.3)	RJ (16.7)	TN, GJ, MH (75.0)
High	JH, BR, OR, CH, CT, AS, MN, AR (42.9)	MP, UP (33.3)	(0.0)
Elderly Population			
Low	CH, AS, AN, MG, MN, AR, MZ (33.3)	(0.0)	DL (25.0)
Middle	WB, JK, BR, HR, JH, CT, TR (33.3)	RJ, MP, UP (50.0)	GJ (25.0)
High	KR, KL, OR, UT, HP, PY, GA (33.3)	AP, TG, PB (50.0)	MH, TN (50.0)
Ailment			
Low	BR, JH, UT, CH, CT, AS, MG, MN (38.1)	TG (16.7)	(0.0)
Middle	JK, KR, PY, AR, MZ (23.8)	MP, UP, PB (50.0)	TN, GJ, DL (75.0)
High	WB, KL, HR, OR, HP, AN, TR, GA (38.1)	RJ, AP (33.3)	MH (25.0)
Literacy			
Low	JK, BR, JH, CT, AS, ML, AR	RJ, MP, UP, AP, TG	(0.0)

	(33.3)	(75.0)	
Middle	WB, KA, HR, OR, UT, MN (28.6)	PB(25.0)	MH, GJ, TN (75.0)
High	KL, HP, CH, GA, AN, TR, PY, MZ (38.1)	(0.0)	DL(25.0)
Scheduled caste/Scheduled tribe population			
Low	OR, CT, HP, CH, MG, AR, MZ (33.3)	PB (16.7)	TN (25.0)
Middle	HR, AN, TR, PY, GA, MN (28.6)	RJ, MP, AP, TG (66.7)	GJ (25.0)
High	WB, JK, KR, BR, KL, JH, UT, AS (38.1)	UP (16.7)	TN, MH (50.0)
Muslim Population			
Low	JK, BR, KL, HR, CH, AS, AN, PY, GA (42.9)	(0.0)	DL, TN (50.0)
Middle	WB, KR, UT, HP, MN (23.8)	RJ, UP, AP, PB, TG (83.3)	MH, GJ (50.0)
High	OR, JH, CT, TR, MG, AR, MZ (33.3)	MP (16.7)	(0.0)
Total Observations	21	6	4

Note: State and union territories of Nagaland, Sikkim, Lakshadweep, Dadar & Nagar Haveli, and Daman & Diu had zero COVID-19 positive case; Figures in the parenthesis indicate the percent distribution of states and union territories by selected characteristics across low, moderate, and high burden COVID-19 states; Appendix-1 provide details of abbreviated names of the state and union territories used above.

Table 3: Sub-national analysis of COVID-19 related socioeconomic and disease exposure patterns in India

Characteristics	Low Burden COVID-19 Cluster	Moderate Burden COVID-19 Cluster	High Burden COVID-19 Cluster
Population Density			
Low	JK, UT, CH, HP, AN, ML, MN, AR, MZ (42.9)	RJ (16.7)	(0.0)
Middle	KR, OD, JH, AS, TR, GA (28.6)	MP, AP, PB, TG (66.6)	MH, GJ(50.0)
High	WB, BR, KL, HR, CH, PY (28.5)	UP (16.7)	DL, TN(50.0)
Slum Population			
Low	BR, KL, JH, AS, HP, ML, GA, MN, AR(42.9)	(0.0)	(0.0)
Middle	JK, KR, OR, UT, AN, TR (28.6)	RJ, UP, PB (50.0)	GJ (25.0)
High	WB, HR, CH, CT, PY, MZ (28.5)	MP, AP, TG (50.0))	MH, DL, TN (75.0)
Urban Population			
Low	JK, BR, HR, OR, CT, AS, TR, MN, AR (42.9)	RJ, UP (33.3)	(0.0)
Middle	WB, JH, UT, HP, AN, MZ (28.6)	AP, PB, TG (50.0)	MH, GJ (50.0)
High	KR, KL, CH, MG, PY, GA (28.5)	MP (16.7)	DL, TN (50.0)
Joint Families			
Low	CH,AS,AN,TR,MG,PY,GA,AR (38.1)	TG(16.7)	TN (25.0)
Middle	WB,JK,OR,CT,UT,MN,MZ (33.3)	UP, AP(33.3)	DL (25.0)
High	KR, BR, KL, HR, JH, HP(28.6)	RJ, MP, PB (50.0)	MH, GJ (50.0)
Average person living per room			
Low	JK, KL, UT AS, HP, AN, ML, GA, MZ (43.0)	(0.0)	(0.0)
Middle	WB, KR, HR, CH, CT, TR, PY, MN (38.0)	AP, PB, TG (50.0)	TN (25.0)
High	BR, OR, JH, AR (19.0)	RJ, MP, UP, (50.0)	MH, GJ, DL (75.0)
Hand Washing			
Low	WB, BR, OR, JH, AS, TR (28.6)	RJ, UP, AP (50.0)	TN (25.0)
Middle	JK, KR, UT, CT, MG, MN, MZ (33.3)	MP, TG (33.3)	GJ (25.0)
High	KL, HR, CH, AN, GA, PY, AR (38.1)	PB (16.7)	MH, DL (50.0)

Note: Figures in the parenthesis indicate the percent distribution of states and union territories by selected characteristics across low, moderate, and high burden COVID-19 states; Appendix-1 provide details of abbreviated names of the state and union territories used above.

Table 4: Sub-national analysis of COVID-19 related public health resilience patterns in India

Characteristics	Low Burden COVID-19 Cluster	Moderate Burden COVID-19 Cluster	High Burden COVID-19 Cluster
Health Index			
Low	WB, KR, BR, JH, CT (23.8)	MP, UP, AP, TG (66.7)	MH, GJ (50.0)
Moderate	JK, KL, HR, OR, UT, AS, MG, MN (38.1)	RJ, PB (33.3)	TN (25.0)
High	CH, HP, AN, TR, PY, GA, AR, MZ (38.1)	(0.0)	DL(25.0)
Mean Per capita Health Expenditure			
Low	WB,KR,BR,OR,JH(28.6)	MP,AP,UP,PB(67.0)	MH,GJ(50.0)
Moderate	KL,UT,CT,AS,TR,MG,MN(33.3)	RJ,TG(33.0)	DL,TN(50.0)
High	JK,CH,HP,AN,PY,GA,AR,MZ(38.0)	(0.0)	(0.0)
Media Exposure			
Low	WB, JH,OR,BR, AS, TR, AR (33.3)	RJ, UP, MP(50.0)	(0.0)
Moderate	JK,HR,UT,CH,AN,MG(28.6)	AP, TG (33.3)	MH, GJ (50.0)
High	KR,KL,CH,HP,PY,GA,MN,MZ (38.1)	PB(16.7)	DL, TN (50.0)

Note: Figures in the parenthesis indicate the percent distribution of states and union territories by selected characteristics across low, moderate, and high burden COVID-19 states; Appendix-1 provide details of abbreviated names of the state and union territories used above.

Table 5: Estimated negative binomial regression coefficients predicting the COVID-19 risk by selected susceptibility, exposure and resilient characteristics across the states/union territories in India

Characteristics	Coefficient	Standard error	z-value
Aging	0.701***	0.250	3.35
Any ailment	0.005***	0.001	0.14
Interstate migration	0.000***	0.000	4.80
International migration	0.000**	0.000	2.34
Literate	-0.306***	0.047	-6.53
Scheduled Caste/Scheduled Tribe population	-0.019	0.012	-1.58
Muslim Population	-0.027	0.016	-1.68
Casual labor in agriculture	-0.103	0.104	-0.98
Casual labor in non-agriculture	0.108*	0.059	1.85
Poverty	0.000	0.030	0.02
Joint/extended family	0.090**	0.041	2.20
Average no. of person living per room	-1.069	0.762	-1.40
Household use for multi-purpose	0.174***	0.063	2.77
Handwash before meal	-0.041*	0.022	-1.84
Handwash before defecation	-0.006	0.017	-0.36
Drinking water outside premises	-0.047***	0.010	-4.53
Population density	0.000**	0.000	1.90
Proportion of urban population	0.034***	0.013	2.71
Proportion of slum population	0.346	0.107	3.23
Health infrastructure index	0.622***	0.186	3.35
Health expenditure	0.000	0.000	0.14
Mass media exposure	-1.236***	0.505	-2.45
lnalpha	-0.755	0.275	
alpha	.469	.129	
N	36		
LR chi2(21)	90.22***		
Log-likelihood	-201.5395		
Likelihood-ratio test of alpha=0: chibar2(01) = 1359.91 Prob>=chibar2 = 0.000			

***p<0.001, **p<0.05 & *p<0.10

Appendix 1: Classification of states and union territories according to the COVID-19 confirmed positive cases in India

Low Burden COVID-19	Medium Burden COVID-19	High Burden COVID-19
West Bengal (WB)	Madhya Pradesh (MP)	Maharashtra (MH)
Jammu & Kashmir (JK)	Rajasthan (RJ)	Gujarat (GJ)
Karnataka (KR)	Tamil Nadu (TN)	Delhi (DL)
Kerala (KL)	Uttar Pradesh (UP)	
Punjab (PJ)	Andhra Pradesh (AP)	
Bihar (BH)	Telangana (TG)	
Haryana (HR)		
Odisha (OR)		
Jharkhand (JH)		
Chandigarh (CD)		
Uttarakhand (UT)		
Assam (AS)		
Chhattisgarh (CT)		
Himachal Pradesh (HP)		
Andaman and Nicobar Islands (ANI)		
Puducherry (PY)		
Goa (GA)		
Manipur (MN)		
Tripura (TR)		
Arunachal Pradesh (AR)		
Mizoram (MZ)		

Figure 1: Prevalence of COVID-19 positive cases and recovered cases of COVID-19 across states and union territories in India




Figure 2: Average weekly confirmed cases and positivity rate in low, medium and high COVID-19 burden states from March 16, 2020 to May 03, 2020, India




Figure 3: Average weekly new recovered cases and recovery rate in low, medium and high COVID-19 burden states from March 16, 2020 to May 03, 2020, India




Figure 4: Average weekly deaths and fatality rate in low, medium and high COVID-19 burden states from March 16, 2020 to May 03, 2020, India




Figure 5: The estimated indices of susceptibility, exposure, resilience and composite vulnerability index related to COVID-19 risk across states and union territories in India

