
Cosmology and linear-bias correlations: $0.4 \le z < 0.6$


Model redshift $z = 0.5$ | $\rho_{ij} = C_{ij}/\sqrt{C_{ii}C_{jj}}$ | Baseline $k_{\rm max} = 0.20\,h_{\rm ref}\,{\rm Mpc}^{-1}$

ALL EIGHT BINS: select this bin's biases from the full marginalized covariance


A: correlated DESI autos




B: DESI autos + crosses




C: B + PFS autos





D: C + PFS crosses





THIS BIN ONLY: marginalize its cosmology and nuisance parameters; insert each prior once











$q_A = \ln(10^{10} A_s)$. Subsample labels (LRG-1, ELG-1, QSO-1, PFS-ELG-1, ...; Sec. 3.1) denote $b_1\sigma_8$ of that subsample at the model redshift. All other fitted parameters are marginalized.

h, ω_b, n_s fixed. Cosmology priors: $\sigma(M_\nu) = 5\,{\rm eV}$, $\sigma(\Omega_m) = 1$; flat q_A and linear-bias priors. Each bin-only matrix is a separate forecast.


PFS baseline: 0.6–1.6 (Takada et al., Table 2). Same finite nuisance priors in all cases, in the amplitude-scaled nuisance coordinates of Sec. 3.3. Values shown to two decimals; full precision in the CSV/JSON companion.

Cosmology and linear-bias correlations: $0.6 \le z < 0.8$


Model redshift $z = 0.7$ | $\rho_{ij} = C_{ij}/\sqrt{C_{ii}C_{jj}}$ | Baseline $k_{\rm max} = 0.20\,h_{\rm ref}\,{\rm Mpc}^{-1}$

ALL EIGHT BINS: select this bin's biases from the full marginalized covariance


A: correlated DESI autos




B: DESI autos + crosses




C: B + PFS autos





D: C + PFS crosses







THIS BIN ONLY: marginalize its cosmology and nuisance parameters; insert each prior once









$q_A = \ln(10^{10} A_s)$. Subsample labels (LRG-1, ELG-1, QSO-1, PFS-ELG-1, ...; Sec. 3.1) denote $b_1\sigma_8$ of that subsample at the model redshift. All other fitted parameters are marginalized.

h, ω_b, n_s fixed. Cosmology priors: $\sigma(M_\nu) = 5\,{\rm eV}$, $\sigma(\Omega_m) = 1$; flat q_A and linear-bias priors. Each bin-only matrix is a separate forecast.

PFS baseline: 0.6–1.6 (Takada et al., Table 2). Same finite nuisance priors in all cases, in the amplitude-scaled nuisance coordinates of Sec. 3.3. Values shown to two decimals; full precision in the CSV/JSON companion.

Cosmology and linear-bias correlations: $0.8 \le z < 1.0$

Model redshift $z = 0.9$ | $\rho_{ij} = C_{ij}/\sqrt{C_{ii}C_{jj}}$ | Baseline $k_{\rm max} = 0.20\,h_{\rm ref}\,{\rm Mpc}^{-1}$

ALL EIGHT BINS: select this bin's biases from the full marginalized covariance

A: correlated DESI autos

	q_A	M_ν	Ω_m	ELG-1	LRG-3	QSO-1
q_A	1.00	0.77	0.49	-0.53	-0.41	-0.21
M_ν	0.77	1.00	0.86	-0.27	-0.22	-0.10
Ω_m	0.49	0.86	1.00	-0.01	0.06	0.03
ELG-1	-0.53	-0.27	-0.01	1.00	0.68	0.29
LRG-3	-0.41	-0.22	0.06	0.68	1.00	0.28
QSO-1	-0.21	-0.10	0.03	0.29	0.28	1.00

B: DESI autos + crosses

	q_A	M_ν	Ω_m	ELG-1	LRG-3	QSO-1
q_A	1.00	0.78	0.49	-0.52	-0.44	-0.32
M_ν	0.78	1.00	0.85	-0.28	-0.24	-0.18
Ω_m	0.49	0.85	1.00	0.01	0.05	0.00
ELG-1	-0.52	-0.28	0.01	1.00	0.84	0.51
LRG-3	-0.44	-0.24	0.05	0.84	1.00	0.52
QSO-1	-0.32	-0.18	0.00	0.51	0.52	1.00

C: B + PFS autos

	q_A	M_ν	Ω_m	ELG-1	LRG-3	QSO-1	PFS-ELG-2
q_A	1.00	0.78	0.49	-0.52	-0.44	-0.32	-0.25
M_ν	0.78	1.00	0.85	-0.28	-0.24	-0.18	-0.14
Ω_m	0.49	0.85	1.00	0.01	0.05	0.00	-0.00
ELG-1	-0.52	-0.28	0.01	1.00	0.84	0.51	0.34
LRG-3	-0.44	-0.24	0.05	0.84	1.00	0.51	0.32
QSO-1	-0.32	-0.18	0.00	0.51	0.51	1.00	0.20
PFS-ELG-2	-0.25	-0.14	-0.00	0.34	0.32	0.20	1.00

D: C + PFS crosses

	q_A	M_ν	Ω_m	ELG-1	LRG-3	QSO-1	PFS-ELG-2
q_A	1.00	0.78	0.50	-0.52	-0.44	-0.31	-0.32
M_ν	0.78	1.00	0.85	-0.29	-0.25	-0.18	-0.19
Ω_m	0.50	0.85	1.00	-0.01	0.04	-0.00	-0.02
ELG-1	-0.52	-0.29	-0.01	1.00	0.84	0.51	0.46
LRG-3	-0.44	-0.25	0.04	0.84	1.00	0.51	0.47
QSO-1	-0.31	-0.18	-0.00	0.51	0.51	1.00	0.33
PFS-ELG-2	-0.32	-0.19	-0.02	0.46	0.47	0.33	1.00

THIS BIN ONLY: marginalize its cosmology and nuisance parameters; insert each prior once

	q_A	M_ν	Ω_m	ELG-1	LRG-3	QSO-1
q_A	1.00	0.77	0.46	-0.75	-0.65	-0.42
M_ν	0.77	1.00	0.86	-0.37	-0.33	-0.19
Ω_m	0.46	0.86	1.00	0.02	0.09	0.06
ELG-1	-0.75	-0.37	0.02	1.00	0.84	0.52
LRG-3	-0.65	-0.33	0.09	0.84	1.00	0.51
QSO-1	-0.42	-0.19	0.06	0.52	0.51	1.00

	q_A	M_ν	Ω_m	ELG-1	LRG-3	QSO-1
q_A	1.00	0.79	0.48	-0.73	-0.66	-0.56
M_ν	0.79	1.00	0.86	-0.38	-0.35	-0.31
Ω_m	0.48	0.86	1.00	0.03	0.08	0.02
ELG-1	-0.73	-0.38	0.03	1.00	0.93	0.72
LRG-3	-0.66	-0.35	0.08	0.93	1.00	0.72
QSO-1	-0.56	-0.31	0.02	0.72	0.72	1.00

	q_A	M_ν	Ω_m	ELG-1	LRG-3	QSO-1	PFS-ELG-2
q_A	1.00	0.79	0.48	-0.73	-0.67	-0.56	-0.47
M_ν	0.79	1.00	0.86	-0.38	-0.36	-0.31	-0.25
Ω_m	0.48	0.86	1.00	0.03	0.07	0.01	0.01
ELG-1	-0.73	-0.38	0.03	1.00	0.92	0.71	0.56
LRG-3	-0.67	-0.36	0.07	0.92	1.00	0.71	0.54
QSO-1	-0.56	-0.31	0.01	0.71	0.71	1.00	0.43
PFS-ELG-2	-0.47	-0.25	0.01	0.56	0.54	0.43	1.00

	q_A	M_ν	Ω_m	ELG-1	LRG-3	QSO-1	PFS-ELG-2
q_A	1.00	0.80	0.49	-0.74	-0.67	-0.56	-0.57
M_ν	0.80	1.00	0.86	-0.41	-0.38	-0.32	-0.33
Ω_m	0.49	0.86	1.00	0.00	0.04	-0.00	-0.03
ELG-1	-0.74	-0.41	0.00	1.00	0.93	0.72	0.68
LRG-3	-0.67	-0.38	0.04	0.93	1.00	0.71	0.68
QSO-1	-0.56	-0.32	-0.00	0.72	0.71	1.00	0.55
PFS-ELG-2	-0.57	-0.33	-0.03	0.68	0.68	0.55	1.00

ρ 1.0 0.5 0.0 −0.5 −1.0

$q_A = \ln(10^{10} A_s)$. Subsample labels (LRG-1, ELG-1, QSO-1, PFS-ELG-1, ...; Sec. 3.1) denote $b_1\sigma_8$ of that subsample at the model redshift. All other fitted parameters are marginalized.

h, ω_b, n_s fixed. Cosmology priors: $\sigma(M_\nu) = 5\,{\rm eV}$, $\sigma(\Omega_m) = 1$; flat q_A and linear-bias priors. Each bin-only matrix is a separate forecast.


PFS baseline: 0.6–1.6 (Takada et al., Table 2). Same finite nuisance priors in all cases, in the amplitude-scaled nuisance coordinates of Sec. 3.3. Values shown to two decimals; full precision in the CSV/JSON companion.

Cosmology and linear-bias correlations: $1.0 \leq z < 1.2$


Model redshift $z = 1.1$ | $\rho_{ij} = C_{ij}/\sqrt{C_{ii}C_{jj}}$ | Baseline $k_{\rm max} = 0.20\,h_{\rm ref}\,{\rm Mpc}^{-1}$

ALL EIGHT BINS: select this bin's biases from the full marginalized covariance


A: correlated DESI autos




B: DESI autos + crosses




C: B + PFS autos





D: C + PFS crosses





THIS BIN ONLY: marginalize its cosmology and nuisance parameters; insert each prior once











$q_A = \ln(10^{10} A_s)$. Subsample labels (LRG-1, ELG-1, QSO-1, PFS-ELG-1, ...; Sec. 3.1) denote $b_1\sigma_8$ of that subsample at the model redshift. All other fitted parameters are marginalized.

h, ω_b, n_s fixed. Cosmology priors: $\sigma(M_\nu) = 5\,{\rm eV}$, $\sigma(\Omega_m) = 1$; flat q_A and linear-bias priors. Each bin-only matrix is a separate forecast.


PFS baseline: 0.6–1.6 (Takada et al., Table 2). Same finite nuisance priors in all cases, in the amplitude-scaled nuisance coordinates of Sec. 3.3. Values shown to two decimals; full precision in the CSV/JSON companion.

Cosmology and linear-bias correlations: $1.2 \le z < 1.4$


Model redshift $z = 1.3$ | $\rho_{ij} = C_{ij}/\sqrt{C_{ii}C_{jj}}$ | Baseline $k_{\rm max} = 0.20\,h_{\rm ref}\,{\rm Mpc}^{-1}$

ALL EIGHT BINS: select this bin's biases from the full marginalized covariance


A: correlated DESI autos




B: DESI autos + crosses




C: B + PFS autos





D: C + PFS crosses





THIS BIN ONLY: marginalize its cosmology and nuisance parameters; insert each prior once











$q_A = \ln(10^{10} A_s)$. Subsample labels (LRG-1, ELG-1, QSO-1, PFS-ELG-1, ...; Sec. 3.1) denote $b_1\sigma_8$ of that subsample at the model redshift. All other fitted parameters are marginalized.

h, ω_b, n_s fixed. Cosmology priors: $\sigma(M_\nu) = 5\,{\rm eV}$, $\sigma(\Omega_m) = 1$; flat q_A and linear-bias priors. Each bin-only matrix is a separate forecast.


PFS baseline: 0.6–1.6 (Takada et al., Table 2). Same finite nuisance priors in all cases, in the amplitude-scaled nuisance coordinates of Sec. 3.3. Values shown to two decimals; full precision in the CSV/JSON companion.

Cosmology and linear-bias correlations: $1.4 \le z < 1.6$


Model redshift $z = 1.5$ | $\rho_{ij} = C_{ij}/\sqrt{C_{ii}C_{jj}}$ | Baseline $k_{\rm max} = 0.20\,h_{\rm ref}\,{\rm Mpc}^{-1}$

ALL EIGHT BINS: select this bin's biases from the full marginalized covariance


A: correlated DESI autos




B: DESI autos + crosses




C: B + PFS autos





D: C + PFS crosses





THIS BIN ONLY: marginalize its cosmology and nuisance parameters; insert each prior once











$q_A = \ln(10^{10} A_s)$. Subsample labels (LRG-1, ELG-1, QSO-1, PFS-ELG-1, ...; Sec. 3.1) denote $b_1\sigma_8$ of that subsample at the model redshift. All other fitted parameters are marginalized.

h, ω_b, n_s fixed. Cosmology priors: $\sigma(M_\nu) = 5\,{\rm eV}$, $\sigma(\Omega_m) = 1$; flat q_A and linear-bias priors. Each bin-only matrix is a separate forecast.


PFS baseline: 0.6–1.6 (Takada et al., Table 2). Same finite nuisance priors in all cases, in the amplitude-scaled nuisance coordinates of Sec. 3.3. Values shown to two decimals; full precision in the CSV/JSON companion.

Cosmology and linear-bias correlations: $1.6 \leq z < 2.0$


Model redshift $z = 1.8$ | $\rho_{ij} = C_{ij}/\sqrt{C_{ii}C_{jj}}$ | Baseline $k_{\max} = 0.20\,h_{\rm ref}\,{\rm Mpc}^{-1}$

ALL EIGHT BINS: select this bin's biases from the full marginalized covariance


A: correlated DESI autos




B: DESI autos + crosses




C: B + PFS autos





D: C + PFS crosses





THIS BIN ONLY: marginalize its cosmology and nuisance parameters; insert each prior once











$q_A = \ln(10^{10} A_s)$. Subsample labels (LRG-1, ELG-1, QSO-1, PFS-ELG-1, ...; Sec. 3.1) denote $b_1\sigma_8$ of that subsample at the model redshift. All other fitted parameters are marginalized.

h, ω_b, n_s fixed. Cosmology priors: $\sigma(M_\nu) = 5\,{\rm eV}$, $\sigma(\Omega_m) = 1$; flat q_A and linear-bias priors. Each bin-only matrix is a separate forecast.

PFS baseline: 0.6–1.6 (Takada et al., Table 2). Same finite nuisance priors in all cases, in the amplitude-scaled nuisance coordinates of Sec. 3.3. Values shown to two decimals; full precision in the CSV/JSON companion.

Cosmology and linear-bias correlations: $2.0 \leq z < 2.1$

Model redshift $z = 2.05$ | $\rho_{ij} = C_{ij}/\sqrt{C_{ii}C_{jj}}$ | Baseline $k_{\rm max} = 0.20\,h_{\rm ref}\,{\rm Mpc}^{-1}$

ALL EIGHT BINS: select this bin's biases from the full marginalized covariance

A: correlated DESI autos

	q_A	M_ν	Ω_m	QSO-6
q_A	1.00	0.77	0.49	-0.22
M_ν	0.77	1.00	0.86	-0.12
Ω_m	0.49	0.86	1.00	-0.01
QSO-6	-0.22	-0.12	-0.01	1.00

B: DESI autos + crosses


	q_A	M_ν	Ω_m	QSO-6
q_A	1.00	0.78	0.49	-0.21
M_ν	0.78	1.00	0.85	-0.12
Ω_m	0.49	0.85	1.00	-0.01
QSO-6	-0.21	-0.12	-0.01	1.00

C: B + PFS autos

	q_A	M_ν	Ω_m	QSO-6
q_A	1.00	0.78	0.49	-0.21
M_ν	0.78	1.00	0.85	-0.12
Ω_m	0.49	0.85	1.00	-0.01
QSO-6	-0.21	-0.12	-0.01	1.00

D: C + PFS crosses

	q_A	M_ν	Ω_m	QSO-6
q_A	1.00	0.78	0.50	-0.20
M_ν	0.78	1.00	0.85	-0.12
Ω_m	0.50	0.85	1.00	-0.01
QSO-6	-0.20	-0.12	-0.01	1.00



THIS BIN ONLY: marginalize its cosmology and nuisance parameters; insert each prior once

	q_A	M_ν	Ω_m	QSO-6
q_A	1.00	0.72	0.55	-0.75
M_ν	0.72	1.00	0.86	-0.44
Ω_m	0.55	0.86	1.00	-0.12
QSO-6	-0.75	-0.44	-0.12	1.00

	q_A	M_ν	Ω_m	QSO-6
q_A	1.00	0.72	0.55	-0.75
M_ν	0.72	1.00	0.86	-0.44
Ω_m	0.55	0.86	1.00	-0.12
QSO-6	-0.75	-0.44	-0.12	1.00

	q_A	M_ν	Ω_m	QSO-6
q_A	1.00	0.72	0.55	-0.75
M_ν	0.72	1.00	0.86	-0.44
Ω_m	0.55	0.86	1.00	-0.12
QSO-6	-0.75	-0.44	-0.12	1.00

	q_A	M_ν	Ω_m	QSO-6
q_A	1.00	0.72	0.55	-0.75
M_ν	0.72	1.00	0.86	-0.44
Ω_m	0.55	0.86	1.00	-0.12
QSO-6	-0.75	-0.44	-0.12	1.00

$q_A = \ln(10^{10} A_s)$. Subsample labels (LRG-1, ELG-1, QSO-1, PFS-ELG-1, ...; Sec. 3.1) denote $b_1\sigma_8$ of that subsample at the model redshift. All other fitted parameters are marginalized.

h, ω_b, n_s fixed. Cosmology priors: $\sigma(M_\nu) = 5\,{\rm eV}$, $\sigma(\Omega_m) = 1$; flat q_A and linear-bias priors. Each bin-only matrix is a separate forecast.

PFS baseline: 0.6–1.6 (Takada et al., Table 2). Same finite nuisance priors in all cases, in the amplitude-scaled nuisance coordinates of Sec. 3.3. Values shown to two decimals; full precision in the CSV/JSON companion.