


Appendix S9. Summary of divergence time estimates in millions of years for shared crown clades (*Adunator* root calibration, narrow priors as in Fig. 4), showing comparisons (where available) with clock estimates for comparable clades from recent studies that sampled across Soricidae. Point estimates are median values in this study, otherwise reported as given in the literature: Brace et al. (2016: fig. 2), Chen et al. (2020: fig. 3, excluding nodes they used as calibrations), Dubey et al. (2007: fig. 2); He et al. (2010: fig. 3 and table 3). We do not use estimates from Sato et al. (2016) due to the issues noted by Springer et al. (2018). "Clade letter" column refers to lettered nodes in topology shown below.

Study	clade	clade letter	point estimate	95% credibility min	95% credibility max
this study	Soricoidea	A	60.92	50.71	75.09
Brace 2016	Soricidae	B	48.3	39.9	56.1
this study	Soricidae	B	51.06	41.32	64.34
Brace 2016	Soricinae	C	37.7	29.6	45.6
He 2010	Soricinae	C	21.12	18.16	25.96
this study	Soricinae	C	37.3	28.32	48.28
He 2010	Soricinei	D	19.73	16.15	24.73
this study	Soricinei	D	34.60	26.34	44.94
Brace 2016	Neomyini	E	29.9	22.5	37.7
this study	Neomyini	E	31.07	23.67	40.84
Dubey 2007	Neomyina	F	13.8	-	-
this study	Neomyina	F	28.04	20.88	37.19
Brace 2016	Nectogalita	G	16.6	11.0	22.8
He 2010	Nectogalita	G	6.63	4.81	8.84
this study	Nectogalita	G	20.52	14.87	27.7
this study	<i>Soriculus-Episoriculus, Nectogale Chimarroale</i>	H	18.81	13.48	25.86
He 2010	<i>Chimarroale-Nectogale</i>	I	3.71	2.10	5.46
this study	<i>Chimarroale-Nectogale</i>	I	13.37	8.31	19.65
this study	<i>Episoriculus-Soriculus</i>	J	16.48	10.97	23.02
He 2010	<i>Anourosorex</i>	K	0.66	0.18	1.35
this study	<i>Anourosorex</i>	K	2.75	0.69	5.55
Chen 2020	Blarininei	L	13.67	-	-
Dubey 2007	Blarininei	L	12.1	-	-
this study	Blarininei	L	26.82	18.56	36.29
Chen 2020	Blarinini	M	10.72	-	-
He 2010	Blarinini	M	15.47	15.04	16.49
this study	Blarinini	M	16.16	9.93	23.48
Chen 2020	<i>Sorex</i>	N	7.97	-	-
Dubey 2007	<i>Sorex</i>	N	13.9	-	-
He 2010	<i>Sorex</i>	N	10.03	6.7	14.21
this study	<i>Sorex</i>	N	21.86	15.52	30.02
this study	<i>S.alpinus-minutus-caecutiens-coronatus-araneus</i>	O	14.72	9.74	20.58
this study	<i>S.minutus-caecutiens-coronatus-araneus</i>	P	11.04	7.08	16.02
this study	<i>S.coronatus-araneus</i>	Q	3.37	1.04	6.18
this study	<i>S.cinereus-monticolus</i>	R	8.64	4.15	13.84

Study	clade	clade letter	point estimate	95% credibility min	95% credibility max
Dubey 2007	Crocidae	S	16.5	-	-
this study	Crocidae	S	39.97	30.9	51.53
Dubey 2007	Crocidae	T	10.8	-	-
this study	Crocidae	T	23.69	17.3	31.8
Dubey 2007	<i>Crocidae-Suncus etruscus</i>	U	9.3	-	-
this study	Crocidae	U	18.93	13.69	25.57
Brace 2016	<i>Crocidae-Suncus murinus</i>	V	17.4	10.8	26.1
this study	<i>Crocidae-Suncus murinus</i>	V	16.8	11.69	22.68
Dubey 2007	<i>Crocidae</i>	W	8	-	-
this study	<i>Crocidae</i>	W	13.98	9.51	19.44
this study	<i>"Diplomesodon"-C.suaveolens-lasiura</i>	X	12.24	7.86	17.34
this study	<i>C.lasiura-suaveolens</i>	Y	10.11	5.95	14.97
this study	<i>C.olivieri-russula</i>	Z	10.62	6.34	15.61
this study	Myosoricini	AA	25.06	16.27	35.06

Below is a guide to divergence estimates for lettered crown nodes in the above table. Topology is a closeup of Soricoidea from the timetree shown in Fig. 4 (*Adunator* root calibration, narrow priors, as explained in Methods). Red indicates Soricinae, Green Crocidurinae, pink Heterosoricidae. Numbers at top and bottom indicate millions of years before present.



- Brace S, Thomas JA, Dalén L, Burger J, MacPhee RDE, Barnes I, Turvey ST. 2016 Evolutionary history of the Nesophontidae, the last unplaced Recent mammal family. *Mol. Biol. Evol.* **33**, 3095–3103.
- Chen S *et al.* 2020 Multilocus phylogeny and cryptic diversity of white-toothed shrews (Mammalia, Eulipotyphla, Crocidura) in China. *BMC Evol. Biol.* **20**, 1–14.
- Dubey S, Salamin N, Ohdachi SD, Barrière P, Vogel P. 2007 Molecular phylogenetics of shrews (Mammalia: Soricidae) reveal timing of transcontinental colonizations. *Mol. Phylogenet. Evol.* **44**, 126–137. (doi:10.1016/j.ympev.2006.12.002)
- He K, Li Y-J, Brandley MC, Lin L-K, Wang Y-X, Zhang Y-P, Jiang X-L. 2010 A multi-locus phylogeny of Nectogalini shrews and influences of the paleoclimate on speciation and evolution. *Mol. Phylogenet. Evol.* **56**, 734–746.
- Sato JJ *et al.* 2016 Molecular phylogenetic analysis of nuclear genes suggests a Cenozoic over-water dispersal origin for the Cuban solenodon. *Sci. Rep.* **6**, 31173.
- Springer MS, Murphy WJ, Roca AL. 2018 Appropriate fossil calibrations and tree constraints uphold the Mesozoic divergence of solenodons from other extant mammals. *Mol. Phylogenet. Evol.* **121**, 158–165.