Supplementary table 1 Normality testing of numerical variables (method: Shapiro-Wilk).

Virus Infection Status	Positive		Negative	
Variables	W	P-value	W	P-value
CD3 absolute count (/U)	0.6116	<0.0001	0.8277	<0.0001
CD4 absolute count (/U)	0.4658	<0.0001	0.8521	0.0001
CD8 absolute count (/U)	0.8326	<0.0001	0.7518	<0.0001
CRP (mg/L)	0.8641	<0.0001	0.7233	<0.0001
D-dimer (mg/L)	0.9533	0.0058	0.8758	0.0005
WBC count (10^9/L)	0.8585	<0.0001	0.6215	<0.0001
Lymphocyte count (10^9/L)	0.6208	<0.0001	0.8793	0.0006
Neutrophil count (10^9/L)	0.8409	<0.0001	0.6017	<0.0001
NLR	0.4241	<0.0001	0.4331	<0.0001
RBC count (10^12/L)	0.9796	0.2380	0.9467	0.0636
HB (g/L)	0.9788	0.2127	0.9467	0.0636
Platelet count (10^9/L)	0.8988	<0.0001	0.9842	0.8492
eGFR (ml/min/1.73m2)	0.9533	0.0057	0.9581	0.1536
Creatinine (umol/L)	0.5954	<0.0001	0.7948	<0.0001
BUN (mmol/L)	0.7794	<0.0001	0.5987	<0.0001
Uric acid (umol/L)	0.9471	0.0025	0.9528	0.1018
TBIL (umol/L)	0.2955	<0.0001	0.5103	<0.0001
TP (g/L)	0.9874	0.6350	0.9577	0.1489
Prealbumin (mg/L)	0.9776	0.1788	0.9644	0.2501
ALT (IU/L)	0.9428	0.0015	0.9585	0.1588
AST (IU/L)	0.7648	<0.0001	0.9562	0.1325
PCT (ng/MI)	0.2767	<0.0001	0.1834	<0.0001
Patient age (year)	0.8916	<0.0001	0.9314	0.0199

* Normal distribution: P>0.05

Supplementary table 1 Normal distribution assessment was performed using the Shapiro-Wilk test on 23 numerical variables across two groups: triple co-infection and double co-infection. W: test statistic. A p-value >= 0.05 was used as the threshold to indicate normality. The RBC count, HB, TP, and prealbumin in both groups all follow a normal distribution. The other variables in both groups deviated significantly from normality ($p < 0.05$). NLR: Neutrophil-to-lymphocyte ratio. HB: Hemoglobine. GFR: Estimated glomerular filtration rate. TBIL: Total bilirubin. TP: