

Michael Goforth
CAAM 550
HW 8
10/27/2021

Problem 1

part a

See Jupyter notebook for code and results.

part b

See Jupyter notebook for code and results.

part c

See Jupyter notebook for code and results.

part d

See Jupyter notebook for code and results.

part e

See Jupyter notebook for code and results.

Problem 2

part a

See Jupyter notebook for code and results.

part b

See Jupyter notebook for code and results. The root approximation is about 1.875.

Problem 3

part a

$$\sum_{k=1}^n \sum_{l=1}^m a_{kl} p_k(x_i) q_l(y_j) = f(x_1, y_j), \quad i = 1, \dots, n, \quad j = 1, \dots, m$$

$$p_i(x) = L_i(x) = \prod_{\substack{j=1 \\ j \neq i}}^n \frac{x - x_j}{x_i - x_j}, \quad i = 1, \dots, n$$

$$q_i(y) = L_i(y) = \prod_{\substack{j=1 \\ j \neq i}}^m \frac{y - y_j}{y_i - y_j}, \quad i = 1, \dots, m$$

For any $(x_i, y_j) \in [a, b] \times [c, d]$, $i = 1, \dots, n$, $j = 1, \dots, m$,

$$p_k(x_i) = \begin{cases} 1, & \text{if } k = i \\ 0, & \text{if } k \neq i \end{cases}$$

$$q_l(y_j) = \begin{cases} 1, & \text{if } l = j \\ 0, & \text{if } l \neq j \end{cases}$$

$$f(x_1, y_j) = \sum_{k=1}^n \sum_{l=1}^m a_{kl} p_k(x_i) q_l(y_j) = a_{kl}$$

See Jupyter notebook for code and further results.

part b

$$\sum_{k=1}^n \sum_{l=1}^m a_{kl} p_k(x_i) q_l(y_j) = f(x_1, y_j), \quad i = 1, \dots, n, \quad j = 1, \dots, m$$

$$p_i(x) = N_i(x) = \prod_{j=1}^{i-1} (x - x_j), \quad i = 1, \dots, n$$

$$q_i(y) = N_i(y) = \prod_{j=1}^{i-1} (y - y_j), \quad i = 1, \dots, m$$

$$N_1(x) = N_1(y) = 1$$

$$p_i(x_k) = \prod_{j=1}^{i-1} (x_k - x_j) = 0 \text{ if } k > i$$

$$q_i(y_l) = \prod_{j=1}^{i-1} (y_l - y_j) = 0 \text{ if } l > i$$

$$a_{kl} = \frac{f(x_k, y_l) - \sum_{i=1}^{k-1} \sum_{j=1}^{l-1} a_{ij} p_i(x_k) q_j(y_l)}{p_k q_l}$$

By starting with a matrix $A \in \mathbb{R}^{n \times m}$ of all zeroes, the equations above can be used to find elements of A one by one starting at the top left and working down the matrix one row at a time.

Problem 4

part a

$\prod_n f = p_n$ where p_n is a polynomial of degree n . Then since the polynomial of degree $n-1$ going through n different points is unique, $\prod_n p_n = p_n$. Therefore $(\prod_n)^2 f = p_n$ and therefore $\prod_n = (\prod_n)^2$ and \prod_n is a projector.

part b

$\prod_0 f = p_0 = f(x_1)$ for any function f and given point x_1 . Therefore the $\|\prod_0 f\|_{L^\infty([a,b])} = f(x_1)$. Then

$$\max_{0 \neq f \in \mathbb{C}([a,b])} \frac{\|\prod_0 f\|_{L^\infty([a,b])}}{\|f\|_{L^\infty([a,b])}} = 1$$

since $\|f\|_{L^\infty([a,b])} \geq |f(x_1)|$.

Similarly, because a polynomial of degree 1 is a line, if $x_0 = a$ and $x_1 = b$, then $\|\prod_1 f\|_{L^\infty([a,b])} = \max|f(a)|, |f(b)|$. Again since $\|f\|_{L^\infty([a,b])} \geq \max|f(a)|, |f(b)|$,

$$\|\prod_1\|_{L^\infty([a,b])} = \max_{0 \neq f \in \mathbb{C}([a,b])} \frac{\|\prod_1 f\|_{L^\infty([a,b])}}{\|f\|_{L^\infty([a,b])}} = 1$$

part c

$$\|Pi_n\|_{L^\infty([a,b])} = \max_{\|f\|_{L^\infty([a,b])}=1} \|Pi_n f\|_{L^\infty([a,b])}$$

$$\|Pi_n\|_{L^\infty([a,b])} = \max_{\|f\|_{L^\infty([a,b])}=1} \left\| \sum_{j=0}^n f(x_j) L_j(x) \right\|_{L^\infty([a,b])}$$

$$\|Pi_n\|_{L^\infty([a,b])} = \max_{\|f\|_{L^\infty([a,b])}=1} \left[\max_{x \in [a,b]} \left| \sum_{j=0}^n f(x_j) L_j(x) \right| \right]$$

Because $\|f\|_{L^\infty([a,b])} = 1$, a function can be constructed with $f(x_j) = \pm 1$ and $\max_{x \in [a,b]} |f(x)| = 1$ (chosen so that the sign of all the terms is the same) which maximizes the sum in the brackets. Therefore

$$\|Pi_n\|_{L^\infty([a,b])} = \max_{x \in [a,b]} \sum_{j=0}^n |L_j(x)|$$

part d

Using the triangle inequality

$$\|f - p_n\|_{L^\infty([a,b])} \leq \|f - p_*\|_{L^\infty([a,b])} + \|p_* - p_n\|_{L^\infty([a,b])}$$

Because Π_n is a projector (shown in part a),

$$\begin{aligned} \|f - p_n\|_{L^\infty([a,b])} &\leq \|f - p_*\|_{L^\infty([a,b])} + \|\Pi_n(p_* - f)\|_{L^\infty([a,b])} \\ \|f - p_n\|_{L^\infty([a,b])} &\leq \|f - p_*\|_{L^\infty([a,b])} + \|\Pi_n\|_{L^\infty([a,b])} \|p_* - f\|_{L^\infty([a,b])} \\ \|f - p_n\|_{L^\infty([a,b])} &\leq (1 + \|\Pi_n\|_{L^\infty([a,b])}) \|f - p_*\|_{L^\infty([a,b])} \end{aligned}$$

part e

Consider the polynomial p_* that minimizes $\|f - p_n\|_{L^\infty([a,b])}$, then from part d above

$$\|f - p_n\|_{L^\infty([a,b])} \leq (1 + \|\Pi_n\|_{L^\infty([a,b])}) \|f - p_*\|_{L^\infty([a,b])}$$

The Lebesgue Constant is acting as a measure of the error of the calculated interpolant and the best possible interpolant, the smaller the constant, the closer the calculated interpolation is to the best interpolation.

Problem 5**part a**

See Jupyter notebook for code and results.

part b

See Jupyter notebook for code and results.

part c

See Jupyter notebook for code and results.

part d

See Jupyter notebook for code and results. The error bound for linear spline interpolation holds for the bounded twice differentiable functions ($\sin(x)$, $\sin(30x)$, and e^x) as written. However the second derivative of $x^{4/3}$, and $\sqrt[3]{x} - 1/2$ are unbounded so the error bound does not apply to these functions. Finally the second derivative of $|x - 0.567|$ is undefined at $x = 0.567$ so the error for this function also does not follow the error bound.

The error bound of $\sin(3x)$ and $\sin(30x)$ both converge $O(h^2)$, but $\sin(30x)$ converges slower due to the larger maximum value of its second derivative on the interval $[0,1]$.

part e

See Jupyter notebook for code and results. Order of error for the functions:

$$f(x) = \sin(3x) \Rightarrow O(4)$$

$$f(x) = \sin(30x) \Rightarrow O(4)$$

$$f(x) = e^x \Rightarrow O(4)$$

$$f(x) = x^{4/3} \Rightarrow O(2.66)$$

$$f(x) = |x - 0.567| \Rightarrow O(2)$$

$$f(x) = \sqrt[3]{x - 1/2} \Rightarrow O(2/3)$$

I determined these values by making plots of L_inf error vs n and h^p vs n for each function and incrementing n up and down until the plot approached linear.