



21:10:30 05-04-22 Example result file.

Experiment Information

Parameter	F1	F2
Nucleus	N/A	¹H
Transmitter Frequency (MHz)	N/A	400
Sweep Width (Hz)	50	5000
Sweep Width (ppm)	N/A	12.5
Transmitter Offset (Hz)	0	2000
Transmitter Offset (ppm)	N/A	5

Estimation Result

Osc.	а	φ (°)	f_1 (Hz)	f ₂ (Hz)	<i>f</i> ₂ (ppm)	$\eta_1 \; (s^{-1})$	$\eta_2~(\mathrm{s}^{-1})$	ſ
1	$0.9906 \\ \pm 9.906 \times 10^{-3}$	-0.51936 $\pm -5.1936 \times 10^{-3}$	20.002 ±0.20002	3.02×10^{3} ± 30.2	7.55 $\pm 7.55 \times 10^{-2}$	$4.9951 \\ \pm 4.9951 \times 10^{-2}$	$4.9953 \\ \pm 4.9953 \times 10^{-2}$	1
2	3.0016 $\pm 3.0016 \times 10^{-2}$	-0.45698 $\pm -4.5698 \times 10^{-3}$	9.9953 $\pm 9.9953 \times 10^{-2}$	3.01×10^{3} ± 30.1	7.525 $\pm 7.525 \times 10^{-2}$	$4.9953 \\ \pm 4.9953 \times 10^{-2}$	4.995 $\pm 4.995 \times 10^{-2}$	3.0604
3	$5.9962 \\ \pm 5.9962 \times 10^{-2}$	0.26285 $\pm 2.6285 \times 10^{-3}$	-4.2529×10^{-4} $\pm -4.2529 \times 10^{-6}$	$3 \times 10^3 \\ \pm 30$	$7.5 \pm 7.5 \times 10^{-2}$	$4.9926 \\ \pm 4.9926 \times 10^{-2}$	$5.0016 \\ \pm 5.0016 \times 10^{-2}$	6.0223
4	$\begin{array}{c} 2.9985 \\ \pm 2.9985 \times 10^{-2} \end{array}$	-0.42287 $\pm -4.2287 \times 10^{-3}$	-9.9949 $\pm -9.9949 \times 10^{-2}$	2.99×10^{3} ± 29.9	7.475 $\pm 7.475 \times 10^{-2}$	$4.9961 \\ \pm 4.9961 \times 10^{-2}$	$5.0025 \\ \pm 5.0025 \times 10^{-2}$	2.9784

Estimation performed using NMR-EsPy.

Author: Simon Hulse For more information:



https://foroozandehgroup.github.io/NMR-EsPy



https://github.com/foroozandehgroup/NMR-EsPy



simon.hulse@chem.ox.ac.uk

If used in a publication, please cite:

Simon G. Hulse, Mohammadali Foroozandeh. "Newton meets Ockham: Parameter estimation and model selection of NMR data with NMR-EsPy". J. Magn. Reson. 338 (2022) 107173.

https://doi.org/10.1016/j.jmr.2022.107173