



10:17:17 06-04-22 Example result file.

Experiment Information

Parameter	F1	F2
Nucleus	N/A	¹H
Transmitter Frequency (MHz)	N/A	400
Sweep Width (Hz)	50	5000
Sweep Width (ppm)	N/A	12.5
Transmitter Offset (Hz)	0	2000
Transmitter Offset (ppm)	N/A	5

Estimation Result

Osc.	а	φ (°)	f ₁ (Hz)	f ₂ (Hz)	<i>f</i> ₂ (ppm)	$\eta_1 \; (s^{-1})$	$\eta_2~(\mathrm{s}^{-1})$	ſ
1	$0.99215 \\ \pm 9.9215 \times 10^{-3}$	0.53866 $\pm 5.3866 \times 10^{-3}$	20.007 ±0.20007	3.02×10^{3} ± 30.2	7.55 $\pm 7.55 \times 10^{-2}$	$5.0039 \\ \pm 5.0039 \times 10^{-2}$	5.0024 ±5.0024 × 10 ⁻²	1
2	3.0048 $\pm 3.0048 \times 10^{-2}$	-0.5152 $\pm -5.152 \times 10^{-3}$	$10 \\ \pm 0.1$	3.01×10^{3} ± 30.1	7.525 $\pm 7.525 \times 10^{-2}$	$5.0001 \\ \pm 5.0001 \times 10^{-2}$	4.999 $\pm 4.999 \times 10^{-2}$	3.0592
3	$6.0006 \\ \pm 6.0006 \times 10^{-2}$	$0.13432\\\pm 1.3432\times 10^{-3}$	9.529×10^{-3} $\pm 9.529 \times 10^{-5}$	$\begin{array}{c} 3 \times 10^3 \\ \pm 30 \end{array}$	$7.5 \pm 7.5 \times 10^{-2}$	$4.9978 \\ \pm 4.9978 \times 10^{-2}$	4.9984 $\pm 4.9984 \times 10^{-2}$	6.015
4	$\begin{array}{c} 2.9991 \\ \pm 2.9991 \times 10^{-2} \end{array}$	-0.26666 $\pm -2.6666 \times 10^{-3}$	$-10.007 \\ \pm -0.10007$	2.99×10^{3} ± 29.9	7.475 $\pm 7.475 \times 10^{-2}$	4.9987 $\pm 4.9987 \times 10^{-2}$	$5.0059 \\ \pm 5.0059 \times 10^{-2}$	2.9756

Estimation performed using NMR-EsPy.

Author: Simon Hulse For more information:



https://foroozandehgroup.github.io/NMR-EsPy



https://github.com/foroozandehgroup/NMR-EsPy



simon.hulse@chem.ox.ac.uk

If used in a publication, please cite:

Simon G. Hulse, Mohammadali Foroozandeh. "Newton meets Ockham: Parameter estimation and model selection of NMR data with NMR-EsPy". J. Magn. Reson. 338 (2022) 107173.

https://doi.org/10.1016/j.jmr.2022.107173