

I. (a) For ridge regression:  $\hat{\beta}_{\text{ridge}} = (\mathbf{Z}^T \mathbf{Z} + \lambda \mathbf{I})^{-1} \mathbf{Z}^T \mathbf{y}$  ( $\lambda > 0$ )

$$\|\hat{\beta}_{\text{ridge}}\|_2^2 = \hat{\beta}_{\text{ridge}}^T \hat{\beta}_{\text{ridge}} \\ = \mathbf{y}^T \mathbf{Z} (\mathbf{Z}^T \mathbf{Z} + \lambda \mathbf{I})^{-2} \mathbf{Z}^T \mathbf{y}$$

$$\frac{\partial \|\hat{\beta}_{\text{ridge}}\|_2^2}{\partial \lambda} = -2 \mathbf{I} [\mathbf{y}^T \mathbf{Z} (\mathbf{Z}^T \mathbf{Z} + \lambda \mathbf{I})^{-3} \mathbf{Z}^T \mathbf{y}]$$

Since  $\mathbf{Z}^T \mathbf{Z} + \lambda \mathbf{I} > 0$ ,  $\frac{\partial \|\hat{\beta}_{\text{ridge}}\|_2^2}{\partial \lambda}$  is negative, which means

$\|\hat{\beta}_{\text{ridge}}\|_2$  increases as  $\lambda \rightarrow 0$ .

(b) For Lasso regression:  $\hat{\beta}_{\text{lasso}} = \underset{\beta}{\operatorname{argmin}} \left\{ \|\mathbf{Y} - \mathbf{X}\beta\|_2^2 + \lambda \|\beta\|_1 \right\}$

Let  $\lambda_1 < \lambda_2$ . For  $\lambda_1$ , the solution  $\hat{\beta}^{(1)}$  satisfy

$$\|\mathbf{Y} - \mathbf{X}\hat{\beta}^{(1)}\|_2^2 + \lambda_1 \|\hat{\beta}^{(1)}\|_1 \leq \|\mathbf{Y} - \mathbf{X}\hat{\beta}^{(2)}\|_2^2 + \lambda_1 \|\hat{\beta}^{(2)}\|_1 \quad (1)$$

Also for  $\lambda_2$ , the solution  $\hat{\beta}^{(2)}$  satisfy

$$\|\mathbf{Y} - \mathbf{X}\hat{\beta}^{(2)}\|_2^2 + \lambda_2 \|\hat{\beta}^{(2)}\|_1 \leq \|\mathbf{Y} - \mathbf{X}\hat{\beta}^{(1)}\|_2^2 + \lambda_2 \|\hat{\beta}^{(1)}\|_1 \quad (2)$$

(1) + (2), then we have

$$\lambda_1 \|\hat{\beta}^{(1)}\|_1 + \lambda_2 \|\hat{\beta}^{(2)}\|_1 \leq \lambda_1 \|\hat{\beta}^{(2)}\|_1 + \lambda_2 \|\hat{\beta}^{(1)}\|_1$$

$$(\lambda_2 - \lambda_1) \left[ \|\hat{\beta}^{(2)}\|_1 - \|\hat{\beta}^{(1)}\|_1 \right] \leq 0$$

Since  $\lambda_2 - \lambda_1 > 0$ , so it is satisfy

$$\|\hat{\beta}^{(2)}\|_1 < \|\hat{\beta}^{(1)}\|_1$$

Thus, when  $\lambda \rightarrow 0$ ,  $\|\hat{\beta}_{\text{lasso}}\|_1$  increases

2.

The solution of Ridge Regression is  $\hat{\beta} = (\mathbf{Z}^T \mathbf{Z} + \lambda \mathbf{I})^{-1} \mathbf{Z}^T \mathbf{y}$

let  $x_i := \mathbf{z}_i$ , and the equation of question is

$$\hat{f}(\mathbf{x}) = (\sum_{i=1}^n c_i x_i)^T \mathbf{x}, \text{ define } \mathbf{C} = [c_1, c_2, \dots, c_n]^T$$

$$\text{Set } (\sum_{i=1}^n c_i x_i)^T = \hat{\beta}^T, (\mathbf{C}^T \mathbf{Z})^T = \hat{\beta}^T$$

$$\mathbf{Z}^T \mathbf{C} = (\mathbf{Z}^T \mathbf{Z} + \lambda \mathbf{I})^{-1} \mathbf{Z}^T \mathbf{y}$$

According to the Hint:

$$\mathbf{Z}^T \mathbf{C} = \mathbf{Z}^T (\mathbf{Z} \mathbf{Z}^T + \lambda \mathbf{I})^{-1} \mathbf{y}$$

$$\mathbf{Z} \mathbf{Z}^T \mathbf{C} = \mathbf{Z} \mathbf{Z}^T (\mathbf{Z} \mathbf{Z}^T + \lambda \mathbf{I})^{-1} \mathbf{y}$$

$$\mathbf{C} = (\mathbf{Z} \mathbf{Z}^T + \lambda \mathbf{I})^{-1} \mathbf{y}, c_i \text{ is the } i\text{th term in vector } \mathbf{C}$$

## Question (a)

The original code as follows:

```
import numpy as np
np.random.seed(123)
def genData(n):
    mu = np.array([-2.0, 3.0])
    A = np.array([[1.0, 2.0], [3.0, 4.0]])
    X = np.random.normal(size = n * 2)
    X = X.reshape((2, n))
    X = (X + mu).dot(A)
    return(X.T)
```

```
genData(10)
```

```
-----  
ValueError
```

```
Traceback (most recent call last)
```

```
Cell In[2], line 1
----> 1 genData(10)
```

```
Cell In[1], line 8, in genData(n)
    6 X = np.random.normal(size = n * 2)
    7 X = X.reshape((2, n))
----> 8 X = (X + mu).dot(A)
    9 return(X.T)
```

```
ValueError: operands could not be broadcast together with shapes (2,10) (2,)
```

## Result:

Theoretically: Return a NumPy array of shape (10, 2) representing 10 samples, each with two features.

## Problem:

The shape of X is (2,10), however, the shape of mu is (2,). NumPy cannot directly add arrays of these two shapes. To solve this problem, we need to make sure that the shape of mu is correctly broadcast to X.

## New errors:

Although I have fixed one error, it seems that a new problem has arisen. The error message should be that the matrix multiplication dimensions do not match. Next, I will fix this problem under the prompt of this error report.

```
import numpy as np
np.random.seed(123)
def genData(n):
    mu = np.array([-2.0, 3.0]).reshape(2,1) # Reshape mu into the shape of (2,1)
    A = np.array([[1.0, 2.0], [3.0, 4.0]])
    X = np.random.normal(size = n * 2)
    X = X.reshape((2, n))
    X = (X + mu).dot(A)
    return(X.T)
```

```
genData(10)
```

```
-----
```

```
ValueError
```

```
Traceback (most recent call last)
```

```
Cell In[2], line 1
----> 1 genData(10)
```

```
Cell In[1], line 8, in genData(n)
    6 X = np.random.normal(size = n * 2)
    7 X = X.reshape((2, n))
----> 8 X = (X + mu).dot(A)
    9 return(X.T)
```

```
ValueError: shapes (2,10) and (2,2) not aligned: 10 (dim 1) != 2 (dim 0)
```

```
import numpy as np
np.random.seed(123)
def genData(n):
    mu = np.array([-2.0, 3.0])
    A = np.array([[1.0, 2.0], [3.0, 4.0]])
    X = np.random.normal(size = n * 2)
    X = X.reshape((2, n))
    X = A.T @ X + mu.reshape(2, 1) # Reshape mu into the shape of (2,1) and fix the
    problem of wrong dimensions
    return(X.T)
```

```
np.random.seed(123)
genData(10)
```

```
array([[-5.12228906, -1.88680581],
       [-1.28678146,  4.61585502],
       [ 2.75714738,  9.5315155 ],
       [-5.4230007 , -2.56819741],
       [-3.91054613,  0.06687166],
       [-1.65161729,  4.56546797],
       [ 2.191111 ,  6.97036184],
       [ 4.13144564, 10.8893191 ],
       [ 2.27809795,  9.54808811],
       [-1.7081812 ,  2.81126479]])
```

The main changes:

code line "X = (X + mu).dot(A)" change to "X = A.T @ X + mu.reshape(2, 1)"

## Question (b)

---

The result of \$x\\_1\$ and \$x\\_2\$

```
$x_1$: array([-5.12228906, -1.28678146, 2.75714738, -5.4230007 , -3.91054613, -1.65161729, 2.191111 ,
4.13144564, 2.27809795, -1.7081812 ])
$x_2$: array([-1.88680581, 4.61585502, 9.5315155 , -2.56819741, 0.06687166, 4.56546797, 6.97036184,
10.8893191 , 9.54808811, 2.81126479])
```

## Question (c)

---

```
np.random.seed(123)
X = genData(10000)

mu_hat = np.mean(X, axis=0)
center = X - mu_hat
sigma_hat = (center.T @ center) / (X.shape[0] - 1)

print("Sample mean: ", mu_hat)
print("Sample covariance matrix: ", sigma_hat)
```

```
Sample mean:  [-1.9591595   3.06092859]
Sample covariance matrix:  [[10.07612824 14.10999769]]
```

```
[14.10999769 20.15635737]]
```

$$\hat{\mu} = \left( -1.9591595, 3.06092859 \right)^T$$

$$\hat{\Sigma} = \begin{pmatrix} 10.07612824 & 14.10999769 \\ 14.10999769 & 20.15635737 \end{pmatrix}$$

## Question (d)

```
np.random.seed(123)
Y = genData(1000)
Y_sq = np.sum(Y**2, axis=1)
dist = Y_sq[:, None] + Y_sq[None, :] - 2 * Y @ Y.T
K = np.exp(-dist)

print(K)
```

```
[[1.0000000e+00 3.66335577e-55 2.42445105e-47 ... 4.25511028e-03
 3.12190047e-01 7.51012396e-07]
 [3.66335577e-55 1.0000000e+00 4.71361082e-01 ... 7.80809228e-35
 9.15395439e-64 7.05305182e-25]
 [2.42445105e-47 4.71361082e-01 1.0000000e+00 ... 1.08316440e-28
 1.79774847e-55 1.00198269e-19]
 ...
 [4.25511028e-03 7.80809228e-35 1.08316440e-28 ... 1.0000000e+00
 2.36317721e-05 1.32376993e-01]
 [3.12190047e-01 9.15395439e-64 1.79774847e-55 ... 2.36317721e-05
 1.0000000e+00 3.08800186e-10]
 [7.51012396e-07 7.05305182e-25 1.00198269e-19 ... 1.32376993e-01
 3.08800186e-10 1.0000000e+00]]
```

```
eigenvalues = np.linalg.eigvals(K)
largest_3 = np.sort(eigenvalues)[-3:][::-1]
print("The largest 3 eigenvalues are: ", largest_3)
```

```
The largest 3 eigenvalues are: [105.14253068+0.j 92.05784395+0.j
79.63704892+0.j]
```

4. (a) Let  $f(x) = (x-1)^2$   $g(x) = \lambda|x|$ . Both  $f(x), g(x)$  are convex.

Hence,  $(x-1)^2 + \lambda|x|$  is convex,  $x^*(\lambda)$  is global minimizer.

For any  $x \neq x^*(\lambda)$ ,  $(x^*(\lambda)-1)^2 + \lambda|x^*(\lambda)| < (x-1)^2 + \lambda|x|$  holds

(b)

For  $x > 0$ , we have  $h(x) = (x-1)^2 + \lambda x$ .

$$h'(x) = 2(x-1) + \lambda = 0, \quad x = \frac{2-\lambda}{2} \geq 0, \quad \lambda < 2$$

For  $x < 0$ , we have  $h(x) = (x-1)^2 - \lambda x$

$$h'(x) = 2(x-1) - \lambda = 0, \quad x = \frac{2+\lambda}{2} < 0, \quad \lambda < -2 \text{ which has a contradiction}$$

with  $\lambda > 0$ . Thus it

$$\text{For } x=0 \quad \lim_{x \rightarrow 0^+} h'(x) = -2 + \lambda, \quad \lim_{x \rightarrow 0^-} h'(x) = -2 - \lambda < 0$$

If  $x=0$  is the global minimizer, then have  $\lim_{x \rightarrow 0^+} h(x) = -2 + \lambda \geq 0$



$\lambda \geq 2$ ,  $\lim_{x \rightarrow 0^-} h(x) = -2 - \lambda < 0$  always holds

In conclusion:

① If  $0 < \lambda < 2$ ,  $x^*(\lambda) = \frac{2-\lambda}{2}$

② If  $\lambda \geq 2$ ,  $x^*(\lambda) = 0$

S.



6.

$$(a) \text{ Let } k(x,y) = x^3y^3 + xy := k_1(x,y) + k_2(x,y)$$

For  $k_1(x,y)$ , let  $x_1 \dots x_n, c_1 \dots c_n \in \mathbb{R}$ . We have

Obviously,  $k_1(x,y) = k_1(y,x)$ . For any positive integer  $n$ , let  $x_1 \dots x_n, c_1 \dots c_n \in \mathbb{R}$ . we have

$$\sum_{i=1}^n \sum_{j=1}^n c_i c_j x_i^3 x_j^3 = \left[ \sum_{i=1}^n c_i x_i^3 \right]^2 \geq 0, \quad k_1(x,y) \text{ is a semi-positive kernel.}$$

For  $k_2(x,y)$ . Similarly,  $k_2(x,y) = k_2(y,x)$

$$\sum_{i=1}^n \sum_{j=1}^n c_i c_j xy = \left[ \sum_{i=1}^n c_i x_i \right]^2 \geq 0, \quad k_2(x,y) \text{ is a semi-positive kernel}$$

$k(x,y) = k_1(x,y) + k_2(x,y)$  is a semi-positive kernel

$$(b) \text{ Let } G_1(x,y) = \langle x, y \rangle^3, \quad G_2(x,y) = \langle x, y \rangle \quad \text{let } x_1 \dots x_n, c_1 \dots c_n \in \mathbb{R}.$$

Consider  $G_2(x,y)$ .  $G_2(x,y) = G_2(y,x)$

$$\sum_{i=1}^n \sum_{j=1}^n c_i c_j \langle x_i, x_j \rangle = \left\langle \sum_{i=1}^n c_i x_i, \sum_{j=1}^n c_j x_j \right\rangle = \left\| \sum_{i=1}^n c_i x_i \right\|^2 \geq 0$$

$G_2(x,y)$  is a semi-positive kernel

Thus, the Gram matrix  $G_2$  of  $G_2(x,y)$  is semi-positive.

Suppose the  $G_1(x,y)$ 's Gram matrix is  $G_1$ .

$$G_1 = G_2 \circ G_2 \circ G_2 \text{ is also semi-positive (Schur theorem)}$$

where  $\circ$  is Hadamard product.  $G_1(x,y)$  is a semi-positive kernel

Hence,  $G(x,y) = G_1(x,y) + G_2(x,y)$  is a semi-positive kernel

(c) In  $(-1, 1)$ , using Geometric series expansion  $\frac{1}{1-xy} = \sum_{k=0}^{\infty} (xy)^k$ .

$$k(x,y) = k(y,x) \cdot k(x,y) = \frac{1+xy}{1-xy} = (1+xy) \sum_{k=0}^{\infty} (xy)^k = \sum_{k=0}^{\infty} (xy)^k + \sum_{k=0}^{\infty} (xy)^{k+1}$$

For any positive integer  $n$ , let  $x_1 \dots x_n \in (-1, 1)$ ,  $c_1 \dots c_n \in \mathbb{R}$

$$\begin{aligned} \sum_{i=1}^n \sum_{j=1}^n c_i c_j \left( \sum_{k=0}^{\infty} (x_i x_j)^k + \sum_{k=0}^{\infty} (x_i x_j)^{k+1} \right) &= \sum_{i=1}^n \sum_{j=1}^n c_i c_j \left( 1 + 2 \sum_{k=1}^{\infty} (x_i x_j)^k \right) \\ &= \left( \sum_{i=1}^n c_i \right)^2 + 2 \sum_{k=1}^{\infty} \left( \sum_{i=1}^n c_i x_i^k \right)^2 \geq 0 \end{aligned}$$

Hence,  $k(x,y)$  is a semi-positive kernel

(d)

$$k(x, u) = WS(x-u) = \frac{e^{i(x-u)} + e^{-i(x-u)}}{2} \quad \text{for all } x \in C$$

$$k(u, x) = WS(u-x) = WS(-(x-u)) = WS(x-u) = k(x, u)$$

For any positive integer  $n$ . Let  $x_1, \dots, x_n \in C$ ,  $c_1, \dots, c_n \in R$

$$\begin{aligned} & \sum_{i=1}^n \sum_{j=1}^n c_i g_j \cdot \frac{e^{i(x_i - x_j)} + e^{-i(x_i - x_j)}}{2} = \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n c_i g_j e^{i(x_i - x_j)} + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n c_i g_j e^{-i(x_i - x_j)} \\ &= \frac{1}{2} \left( \sum_{i=1}^n c_i e^{ix_i} \right) \left( \sum_{j=1}^n g_j e^{-ix_j} \right) + \frac{1}{2} \left( \sum_{i=1}^n c_i e^{-ix_i} \right) \left( \sum_{j=1}^n g_j e^{ix_j} \right) \\ &= \frac{1}{2} \left( \sum_{i=1}^n c_i e^{ix_i} \right) \overline{\left( \sum_{j=1}^n g_j e^{-ix_j} \right)} + \overline{\frac{1}{2} \left( \sum_{i=1}^n c_i e^{-ix_i} \right) \left( \sum_{j=1}^n g_j e^{ix_j} \right)} \\ &= \boxed{\left| \sum_{i=1}^n c_i e^{ix_i} \right|^2 \geq 0} \end{aligned}$$

Hence,  $k(u, x)$  is a semi-positive kernel