

The YEASTRACT+ Regulation File: Publications and Links

Michael Volk

October 4, 2026

The YEASTRACT+ regulation file for *Saccharomyces cerevisiae* holds 356,181 rows of evidence that a transcription factor regulates a target gene, credited to 1,671 papers. Table 4 lists every paper, most rows first, with its DOI, PubMed and PubMed Central links. 84 of the papers are also studies in the SPELL expression archive, and one more reanalyzes a SPELL study. Table 3 ranks the papers outside SPELL: 2 of them carry half of the rows credited outside the archive and 89 carry 95 percent.

Tables generated by `experiments/037-yeastract/scripts/yeastract_publication_tables.py`

Manifest: `experiments/037-yeastract/results/yeastract_publications.csv`

SPELL manifest: `experiments/015-spell/results/spell_publications.csv`

Working notes: `notes/experiments.037-yeastract.publications.md`

Section status: ✓ ready ■ author-reviewed, keep stable ✗ not done, agents may edit freely

Contents

1	What the file holds ✗	1
1.1	Where each field comes from ✗	2
2	Overlap with the SPELL archive ✗	2
3	Papers outside SPELL, by rows ✗	2
4	All publications ✗	11

1 What the file holds ✗

The source is the *S. cerevisiae* member of the YEASTRACT+ 2022 regulation archive, which holds one flat file for each of eleven yeast species:

archive: `YeasttractPlus_2022_all_regs.zip`
sha256: `2f2e9f8b3c905ae333814f04161d0ffd2fabf00cc9878c1bd86a8f5e366ebb3e`
member: `YeasttractPlus_2022_Scerevisiae_regs_flatfile.tsv.gz`

The member is a tab-separated table of 356,181 rows and eleven columns with no header row, and no row is repeated. The column meanings used here are read off the values: the systematic and standard names of a transcription factor, the systematic and standard names of a target gene, a PMID, a sign (**Positive**, **Negative** or **N/A**), an evidence label (**Direct**, **Indirect** or **N/A**), an assay, a condition group, a condition subgroup and a free-text condition.

A row is one piece of evidence, and one factor-target pair recurs once for each paper, assay and condition that reports it. The 356,181 rows name 226 transcription factors and 6,943 targets in 215,398 distinct pairs. 264,772 rows are labeled **Indirect**, 91,155 are labeled **Direct** and 254 carry no label (Table 1).

Table 1: Papers and rows in the YEASTRACT+ flat file by relation to the SPELL archive. *Rows* counts flat-file rows credited to the papers and *%* is their share of all rows. *Direct* and *indirect* count rows by the evidence label in column 7 of the file; rows labeled **N/A** are in neither. *DOI* and *PMC id* count the papers whose PubMed record carries one. *Years* is the range of publication years.

	papers	rows	%	direct	indirect	DOI	PMC id	years
same PMID in SPELL	84	75,342	21.2	3,493	71,846	84	55	1997–2019
reanalysis of a SPELL study	1	66,920	18.8	0	66,920	1	1	2010–2010
not in SPELL	1,586	213,919	60.1	87,662	126,006	1,547	920	1976–2022
all	1,671	356,181	100.0	91,155	264,772	1,632	976	1976–2022

1.1 Where each field comes from ✗

The PMID is column 5 of the file, and the row, factor, pair, evidence and condition counts of a paper are taken over the rows that carry its PMID. The file gives no citation, so the authors, title, journal, DOI and PMC id come from the PubMed record for that PMID, retrieved through the NCBI E-utilities `esummary` endpoint and stored beside the manifest in `experiments/037-yeastract/results/yeastract_pubmed_esummary.json`.

PubMed returns no record for 1 of the 1,671 PMIDs, and that row is listed by its id alone. 1,632 records carry a DOI and 38 do not; those are linked by their PubMed record. 976 records carry a PMC id, which means PubMed Central holds a full text. Whether that copy is in the open-access subset served by the PMC OA API was not checked.

For 106 papers the most frequent value of the assay column is empty, and those papers show a dash in the assay column of Table 4.

2 Overlap with the SPELL archive ✗

The SPELL archive distributed by the *Saccharomyces* Genome Database holds 603 expression studies, one PMID each. 84 of the 1,671 YEASTRACT+ papers have a PMID in that archive. They carry 75,342 rows, 21.2 percent of the file, and 71,846 of those rows are labeled **Indirect** against 3,493 labeled **Direct**. For these papers the archive holds an expression matrix from the same publication. 24 of the 84 SPELL studies name no GEO series.

A PMID match misses a paper that analyzes another paper’s measurements, because the file credits the analysis. One such paper has been identified. Reimand 2010 (PMID 20385592) carries 66,920 rows, 18.8 percent of the file, and its abstract states the source: “Recently, Hu and colleagues published a comprehensive study covering 269 TF knockout mutants for the yeast *Saccharomyces cerevisiae*. However, the information that can be extracted from this valuable dataset is limited by the method employed to process the microarray data. Here, we present a reanalysis of the original data using improved statistical techniques freely available from the BioConductor project.” The SPELL study Hu 2007 (PMID 17417638, GSE4654) has 269 conditions. Table 4 marks the Reimand 2010 row *via* Hu 2007.

The remaining 1,586 papers were not read for the origin of their data, so other reanalyses may be among them. They carry 213,919 rows, 60.1 percent of the file.

3 Papers outside SPELL, by rows ✗

1,587 papers have a PMID that is not in the SPELL archive, and they carry 280,839 rows, 78.8 percent of the file. The rows are concentrated in a few papers. The largest, Salin 2008, carries 74,129 rows (20.8 percent of the file) and the second, Reimand 2010, carries 66,920 (18.8 percent). Ranked by rows, 2 papers reach half of the rows outside SPELL, 20 reach 80 percent, 47 reach 90 percent, 89 reach 95 percent and 457 reach 99 percent.

Table 2 gives the same distribution over all 1,671 papers. 1,077 papers contribute five rows or fewer and together hold 2,415 rows; 384 contribute a single row.

Table 2: Papers by the number of flat-file rows each contributes. *Rows* sums over the papers in the band and *%* is the share of all rows. *Direct* and *indirect* count rows by evidence label. *SPELL* counts the papers in the band that are in the SPELL archive, by PMID or as a reanalysis; a dash is zero.

rows per paper	papers	rows	%	direct	indirect	SPELL
1	384	384	0.1	136	227	6
2–5	693	2,031	0.6	557	1,292	13
6–10	212	1,581	0.4	301	1,231	4
11–100	228	7,405	2.1	1,303	6,100	22
101–1,000	112	41,214	11.6	9,161	32,053	28
1,001–10,000	35	84,333	23.7	44,046	40,287	9
more than 10,000	7	219,233	61.6	35,651	183,582	3
all	1,671	356,181	100.0	91,155	264,772	85

Table 3 lists the 89 papers that reach 95 percent, which carry 266,921 rows. In 51 of them most rows are labeled **Indirect**, and in 38 at least half are labeled **Direct**. The 1,498 papers below the cut carry 13,918 rows and appear in Table 4 only.

Table 3: Papers whose PMID is not in the SPELL archive, most rows first, down to the paper that brings the cumulative share to 95 percent. # is the row number in Table 4. *cum.* % is the cumulative share of the rows credited to papers outside SPELL. *TFs* is the number of distinct transcription factors and *pairs* the number of distinct factor-target pairs in the paper’s rows. *Direct* and *indirect* count rows by evidence label; a dash is zero. *cond.* is the number of distinct free-text conditions and *top assay* the most frequent value of the assay column. *SPELL* names the archive study a reanalysis draws on (Sec. 2).

rank	#	Publication	rows	cum. %	TFs	pairs	direct	indirect	cond.	top assay	SPELL	link
1	1	Salin 2008 Structure and properties of transcriptional networks driving selenite stress response in yeasts. <i>BMC Genomics</i> 2008.	74,129	26.4	4	22,523	365	73,764	7	Microarray analysis - wt vs TF mutant	–	doi
2	2	Reimand 2010 Comprehensive reanalysis of transcription factor knockout expression data in <i>Saccharomyces cerevisiae</i> reveals many new targets. <i>Nucleic Acids Res</i> 2010.	66,920	50.2	178	66,920	–	66,920	1	Microarray analysis - wt vs TF mutant	via Hu 2007 GSE4654	doi
3	5	Rossi 2021 A high-resolution protein architecture of the budding yeast genome. <i>Nature</i> 2021.	13,776	55.1	86	13,776	13,776	–	1	ChIP-exo	–	doi
4	6	Holland 2019 Predictive models of eukaryotic transcriptional regulation reveals changes in transcription factor roles and promoter usage between metabolic conditions. <i>Nucleic Acids Res</i> 2019.	11,484	59.2	20	4,729	11,484	–	4	ChIP-exo	–	doi
5	7	Harbison 2004 Transcriptional regulatory code of a eukaryotic genome. <i>Nature</i> 2004.	10,026	62.8	145	10,008	10,026	–	4	Chip-on-chip	–	doi
6	8	Lee 2002 Transcriptional regulatory networks in <i>Saccharomyces cerevisiae</i> . <i>Science</i> 2002.	8,335	65.8	99	8,335	8,335	–	1	Chip-on-chip	–	doi
7	9	Staschke 2010 Integration of general amino acid control and target of rapamycin (TOR) regulatory pathways in nitrogen assimilation in yeast. <i>J Biol Chem</i> 2010.	6,155	67.9	4	3,601	–	6,155	19	Microarray analysis - wt vs TF mutant	–	doi
8	11	Bergenholtm 2018 Reconstruction of a Global Transcriptional Regulatory Network for Control of Lipid Metabolism in Yeast by Using Chromatin Immunoprecipitation with Lambda Exonuclease Digestion. <i>mSystems</i> 2018.	5,050	69.7	5	2,598	5,050	–	4	ChIP-exo	–	doi
9	12	Ostrow 2014 Fkh1 and Fkh2 bind multiple chromosomal elements in the <i>S. cerevisiae</i> genome with distinct specificities and cell cycle dynamics. <i>PLoS One</i> 2014.	3,811	71.1	2	3,811	3,811	–	1	Chip-on-chip	–	doi
10	14	Workman 2006 A systems approach to mapping DNA damage response pathways. <i>Science</i> 2006.	3,532	72.4	86	3,532	3,532	–	1	Chip-on-chip	–	doi
11	15	Ni 2009 Dynamic and complex transcription factor binding during an inducible response in yeast. <i>Genes Dev</i> 2009.	3,296	73.5	9	2,187	3,296	–	3	Chip-on-chip	–	doi

Table 3, continued

rank	#	Publication	rows	cum. %	TFs	pairs	direct	indirect	cond.	top assay	SPELL	link
12	16	Van Dalfsen 2018 Global Proteome Remodeling during ER Stress Involves Hac1-Driven Expression of Long Undecoded Transcript Isoforms. <i>Dev Cell</i> 2018.	3,019	74.6	1	1,185	–	3,019	3	Ribo-seq: WT vs TF mutant	–	doi
13	17	Lavoie 2010 Evolutionary tinkering with conserved components of a transcriptional regulatory network. <i>PLoS Biol</i> 2010.	2,669	75.6	6	2,669	2,669	–	1	Chip-on-chip	–	doi
14	18	Hong 2018 An incoherent feedforward loop facilitates adaptive tuning of gene expression. <i>Elife</i> 2018.	2,478	76.4	1	2,319	2	2,476	3	RNA-seq - wt vs TF mutant	–	doi
15	19	Horak 2002 Complex transcriptional circuitry at the G1/S transition in <i>Saccharomyces cerevisiae</i> . <i>Genes Dev</i> 2002.	2,128	77.2	10	2,128	2,128	–	2	Chip-on-chip	–	doi
16	20	Kuang 2017 Msn2/4 regulate expression of glycolytic enzymes and control transition from quiescence to growth. <i>Elife</i> 2017.	2,049	77.9	2	2,049	2,049	–	1	ChIP-seq	–	doi
17	21	Venters 2009 A canonical promoter organization of the transcription machinery and its regulators in the <i>Saccharomyces</i> genome. <i>Genome Res</i> 2009.	2,002	78.6	8	2,002	2,002	–	1	ChIP	–	doi
18	22	Rong-Mullins 2018 Transcriptional Profiling of <i>Saccharomyces cerevisiae</i> Reveals the Impact of Variation of a Single Transcription Factor on Differential Gene Expression in 4NQO, Fermentable, and Nonfermentable Carbon Sources. <i>G3 (Bethesda)</i> 2018.	1,883	79.3	1	1,109	–	1,883	2	RNA-seq - wt vs TF mutant	–	doi
19	24	Nocedal 2017 Gene regulatory network plasticity predates a switch in function of a conserved transcription regulator. <i>Elife</i> 2017.	1,805	80.0	1	1,765	1,805	–	2	ChIP-seq	–	doi
20	26	Shu 2020 Rph1 coordinates transcription of ribosomal protein genes and ribosomal RNAs to control cell growth under nutrient stress conditions. <i>Nucleic Acids Res</i> 2020.	1,774	80.6	1	1,573	–	1,774	2	RNA-seq - wt vs TF overexpression	–	doi
21	27	Kasahara 2007 Assembly of regulatory factors on rRNA and ribosomal protein genes in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2007.	1,689	81.2	3	1,689	1,689	–	2	Chip-on-chip	–	doi
22	30	Vizoso-Vázquez 2018 Ixr1 Regulates Ribosomal Gene Transcription and Yeast Response to Cisplatin. <i>Sci Rep</i> 2018.	1,522	81.7	1	1,100	–	1,522	4	Microarray analysis - wt vs TF mutant	–	doi
23	31	Borneman 2006 Target hub proteins serve as master regulators of development in yeast. <i>Genes Dev</i> 2006.	1,512	82.3	6	1,512	1,512	–	4	Chip-on-chip	–	doi
24	32	Zhang 2016 Transcriptomic profiling of chemical exposure reveals roles of Yap1 in protecting yeast cells from oxidative and other types of stresses. <i>Yeast</i> 2016.	1,405	82.8	1	1,405	–	1,405	5	RNA-seq - wt vs TF mutant	–	doi

Table 3, continued

rank	#	Publication	rows	cum. %	TFs	pairs	direct	indirect	cond.	top assay	SPELL	link
25	33	Eastmond 2006 Genome-wide analysis reveals new roles for the activation domains of the <i>Saccharomyces cerevisiae</i> heat shock transcription factor (Hsf1) during the transient heat shock response. <i>J Biol Chem</i> 2006.	1,366	83.3	3	1,366	–	1,366	1	Microarray analysis - wt vs TF mutant	–	doi
26	35	Borneman 2007 Transcription factor binding site identification in yeast: a comparison of high-density oligonucleotide and PCR-based microarray platforms. <i>Funct Integr Genomics</i> 2007.	1,323	83.7	3	1,323	1,323	–	1	Chip-on-chip	–	doi
27	36	Read 2016 A trans-acting Variant within the Transcription Factor RIM101 Interacts with Genetic Background to Determine its Regulatory Capacity. <i>PLoS Genet</i> 2016.	1,271	84.2	1	1,271	–	1,271	2	RNA-seq - wt vs TF mutant	–	doi
28	39	Srinivasan 2020 Genome-scale reconstruction of Gcn4/ATF4 networks driving a growth program. <i>PLoS Genet</i> 2020.	1,220	84.6	1	1,111	308	912	1	RNA-seq - wt vs TF mutant	–	doi
29	40	Huebert 2012 Dynamic changes in nucleosome occupancy are not predictive of gene expression dynamics but are linked to transcription and chromatin regulators. <i>Mol Cell Biol</i> 2012.	1,203	85.0	1	1,203	1,203	–	1	Chip-on-chip	–	doi
30	41	Rossouw 2012 Transcriptional regulation and the diversification of metabolism in wine yeast strains. <i>Genetics</i> 2012.	1,126	85.4	1	1,126	–	1,126	1	Microarray analysis - wt vs TF overexpression	–	doi
31	42	Chow 2019 Filamentation Regulatory Pathways Control Adhesion-Dependent Surface Responses in Yeast. <i>Genetics</i> 2019.	1,011	85.8	2	1,001	–	1,011	2	RNA-seq - wt vs TF mutant	–	doi
32	45	Cromie 2017 Transcriptional Profiling of Biofilm Regulators Identified by an Overexpression Screen in <i>Saccharomyces cerevisiae</i> . <i>G3 (Bethesda)</i> 2017.	938	86.1	3	938	–	938	1	RNA-seq - wt vs TF overexpression	–	doi
33	46	Zhu 2015 High-Resolution Global Analysis of the Influences of Bas1 and Ino4 Transcription Factors on Meiotic DNA Break Distributions in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2015.	936	86.5	2	936	234	702	1	RNA-seq - wt vs TF mutant	–	doi
34	47	Lefrançois 2009 Efficient yeast ChIP-Seq using multiplex short-read DNA sequencing. <i>BMC Genomics</i> 2009.	926	86.8	1	926	926	–	2	ChIP-seq	–	doi
35	48	Miles 2013 Xbp1 directs global repression of budding yeast transcription during the transition to quiescence and is important for the longevity and reversibility of the quiescent state. <i>PLoS Genet</i> 2013.	853	87.1	1	853	7	846	6	RNA-seq - wt vs TF mutant	–	doi

Table 3, continued

rank	#	Publication	rows	cum. %	TFs	pairs	direct	indirect	cond.	top assay	SPELL	link
36	49	Eckert-Boulet 2004 Transcriptional profiling of extracellular amino acid sensing in <i>Saccharomyces cerevisiae</i> and the role of Stp1p and Stp2p. <i>Yeast</i> 2004.	840	87.4	2	569	–	840	2	Microarray analysis - wt vs TF mutant	–	doi
37	51	Walvekar 2018 Methionine coordinates a hierarchically organized anabolic program enabling proliferation. <i>Mol Biol Cell</i> 2018.	835	87.7	1	835	–	835	1	RNA-seq - wt vs TF mutant	–	doi
38	52	Weidberg 2018 MitoCPR-A surveillance pathway that protects mitochondria in response to protein import stress. <i>Science</i> 2018.	805	88.0	1	803	–	805	3	RNA-seq - wt vs TF mutant	–	doi
39	53	Lardenois 2015 Global alterations of the transcriptional landscape during yeast growth and development in the absence of Ume6-dependent chromatin modification. <i>Mol Genet Genomics</i> 2015.	788	88.3	1	788	3	785	6	Microarray analysis - wt vs TF mutant	–	doi
40	54	Marion 2004 Sfp1 is a stress- and nutrient-sensitive regulator of ribosomal protein gene expression. <i>Proc Natl Acad Sci U S A</i> 2004.	766	88.5	1	766	92	674	5	Microarray analysis - wt vs TF mutant	–	doi
41	56	Kim 2019 Tailoring the <i>Saccharomyces cerevisiae</i> endoplasmic reticulum for functional assembly of terpene synthesis pathway. <i>Metab Eng</i> 2019.	739	88.8	1	739	–	739	1	RNA-seq - wt vs TF overexpression	–	doi
42	57	Albert 2019 Sfp1 regulates transcriptional networks driving cell growth and division through multiple promoter-binding modes. <i>Genes Dev</i> 2019.	713	89.1	1	311	713	–	3	ChIP-seq	–	doi
43	59	Jorgensen 2004 A dynamic transcriptional network communicates growth potential to ribosome synthesis and critical cell size. <i>Genes Dev</i> 2004.	656	89.3	2	643	–	656	6	Microarray analysis - wt vs TF mutant	–	doi
44	60	Gallagher 2014 Divergence in a master variator generates distinct phenotypes and transcriptional responses. <i>Genes Dev</i> 2014.	644	89.5	1	644	644	–	3	ChIP-seq	–	doi
45	61	Lahav 2007 Role of transcription factor Kar4 in regulating downstream events in the <i>Saccharomyces cerevisiae</i> pheromone response pathway. <i>Mol Cell Biol</i> 2007.	609	89.7	2	609	2	607	4	Microarray analysis - wt vs TF mutant	–	doi
46	62	Hazelwood 2009 Identity of the growth-limiting nutrient strongly affects storage carbohydrate accumulation in anaerobic chemostat cultures of <i>Saccharomyces cerevisiae</i> . <i>Appl Environ Microbiol</i> 2009.	598	90.0	2	540	–	598	2	Microarray analysis - wt vs TF mutant	–	doi
47	63	McIsaac 2013 Synthetic gene expression perturbation systems with rapid, tunable, single-gene specificity in yeast. <i>Nucleic Acids Res</i> 2013.	581	90.2	1	581	–	581	1	Microarray analysis - wt vs TF overexpression	–	doi

Table 3, continued

rank	#	Publication	rows	cum. %	TFs	pairs	direct	indirect	cond.	top assay	SPELL	link
48	65	Banerjee 2008 Responses of pathogenic and nonpathogenic yeast species to steroids reveal the functioning and evolution of multidrug resistance transcriptional networks. <i>Eukaryot Cell</i> 2008.	572	90.4	2	572	–	572	2	Microarray analysis - wt vs TF mutant	–	doi
49	70	Iyer 2001 Genomic binding sites of the yeast cell-cycle transcription factors SBF and MBF. <i>Nature</i> 2001.	532	90.6	2	353	532	–	3	Chip-on-chip	–	doi
50	71	Pfanzagl 2018 A constitutive active allele of the transcription factor Msn2 mimicking low PKA activity dictates metabolic remodeling in yeast. <i>Mol Biol Cell</i> 2018.	530	90.7	1	530	–	530	1	Microarray analysis - wt vs TF mutant	–	doi
51	72	Knott 2012 Forkhead transcription factors establish origin timing and long-range clustering in <i>S. cerevisiae</i> . <i>Cell</i> 2012.	513	90.9	1	513	467	46	2	ChIP-seq	–	doi
52	73	Kruegel 2011 Elevated proteasome capacity extends replicative lifespan in <i>Saccharomyces cerevisiae</i> . <i>PLoS Genet</i> 2011.	500	91.1	1	500	–	500	1	Proteomics - wt vs TF mutant	–	doi
53	74	Larochelle 2006 Oxidative stress-activated zinc cluster protein Stb5 has dual activator/repressor functions required for pentose phosphate pathway regulation and NADPH production. <i>Mol Cell Biol</i> 2006.	476	91.3	1	476	–	476	1	Microarray analysis - wt vs TF mutant	–	doi
54	76	Tan 2008 A systems approach to delineate functions of paralogous transcription factors: role of the Yap family in the DNA damage response. <i>Proc Natl Acad Sci U S A</i> 2008.	457	91.4	5	457	457	–	2	Chip-on-chip	–	doi
55	77	Fendt 2010 Tradeoff between enzyme and metabolite efficiency maintains metabolic homeostasis upon perturbations in enzyme capacity. <i>Mol Syst Biol</i> 2010.	453	91.6	1	453	–	453	1	Microarray analysis - wt vs TF mutant	–	doi
56	78	Ouyang 2018 Integrated analysis of the yeast NADPH-regulator Stb5 reveals distinct differences in NADPH requirements and regulation in different states of yeast metabolism. <i>FEMS Yeast Res</i> 2018.	438	91.7	1	185	382	56	4	ChIP-exo	–	doi
57	79	Zeitlinger 2003 Program-specific distribution of a transcription factor dependent on partner transcription factor and MAPK signaling. <i>Cell</i> 2003.	407	91.9	3	407	407	–	4	Chip-on-chip	–	doi
58	81	Carrillo 2012 Characterizing the roles of Met31 and Met32 in coordinating Met4-activated transcription in the absence of Met30. <i>Mol Biol Cell</i> 2012.	404	92.0	2	362	363	41	1	Chip-on-chip	–	doi
59	83	Pagani 2007 Disruption of iron homeostasis in <i>Saccharomyces cerevisiae</i> by high zinc levels: a genome-wide study. <i>Mol Microbiol</i> 2007.	397	92.2	1	396	–	397	2	Microarray analysis - wt vs TF mutant	–	doi

Table 3, continued

rank	#	Publication	rows	cum. %	TFs	pairs	direct	indirect	cond.	top assay	SPELL	link
60	86	Raghevendran 2006 Hap4 is not essential for activation of respiration at low specific growth rates in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2006.	385	92.3	1	385	–	385	1	Microarray analysis - wt vs TF mutant	–	doi
61	87	Lieb 2001 Promoter-specific binding of Rap1 revealed by genome-wide maps of protein-DNA association. <i>Nat Genet</i> 2001.	374	92.4	1	374	374	–	1	Chip-on-chip	–	doi
62	88	Kelley 2009 Genome-wide fitness and expression profiling implicate Mga2 in adaptation to hydrogen peroxide. <i>PLoS Genet</i> 2009.	373	92.6	5	309	–	373	2	Microarray analysis - wt vs TF mutant	–	doi
63	89	Kleinschmidt 2005 Transcriptional profiling of <i>Saccharomyces cerevisiae</i> cells under adhesion-inducing conditions. <i>Mol Genet Genomics</i> 2005.	358	92.7	1	358	–	358	2	Microarray analysis - wt vs TF mutant	–	doi
64	90	Ernst 2007 Reconstructing dynamic regulatory maps. <i>Mol Syst Biol</i> 2007.	357	92.8	2	357	357	–	5	Chip-on-chip	–	doi
65	92	Gertz 2005 Discovery, validation, and genetic dissection of transcription factor binding sites by comparative and functional genomics. <i>Genome Res</i> 2005.	351	93.0	2	351	1	350	3	Microarray analysis - wt vs TF mutant	–	doi
66	93	Börlin 2020 The transcription factor Leu3 shows differential binding behavior in response to changing leucine availability. <i>FEMS Microbiol Lett</i> 2020.	339	93.1	1	107	339	–	5	ChIP-seq	–	doi
67	94	James Theoga Raj 2019 The copper-sensing transcription factor Mac1, the histone deacetylase Hst1, and nicotinic acid regulate de novo NAD(+) biosynthesis in budding yeast. <i>J Biol Chem</i> 2019.	327	93.2	1	316	1	326	1	RNA-seq - wt vs TF mutant	–	doi
68	95	Reja 2015 Molecular mechanisms of ribosomal protein gene coregulation. <i>Genes Dev</i> 2015.	312	93.3	4	312	312	–	1	ChIP-exo	–	doi
69	98	Schüller 2007 Membrane-active compounds activate the transcription factors Pdr1 and Pdr3 connecting pleiotropic drug resistance and membrane lipid homeostasis in <i>saccharomyces cerevisiae</i> . <i>Mol Biol Cell</i> 2007.	297	93.4	4	297	–	297	3	Microarray analysis - wt vs TF mutant	–	doi
70	99	Rawal 2018 Gcn4 Binding in Coding Regions Can Activate Internal and Canonical 5' Promoters in Yeast. <i>Mol Cell</i> 2018.	279	93.5	1	279	279	–	1	ChIP-seq	–	doi
71	102	Pimentel 2012 The role of the Yap5 transcription factor in remodeling gene expression in response to Fe bioavailability. <i>PLoS One</i> 2012.	259	93.6	1	241	1	258	10	Microarray analysis - wt vs TF mutant	–	doi
72	103	Simon 2001 Serial regulation of transcriptional regulators in the yeast cell cycle. <i>Cell</i> 2001.	252	93.7	8	252	252	–	2	Chip-on-chip	–	doi

Table 3, continued

rank	#	Publication	rows	cum. %	TFs	pairs	direct	indirect	cond.	top assay	SPELL	link
73	104	Tang 2006 Inferring direct regulatory targets from expression and genome location analyses: a comparison of transcription factor deletion and overexpression. <i>BMC Genomics</i> 2006.	250	93.8	1	250	–	250	1	Microarray analysis - wt vs TF overexpression	–	doi
74	105	Tuch 2008 The evolution of combinatorial gene regulation in fungi. <i>PLoS Biol</i> 2008.	250	93.9	1	250	250	–	2	Chip-on-chip	–	doi
75	106	Hanlon 2011 The stress response factors Yap6, Cin5, Phd1, and Skn7 direct targeting of the conserved co-repressor Tup1-Ssn6 in <i>S. cerevisiae</i> . <i>PLoS One</i> 2011.	249	94.0	1	249	249	–	1	Chip-on-chip	–	doi
76	107	Wade 2004 The transcription factor Ifh1 is a key regulator of yeast ribosomal protein genes. <i>Nature</i> 2004.	248	94.1	4	248	239	9	4	Chip-on-chip	–	doi
77	108	Chavel 2010 Multiple signals converge on a differentiation MAPK pathway. <i>PLoS Genet</i> 2010.	246	94.1	16	241	1	245	20	Microarray analysis - wt vs TF mutant	–	doi
78	109	Nikolov 2022 SWI/SNF and the histone chaperone Rtt106 drive expression of the Pleiotropic Drug Resistance network genes. <i>Nat Commun</i> 2022.	231	94.2	2	231	–	231	1	RNA-seq - wt vs TF mutant	–	doi
79	110	Tachibana 2005 Combined global localization analysis and transcriptome data identify genes that are directly coregulated by Adr1 and Cat8. <i>Mol Cell Biol</i> 2005.	226	94.3	2	226	222	4	1	ChIP	–	doi
80	111	Váchová 2004 Sok2p transcription factor is involved in adaptive program relevant for long term survival of <i>Saccharomyces cerevisiae</i> colonies. <i>J Biol Chem</i> 2004.	224	94.4	1	110	–	224	13	Microarray analysis - wt vs TF mutant	–	doi
81	112	Hickman 2011 Coordinated regulation of sulfur and phospholipid metabolism reflects the importance of methylation in the growth of yeast. <i>Mol Biol Cell</i> 2011.	224	94.5	2	224	–	224	1	Microarray analysis - wt vs TF mutant	–	doi
82	113	Loll-Krippléber 2017 P-body proteins regulate transcriptional rewiring to promote DNA replication stress resistance. <i>Nat Commun</i> 2017.	224	94.5	2	224	–	224	1	RNA-seq - wt vs TF overexpression	–	doi
83	114	Vyas 2005 Repressors Nrg1 and Nrg2 regulate a set of stress-responsive genes in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2005.	216	94.6	2	216	–	216	1	Microarray analysis - wt vs TF mutant	–	doi
84	115	Servant 2012 Tye7 regulates yeast Ty1 retrotransposon sense and antisense transcription in response to adenylc nucleotides stress. <i>Nucleic Acids Res</i> 2012.	209	94.7	2	209	1	208	2	Microarray analysis - wt vs TF mutant	–	doi
85	116	Westholm 2008 Combinatorial control of gene expression by the three yeast repressors Mig1, Mig2 and Mig3. <i>BMC Genomics</i> 2008.	206	94.8	3	206	–	206	1	Microarray analysis - wt vs TF mutant	–	doi

Table 3, continued

rank	#	Publication	rows	cum. %	TFs	pairs	direct	indirect	cond.	top assay	SPELL	link
86	118	Roque 2016 Genome-wide recruitment profiling of transcription factor Crz1 in response to high pH stress. <i>BMC Genomics</i> 2016.	203	94.8	1	203	203	–	1	ChIP-seq	–	doi
87	119	Maity 2016 Oxidative Homeostasis Regulates the Response to Reductive Endoplasmic Reticulum Stress through Translation Control. <i>Cell Rep</i> 2016.	195	94.9	1	195	–	195	1	iTRAQ-MS - wt vs TF mutant	–	doi
88	120	Elfving 2014 A dynamic interplay of nucleosome and Msn2 binding regulates kinetics of gene activation and repression following stress. <i>Nucleic Acids Res</i> 2014.	193	95.0	1	193	62	131	2	Microarray analysis - wt vs TF overexpression	–	doi
89	121	Pincus 2018 Genetic and epigenetic determinants establish a continuum of Hsf1 occupancy and activity across the yeast genome. <i>Mol Biol Cell</i> 2018.	189	95.0	1	70	129	60	5	ChIP-seq	–	doi

4 All publications ✗

Table 4 has one row per PMID, ordered by the number of flat-file rows credited to it. Authors are given in full up to six names; a longer list is cut to the first three and the last.

Table 4: Every paper credited in the YEASTRACT+ flat file, most rows first; ties are ordered by PMID. The bold label is the first author and year. *Links* gives the DOI, the PubMed record and, where one exists, the PubMed Central copy. *SPELL* is the GEO series of the SPELL study with the same PMID, *no GEO* where that study names none, *via* a study where the paper reanalyzes it, and a dash where the paper is not in the archive. *rows* is the number of flat-file rows credited to the paper, *TFs* the distinct transcription factors and *pairs* the distinct factor-target pairs in them. *Direct* and *indirect* count rows by evidence label; a dash is zero. *cond.* is the number of distinct free-text conditions and *top assay* the most frequent value of the assay column, a dash where that value is empty. Every link is clickable.

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1	Salin 2008 Salin H, Fardeau V, Piccini E, et al., Devaux F. Structure and properties of transcriptional networks driving selenite stress response in yeasts. <i>BMC Genomics</i> 2008;9:333.	doi:10.1186/1471-2164-9-333 PMID 18627600 PMC2515152	–	74,129	4	22,523	365	73,764	7	Microarray analysis - wt vs TF mutant
2	Reimand 2010 Reimand J, Vaquerizas JM, Todd AE, Vilo J, Luscombe NM. Comprehensive reanalysis of transcription factor knockout expression data in <i>Saccharomyces cerevisiae</i> reveals many new targets. <i>Nucleic Acids Res</i> 2010;38(14):4768–77.	doi:10.1093/nar/gkq232 PMID 20385592 PMC2919724	via Hu 2007 GSE4654	66,920	178	66,920	–	66,920	1	Microarray analysis - wt vs TF mutant
3	Moxley 2009 Moxley JF, Jewett MC, Antoniewicz MR, et al., Stephanopoulos G. Linking high-resolution metabolic flux phenotypes and transcriptional regulation in yeast modulated by the global regulator Gcn4p. <i>Proc Natl Acad Sci U S A</i> 2009;106(16):6477–82.	doi:10.1073/pnas.0811091106 PMID 19346491 PMC2672541	GSE4709	25,142	5	25,142	–	25,142	2	Microarray analysis - wt vs TF mutant
4	Chua 2006 Chua G, Morris QD, Sopko R, et al., Hughes TR. Identifying transcription factor functions and targets by phenotypic activation. <i>Proc Natl Acad Sci U S A</i> 2006;103(32):12045–50.	doi:10.1073/pnas.0605140103 PMID 16880382 PMC1567694	GSE5499	17,756	46	17,393	–	17,756	2	Microarray analysis - wt vs TF overexpression
5	Rossi 2021 Rossi MJ, Kuntala PK, Lai WKM, et al., Pugh BF. A high-resolution protein architecture of the budding yeast genome. <i>Nature</i> 2021;592(7853):309–314.	doi:10.1038/s41586-021-03314-8 PMID 33692541 PMC8035251	–	13,776	86	13,776	13,776	–	1	ChIP-exo
6	Holland 2019 Holland P, Bergenholm D, Börlin CS, Liu G, Nielsen J. Predictive models of eukaryotic transcriptional regulation reveals changes in transcription factor roles and promoter usage between metabolic conditions. <i>Nucleic Acids Res</i> 2019; 47(10):4986–5000.	doi:10.1093/nar/gkz253 PMID 30976803 PMC6547448	–	11,484	20	4,729	11,484	–	4	ChIP-exo
7	Harbison 2004 Harbison CT, Gordon DB, Lee TI, et al., Young RA. Transcriptional regulatory code of a eukaryotic genome. <i>Nature</i> 2004;431(7004):99–104.	doi:10.1038/nature02800 PMID 15343339 PMC3006441	–	10,026	145	10,008	10,026	–	4	Chip-on-chip
8	Lee 2002 Lee TI, Rinaldi NJ, Robert F, et al., Young RA. Transcriptional regulatory networks in <i>Saccharomyces cerevisiae</i> . <i>Science</i> 2002;298(5594):799–804.	doi:10.1126/science.1075090 PMID 12399584	–	8,335	99	8,335	8,335	–	1	Chip-on-chip
9	Staschke 2010 Staschke KA, Dey S, Zaborske JM, et al., Wek RC. Integration of general amino acid control and target of rapamycin (TOR) regulatory pathways in nitrogen assimilation in yeast. <i>J Biol Chem</i> 2010;285(22):16893–911.	doi:10.1074/jbc.M110.121947 PMID 20233714 PMC2878067	–	6,155	4	3,601	–	6,155	19	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
10	Madhani 1999 Madhani HD, Galitski T, Lander ES, Fink GR. Effectors of a developmental mitogen-activated protein kinase cascade revealed by expression signatures of signaling mutants. <i>Proc Natl Acad Sci U S A</i> 1999;96(22):12530–5.	doi:10.1073/pnas.96.22.12530 PMID 10535956 PMC22972	no GEO	5,628	6	5,628	45	5,583	11	Microarray analysis - wt vs TF mutant
11	Bergenholtm 2018 Bergenholtm D, Liu G, Holland P, Nielsen J. Reconstruction of a Global Transcriptional Regulatory Network for Control of Lipid Metabolism in Yeast by Using Chromatin Immunoprecipitation with Lambda Exonuclease Digestion. <i>mSystems</i> 2018;3(4).	doi:10.1128/mSystems.00215-17 PMID 30073202 PMC6068829	–	5,050	5	2,598	5,050	–	4	ChIP-exo
12	Ostrow 2014 Ostrow AZ, Nellimoottil T, Knott SR, Fox CA, Tavaré S, Aparicio OM. Fkh1 and Fkh2 bind multiple chromosomal elements in the <i>S. cerevisiae</i> genome with distinct specificities and cell cycle dynamics. <i>PLoS One</i> 2014;9(2):e87647.	doi:10.1371/journal.pone.0087647 PMID 24504085 PMC3913637	–	3,811	2	3,811	3,811	–	1	Chip-on-chip
13	Fendt 2010 Fendt SM, Oliveira AP, Christen S, Picotti P, Dechant RC, Sauer U. Unraveling condition-dependent networks of transcription factors that control metabolic pathway activity in yeast. <i>Mol Syst Biol</i> 2010;6:432.	doi:10.1038/msb.2010.91 PMID 21119627 PMC3010106	GSE24056	3,536	4	3,536	–	3,536	1	Microarray analysis - wt vs TF mutant
14	Workman 2006 Workman CT, Mak HC, McCuine S, et al., Ideker T. A systems approach to mapping DNA damage response pathways. <i>Science</i> 2006;312(5776):1054–9.	doi:10.1126/science.1122088 PMID 16709784 PMC2811083	–	3,532	86	3,532	3,532	–	1	Chip-on-chip
15	Ni 2009 Ni L, Bruce C, Hart C, et al., Snyder M. Dynamic and complex transcription factor binding during an inducible response in yeast. <i>Genes Dev</i> 2009;23(11):1351–63.	doi:10.1101/gad.1781909 PMID 19487574 PMC2701586	–	3,296	9	2,187	3,296	–	3	Chip-on-chip
16	Van Dalfsen 2018 Van Dalfsen KM, Hodapp S, Keskin A, et al., Brar GA. Global Proteome Remodeling during ER Stress Involves Hac1-Driven Expression of Long Undecoded Transcript Isoforms. <i>Dev Cell</i> 2018;46(2):219–235.e8.	doi:10.1016/j.devcel.2018.06.016 PMID 30016623 PMC6140797	–	3,019	1	1,185	–	3,019	3	Ribo-seq: WT vs TF mutant
17	Lavoie 2010 Lavoie H, Hogues H, Mallick J, Sellam A, Nantel A, Whiteway M. Evolutionary tinkering with conserved components of a transcriptional regulatory network. <i>PLoS Biol</i> 2010;8(3):e1000329.	doi:10.1371/journal.pbio.1000329 PMID 20231876 PMC2834713	–	2,669	6	2,669	2,669	–	1	Chip-on-chip
18	Hong 2018 Hong J, Brandt N, Abdul-Rahman F, Yang A, Hughes T, Gresham D. An incoherent feedforward loop facilitates adaptive tuning of gene expression. <i>Elife</i> 2018;7.	doi:10.7554/eLife.32323 PMID 29620523 PMC5903863	–	2,478	1	2,319	2	2,476	3	RNA-seq - wt vs TF mutant
19	Horak 2002 Horak CE, Luscombe NM, Qian J, et al., Snyder M. Complex transcriptional circuitry at the G1/S transition in <i>Saccharomyces cerevisiae</i> . <i>Genes Dev</i> 2002;16(23):3017–33.	doi:10.1101/gad.1039602 PMID 12464632 PMC187492	–	2,128	10	2,128	2,128	–	2	Chip-on-chip

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
20	Kuang 2017 Kuang Z, Pinglay S, Ji H, Boeke JD. Msn2/4 regulate expression of glycolytic enzymes and control transition from quiescence to growth. <i>Elife</i> 2017;6.	doi:10.7554/eLife.29938 PMID 28949295 PMC5634782	–	2,049	2	2,049	2,049	–	1	ChIP-seq
21	Venters 2009 Venters BJ, Pugh BF. A canonical promoter organization of the transcription machinery and its regulators in the <i>Saccharomyces</i> genome. <i>Genome Res</i> 2009;19(3):360–71.	doi:10.1101/gr.084970.108 PMID 19124666 PMC2661807	–	2,002	8	2,002	2,002	–	1	ChIP
22	Rong-Mullins 2018 Rong-Mullins X, Ayers MC, Summers M, Gallagher JEG. Transcriptional Profiling of <i>Saccharomyces cerevisiae</i> Reveals the Impact of Variation of a Single Transcription Factor on Differential Gene Expression in 4NQO, Fermentable, and Nonfermentable Carbon Sources. <i>G3 (Bethesda)</i> 2018;8(2):607–619.	doi:10.1534/g3.117.300138 PMID 29208650 PMC5919752	–	1,883	1	1,109	–	1,883	2	RNA-seq - wt vs TF mutant
23	Capaldi 2008 Capaldi AP, Kaplan T, Liu Y, et al., O’Shea EK. Structure and function of a transcriptional network activated by the MAPK Hog1. <i>Nat Genet</i> 2008;40(11):1300–6.	doi:10.1038/ng.235 PMID 18931682 PMC2825711	GSE12270	1,861	8	1,861	173	1,688	3	Microarray analysis - wt vs TF mutant
24	Nocedal 2017 Nocedal I, Mancera E, Johnson AD. Gene regulatory network plasticity predates a switch in function of a conserved transcription regulator. <i>Elife</i> 2017;6.	doi:10.7554/eLife.23250 PMID 28327289 PMC5391208	–	1,805	1	1,765	1,805	–	2	ChIP-seq
25	Auld 2006 Auld KL, Brown CR, Casolari JM, Komili S, Silver PA. Genomic association of the proteasome demonstrates overlapping gene regulatory activity with transcription factor substrates. <i>Mol Cell</i> 2006;21(6):861–71.	doi:10.1016/j.molcel.2006.02.020 PMID 16543154	GSE4272	1,784	2	1,784	1,784	–	5	Chip-on-chip
26	Shu 2020 Shu WJ, Chen R, Yin ZH, Li F, Zhang H, Du HN. Rph1 coordinates transcription of ribosomal protein genes and ribosomal RNAs to control cell growth under nutrient stress conditions. <i>Nucleic Acids Res</i> 2020;48(15):8360–8373.	doi:10.1093/nar/gkaa558 PMID 32619236 PMC7470948	–	1,774	1	1,573	–	1,774	2	RNA-seq - wt vs TF overexpression
27	Kasahara 2007 Kasahara K, Ohtsuki K, Ki S, et al., Kokubo T. Assembly of regulatory factors on rRNA and ribosomal protein genes in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2007;27(19):6686–705.	doi:10.1128/MCB.00876-07 PMID 17646381 PMC2099245	–	1,689	3	1,689	1,689	–	2	Chip-on-chip
28	Lewis 2012 Lewis JA, Gasch AP. Natural variation in the yeast glucose-signaling network reveals a new role for the Mig3p transcription factor. <i>G3 (Bethesda)</i> 2012;2(12):1607–12.	doi:10.1534/g3.112.004127 PMID 23275883 PMC3516482	GSE40153	1,547	1	1,547	–	1,547	2	Microarray analysis - wt vs TF mutant
29	Haugen 2004 Haugen AC, Kelley R, Collins JB, et al., Van Houten B. Integrating phenotypic and expression profiles to map arsenic-response networks. <i>Genome Biol</i> 2004;5(12):R95.	doi:10.1186/gb-2004-5-12-r95 PMID 15575969 PMC545798	no GEO	1,528	4	1,528	–	1,528	1	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
30	Vizoso-Vázquez 2018 Vizoso-Vázquez Á, Lamas-Maceiras M, González-Siso MI, Cerdán ME. Ixr1 Regulates Ribosomal Gene Transcription and Yeast Response to Cisplatin. <i>Sci Rep</i> 2018;8(1):3090.	doi:10.1038/s41598-018-21439-1 PMID 29449612 PMC5814428	–	1,522	1	1,100	–	1,522	4	Microarray analysis - wt vs TF mutant
31	Borneman 2006 Borneman AR, Leigh-Bell JA, Yu H, Bertone P, Gerstein M, Snyder M. Target hub proteins serve as master regulators of development in yeast. <i>Genes Dev</i> 2006;20(4):435–48.	doi:10.1101/gad.1389306 PMID 16449570 PMC1369046	–	1,512	6	1,512	1,512	–	4	Chip-on-chip
32	Zhang 2016 Zhang C, Li Z, Zhang X, Yuan L, Dai H, Xiao W. Transcriptomic profiling of chemical exposure reveals roles of Yap1 in protecting yeast cells from oxidative and other types of stresses. <i>Yeast</i> 2016;33(1):5–19.	doi:10.1002/yea.3135 PMID 26389527	–	1,405	1	1,405	–	1,405	5	RNA-seq - wt vs TF mutant
33	Eastmond 2006 Eastmond DL, Nelson HC. Genome-wide analysis reveals new roles for the activation domains of the <i>Saccharomyces cerevisiae</i> heat shock transcription factor (Hsf1) during the transient heat shock response. <i>J Biol Chem</i> 2006;281(43):32909–21.	doi:10.1074/jbc.M602454200 PMID 16926161 PMC2243236	–	1,366	3	1,366	–	1,366	1	Microarray analysis - wt vs TF mutant
34	Zheng 2010 Zheng W, Zhao H, Mancera E, Steinmetz LM, Snyder M. Genetic analysis of variation in transcription factor binding in yeast. <i>Nature</i> 2010;464(7292):1187–91.	doi:10.1038/nature08934 PMID 20237471 PMC2941147	GSE19634	1,330	1	1,330	1,330	–	3	ChIP-seq
35	Borneman 2007 Borneman AR, Zhang ZD, Rozowsky J, Seringhaus MR, Gerstein M, Snyder M. Transcription factor binding site identification in yeast: a comparison of high-density oligonucleotide and PCR-based microarray platforms. <i>Funct Integr Genomics</i> 2007;7(4):335–45.	doi:10.1007/s10142-007-0054-7 PMID 17638031	–	1,323	3	1,323	1,323	–	1	Chip-on-chip
36	Read 2016 Read T, Richmond PA, Dowell RD. A trans-acting Variant within the Transcription Factor RIM101 Interacts with Genetic Background to Determine its Regulatory Capacity. <i>PLoS Genet</i> 2016;12(1):e1005746.	doi:10.1371/journal.pgen.1005746 PMID 26751950 PMC4709078	–	1,271	1	1,271	–	1,271	2	RNA-seq - wt vs TF mutant
37	Vizoso-Vázquez 2012 Vizoso-Vázquez A, Lamas-Maceiras M, Becerra M, González-Siso MI, Rodríguez-Belmonte E, Cerdán ME. Ixr1p and the control of the <i>Saccharomyces cerevisiae</i> hypoxic response. <i>Appl Microbiol Biotechnol</i> 2012;94(1):173–84.	doi:10.1007/s00253-011-3785-2 PMID 22189861	GSE30046	1,257	1	1,257	–	1,257	3	Microarray analysis - wt vs TF mutant
38	Lai 2005 Lai LC, Kosorukoff AL, Burke PV, Kwast KE. Dynamical remodeling of the transcriptome during short-term anaerobiosis in <i>Saccharomyces cerevisiae</i> : differential response and role of Msn2 and/or Msn4 and other factors in galactose and glucose media. <i>Mol Cell Biol</i> 2005;25(10):4075–91.	doi:10.1128/MCB.25.10.4075-4091.2005 PMID 15870279 PMC1087712	GSE1879	1,228	2	1,228	–	1,228	5	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
39	Srinivasan 2020 Srinivasan R, Walvekar AS, Rashida Z, Seshasayee A, Laxman S. Genome-scale reconstruction of Gcn4/ATF4 networks driving a growth program. <i>PLoS Genet</i> 2020;16(12):e1009252.	doi:10.1371/journal.pgen.1009252 PMID 33378328 PMC7773203	–	1,220	1	1,111	308	912	1	RNA-seq - wt vs TF mutant
40	Huebert 2012 Huebert DJ, Kuan PF, Keleş S, Gasch AP. Dynamic changes in nucleosome occupancy are not predictive of gene expression dynamics but are linked to transcription and chromatin regulators. <i>Mol Cell Biol</i> 2012;32(9):1645–53.	doi:10.1128/MCB.06170-11 PMID 22354995 PMC3347246	–	1,203	1	1,203	1,203	–	1	Chip-on-chip
41	Rossouw 2012 Rossouw D, Jacobson D, Bauer FF. Transcriptional regulation and the diversification of metabolism in wine yeast strains. <i>Genetics</i> 2012;190(1):251–61.	doi:10.1534/genetics.111.132720 PMID 22042577 PMC3249374	–	1,126	1	1,126	–	1,126	1	Microarray analysis - wt vs TF overexpression
42	Chow 2019 Chow J, Starr I, Jamalzadeh S, et al., Cullen PJ. Filamentation Regulatory Pathways Control Adhesion-Dependent Surface Responses in Yeast. <i>Genetics</i> 2019;212(3):667–690.	doi:10.1534/genetics.119.302004 PMID 31053593 PMC6614897	–	1,011	2	1,001	–	1,011	2	RNA-seq - wt vs TF mutant
43	Tai 2017 Tai A, Kamei Y, Mukai Y. The forkhead-like transcription factor (Fhl1p) maintains yeast replicative lifespan by regulating ribonucleotide reductase 1 (RNR1) gene transcription. <i>Biochem Biophys Res Commun</i> 2017;488(1):218–223.	doi:10.1016/j.bbrc.2017.05.038 PMID 28495531	GSE95591	982	1	977	2	980	1	Microarray analysis - wt vs TF mutant
44	Lu 2005 Lu L, Roberts GG, Oszust C, Hudson AP. The YJR127C/ZMS1 gene product is involved in glycerol-based respiratory growth of the yeast <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 2005;48(4):235–46.	doi:10.1007/s00294-005-0023-4 PMID 16208474	GSE29530	960	1	957	–	960	3	Microarray analysis - wt vs TF mutant
45	Cromie 2017 Cromie GA, Tan Z, Hays M, Sirr A, Jeffery EW, Dudley AM. Transcriptional Profiling of Biofilm Regulators Identified by an Overexpression Screen in <i>Saccharomyces cerevisiae</i> . <i>G3 (Bethesda)</i> 2017;7(8):2845–2854.	doi:10.1534/g3.117.042440 PMID 28673928 PMC5555487	–	938	3	938	–	938	1	RNA-seq - wt vs TF overexpression
46	Zhu 2015 Zhu X, Keeney S. High-Resolution Global Analysis of the Influences of Bas1 and Ino4 Transcription Factors on Meiotic DNA Break Distributions in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2015;201(2):525–42.	doi:10.1534/genetics.115.178293 PMID 26245832 PMC4596667	–	936	2	936	234	702	1	RNA-seq - wt vs TF mutant
47	Lefrançois 2009 Lefrançois P, Euskirchen GM, Auerbach RK, et al., Snyder M. Efficient yeast ChIP-Seq using multiplex short-read DNA sequencing. <i>BMC Genomics</i> 2009;10:37.	doi:10.1186/1471-2164-10-37 PMID 19159457 PMC2656530	–	926	1	926	926	–	2	ChIP-seq

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
48	Miles 2013 Miles S, Li L, Davison J, Breeden LL. Xbp1 directs global repression of budding yeast transcription during the transition to quiescence and is important for the longevity and reversibility of the quiescent state. <i>PLoS Genet</i> 2013;9(10): e1003854.	doi:10.1371/journal.pgen.1003854 PMID 24204289 PMC3814307	–	853	1	853	7	846	6	RNA-seq - wt vs TF mutant
49	Eckert-Boulet 2004 Eckert-Boulet N, Nielsen PS, Friis C, et al., Regenberg B. Transcriptional profiling of extracellular amino acid sensing in <i>Saccharomyces cerevisiae</i> and the role of Stp1p and Stp2p. <i>Yeast</i> 2004;21(8):635–48.	doi:10.1002/yea.1120 PMID 15197729	–	840	2	569	–	840	2	Microarray analysis - wt vs TF mutant
50	Chumnanpuen 2013 Chumnanpuen P, Nookaew I, Nielsen J. Integrated analysis, transcriptome-lipidome, reveals the effects of INO-level (INO2 and INO4) on lipid metabolism in yeast. <i>BMC Syst Biol</i> 2013; 7 Suppl 3(Suppl 3):S7.	doi:10.1186/1752-0509-7-S3-S7 PMID 24456840 PMC3852131	GSE36298	837	3	837	–	837	2	Microarray analysis - wt vs TF mutant
51	Walvekar 2018 Walvekar AS, Srinivasan R, Gupta R, Laxman S. Methionine coordinates a hierarchically organized anabolic program enabling proliferation. <i>Mol Biol Cell</i> 2018;29(26):3183–3200.	doi:10.1091/mbc.E18-08-0515 PMID 30354837 PMC6340205	–	835	1	835	–	835	1	RNA-seq - wt vs TF mutant
52	Weidberg 2018 Weidberg H, Amon A. MitoCPR-A surveillance pathway that protects mitochondria in response to protein import stress. <i>Science</i> 2018;360(6385).	doi:10.1126/science.aan4146 PMID 29650645 PMC6528467	–	805	1	803	–	805	3	RNA-seq - wt vs TF mutant
53	Lardenois 2015 Lardenois A, Becker E, Walther T, et al., Primig M. Global alterations of the transcriptional landscape during yeast growth and development in the absence of Ume6-dependent chromatin modification. <i>Mol Genet Genomics</i> 2015;290(5): 2031–46.	doi:10.1007/s00438-015-1051-5 PMID 25957495 PMC5469029	–	788	1	788	3	785	6	Microarray analysis - wt vs TF mutant
54	Marion 2004 Marion RM, Regev A, Segal E, et al., O’Shea EK. Sfp1 is a stress- and nutrient-sensitive regulator of ribosomal protein gene expression. <i>Proc Natl Acad Sci U S A</i> 2004;101(40): 14315–22.	doi:10.1073/pnas.0405353101 PMID 15353587 PMC521938	–	766	1	766	92	674	5	Microarray analysis - wt vs TF mutant
55	Ouyang 2011 Ouyang X, Tran QT, Goodwin S, Wible RS, Sutter CH, Sutter TR. Yap1 activation by H2O2 or thiol-reactive chemicals elicits distinct adaptive gene responses. <i>Free Radic Biol Med</i> 2011;50(1):1–13.	doi:10.1016/j.freeradbiomed.2010.10.697 PMID 20971184	GSE19213	741	1	411	–	741	3	Microarray analysis - wt vs TF mutant
56	Kim 2019 Kim JE, Jang IS, Son SH, et al., Lee JY. Tailoring the <i>Saccharomyces cerevisiae</i> endoplasmic reticulum for functional assembly of terpene synthesis pathway. <i>Metab Eng</i> 2019;56: 50–59.	doi:10.1016/j.ymben.2019.08.013 PMID 31445083	–	739	1	739	–	739	1	RNA-seq - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
57	Albert 2019 Albert B, Tomassetti S, Gloor Y, et al., Shore D. Sfp1 regulates transcriptional networks driving cell growth and division through multiple promoter-binding modes. <i>Genes Dev</i> 2019;33(5–6):288–293.	doi:10.1101/gad.322040.118 PMID 30804227 PMC6411004	–	713	1	311	713	–	3	ChIP-seq
58	Cipollina 2008 Cipollina C, van den Brink J, Daran-Lapujade P, Pronk JT, Porro D, de Winde JH. <i>Saccharomyces cerevisiae</i> SFP1: at the crossroads of central metabolism and ribosome biogenesis. <i>Microbiology (Reading)</i> 2008;154(Pt 6):1686–1699.	doi:10.1099/mic.0.2008/017392-0 PMID 18524923	GSE9644	696	1	696	–	696	1	Microarray analysis - wt vs TF mutant
59	Jorgensen 2004 Jorgensen P, Rupes I, Sharom JR, Schnepel L, Broach JR, Tyers M. A dynamic transcriptional network communicates growth potential to ribosome synthesis and critical cell size. <i>Genes Dev</i> 2004;18(20):2491–505.	doi:10.1101/gad.1228804 PMID 15466158 PMC529537	–	656	2	643	–	656	6	Microarray analysis - wt vs TF mutant
60	Gallagher 2014 Gallagher JE, Zheng W, Rong X, et al., Snyder MP. Divergence in a master variator generates distinct phenotypes and transcriptional responses. <i>Genes Dev</i> 2014;28(4):409–21.	doi:10.1101/gad.228940.113 PMID 24532717 PMC3937518	–	644	1	644	644	–	3	ChIP-seq
61	Lahav 2007 Lahav R, Gammie A, Tavazoie S, Rose MD. Role of transcription factor Kar4 in regulating downstream events in the <i>Saccharomyces cerevisiae</i> pheromone response pathway. <i>Mol Cell Biol</i> 2007;27(3):818–29.	doi:10.1128/MCB.00439-06 PMID 17101777 PMC1800688	–	609	2	609	2	607	4	Microarray analysis - wt vs TF mutant
62	Hazelwood 2009 Hazelwood LA, Walsh MC, Luttik MA, Daran-Lapujade P, Pronk JT, Daran JM. Identity of the growth-limiting nutrient strongly affects storage carbohydrate accumulation in anaerobic chemostat cultures of <i>Saccharomyces cerevisiae</i> . <i>Appl Environ Microbiol</i> 2009;75(21):6876–85.	doi:10.1128/AEM.01464-09 PMID 19734328 PMC2772451	–	598	2	540	–	598	2	Microarray analysis - wt vs TF mutant
63	McIsaac 2013 McIsaac RS, Oakes BL, Wang X, Dummit KA, Botstein D, Noyes MB. Synthetic gene expression perturbation systems with rapid, tunable, single-gene specificity in yeast. <i>Nucleic Acids Res</i> 2013;41(4):e57.	doi:10.1093/nar/gks1313 PMID 23275543 PMC3575806	–	581	1	581	–	581	1	Microarray analysis - wt vs TF overexpression
64	Yoshimoto 2002 Yoshimoto H, Saltsman K, Gasch AP, et al., Cyert MS. Genome-wide analysis of gene expression regulated by the calcineurin/Crz1p signaling pathway in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2002;277(34):31079–88.	doi:10.1074/jbc.M202718200 PMID 12058033	no GEO	575	1	350	–	575	4	Microarray analysis - wt vs TF mutant
65	Banerjee 2008 Banerjee D, Lelandais G, Shukla S, et al., Prasad R. Responses of pathogenic and nonpathogenic yeast species to steroids reveal the functioning and evolution of multidrug resistance transcriptional networks. <i>Eukaryot Cell</i> 2008;7(1):68–77.	doi:10.1128/EC.00256-07 PMID 17993571 PMC2224153	–	572	2	572	–	572	2	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
66	Rudra 2005 Rudra D, Zhao Y, Warner JR. Central role of Ifh1p-Fhl1p interaction in the synthesis of yeast ribosomal proteins. <i>EMBO J</i> 2005;24(3):533–42.	doi:10.1038/sj.emboj.7600553 PMID 15692568 PMC548658	no GEO	566	2	566	14	552	3	Microarray analysis - wt vs TF mutant
67	Berry 2008 Berry DB, Gasch AP. Stress-activated genomic expression changes serve a preparative role for impending stress in yeast. <i>Mol Biol Cell</i> 2008;19(11):4580–7.	doi:10.1091/mbc.e07-07-0680 PMID 18753408 PMC2575158	GSE8335	564	2	507	–	564	3	Microarray analysis - wt vs TF mutant
68	Barbara 2007 Barbara KE, Haley TM, Willis KA, Santangelo GM. The transcription factor Gcr1 stimulates cell growth by participating in nutrient-responsive gene expression on a global level. <i>Mol Genet Genomics</i> 2007;277(2):171–88.	doi:10.1007/s00438-006-0182-0 PMID 17124610	GSE5027	555	1	291	–	555	4	Microarray analysis - wt vs TF mutant
69	Smith 2007 Smith JJ, Ramsey SA, Marelli M, et al., Aitchison JD. Transcriptional responses to fatty acid are coordinated by combinatorial control. <i>Mol Syst Biol</i> 2007;3:115.	doi:10.1038/msb4100157 PMID 17551510 PMC1911199	GSE5862	546	4	546	–	546	1	Microarray analysis - wt vs TF mutant
70	Iyer 2001 Iyer VR, Horak CE, Scafe CS, Botstein D, Snyder M, Brown PO. Genomic binding sites of the yeast cell-cycle transcription factors SBF and MBF. <i>Nature</i> 2001;409(6819):533–8.	doi:10.1038/35054095 PMID 11206552	–	532	2	353	532	–	3	Chip-on-chip
71	Pfanzagl 2018 Pfanzagl V, Görner W, Radolf M, et al., Schüller C. A constitutive active allele of the transcription factor Msn2 mimicking low PKA activity dictates metabolic remodeling in yeast. <i>Mol Biol Cell</i> 2018;29(23):2848–2862.	doi:10.1091/mbc.E18-06-0389 PMID 30256697 PMC6249869	–	530	1	530	–	530	1	Microarray analysis - wt vs TF mutant
72	Knott 2012 Knott SR, Peace JM, Ostrow AZ, et al., Aparicio OM. Forkhead transcription factors establish origin timing and long-range clustering in <i>S. cerevisiae</i> . <i>Cell</i> 2012;148(1–2):99–111.	doi:10.1016/j.cell.2011.12.012 PMID 22265405 PMC3266545	–	513	1	513	467	46	2	ChIP-seq
73	Kruegel 2011 Kruegel U, Robison B, Dange T, et al., Schmidt M. Elevated proteasome capacity extends replicative lifespan in <i>Saccharomyces cerevisiae</i> . <i>PLoS Genet</i> 2011;7(9):e1002253.	doi:10.1371/journal.pgen.1002253 PMID 21931558 PMC3169524	–	500	1	500	–	500	1	Proteomics - wt vs TF mutant
74	Larochelle 2006 Larochelle M, Drouin S, Robert F, Turcotte B. Oxidative stress-activated zinc cluster protein Stb5 has dual activator/repressor functions required for pentose phosphate pathway regulation and NADPH production. <i>Mol Cell Biol</i> 2006;26(17):6690–701.	doi:10.1128/MCB.02450-05 PMID 16914749 PMC1592823	–	476	1	476	–	476	1	Microarray analysis - wt vs TF mutant
75	Berthelet 2010 Berthelet S, Usher J, Shulist K, et al., Baetz K. Functional genomics analysis of the <i>Saccharomyces cerevisiae</i> iron responsive transcription factor Aft1 reveals iron-independent functions. <i>Genetics</i> 2010;185(3):1111–28.	doi:10.1534/genetics.110.117531 PMID 20439772 PMC2900968	GSE20531	471	1	471	1	470	2	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
76	Tan 2008 Tan K, Feizi H, Luo C, Fan SH, Ravasi T, Ideker TG. A systems approach to delineate functions of paralogous transcription factors: role of the Yap family in the DNA damage response. <i>Proc Natl Acad Sci U S A</i> 2008;105(8):2934–9.	doi:10.1073/pnas.0708670105 PMID 18287073 PMC2268563	–	457	5	457	457	–	2	Chip-on-chip
77	Fendt 2010 Fendt SM, Buescher JM, Rudroff F, Picotti P, Zamboni N, Sauer U. Tradeoff between enzyme and metabolite efficiency maintains metabolic homeostasis upon perturbations in enzyme capacity. <i>Mol Syst Biol</i> 2010;6:356.	doi:10.1038/msb.2010.11 PMID 20393576 PMC2872607	–	453	1	453	–	453	1	Microarray analysis - wt vs TF mutant
78	Ouyang 2018 Ouyang L, Holland P, Lu H, Bergenholm D, Nielsen J. Integrated analysis of the yeast NADPH-regulator Stb5 reveals distinct differences in NADPH requirements and regulation in different states of yeast metabolism. <i>FEMS Yeast Res</i> 2018;18(8).	doi:10.1093/femsyr/foy091 PMID 30107458	–	438	1	185	382	56	4	ChIP-exo
79	Zeitlinger 2003 Zeitlinger J, Simon I, Harbison CT, et al., Young RA. Program-specific distribution of a transcription factor dependent on partner transcription factor and MAPK signaling. <i>Cell</i> 2003;113(3):395–404.	doi:10.1016/s0092-8674(03)00301-5 PMID 12732146	–	407	3	407	407	–	4	Chip-on-chip
80	Thorsen 2007 Thorsen M, Lagniel G, Kristiansson E, et al., Tamás MJ. Quantitative transcriptome, proteome, and sulfur metabolite profiling of the <i>Saccharomyces cerevisiae</i> response to arsenite. <i>Physiol Genomics</i> 2007;30(1):35–43.	doi:10.1152/physiolgenomics.00236.2006 PMID 17327492	GSE6129	405	2	389	–	405	2	Microarray analysis - wt vs TF mutant
81	Carrillo 2012 Carrillo E, Ben-Ari G, Wildenhain J, Tyers M, Grammentz D, Lee TA. Characterizing the roles of Met31 and Met32 in coordinating Met4-activated transcription in the absence of Met30. <i>Mol Biol Cell</i> 2012;23(10):1928–42.	doi:10.1091/mbc.E11-06-0532 PMID 22438580 PMC3350556	–	404	2	362	363	41	1	Chip-on-chip
82	Goossens 2015 Goossens KV, Ielasi FS, Nookaew I, et al., Willaert RG. Molecular mechanism of flocculation self-recognition in yeast and its role in mating and survival. <i>mBio</i> 2015;6(2).	doi:10.1128/mBio.00427-15 PMID 25873380 PMC4453552	GSE64468	403	1	403	–	403	2	Microarray analysis - wt vs TF mutant
83	Pagani 2007 Pagani MA, Casamayor A, Serrano R, Atrian S, Ariño J. Disruption of iron homeostasis in <i>Saccharomyces cerevisiae</i> by high zinc levels: a genome-wide study. <i>Mol Microbiol</i> 2007; 65(2):521–37.	doi:10.1111/j.1365-2958.2007.05807.x PMID 17630978	–	397	1	396	–	397	2	Microarray analysis - wt vs TF mutant
84	Jaehnig 2013 Jaehnig EJ, Kuo D, Hombauer H, Ideker TG, Kolodner RD. Checkpoint kinases regulate a global network of transcription factors in response to DNA damage. <i>Cell Rep</i> 2013;4(1):174–88.	doi:10.1016/j.celrep.2013.05.041 PMID 23810556 PMC3855057	GSE40351	396	5	396	–	396	1	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
85	Schawalter 2004 Schawalter SB, Kabani M, Howald I, Choudhury U, Werner M, Shore D. Growth-regulated recruitment of the essential yeast ribosomal protein gene activator Ifh1. <i>Nature</i> 2004; 432(7020):1058–61.	doi:10.1038/nature03200 PMID 15616569	no GEO	393	3	132	4	389	6	Microarray analysis - wt vs TF overexpression
86	Raghevedran 2006 Raghevedran V, Patil KR, Olsson L, Nielsen J. Hap4 is not essential for activation of respiration at low specific growth rates in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2006;281(18): 12308–14.	doi:10.1074/jbc.M512972200 PMID 16522629	–	385	1	385	–	385	1	Microarray analysis - wt vs TF mutant
87	Lieb 2001 Lieb JD, Liu X, Botstein D, Brown PO. Promoter-specific binding of Rap1 revealed by genome-wide maps of protein-DNA association. <i>Nat Genet</i> 2001;28(4):327–34.	doi:10.1038/ng569 PMID 11455386	–	374	1	374	374	–	1	Chip-on-chip
88	Kelley 2009 Kelley R, Ideker T. Genome-wide fitness and expression profiling implicate Mga2 in adaptation to hydrogen peroxide. <i>PLoS Genet</i> 2009;5(5):e1000488.	doi:10.1371/journal.pgen.1000488 PMID 19503593 PMC2676504	–	373	5	309	–	373	2	Microarray analysis - wt vs TF mutant
89	Kleinschmidt 2005 Kleinschmidt M, Grundmann O, Blüthgen N, Mösch HU, Braus GH. Transcriptional profiling of <i>Saccharomyces cerevisiae</i> cells under adhesion-inducing conditions. <i>Mol Genet Genomics</i> 2005;273(5):382–93.	doi:10.1007/s00438-005-1139-4 PMID 15843968	–	358	1	358	–	358	2	Microarray analysis - wt vs TF mutant
90	Ernst 2007 Ernst J, Vainas O, Harbison CT, Simon I, Bar-Joseph Z. Reconstructing dynamic regulatory maps. <i>Mol Syst Biol</i> 2007; 3:74.	doi:10.1038/msb4100115 PMID 17224918 PMC1800355	–	357	2	357	357	–	5	Chip-on-chip
91	Casado 2011 Casado C, González A, Platara M, Ruiz A, Ariño J. The role of the protein kinase A pathway in the response to alkaline pH stress in yeast. <i>Biochem J</i> 2011;438(3):523–33.	doi:10.1042/BJ20110607 PMID 21749328 PMC3253439	GSE27925	356	2	356	–	356	2	Microarray analysis - wt vs TF mutant
92	Gertz 2005 Gertz J, Riles L, Turnbaugh P, Ho SW, Cohen BA. Discovery, validation, and genetic dissection of transcription factor binding sites by comparative and functional genomics. <i>Genome Res</i> 2005;15(8):1145–52.	doi:10.1101/gr.3859605 PMID 16077013 PMC1182227	–	351	2	351	1	350	3	Microarray analysis - wt vs TF mutant
93	Börlin 2020 Börlin CS, Nielsen J, Siewers V. The transcription factor Leu3 shows differential binding behavior in response to changing leucine availability. <i>FEMS Microbiol Lett</i> 2020;367(13).	doi:10.1093/femsle/fnaa107 PMID 32589214 PMC7351879	–	339	1	107	339	–	5	ChIP-seq
94	James Theoga Raj 2019 James Theoga Raj C, Croft T, Venkatakrishnan P, et al., Lin SJ. The copper-sensing transcription factor Mac1, the histone deacetylase Hst1, and nicotinic acid regulate de novo NAD(+) biosynthesis in budding yeast. <i>J Biol Chem</i> 2019;294(14): 5562–5575.	doi:10.1074/jbc.RA118.006987 PMID 30760525 PMC6462523	–	327	1	316	1	326	1	RNA-seq - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
95	Reja 2015 Reja R, Vinayachandran V, Ghosh S, Pugh BF. Molecular mechanisms of ribosomal protein gene coregulation. <i>Genes Dev</i> 2015;29(18):1942–54.	doi:10.1101/gad.268896.115 PMID 26385964 PMC4579351	–	312	4	312	312	–	1	ChIP-exo
96	Yarragudi 2007 Yarragudi A, Parfrey LW, Morse RH. Genome-wide analysis of transcriptional dependence and probable target sites for Abf1 and Rap1 in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Res</i> 2007;35(1):193–202.	doi:10.1093/nar/gkl1059 PMID 17158163 PMC1802568	GSE6073	309	2	309	–	309	1	Microarray analysis - wt vs TF mutant
97	Cipollina 2008 Cipollina C, van den Brink J, Daran-Lapujade P, Pronk JT, Vai M, de Winde JH. Revisiting the role of yeast Sfp1 in ribosome biogenesis and cell size control: a chemostat study. <i>Microbiology (Reading)</i> 2008;154(Pt 1):337–346.	doi:10.1099/mic.0.2007/011767-0 PMID 18174152	GSE5238	299	1	279	–	299	2	Microarray analysis - wt vs TF mutant
98	Schüller 2007 Schüller C, Mamnun YM, Wolfger H, Rockwell N, Thorner J, Kuchler K. Membrane-active compounds activate the transcription factors Pdr1 and Pdr3 connecting pleiotropic drug resistance and membrane lipid homeostasis in <i>saccharomyces cerevisiae</i> . <i>Mol Biol Cell</i> 2007;18(12):4932–44.	doi:10.1091/mbc.e07-06-0610 PMID 17881724 PMC2096591	–	297	4	297	–	297	3	Microarray analysis - wt vs TF mutant
99	Rawal 2018 Rawal Y, Chereji RV, Valabhoju V, et al., Hinnebusch AG. Gcn4 Binding in Coding Regions Can Activate Internal and Canonical 5' Promoters in Yeast. <i>Mol Cell</i> 2018;70(2):297–311.e4.	doi:10.1016/j.molcel.2018.03.007 PMID 29628310 PMC6133248	–	279	1	279	279	–	1	ChIP-seq
100	Sanz 2012 Sanz AB, García R, Rodríguez-Peña JM, et al., Arroyo J. Chromatin remodeling by the SWI/SNF complex is essential for transcription mediated by the yeast cell wall integrity MAPK pathway. <i>Mol Biol Cell</i> 2012;23(14):2805–17.	doi:10.1091/mbc.E12-04-0278 PMID 22621902 PMC3395667	GSE31176	272	1	272	1	271	2	Microarray analysis - wt vs TF mutant
101	Judeh 2013 Judeh T, Johnson C, Kumar A, Zhu D. TEAK: topology enrichment analysis framework for detecting activated biological subpathways. <i>Nucleic Acids Res</i> 2013;41(3):1425–37.	doi:10.1093/nar/gks1299 PMID 23268448 PMC3561980	GSE40530	270	2	270	–	270	1	Microarray analysis - wt vs TF mutant
102	Pimentel 2012 Pimentel C, Vicente C, Menezes RA, Caetano S, Carreto L, Rodrigues-Pousada C. The role of the Yap5 transcription factor in remodeling gene expression in response to Fe bioavailability. <i>PLoS One</i> 2012;7(5):e37434.	doi:10.1371/journal.pone.0037434 PMID 22616008 PMC3353947	–	259	1	241	1	258	10	Microarray analysis - wt vs TF mutant
103	Simon 2001 Simon I, Barnett J, Hannett N, et al., Young RA. Serial regulation of transcriptional regulators in the yeast cell cycle. <i>Cell</i> 2001;106(6):697–708.	doi:10.1016/s0092-8674(01)00494-9 PMID 11572776	–	252	8	252	252	–	2	Chip-on-chip

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
104	Tang 2006 Tang L, Liu X, Clarke ND. Inferring direct regulatory targets from expression and genome location analyses: a comparison of transcription factor deletion and overexpression. <i>BMC Genomics</i> 2006;7:215.	doi:10.1186/1471-2164-7-215 PMID 16923194 PMC1559704	–	250	1	250	–	250	1	Microarray analysis - wt vs TF overexpression
105	Tuch 2008 Tuch BB, Galgoczy DJ, Hernday AD, Li H, Johnson AD. The evolution of combinatorial gene regulation in fungi. <i>PLoS Biol</i> 2008;6(2):e38.	doi:10.1371/journal.pbio.0060038 PMID 18303948 PMC2253631	–	250	1	250	250	–	2	Chip-on-chip
106	Hanlon 2011 Hanlon SE, Rizzo JM, Tatomer DC, Lieb JD, Buck MJ. The stress response factors Yap6, Cin5, Phd1, and Skn7 direct targeting of the conserved co-repressor Tup1-Ssn6 in <i>S. cerevisiae</i> . <i>PLoS One</i> 2011;6(4):e19060.	doi:10.1371/journal.pone.0019060 PMID 21552514 PMC3084262	–	249	1	249	249	–	1	Chip-on-chip
107	Wade 2004 Wade JT, Hall DB, Struhl K. The transcription factor Ifh1 is a key regulator of yeast ribosomal protein genes. <i>Nature</i> 2004; 432(7020):1054–8.	doi:10.1038/nature03175 PMID 15616568	–	248	4	248	239	9	4	Chip-on-chip
108	Chavel 2010 Chavel CA, Dionne HM, Birkaya B, Joshi J, Cullen PJ. Multiple signals converge on a differentiation MAPK pathway. <i>PLoS Genet</i> 2010;6(3):e1000883.	doi:10.1371/journal.pgen.1000883 PMID 20333241 PMC2841618	–	246	16	241	1	245	20	Microarray analysis - wt vs TF mutant
109	Nikolov 2022 Nikolov VN, Malavia D, Kubota T. SWI/SNF and the histone chaperone Rtt106 drive expression of the Pleiotropic Drug Resistance network genes. <i>Nat Commun</i> 2022;13(1):1968.	doi:10.1038/s41467-022-29591-z PMID 35413952 PMC9005695	–	231	2	231	–	231	1	RNA-seq - wt vs TF mutant
110	Tachibana 2005 Tachibana C, Yoo JY, Tagne JB, Kacherovsky N, Lee TI, Young ET. Combined global localization analysis and transcriptome data identify genes that are directly coregulated by Adr1 and Cat8. <i>Mol Cell Biol</i> 2005;25(6):2138–46.	doi:10.1128/MCB.25.6.2138-2146.2005 PMID 15743812 PMC1061606	–	226	2	226	222	4	1	ChIP
111	Váchová 2004 Váchová L, Devaux F, KucEROVÁ H, RíCICOVÁ M, Jacq C, PalkOVÁ Z. Sok2p transcription factor is involved in adaptive program relevant for long term survival of <i>Saccharomyces cerevisiae</i> colonies. <i>J Biol Chem</i> 2004;279(36):37973–81.	doi:10.1074/jbc.M404594200 PMID 15229222	–	224	1	110	–	224	13	Microarray analysis - wt vs TF mutant
112	Hickman 2011 Hickman MJ, Petti AA, Ho-Shing O, et al., Botstein D. Coordinated regulation of sulfur and phospholipid metabolism reflects the importance of methylation in the growth of yeast. <i>Mol Biol Cell</i> 2011;22(21):4192–204.	doi:10.1091/mbc.E11-05-0467 PMID 21900497 PMC3204079	–	224	2	224	–	224	1	Microarray analysis - wt vs TF mutant
113	Loll-Krippelbeier 2017 Loll-Krippelbeier R, Brown GW. P-body proteins regulate transcriptional rewiring to promote DNA replication stress resistance. <i>Nat Commun</i> 2017;8(1):558.	doi:10.1038/s41467-017-00632-2 PMID 28916784 PMC5601920	–	224	2	224	–	224	1	RNA-seq - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
114	Vyas 2005 Vyas VK, Berkey CD, Miyao T, Carlson M. Repressors Nrg1 and Nrg2 regulate a set of stress-responsive genes in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2005;4(11):1882–91.	doi:10.1128/EC.4.11.1882-1891.2005 PMID 16278455 PMID 22379133 PMC1287862	–	216	2	216	–	216	1	Microarray analysis - wt vs TF mutant
115	Servant 2012 Servant G, Pinson B, Tchalikian-Cosson A, et al., Lesage P. Tye7 regulates yeast Ty1 retrotransposon sense and antisense transcription in response to adenylic nucleotides stress. <i>Nucleic Acids Res</i> 2012;40(12):5271–82.	doi:10.1093/nar/gks166 PMID 22379133 PMC3384299	–	209	2	209	1	208	2	Microarray analysis - wt vs TF mutant
116	Westholm 2008 Westholm JO, Nordberg N, Murén E, Ameer A, Komorowski J, Ronne H. Combinatorial control of gene expression by the three yeast repressors Mig1, Mig2 and Mig3. <i>BMC Genomics</i> 2008;9:601.	doi:10.1186/1471-2164-9-601 PMID 19087243 PMC2631581	–	206	3	206	–	206	1	Microarray analysis - wt vs TF mutant
117	Liang 2013 Liang CY, Wang LC, Lo WS. Dissociation of the H3K36 demethylase Rph1 from chromatin mediates derepression of environmental stress-response genes under genotoxic stress in <i>Saccharomyces cerevisiae</i> . <i>Mol Biol Cell</i> 2013;24(20):3251–62.	doi:10.1091/mbc.E12-11-0820 PMID 23985319 PMC3806659	GSE39950	206	1	206	–	206	1	Microarray analysis - wt vs TF mutant
118	Roque 2016 Roque A, Petrezsélyová S, Serra-Cardona A, Ariño J. Genome-wide recruitment profiling of transcription factor Crz1 in response to high pH stress. <i>BMC Genomics</i> 2016;17:662.	doi:10.1186/s12864-016-3006-6 PMID 27544903 PMC4992276	–	203	1	203	203	–	1	ChIP-seq
119	Maity 2016 Maity S, Rajkumar A, Matai L, et al., Chakraborty K. Oxidative Homeostasis Regulates the Response to Reductive Endoplasmic Reticulum Stress through Translation Control. <i>Cell Rep</i> 2016;16(3):851–65.	doi:10.1016/j.celrep.2016.06.025 PMID 27373166	–	195	1	195	–	195	1	iTRAQ-MS - wt vs TF mutant
120	Elfving 2014 Elfving N, Chereji RV, Bharatula V, Björklund S, Morozov AV, Broach JR. A dynamic interplay of nucleosome and Msn2 binding regulates kinetics of gene activation and repression following stress. <i>Nucleic Acids Res</i> 2014;42(9):5468–82.	doi:10.1093/nar/gku176 PMID 24598258 PMC4027177	–	193	1	193	62	131	2	Microarray analysis - wt vs TF overexpression
121	Pincus 2018 Pincus D, Anandhakumar J, Thiru P, Guertin MJ, Erkin AM, Gross DS. Genetic and epigenetic determinants establish a continuum of Hsf1 occupancy and activity across the yeast genome. <i>Mol Biol Cell</i> 2018;29(26):3168–3182.	doi:10.1091/mbc.E18-06-0353 PMID 30332327 PMC6340206	–	189	1	70	129	60	5	ChIP-seq
122	Onda 2004 Onda M, Ota K, Chiba T, Sakaki Y, Ito T. Analysis of gene network regulating yeast multidrug resistance by artificial activation of transcription factors: involvement of Pdr3 in salt tolerance. <i>Gene</i> 2004;332:51–9.	doi:10.1016/j.gene.2004.02.003 PMID 15145054	–	185	8	185	–	185	2	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
123	Soontorngun 2012 Soontorngun N, Baramée S, Tangsombatvichit C, et al., Turcotte B. Genome-wide location analysis reveals an important overlap between the targets of the yeast transcriptional regulators Rds2 and Adr1. <i>Biochem Biophys Res Commun</i> 2012;423(4):632–7.	doi:10.1016/j.bbrc.2012.05.151 PMID 22687600	–	185	2	185	181	4	2	Chip-on-chip
124	Liang 2021 Liang L, Liu R, Freed EF, Eckert CA, Gill RT. Transcriptional Regulatory Networks Involved in C3-C4 Alcohol Stress Response and Tolerance in Yeast. <i>ACS Synth Biol</i> 2021;10(1):19–28.	doi:10.1021/acssynbio.0c00253 PMID 33356165	–	179	4	176	–	179	2	RNA-seq - wt vs TF mutant
125	Zhang 2010 Zhang N, Oliver SG. The transcription activity of Gis1 is negatively modulated by proteasome-mediated limited proteolysis. <i>J Biol Chem</i> 2010;285(9):6465–76.	doi:10.1074/jbc.M109.073288 PMID 20022953 PMC2825442	–	176	1	88	–	176	2	Microarray analysis - wt vs TF overexpression
126	García 2004 García R, Bermejo C, Grau C, et al., Arroyo J. The global transcriptional response to transient cell wall damage in <i>Saccharomyces cerevisiae</i> and its regulation by the cell integrity signaling pathway. <i>J Biol Chem</i> 2004;279(15):15183–95.	doi:10.1074/jbc.M312954200 PMID 14739279	GSE961	174	1	174	–	174	2	Microarray analysis - wt vs TF mutant
127	Nishizawa 2008 Nishizawa M, Komai T, Katou Y, Shirahige K, Ito T, Toh-E A. Nutrient-regulated antisense and intragenic RNAs modulate a signal transduction pathway in yeast. <i>PLoS Biol</i> 2008;6(12):2817–30.	doi:10.1371/journal.pbio.0060326 PMID 19108609 PMC2605928	–	163	2	163	161	2	3	Chip-on-chip
128	Sasaki 2005 Sasaki H, Uemura H. Influence of low glycolytic activities in <i>gcr1</i> and <i>gcr2</i> mutants on the expression of other metabolic pathway genes in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 2005;22(2):111–27.	doi:10.1002/yea.1198 PMID 15645478	–	155	2	118	–	155	2	Microarray analysis - wt vs TF mutant
129	da Silva 2018 da Silva SM, Batista-Nascimento L, Gaspar-Cordeiro A, Vernis L, Pimentel C, Rodrigues-Pousada C. Transcriptional regulation of FeS biogenesis genes: A possible shield against arsenate toxicity activated by Yap1. <i>Biochim Biophys Acta Gen Subj</i> 2018;1862(10):2152–2161.	doi:10.1016/j.bbagen.2018.07.013 PMID 30025855	–	155	2	146	–	155	3	Microarray analysis - wt vs TF mutant
130	Natarajan 2001 Natarajan K, Meyer MR, Jackson BM, et al., Marton MJ. Transcriptional profiling shows that Gcn4p is a master regulator of gene expression during amino acid starvation in yeast. <i>Mol Cell Biol</i> 2001;21(13):4347–68.	doi:10.1128/MCB.21.13.4347-4368.2001 PMID 11390663 PMC87095	–	154	1	154	–	154	2	Microarray analysis - wt vs TF mutant
131	Mittal 2017 Mittal N, Guimaraes JC, Gross T, et al., Zavolan M. The Gcn4 transcription factor reduces protein synthesis capacity and extends yeast lifespan. <i>Nat Commun</i> 2017;8(1):457.	doi:10.1038/s41467-017-00539-y PMID 28878244 PMC5587724	–	150	1	150	150	–	1	ChIP-seq

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
132	Jorgensen 2002 Jorgensen P, Nishikawa JL, Breitkreutz BJ, Tyers M. Systematic identification of pathways that couple cell growth and division in yeast. <i>Science</i> 2002;297(5580):395–400.	doi:10.1126/science.1070850 PMID 12089449	–	146	1	103	–	146	6	Microarray analysis - wt vs TF overexpression
133	Williams 2002 Williams RM, Primig M, Washburn BK, et al., Esposito RE. The Ume6 regulon coordinates metabolic and meiotic gene expression in yeast. <i>Proc Natl Acad Sci U S A</i> 2002;99(21):13431–6.	doi:10.1073/pnas.202495299 PMID 12370439 PMC129690	no GEO	140	1	82	–	140	2	Microarray analysis - wt vs TF mutant
134	Hasan 2002 Hasan R, Leroy C, Isnard AD, Labarre J, Boy-Marcotte E, Toledano MB. The control of the yeast H2O2 response by the Msn2/4 transcription factors. <i>Mol Microbiol</i> 2002;45(1):233–41.	doi:10.1046/j.1365-2958.2002.03011.x PMID 12100562	–	138	3	102	–	138	1	Proteomics - wt vs TF mutant
135	Boer 2005 Boer VM, Daran JM, Almering MJ, de Winde JH, Pronk JT. Contribution of the <i>Saccharomyces cerevisiae</i> transcriptional regulator Leu3p to physiology and gene expression in nitrogen- and carbon-limited chemostat cultures. <i>FEMS Yeast Res</i> 2005;5(10):885–97.	doi:10.1016/j.femsyr.2005.04.003 PMID 15949974	no GEO	137	1	135	–	137	2	Microarray analysis - wt vs TF mutant
136	Bosio 2017 Bosio MC, Fermi B, Spagnoli G, et al., Dieci G. Abf1 and other general regulatory factors control ribosome biogenesis gene expression in budding yeast. <i>Nucleic Acids Res</i> 2017;45(8):4493–4506.	doi:10.1093/nar/gkx058 PMID 28158860 PMC5416754	–	134	3	122	134	–	2	ChIP-seq
137	Mascarenhas 2008 Mascarenhas C, Edwards-Ingram LC, Zeef L, Shenton D, Ashe MP, Grant CM. Gcn4 is required for the response to peroxide stress in the yeast <i>Saccharomyces cerevisiae</i> . <i>Mol Biol Cell</i> 2008;19(7):2995–3007.	doi:10.1091/mbc.e07-11-1173 PMID 18417611 PMC2441660	–	131	1	70	–	131	3	Microarray analysis - wt vs TF mutant
138	Bro 2005 Bro C, Knudsen S, Regenber B, Olsson L, Nielsen J. Improvement of galactose uptake in <i>Saccharomyces cerevisiae</i> through overexpression of phosphoglucomutase: example of transcript analysis as a tool in inverse metabolic engineering. <i>Appl Environ Microbiol</i> 2005;71(11):6465–72.	doi:10.1128/AEM.71.11.6465-6472.2005 PMID 16269670 PMC1287681	–	128	4	128	–	128	1	Microarray analysis - wt vs TF overexpression
139	Nawrocki 2001 Nawrocki A, Fey SJ, Goffeau A, Roepstorff P, Larsen PM. The effects of transcription regulating genes PDR1, pdr1-3 and PDR3 in pleiotropic drug resistance. <i>Proteomics</i> 2001;1(8):1022–32.	<a href="https://doi.org/10.1002/1615-9861(200108)1:8<1022::AID-PROT1022>3.0.CO;2-7">doi:10.1002/1615-9861(200108)1:8<1022::AID-PROT1022>3.0.CO;2-7 PMID 11683503	–	125	2	72	–	125	1	Proteomics - wt vs TF mutant
140	Hickman 2007 Hickman MJ, Winston F. Heme levels switch the function of Hap1 of <i>Saccharomyces cerevisiae</i> between transcriptional activator and transcriptional repressor. <i>Mol Cell Biol</i> 2007;27(21):7414–24.	doi:10.1128/MCB.00887-07 PMID 17785431 PMC2169065	GSE8613	125	1	88	–	125	2	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
141	Lucau-Danila 2005 Lucau-Danila A, Lelandais G, Kozovska Z, et al., Jacq C. Early expression of yeast genes affected by chemical stress. <i>Mol Cell Biol</i> 2005;25(5):1860–8.	doi:10.1128/MCB.25.5.1860-1868.2005 PMID 15713640 PMC549374	–	124	2	124	8	116	2	Microarray analysis - wt vs TF mutant
142	Weiner 2012 Weiner A, Chen HV, Liu CL, et al., Rando OJ. Systematic dissection of roles for chromatin regulators in a yeast stress response. <i>PLoS Biol</i> 2012;10(7):e1001369.	doi:10.1371/journal.pbio.1001369 PMID 22912562 PMC3416867	–	121	7	121	–	121	5	Microarray analysis - wt vs TF mutant
143	Lage 2019 Lage P, Sampaio-Marques B, Ludovico P, Mira NP, Mendes-Ferreira A. Transcriptomic and chemogenomic analyses unveil the essential role of Com2-regulon in response and tolerance of <i>Saccharomyces cerevisiae</i> to stress induced by sulfur dioxide. <i>Microb Cell</i> 2019;6(11):509–523.	doi:10.15698/mic2019.11.697 PMID 31799324 PMC6859422	–	121	1	72	–	121	3	Microarray analysis - wt vs TF mutant
144	Horan 2006 Horan S, Bourges I, Meunier B. Transcriptional response to nitrosative stress in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 2006; 23(7):519–35.	doi:10.1002/yea.1372 PMID 16710843	–	119	3	119	–	119	1	Microarray analysis - wt vs TF mutant
145	Zhu 2000 Zhu G, Spellman PT, Volpe T, et al., Fitcher B. Two yeast forkhead genes regulate the cell cycle and pseudohyphal growth. <i>Nature</i> 2000;406(6791):90–4.	doi:10.1038/35017581 PMID 10894548	GSE3182	118	2	109	4	114	2	Microarray analysis - wt vs TF mutant
146	Dubacq 2006 Dubacq C, Chevalier A, Courbeyrette R, Petat C, Gidrol X, Mann C. Role of the iron mobilization and oxidative stress regulons in the genomic response of yeast to hydroxyurea. <i>Mol Genet Genomics</i> 2006;275(2):114–24.	doi:10.1007/s00438-005-0077-5 PMID 16328372	GSE1914	115	3	115	–	115	2	Microarray analysis - wt vs TF mutant
147	Heise 2010 Heise B, van der Felden J, Kern S, Malcher M, Brückner S, Mösch HU. The TEA transcription factor Tec1 confers promoter-specific gene regulation by Ste12-dependent and -independent mechanisms. <i>Eukaryot Cell</i> 2010;9(4):514–31.	doi:10.1128/EC.00251-09 PMID 20118212 PMC2863411	–	115	2	115	–	115	1	Microarray analysis - wt vs TF mutant
148	Kwast 2002 Kwast KE, Lai LC, Menda N, James DT 3rd, Aref S, Burke PV. Genomic analyses of anaerobically induced genes in <i>Saccharomyces cerevisiae</i> : functional roles of Rox1 and other factors in mediating the anoxic response. <i>J Bacteriol</i> 2002; 184(1):250–65.	doi:10.1128/JB.184.1.250-265.2002 PMID 11741867 PMC134782	–	112	1	106	–	112	3	Microarray analysis - wt vs TF mutant
149	Courel 2005 Courel M, Lallet S, Camadro JM, Blaiseau PL. Direct activation of genes involved in intracellular iron use by the yeast iron-responsive transcription factor Aft2 without its paralog Aft1. <i>Mol Cell Biol</i> 2005;25(15):6760–71.	doi:10.1128/MCB.25.15.6760-6771.2005 PMID 16024809 PMC1190354	–	110	2	63	–	110	2	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
150	Rep 2000 Rep M, Krantz M, Thevelein JM, Hohmann S. The transcriptional response of <i>Saccharomyces cerevisiae</i> to osmotic shock. Hot1p and Msn2p/Msn4p are required for the induction of subsets of high osmolarity glycerol pathway-dependent genes. <i>J Biol Chem</i> 2000;275(12):8290–300.	doi:10.1074/jbc.275.12.8290 PMID 10722658	–	107	3	107	–	107	3	Microarray analysis - wt vs TF mutant
151	Cameroni 2004 Cameroni E, Hulo N, Roosen J, Winderickx J, De Virgilio C. The novel yeast PAS kinase Rim 15 orchestrates G0-associated antioxidant defense mechanisms. <i>Cell Cycle</i> 2004;3(4):462–8.	PMID 15300954	–	107	3	107	–	107	2	Microarray analysis - wt vs TF mutant
152	Dumond 2000 Dumond H, Danielou N, Pinto M, Bolotin-Fukuhara M. A large-scale study of Yap1p-dependent genes in normal aerobic and H ₂ O ₂ -stress conditions: the role of Yap1p in cell proliferation control in yeast. <i>Mol Microbiol</i> 2000;36(4):830–45.	doi:10.1046/j.1365-2958.2000.01845.x PMID 10844671	–	104	1	62	–	104	5	Microarray analysis - wt vs TF mutant
153	Kimata 2006 Kimata Y, Ishiwata-Kimata Y, Yamada S, Kohno K. Yeast unfolded protein response pathway regulates expression of genes for anti-oxidative stress and for cell surface proteins. <i>Genes Cells</i> 2006;11(1):59–69.	doi:10.1111/j.1365-2443.2005.00921.x PMID 16371132	–	104	1	104	–	104	1	Microarray analysis - wt vs TF mutant
154	Anderson 2006 Anderson JB, Ricker N, Sirjusingh C. Antagonism between two mechanisms of antifungal drug resistance. <i>Eukaryot Cell</i> 2006;5(8):1243–51.	doi:10.1128/EC.00048-06 PMID 16896209 PMC1539145	GSE4261	103	1	103	–	103	3	Microarray analysis - wt vs TF mutant
155	Cai 2013 Cai L, McCormick MA, Kennedy BK, Tu BP. Integration of multiple nutrient cues and regulation of lifespan by ribosomal transcription factor Ifh1. <i>Cell Rep</i> 2013;4(6):1063–71.	doi:10.1016/j.celrep.2013.08.016 PMID 24035395 PMC3792855	–	99	1	99	99	–	3	ChIP-seq
156	Wilcox 2002 Wilcox LJ, Balderes DA, Wharton B, Tinkelenberg AH, Rao G, Sturley SL. Transcriptional profiling identifies two members of the ATP-binding cassette transporter superfamily required for sterol uptake in yeast. <i>J Biol Chem</i> 2002;277(36):32466–72.	doi:10.1074/jbc.M204707200 PMID 12077145	–	96	1	88	–	96	2	Microarray analysis - wt vs TF mutant
157	Mira 2010 Mira NP, Becker JD, Sá-Correia I. Genomic expression program involving the Haa1p-regulon in <i>Saccharomyces cerevisiae</i> response to acetic acid. <i>OMICS</i> 2010;14(5):587–601.	doi:10.1089/omi.2010.0048 PMID 20955010 PMC3125556	–	96	1	86	–	96	2	Microarray analysis - wt vs TF mutant
158	Pramila 2002 Pramila T, Miles S, GuhaThakurta D, Jemiolo D, Breeden LL. Conserved homeodomain proteins interact with MADS box protein Mcm1 to restrict ECB-dependent transcription to the M/G1 phase of the cell cycle. <i>Genes Dev</i> 2002;16(23):3034–45.	doi:10.1101/gad.1034302 PMID 12464633 PMC187489	GSE5283	95	3	59	11	84	7	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
159	Lee 2010 Lee TA, Jorgensen P, Bognar AL, Peyraud C, Thomas D, Tyers M. Dissection of combinatorial control by the Met4 transcriptional complex. <i>Mol Biol Cell</i> 2010;21(3):456–69.	doi:10.1091/mbc.e09-05-0420 PMID 19940020 PMC2814790	–	95	5	81	–	95	3	Microarray analysis - wt vs TF mutant
160	Cook 2012 Cook KE, O’Shea EK. Hog1 controls global reallocation of RNA Pol II upon osmotic shock in <i>Saccharomyces cerevisiae</i> . <i>G3 (Bethesda)</i> 2012;2(9):1129–36.	doi:10.1534/g3.112.003251 PMID 22973550 PMC3429927	–	95	2	95	95	–	2	ChIP-seq
161	Slattery 2006 Slattery MG, Liko D, Heideman W. The function and properties of the Azf1 transcriptional regulator change with growth conditions in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2006;5(2):313–20.	doi:10.1128/EC.5.2.313-320.2006 PMID 16467472 PMC1405897	–	86	1	83	–	86	2	Microarray analysis - wt vs TF mutant
162	Miyake 2004 Miyake T, Reese J, Loch CM, Auble DT, Li R. Genome-wide analysis of ARS (autonomously replicating sequence) binding factor 1 (Abf1p)-mediated transcriptional regulation in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2004;279(33):34865–72.	doi:10.1074/jbc.M405156200 PMID 15192094	no GEO	85	1	85	5	80	1	Microarray analysis - wt vs TF mutant
163	Yamamoto 2005 Yamamoto A, Mizukami Y, Sakurai H. Identification of a novel class of target genes and a novel type of binding sequence of heat shock transcription factor in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2005;280(12):11911–9.	doi:10.1074/jbc.M411256200 PMID 15647283	no GEO	85	3	84	4	81	2	Microarray analysis - wt vs TF mutant
164	Lamb 2003 Lamb TM, Mitchell AP. The transcription factor Rim101p governs ion tolerance and cell differentiation by direct repression of the regulatory genes NRG1 and SMP1 in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2003;23(2):677–86.	doi:10.1128/MCB.23.2.677-686.2003 PMID 12509465 PMC151549	–	84	3	63	7	77	6	Microarray analysis - wt vs TF mutant
165	Pincus 2014 Pincus D, Aranda-Díaz A, Zuleta IA, Walter P, El-Samad H. Delayed Ras/PKA signaling augments the unfolded protein response. <i>Proc Natl Acad Sci U S A</i> 2014;111(41):14800–5.	doi:10.1073/pnas.1409588111 PMID 25275008 PMC4205644	–	82	1	82	–	82	1	RNA-seq - wt vs TF overexpression
166	Bai 2020 Bai C, Tesker M, Melamed-Kadosh D, Engelberg D, Admon A. Hog1-induced transcription of RTC3 and HSP12 is robust and occurs in cells lacking Msn2, Msn4, Hot1 and Sko1. <i>PLoS One</i> 2020;15(8):e0237540.	doi:10.1371/journal.pone.0237540 PMID 32804965 PMC7430751	–	82	4	16	–	82	2	RT-PCR - wt vs. TF quadruple mutant
167	Zhang 2009 Zhang N, Wu J, Oliver SG. Gis1 is required for transcriptional reprogramming of carbon metabolism and the stress response during transition into stationary phase in yeast. <i>Microbiology (Reading)</i> 2009;155(Pt 5):1690–1698.	doi:10.1099/mic.0.026377-0 PMID 19383711	–	81	1	50	–	81	4	Microarray analysis - wt vs TF mutant
168	DeRisi 2000 DeRisi J, van den Hazel B, Marc P, et al., Goffeau A. Genome microarray analysis of transcriptional activation in multidrug resistance yeast mutants. <i>FEBS Lett</i> 2000;470(2):156–60.	doi:10.1016/s0014-5793(00)01294-1 PMID 10734226	–	79	2	79	–	79	1	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
169	Bester 2012 Bester MC, Jacobson D, Bauer FF. Many <i>Saccharomyces cerevisiae</i> Cell Wall Protein Encoding Genes Are Coregulated by Mss11, but Cellular Adhesion Phenotypes Appear Only Flo Protein Dependent. <i>G3 (Bethesda)</i> 2012;2(1):131–41.	doi:10.1534/g3.111.001644 PMID 22384390 PMC3276193	GSE29371	76	1	76	2	74	1	Microarray analysis - wt vs TF overexpression
170	Davies 2006 Davies BS, Rine J. A role for sterol levels in oxygen sensing in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2006;174(1):191–201.	doi:10.1534/genetics.106.059964 PMID 16783004 PMC1569814	–	74	2	21	–	74	3	RT-PCR - wt vs TF mutant
171	Hendler 2017 Hendler A, Medina EM, Kishkevich A, et al., Aharoni A. Gene duplication and co-evolution of G1/S transcription factor specificity in fungi are essential for optimizing cell fitness. <i>PLoS Genet</i> 2017;13(5):e1006778.	doi:10.1371/journal.pgen.1006778 PMID 28505153 PMC5448814	–	74	1	68	6	68	3	RNA-seq - wt vs TF mutant
172	Hoppen 2005 Hoppen J, Repenning A, Albrecht A, Geburtig S, Schüller HJ. Comparative analysis of promoter regions containing binding sites of the heterodimeric transcription factor Ino2/Ino4 involved in yeast phospholipid biosynthesis. <i>Yeast</i> 2005;22(8):601–13.	doi:10.1002/yea.1209 PMID 16034810	–	72	3	30	–	72	2	lacZ - wt vs TF mutant
173	Mojzita 2006 Mojzita D, Hohmann S. Pdc2 coordinates expression of the THI regulon in the yeast <i>Saccharomyces cerevisiae</i> . <i>Mol Genet Genomics</i> 2006;276(2):147–61.	doi:10.1007/s00438-006-0130-z PMID 16850348	–	70	2	50	–	70	3	Microarray analysis - wt vs TF mutant
174	Wang 2017 Wang X, Liang Z, Hou J, Shen Y, Bao X. The Absence of the Transcription Factor Yrr1p, Identified from Comparative Genome Profiling, Increased Vanillin Tolerance Due to Enhancements of ABC Transporters Expressing, rRNA Processing and Ribosome Biogenesis in <i>Saccharomyces cerevisiae</i> . <i>Front Microbiol</i> 2017;8:367.	doi:10.3389/fmicb.2017.00367 PMID 28360888 PMC5352702	–	69	1	69	–	69	2	RNA-seq - wt vs TF mutant
175	Parnell 2020 Parnell EJ, Parnell TJ, Yan C, Bai L, Stillman DJ. Ash1 and Tup1 dependent repression of the <i>Saccharomyces cerevisiae</i> HO promoter requires activator-dependent nucleosome eviction. <i>PLoS Genet</i> 2020;16(12):e1009133.	doi:10.1371/journal.pgen.1009133 PMID 33382702 PMC7806131	–	67	2	32	66	1	1	ChIP-qPCR
176	Yin 2004 Yin Z, Stead D, Selway L, et al., Brown AJ. Proteomic response to amino acid starvation in <i>Candida albicans</i> and <i>Saccharomyces cerevisiae</i> . <i>Proteomics</i> 2004;4(8):2425–36.	doi:10.1002/pmic.200300760 PMID 15274137	–	65	1	65	–	65	1	Proteomics - wt vs TF mutant
177	Subramanian 2005 Subramanian M, Qiao WB, Khanam N, et al., Bognar AL. Transcriptional regulation of the one-carbon metabolism regulon in <i>Saccharomyces cerevisiae</i> by Bas1p. <i>Mol Microbiol</i> 2005;57(1):53–69.	doi:10.1111/j.1365-2958.2005.04663.x PMID 15948949	–	65	3	65	–	65	2	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
178	Niu 2008 Niu W, Li Z, Zhan W, Iyer VR, Marcotte EM. Mechanisms of cell cycle control revealed by a systematic and quantitative overexpression screen in <i>S. cerevisiae</i> . <i>PLoS Genet</i> 2008;4(7):e1000120.	doi:10.1371/journal.pgen.1000120 PMID 18617996 PMC2438615	–	65	1	65	–	65	1	Microarray analysis - wt vs TF overexpression
179	Ter Linde 2002 Ter Linde JJ, Steensma HY. A microarray-assisted screen for potential Hap1 and Rox1 target genes in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 2002;19(10):825–40.	doi:10.1002/yea.879 PMID 12112237	–	63	2	63	–	63	1	Microarray analysis - wt vs TF mutant
180	Ulusik 2011 Ulusik I, Kaya A, Fomenko DE, et al., Koc A. Boron stress activates the general amino acid control mechanism and inhibits protein synthesis. <i>PLoS One</i> 2011;6(11):e27772.	doi:10.1371/journal.pone.0027772 PMID 22114689 PMC3219688	GSE29887	63	2	62	1	62	3	Microarray analysis - wt vs TF mutant
181	Goh 2010 Goh WS, Orlov Y, Li J, Clarke ND. Blurring of high-resolution data shows that the effect of intrinsic nucleosome occupancy on transcription factor binding is mostly regional, not local. <i>PLoS Comput Biol</i> 2010;6(1):e1000649.	doi:10.1371/journal.pcbi.1000649 PMID 20098497 PMC2799660	–	62	3	62	62	–	2	ChIP
182	Hou 2014 Hou J, Tang H, Liu Z, Österlund T, Nielsen J, Petranovic D. Management of the endoplasmic reticulum stress by activation of the heat shock response in yeast. <i>FEMS Yeast Res</i> 2014;14(3):481–94.	doi:10.1111/1567-1364.12125 PMID 24237754	GSE39311	62	2	62	–	62	1	Microarray analysis - wt vs TF mutant
183	Liu 2016 Liu G, Bergenholm D, Nielsen J. Genome-Wide Mapping of Binding Sites Reveals Multiple Biological Functions of the Transcription Factor Cst6p in <i>Saccharomyces cerevisiae</i> . <i>mBio</i> 2016;7(3).	doi:10.1128/mBio.00559-16 PMID 27143390 PMC4959655	–	62	1	62	53	9	2	ChIP-exo
184	Wu 2008 Wu CY, Bird AJ, Chung LM, Newton MA, Winge DR, Eide DJ. Differential control of Zap1-regulated genes in response to zinc deficiency in <i>Saccharomyces cerevisiae</i> . <i>BMC Genomics</i> 2008;9:370.	doi:10.1186/1471-2164-9-370 PMID 18673560 PMC2535606	GSE11983	61	1	61	19	42	2	Microarray analysis - wt vs TF overexpression
185	Liu 2019 Liu R, Liang L, Choudhury A, et al., Gill RT. Multiplex navigation of global regulatory networks (MINR) in yeast for improved ethanol tolerance and production. <i>Metab Eng</i> 2019;51:50–58.	doi:10.1016/j.ymben.2018.07.007 PMID 30030154	–	61	3	54	–	61	2	RNA-seq - wt vs TF mutant
186	Causton 2001 Causton HC, Ren B, Koh SS, et al., Young RA. Remodeling of yeast genome expression in response to environmental changes. <i>Mol Biol Cell</i> 2001;12(2):323–37.	doi:10.1091/mbc.12.2.323 PMID 11179418 PMC30946	no GEO	60	2	60	–	60	1	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
187	González 2017 González J, López G, Argueta S, et al., González A. Diversification of Transcriptional Regulation Determines Subfunctionalization of Paralogous Branched Chain Aminotransferases in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2017;207(3):975–991.	doi:10.1534/genetics.117.300290 PMID 28912343 PMC5676234	–	59	5	16	33	26	5	ChIP
188	Muñoz 2020 Muñoz SA, Gullías JF, Valencia-Guillén J, Correa-García S, Bermúdez-Moretti M. Novel function of transcription factor Uga3 as an activator of branched-chain amino acid permease BAP2 gene expression. <i>Microbiology (Reading)</i> 2020;166(1):85–92.	doi:10.1099/mic.0.000863 PMID 31625834	–	59	6	6	8	51	9	lacZ - wt vs TF mutant
189	Boy-Marcotte 1999 Boy-Marcotte E, Lagniel G, Perrot M, et al., Labarre J. The heat shock response in yeast: differential regulations and contributions of the Msn2p/Msn4p and Hsf1p regulons. <i>Mol Microbiol</i> 1999;33(2):274–83.	doi:10.1046/j.1365-2958.1999.01467.x PMID 10411744	–	57	3	57	–	57	1	Proteomics - wt vs TF mutant
190	Lyons 2000 Lyons TJ, Gasch AP, Gaither LA, Botstein D, Brown PO, Eide DJ. Genome-wide characterization of the Zap1p zinc-responsive regulon in yeast. <i>Proc Natl Acad Sci U S A</i> 2000;97(14):7957–62.	doi:10.1073/pnas.97.14.7957 PMID 10884426 PMC16652	no GEO	57	1	46	–	57	2	Microarray analysis - wt vs TF mutant
191	Cai 2008 Cai L, Dalal CK, Elowitz MB. Frequency-modulated nuclear localization bursts coordinate gene regulation. <i>Nature</i> 2008;455(7212):485–90.	doi:10.1038/nature07292 PMID 18818649 PMC2695983	–	55	1	55	–	55	1	GFP - wt vs TF overexpression
192	Epstein 2001 Epstein CB, Waddle JA, Hale W 4th, et al., Butow RA. Genome-wide responses to mitochondrial dysfunction. <i>Mol Biol Cell</i> 2001;12(2):297–308.	doi:10.1091/mbc.12.2.297 PMID 11179416 PMC30944	no GEO	54	2	54	–	54	1	Microarray analysis - wt vs TF mutant
193	Pierce 2003 Pierce M, Benjamin KR, Montano SP, Georgiadis MM, Winter E, Vershon AK. Sum1 and Ndt80 proteins compete for binding to middle sporulation element sequences that control meiotic gene expression. <i>Mol Cell Biol</i> 2003;23(14):4814–25.	doi:10.1128/MCB.23.14.4814-4825.2003 PMID 12832469 PMC162219	–	54	2	54	–	54	2	Microarray analysis - wt vs TF mutant
194	Nordberg 2014 Nordberg N, Olsson I, Carlsson M, Hu GZ, Orzechowski Westholm J, Ronne H. The histone demethylase activity of Rph1 is not essential for its role in the transcriptional response to nutrient signaling. <i>PLoS One</i> 2014;9(7):e95078.	doi:10.1371/journal.pone.0095078 PMID 24999627 PMC4085034	–	53	2	53	–	53	2	Microarray analysis - wt vs TF mutant
195	Deed 2011 Deed NK, van Vuuren HJ, Gardner RC. Effects of nitrogen catabolite repression and di-ammonium phosphate addition during wine fermentation by a commercial strain of <i>S. cerevisiae</i> . <i>Appl Microbiol Biotechnol</i> 2011;89(5):1537–49.	doi:10.1007/s00253-011-3084-y PMID 21246356	–	52	2	52	–	52	1	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
196	Xiao 2018 Xiao W, Duan X, Lin Y, et al., Shui W. Distinct Proteome Remodeling of Industrial <i>Saccharomyces cerevisiae</i> in Response to Prolonged Thermal Stress or Transient Heat Shock. <i>J Proteome Res</i> 2018;17(5):1812–1825.	doi:10.1021/acs.jproteome.7b00842 PMID 29611422	–	52	1	52	–	52	1	SWATH- and DIA-based proteomic workflow
197	Tam 2020 Tam J, van Werven FJ. Regulated repression governs the cell fate promoter controlling yeast meiosis. <i>Nat Commun</i> 2020; 11(1):2271.	doi:10.1038/s41467-020-16107-w PMID 32385261 PMC7210989	–	52	10	10	44	8	10	ChIP
198	Bean 2005 Bean JM, Siggia ED, Cross FR. High functional overlap between MluI cell-cycle box binding factor and Swi4/6 cell-cycle box binding factor in the G1/S transcriptional program in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2005;171(1): 49–61.	doi:10.1534/genetics.105.044560 PMID 15965243 PMC1456534	–	51	2	51	3	48	2	Northern blot - wt vs TF mutant
199	Fardeau 2007 Fardeau V, Lelandais G, Oldfield A, et al., Devaux F. The central role of PDR1 in the foundation of yeast drug resistance. <i>J Biol Chem</i> 2007;282(7):5063–5074.	doi:10.1074/jbc.M610197200 PMID 17158869	–	51	2	51	–	51	1	Microarray analysis - wt vs TF mutant
200	Soontorngun 2007 Soontorngun N, Larochelle M, Drouin S, Robert F, Turcotte B. Regulation of gluconeogenesis in <i>Saccharomyces cerevisiae</i> is mediated by activator and repressor functions of Rds2. <i>Mol Cell Biol</i> 2007;27(22):7895–905.	doi:10.1128/MCB.01055-07 PMID 17875938 PMC2169140	–	51	1	51	51	–	2	ChIP
201	Gurvich 2017 Gurvich Y, Leshkowitz D, Barkai N. Dual role of starvation signaling in promoting growth and recovery. <i>PLoS Biol</i> 2017; 15(12):e2002039.	doi:10.1371/journal.pbio.2002039 PMID 29236696 PMC5728490	–	51	1	29	–	51	3	RNA-seq - wt vs TF mutant
202	Lee 1999 Lee J, Godon C, Lagniel G, et al., Toledano MB. Yap1 and Skn7 control two specialized oxidative stress response regulons in yeast. <i>J Biol Chem</i> 1999;274(23):16040–6.	doi:10.1074/jbc.274.23.16040 PMID 10347154	–	50	2	50	2	48	4	Proteomics - wt vs TF mutant
203	Lenssen 2005 Lenssen E, James N, Pedruzzi I, et al., De Virgilio C. The Ccr4-Not complex independently controls both Msn2-dependent transcriptional activation–via a newly identified Glc7/Bud14 type I protein phosphatase module–and TFIID promoter distribution. <i>Mol Cell Biol</i> 2005;25(1): 488–98.	doi:10.1128/MCB.25.1.488-498.2005 PMID 15601868 PMC538800	–	50	2	50	–	50	1	Microarray analysis - wt vs TF mutant
204	Westergaard 2007 Westergaard SL, Oliveira AP, Bro C, Olsson L, Nielsen J. A systems biology approach to study glucose repression in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biotechnol Bioeng</i> 2007;96(1): 134–45.	doi:10.1002/bit.21135 PMID 16878332	–	50	2	50	–	50	1	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
205	Frey 2011 Frey AG, Eide DJ. Roles of two activation domains in Zap1 in the response to zinc deficiency in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2011;286(8):6844–54.	doi:10.1074/jbc.M110.203927 PMID 21177862 PMC3057842	–	50	1	50	1	49	1	Microarray analysis - wt vs TF mutant
206	Lickwar 2012 Lickwar CR, Mueller F, Hanlon SE, McNally JG, Lieb JD. Genome-wide protein-DNA binding dynamics suggest a molecular clutch for transcription factor function. <i>Nature</i> 2012;484(7393):251–5.	doi:10.1038/nature10985 PMID 22498630 PMC3341663	–	50	1	50	50	–	1	ChIP
207	Becerra 2002 Becerra M, Lombardía-Ferreira LJ, Hauser NC, Hoheisel JD, Tizon B, Cerdán ME. The yeast transcriptome in aerobic and hypoxic conditions: effects of hap1, rox1, rox3 and srb10 deletions. <i>Mol Microbiol</i> 2002;43(3):545–55.	doi:10.1046/j.1365-2958.2002.02724.x PMID 11929514	–	48	3	48	–	48	2	Microarray analysis - wt vs TF mutant
208	Young 2003 Young ET, Dombek KM, Tachibana C, Ideker T. Multiple pathways are co-regulated by the protein kinase Snf1 and the transcription factors Adr1 and Cat8. <i>J Biol Chem</i> 2003;278(28):26146–58.	doi:10.1074/jbc.M301981200 PMID 12676948	–	47	1	47	–	47	1	Microarray analysis - wt vs TF mutant
209	Wimalarathna 2011 Wimalarathna R, Tsai CH, Shen CH. Transcriptional control of genes involved in yeast phospholipid biosynthesis. <i>J Microbiol</i> 2011;49(2):265–73.	doi:10.1007/s12275-011-1130-1 PMID 21538248	–	47	1	47	–	47	1	RT-PCR - wt vs TF mutant
210	Proft 2005 Proft M, Gibbons FD, Copeland M, Roth FP, Struhl K. Genomewide identification of Sko1 target promoters reveals a regulatory network that operates in response to osmotic stress in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2005;4(8):1343–52.	doi:10.1128/EC.4.8.1343-1352.2005 PMID 16087739 PMC1214534	–	46	1	46	41	5	2	ChIP
211	Schroeder 2013 Schroeder EA, Raimundo N, Shadel GS. Epigenetic silencing mediates mitochondria stress-induced longevity. <i>Cell Metab</i> 2013;17(6):954–964.	doi:10.1016/j.cmet.2013.04.003 PMID 23747251 PMC3694503	–	45	1	45	–	45	1	Microarray analysis - wt vs TF mutant
212	Pimentel 2014 Pimentel C, Caetano SM, Menezes R, et al., Rodrigues-Pousada C. Yap1 mediates tolerance to cobalt toxicity in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochim Biophys Acta</i> 2014;1840(6):1977–86.	doi:10.1016/j.bbagen.2014.01.032 PMID 24486411	–	44	2	44	–	44	2	Microarray analysis - wt vs TF mutant
213	Pérez-Sampietro 2016 Pérez-Sampietro M, Serra-Cardona A, Canadell D, Casas C, Ariño J, Herrero E. The yeast Aft2 transcription factor determines selenite toxicity by controlling the low affinity phosphate transport system. <i>Sci Rep</i> 2016;6:32836.	doi:10.1038/srep32836 PMID 27618952 PMC5020356	–	44	1	44	–	44	1	RNA-seq - wt vs TF mutant
214	Mondeel 2019 Mondeel TDGA, Holland P, Nielsen J, Barberis M. ChIP-exo analysis highlights Fkh1 and Fkh2 transcription factors as hubs that integrate multi-scale networks in budding yeast. <i>Nucleic Acids Res</i> 2019;47(15):7825–7841.	doi:10.1093/nar/gkz603 PMID 31299083 PMC6736057	–	44	2	44	44	–	1	ChIP-exo

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
215	Karpichev 2008 Karpichev IV, Durand-Heredia JM, Luo Y, Small GM. Binding characteristics and regulatory mechanisms of the transcription factors controlling oleate-responsive genes in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2008;283(16):10264–75.	doi:10.1074/jbc.M708215200 PMID 18285336 PMC2447635	–	43	3	43	29	14	2	ChIP
216	de Jonge 2017 de Jonge WJ, O’Duibhir E, Lijnzaad P, et al., Holstege FC. Molecular mechanisms that distinguish TFIID housekeeping from regulatable SAGA promoters. <i>EMBO J</i> 2017;36(3):274–290.	doi:10.15252/embj.201695621 PMID 27979920 PMC5286361	GSE81480	42	1	21	21	21	1	ChIP
217	Bergman 2019 Bergman A, Vitay D, Hellgren J, Chen Y, Nielsen J, Siewers V. Effects of overexpression of STB5 in <i>Saccharomyces cerevisiae</i> on fatty acid biosynthesis, physiology and transcriptome. <i>FEMS Yeast Res</i> 2019;19(3).	doi:10.1093/femsyr/foz027 PMID 30924859 PMC6755256	–	42	1	29	–	42	3	RNA-seq - wt vs TF overexpression
218	Abramova 2001 Abramova NE, Cohen BD, Sertil O, Kapoor R, Davies KJ, Lowry CV. Regulatory mechanisms controlling expression of the DAN/TIR mannoprotein genes during anaerobic remodeling of the cell wall in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2001;157(3):1169–77.	doi:10.1093/genetics/157.3.1169 PMID 11238402 PMC1461566	–	41	5	35	–	41	6	Northern blot - wt vs TF mutant
219	Barreto 2006 Barreto L, Garcerá A, Jansson K, Sunnerhagen P, Herrero E. A peroxisomal glutathione transferase of <i>Saccharomyces cerevisiae</i> is functionally related to sulfur amino acid metabolism. <i>Eukaryot Cell</i> 2006;5(10):1748–59.	doi:10.1128/EC.00216-06 PMID 16936141 PMC1595348	–	41	3	10	–	41	8	Northern blot - wt vs TF mutant
220	Bernard 2015 Bernard A, Jin M, Xu Z, Klionsky DJ. A large-scale analysis of autophagy-related gene expression identifies new regulators of autophagy. <i>Autophagy</i> 2015;11(11):2114–2122.	doi:10.1080/15548627.2015.1099796 PMID 26649943 PMC4824583	–	41	11	41	4	37	3	RT-PCR - wt vs TF mutant
221	He 2017 He BZ, Zhou X, O’Shea EK. Evolution of reduced co-activator dependence led to target expansion of a starvation response pathway. <i>Elife</i> 2017;6.	doi:10.7554/eLife.25157 PMID 28485712 PMC5446240	–	41	1	39	–	41	2	RNA-seq - wt vs TF mutant
222	Voth 2007 Voth WP, Yu Y, Takahata S, et al., Stillman DJ. Forkhead proteins control the outcome of transcription factor binding by antiactivation. <i>EMBO J</i> 2007;26(20):4324–34.	doi:10.1038/sj.emboj.7601859 PMID 17898805 PMC2034669	–	40	4	40	40	–	1	Chip-on-chip
223	Martínez-Montañés 2013 Martínez-Montañés F, Rienzo A, Poveda-Huertes D, Pascual-Ahuir A, Proft M. Activator and repressor functions of the Mot3 transcription factor in the osmostress response of <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2013;12(5):636–47.	doi:10.1128/EC.00037-13 PMID 23435728 PMC3647766	–	40	6	40	12	28	3	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
224	Gasmi 2014 Gasmi N, Jacques PE, Klimova N, et al., Turcotte B. The switch from fermentation to respiration in <i>Saccharomyces cerevisiae</i> is regulated by the Ert1 transcriptional activator/repressor. <i>Genetics</i> 2014;198(2):547–60.	doi:10.1534/genetics.114.168609 PMID 25123508 PMC4196612	–	40	3	40	34	6	4	Chip-on-chip
225	Linke 2017 Linke C, Chasapi A, González-Novo A, et al., Barberis M. A Clb/Cdk1-mediated regulation of Fkh2 synchronizes CLB expression in the budding yeast cell cycle. <i>NPJ Syst Biol Appl</i> 2017;3:7.	doi:10.1038/s41540-017-0008-1 PMID 28649434 PMC5460246	–	40	3	11	6	34	7	RT-PCR - wt vs TF mutant
226	Pereira 2008 Pereira Y, Lagniel G, Godat E, Baudouin-Cornu P, Junot C, Labarre J. Chromate causes sulfur starvation in yeast. <i>Toxicol Sci</i> 2008;106(2):400–12.	doi:10.1093/toxsci/kfn193 PMID 18794233	–	39	1	39	–	39	1	Microarray analysis - wt vs TF mutant
227	Ponomarova 2017 Ponomarova O, Gabrielli N, Sévin DC, et al., Patil KR. Yeast Creates a Niche for Symbiotic Lactic Acid Bacteria through Nitrogen Overflow. <i>Cell Syst</i> 2017;5(4):345–357.e6.	doi:10.1016/j.cels.2017.09.002 PMID 28964698 PMC5660601	–	39	2	39	–	39	1	RNA-seq - wt vs TF mutant
228	Devaux 2002 Devaux F, Carvajal E, Moye-Rowley S, Jacq C. Genome-wide studies on the nuclear PDR3-controlled response to mitochondrial dysfunction in yeast. <i>FEBS Lett</i> 2002;515(1–3):25–8.	doi:10.1016/s0014-5793(02)02387-6 PMID 11943188	–	38	2	30	–	38	2	Microarray analysis - wt vs TF mutant
229	Mira 2009 Mira NP, Lourenço AB, Fernandes AR, Becker JD, Sá-Correia I. The RIM101 pathway has a role in <i>Saccharomyces cerevisiae</i> adaptive response and resistance to propionic acid and other weak acids. <i>FEMS Yeast Res</i> 2009;9(2):202–16.	doi:10.1111/j.1567-1364.2008.00473.x PMID 19220866	–	38	1	38	–	38	2	Microarray analysis - wt vs TF mutant
230	Palavecino-Ruiz 2017 Palavecino-Ruiz M, Bermudez-Moretti M, Correa-Garcia S. Unravelling the transcriptional regulation of <i>Saccharomyces cerevisiae</i> UGA genes: the dual role of transcription factor Leu3. <i>Microbiology (Reading)</i> 2017;163(11):1692–1701.	doi:10.1099/mic.0.000560 PMID 29058647	–	37	7	13	1	36	17	lacZ - wt vs TF mutant
231	Amorós 2001 Amorós M, Estruch F. Hsf1p and Msn2/4p cooperate in the expression of <i>Saccharomyces cerevisiae</i> genes HSP26 and HSP104 in a gene- and stress type-dependent manner. <i>Mol Microbiol</i> 2001;39(6):1523–32.	doi:10.1046/j.1365-2958.2001.02339.x PMID 11260469	–	36	3	10	–	36	4	Northern blot - wt vs TF mutant
232	Cox 1999 Cox KH, Pinchak AB, Cooper TG. Genome-wide transcriptional analysis in <i>S. cerevisiae</i> by mini-array membrane hybridization. <i>Yeast</i> 1999;15(8):703–13.	<a href="https://doi.org/10.1002/(SICI)1097-0061(19990615)15:8<703::AID-YEA413>3.0.CO;2-Z">doi:10.1002/(SICI)1097-0061(19990615)15:8<703::AID-YEA413>3.0.CO;2-Z PMID 10392447	–	35	2	33	–	35	1	Microarray analysis - wt vs TF mutant
233	López 2000 López MC, Baker HV. Understanding the growth phenotype of the yeast <i>gcr1</i> mutant in terms of global genomic expression patterns. <i>J Bacteriol</i> 2000;182(17):4970–8.	doi:10.1128/JB.182.17.4970-4978.2000 PMID 10940042 PMC111378	no GEO	35	1	35	–	35	2	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
234	Bai 2015 Bai C, Tesker M, Engelberg D. The yeast Hot1 transcription factor is critical for activating a single target gene, STL1. <i>Mol Biol Cell</i> 2015;26(12):2357–74.	doi:10.1091/mbc.E14-12-1626 PMID 25904326 PMC4462951	–	35	1	35	–	35	5	Microarray analysis - wt vs TF mutant
235	Schuurmans 2008 Schuurmans JM, Boorsma A, Lascaris R, Hellingwerf KJ, Teixeira de Mattos MJ. Physiological and transcriptional characterization of <i>Saccharomyces cerevisiae</i> strains with modified expression of catabolic regulators. <i>FEMS Yeast Res</i> 2008;8(1):26–34.	doi:10.1111/j.1567-1364.2007.00309.x PMID 17892474	–	34	1	34	–	34	1	Microarray analysis - wt vs TF mutant
236	Hazelwood 2010 Hazelwood LA, Walsh MC, Pronk JT, Daran JM. Involvement of vacuolar sequestration and active transport in tolerance of <i>Saccharomyces cerevisiae</i> to hop iso-alpha-acids. <i>Appl Environ Microbiol</i> 2010;76(1):318–28.	doi:10.1128/AEM.01457-09 PMID 19915041 PMC2798648	–	34	4	34	–	34	1	Microarray analysis - wt vs TF mutant
237	Jordá 2020 Jordá T, Romero AM, Perea-García A, Rozès N, Puig S. The lipid composition of yeast cells modulates the response to iron deficiency. <i>Biochim Biophys Acta Mol Cell Biol Lipids</i> 2020; 1865(8):158707.	doi:10.1016/j.bbalip.2020.158707 PMID 32251724	–	34	2	12	–	34	9	RT-PCR - wt vs. TF mutant
238	Denis 1998 Denis V, Boucherie H, Monribot C, Daignan-Fornier B. Role of the myb-like protein bas1p in <i>Saccharomyces cerevisiae</i> : a proteome analysis. <i>Mol Microbiol</i> 1998;30(3):557–66.	doi:10.1046/j.1365-2958.1998.01087.x PMID 9822821	–	33	2	19	–	33	2	lacZ - wt vs TF mutant
239	Haurie 2001 Haurie V, Perrot M, Mini T, Jenö P, Sagliocco F, Boucherie H. The transcriptional activator Cat8p provides a major contribution to the reprogramming of carbon metabolism during the diauxic shift in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2001;276(1):76–85.	doi:10.1074/jbc.M008752200 PMID 11024040	–	33	1	33	–	33	3	Northern blot - wt vs TF mutant
240	Aranda 2004 Aranda A, del Olmo ML. Exposure of <i>Saccharomyces cerevisiae</i> to acetaldehyde induces sulfur amino acid metabolism and polyamine transporter genes, which depend on Met4p and Haa1p transcription factors, respectively. <i>Appl Environ Microbiol</i> 2004;70(4):1913–22.	doi:10.1128/AEM.70.4.1913-1922.2004 PMID 15066780 PMC383134	–	33	7	29	–	33	2	Western blot - wt vs TF mutant
241	Biddick 2008 Biddick RK, Law GL, Young ET. Adr1 and Cat8 mediate coactivator recruitment and chromatin remodeling at glucose-regulated genes. <i>PLoS One</i> 2008;3(1):e1436.	doi:10.1371/journal.pone.0001436 PMID 18197247 PMC2175534	–	33	2	33	–	33	2	RT-PCR - wt vs TF mutant
242	Nunez 2008 Nunez LR, Jesch SA, Gaspar ML, et al., Henry SA. Cell wall integrity MAPK pathway is essential for lipid homeostasis. <i>J Biol Chem</i> 2008;283(49):34204–17.	doi:10.1074/jbc.M806391200 PMID 18842580 PMC2590691	–	33	2	33	–	33	1	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
243	Liu 1999 Liu Z, Butow RA. A transcriptional switch in the expression of yeast tricarboxylic acid cycle genes in response to a reduction or loss of respiratory function. <i>Mol Cell Biol</i> 1999; 19(10):6720–8.	doi:10.1128/MCB.19.10.6720 PMID 10490611 PMC84662	–	32	4	16	3	29	5	Northern blot - wt vs TF mutant
244	Jung 1999 Jung US, Levin DE. Genome-wide analysis of gene expression regulated by the yeast cell wall integrity signalling pathway. <i>Mol Microbiol</i> 1999;34(5):1049–57.	doi:10.1046/j.1365-2958.1999.01667.x PMID 10594829	–	32	1	22	–	32	2	Northern blot - wt vs TF mutant
245	Stewart-Ornstein 2013 Stewart-Ornstein J, Nelson C, DeRisi J, Weissman JS, El-Samad H. Msn2 coordinates a stoichiometric gene expression program. <i>Curr Biol</i> 2013;23(23):2336–45.	doi:10.1016/j.cub.2013.09.043 PMID 24210615 PMC4072881	–	32	1	32	–	32	1	GFP - wt vs TF overexpression
246	Karpichev 1998 Karpichev IV, Small GM. Global regulatory functions of Oaf1p and Pip2p (Oaf2p), transcription factors that regulate genes encoding peroxisomal proteins in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1998;18(11):6560–70.	doi:10.1128/MCB.18.11.6560 PMID 9774671 PMC109241	–	31	2	31	–	31	3	Northern blot - wt vs TF mutant
247	Jesch 2005 Jesch SA, Zhao X, Wells MT, Henry SA. Genome-wide analysis reveals inositol, not choline, as the major effector of Ino2p-Ino4p and unfolded protein response target gene expression in yeast. <i>J Biol Chem</i> 2005;280(10):9106–18.	doi:10.1074/jbc.M411770200 PMID 15611057 PMC1352320	–	31	1	31	–	31	2	Microarray analysis - wt vs TF mutant
248	Petti 2012 Petti AA, McIsaac RS, Ho-Shing O, Bussemaker HJ, Botstein D. Combinatorial control of diverse metabolic and physiological functions by transcriptional regulators of the yeast sulfur assimilation pathway. <i>Mol Biol Cell</i> 2012;23(15):3008–24.	doi:10.1091/mbc.E12-03-0233 PMID 22696679 PMC3408426	–	31	3	31	–	31	1	Microarray analysis - wt vs TF mutant
249	Tadi 1999 Tadi D, Hasan RN, Bussereau F, Boy-Marcotte E, Jacquet M. Selection of genes repressed by cAMP that are induced by nutritional limitation in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 1999; 15(16):1733–45.	<a href="https://doi.org/10.1002/(SICI)1097-0061(199912)15:16<1733::AID-YEA490>3.0.CO;2-L">doi:10.1002/(SICI)1097-0061(199912)15:16<1733::AID-YEA490>3.0.CO;2-L PMID 10590462	–	30	2	30	–	30	1	lacZ - wt vs TF mutant
250	Miura 2001 Miura F, Yada T, Nakai K, Sakaki Y, Ito T. Differential display analysis of mutants for the transcription factor Pdr1p regulating multidrug resistance in the budding yeast. <i>FEBS Lett</i> 2001;505(1):103–8.	doi:10.1016/s0014-5793(01)02792-2 PMID 11557050	–	30	1	20	1	29	1	ATAC-PCR - wt vs TF overexpression
251	Woods 2016 Woods K, Höfken T. The zinc cluster proteins Upc2 and Ecm22 promote filamentation in <i>Saccharomyces cerevisiae</i> by sterol biosynthesis-dependent and -independent pathways. <i>Mol Microbiol</i> 2016;99(3):512–27.	doi:10.1111/mmi.13244 PMID 26448198	–	30	8	30	–	30	10	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
252	Chen 2017 Chen X, Wang Z, Guo X, Liu S, He X. Regulation of general amino acid permeases Gap1p, GATA transcription factors Gln3p and Gat1p on 2-phenylethanol biosynthesis via Ehrlich pathway. <i>J Biotechnol</i> 2017;242:83–91.	doi:10.1016/j.jbiotec.2016.11.028 PMID 27908775	–	30	2	10	–	30	6	RT-PCR - wt vs TF mutant
253	Urnov 2002 Urnov FD. A feel for the template: zinc finger protein transcription factors and chromatin. <i>Biochem Cell Biol</i> 2002; 80(3):321–33.	doi:10.1139/o02-084 PMID 12123285	–	29	1	29	29	–	1	Chip-on-chip
254	Tanaka 2015 Tanaka N, Mukai Y. Yeast Cyc8p and Tup1p proteins function as coactivators for transcription of Stp1/2p-dependent amino acid transporter genes. <i>Biochem Biophys Res Commun</i> 2015;468(1–2):32–8.	doi:10.1016/j.bbrc.2015.11.001 PMID 26546823	–	29	4	29	2	27	4	RT-PCR - wt vs TF mutant
255	Ogawa 2000 Ogawa N, DeRisi J, Brown PO. New components of a system for phosphate accumulation and polyphosphate metabolism in <i>Saccharomyces cerevisiae</i> revealed by genomic expression analysis. <i>Mol Biol Cell</i> 2000;11(12):4309–21.	doi:10.1091/mbc.11.12.4309 PMID 11102525 PMC15074	no GEO	28	1	28	–	28	2	Microarray analysis - wt vs TF mutant
256	Shamji 2000 Shamji AF, Kuruvilla FG, Schreiber SL. Partitioning the transcriptional program induced by rapamycin among the effectors of the Tor proteins. <i>Curr Biol</i> 2000;10(24):1574–81.	doi:10.1016/s0960-9822(00)00866-6 PMID 11137008	–	28	2	28	–	28	1	Microarray analysis - wt vs TF mutant
257	Aguilera 2005 Aguilera J, Rodríguez-Vargas S, Prieto JA. The HOG MAP kinase pathway is required for the induction of methylglyoxal-responsive genes and determines methylglyoxal resistance in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2005; 56(1):228–39.	doi:10.1111/j.1365-2958.2005.04533.x PMID 15773992	–	27	4	27	–	27	2	Microarray analysis - wt vs TF mutant
258	Viladevall 2004 Viladevall L, Serrano R, Ruiz A, et al., Ariño J. Characterization of the calcium-mediated response to alkaline stress in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2004;279(42): 43614–24.	doi:10.1074/jbc.M403606200 PMID 15299026	–	26	3	26	–	26	6	Microarray analysis - wt vs TF mutant
259	Galdieri 2016 Galdieri L, Zhang T, Rogerson D, Vancura A. Reduced Histone Expression or a Defect in Chromatin Assembly Induces Respiration. <i>Mol Cell Biol</i> 2016;36(7):1064–77.	doi:10.1128/MCB.00770-15 PMID 26787838 PMC4800798	–	26	2	26	–	26	1	RT-PCR - wt vs TF mutant
260	Zhang 2018 Zhang P, Li B, Wen P, et al., Hu X. Metabolic Engineering of Four GATA Factors to Reduce Urea and Ethyl Carbamate Formation in a Model Rice Wine System. <i>J Agric Food Chem</i> 2018;66(41):10881–10889.	doi:10.1021/acs.jafc.8b04370 PMID 30246534	–	26	3	6	–	26	13	RT-PCR - wt vs TF mutant
261	Doolin 2001 Doolin MT, Johnson AL, Johnston LH, Butler G. Overlapping and distinct roles of the duplicated yeast transcription factors Ace2p and Swi5p. <i>Mol Microbiol</i> 2001;40(2):422–32.	doi:10.1046/j.1365-2958.2001.02388.x PMID 11309124	–	25	2	25	–	25	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
262	Rutherford 2001 Rutherford JC, Jaron S, Ray E, Brown PO, Winge DR. A second iron-regulatory system in yeast independent of Aft1p. <i>Proc Natl Acad Sci U S A</i> 2001;98(25):14322–7.	doi:10.1073/pnas.261381198 PMID 11734641 PMC64680	–	25	2	25	6	19	4	Microarray analysis - wt vs TF overexpression
263	Cormier 2010 Cormier L, Barbey R, Kuras L. Transcriptional plasticity through differential assembly of a multiprotein activation complex. <i>Nucleic Acids Res</i> 2010;38(15):4998–5014.	doi:10.1093/nar/gkq257 PMID 20392822 PMC2926612	–	25	4	13	18	7	15	ChIP
264	Galello 2017 Galello F, Pautasso C, Reca S, et al., Rossi S. Transcriptional regulation of the protein kinase a subunits in <i>Saccharomyces cerevisiae</i> during fermentative growth. <i>Yeast</i> 2017;34(12):495–508.	doi:10.1002/yea.3252 PMID 28812308	–	25	3	11	2	23	4	RT-PCR - wt vs TF mutant
265	Boy-Marcotte 1998 Boy-Marcotte E, Perrot M, Bussereau F, Boucherie H, Jacquet M. Msn2p and Msn4p control a large number of genes induced at the diauxic transition which are repressed by cyclic AMP in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1998;180(5):1044–52.	doi:10.1128/JB.180.5.1044-1052.1998 PMID 9495741 PMC106990	–	24	2	24	–	24	1	Proteomics - wt vs TF mutant
266	Shapira 2004 Shapira M, Segal E, Botstein D. Disruption of yeast forkhead-associated cell cycle transcription by oxidative stress. <i>Mol Biol Cell</i> 2004;15(12):5659–69.	doi:10.1091/mbc.e04-04-0340 PMID 15371544 PMC532044	no GEO	24	3	24	–	24	1	Microarray analysis - wt vs TF mutant
267	Watanabe 2011 Watanabe D, Wu H, Noguchi C, Zhou Y, Akao T, Shimoi H. Enhancement of the initial rate of ethanol fermentation due to dysfunction of yeast stress response components Msn2p and/or Msn4p. <i>Appl Environ Microbiol</i> 2011;77(3):934–41.	doi:10.1128/AEM.01869-10 PMID 21131516 PMC3028739	–	24	2	24	–	24	1	RT-PCR - wt vs TF mutant
268	Caballero-Molada 2018 Caballero-Molada M, Planes MD, Benlloch H, Atares S, Naranjo MA, Serrano R. The Gcn2-eIF2 α pathway connects iron and amino acid homeostasis in <i>Saccharomyces cerevisiae</i> . <i>Biochem J</i> 2018;475(8):1523–1534.	doi:10.1042/BCJ20170871 PMID 29626156	–	24	2	13	–	24	3	RT-PCR - wt vs TF mutant
269	Vandermeulen 2022 Vandermeulen MD, Cullen PJ. Gene by Environment Interactions reveal new regulatory aspects of signaling network plasticity. <i>PLoS Genet</i> 2022;18(1):e1009988.	doi:10.1371/journal.pgen.1009988 PMID 34982769 PMC8759647	–	24	5	12	–	24	4	RT-PCR - wt vs. TF mutant
270	Rep 2001 Rep M, Proft M, Remize F, et al., Hohmann S. The <i>Saccharomyces cerevisiae</i> Sko1p transcription factor mediates HOG pathway-dependent osmotic regulation of a set of genes encoding enzymes implicated in protection from oxidative damage. <i>Mol Microbiol</i> 2001;40(5):1067–83.	doi:10.1046/j.1365-2958.2001.02384.x PMID 11401713	–	23	7	15	–	23	5	Northern blot - wt vs TF mutant
271	Santiago 2003 Santiago TC, Mamoun CB. Genome expression analysis in yeast reveals novel transcriptional regulation by inositol and choline and new regulatory functions for Opi1p, Ino2p, and Ino4p. <i>J Biol Chem</i> 2003;278(40):38723–30.	doi:10.1074/jbc.M303008200 PMID 12871953	–	23	2	23	–	23	1	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
272	Hansen 2013 Hansen AS, O'Shea EK. Promoter decoding of transcription factor dynamics involves a trade-off between noise and control of gene expression. <i>Mol Syst Biol</i> 2013;9:704.	doi:10.1038/msb.2013.56 PMID 24189399 PMC4039373	–	23	1	23	–	23	1	Microarray analysis - wt vs TF mutant
273	Natkańska 2017 Natkańska U, Skoneczna A, Sieńko M, Skoneczny M. The budding yeast orthologue of Parkinson's disease-associated DJ-1 is a multi-stress response protein protecting cells against toxic glycolytic products. <i>Biochim Biophys Acta Mol Cell Res</i> 2017;1864(1):39–50.	doi:10.1016/j.bbamcr.2016.10.016 PMID 27984092	–	23	7	7	–	23	11	lacZ - wt vs TF mutant
274	Ravi 2019 Ravi C, Gowsalya R, Nachiappan V. Impaired GCR1 transcription resulted in defective inositol levels, vacuolar structure and autophagy in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 2019;65(4):995–1014.	doi:10.1007/s00294-019-00954-2 PMID 30879088	–	23	4	9	–	23	6	RT-PCR - wt vs TF mutant
275	Becerra 2011 Becerra M, Lombardía LJ, Lamas-Maceiras M, Canto E, Rodríguez-Belmonte E, Cerdán ME. Comparative transcriptome analysis of yeast strains carrying slt2, rlm1, and pop2 deletions. <i>Genome</i> 2011;54(2):99–109.	doi:10.1139/G10-101 PMID 21326366	–	22	1	22	4	18	1	Microarray analysis - wt vs TF mutant
276	Chubukov 2012 Chubukov V, Zuleta IA, Li H. Regulatory architecture determines optimal regulation of gene expression in metabolic pathways. <i>Proc Natl Acad Sci U S A</i> 2012;109(13):5127–32.	doi:10.1073/pnas.1114235109 PMID 22416120 PMC3324031	–	22	3	22	–	22	3	GFP - wt vs TF mutant
277	McIsaac 2012 McIsaac RS, Petti AA, Bussemaker HJ, Botstein D. Perturbation-based analysis and modeling of combinatorial regulation in the yeast sulfur assimilation pathway. <i>Mol Biol Cell</i> 2012;23(15):2993–3007.	doi:10.1091/mbc.E12-03-0232 PMID 22696683 PMC3408425	–	22	2	22	–	22	1	Microarray analysis - wt vs TF overexpression
278	Shin 2021 Shin M, Park H, Kim S, et al., Kim SR. Transcriptomic Changes Induced by Deletion of Transcriptional Regulator GCR2 on Pentose Sugar Metabolism in <i>Saccharomyces cerevisiae</i> . <i>Front Bioeng Biotechnol</i> 2021;9:654177.	doi:10.3389/fbioe.2021.654177 PMID 33842449 PMC8027353	–	22	1	22	–	22	2	RNA-seq
279	Coffman 1997 Coffman JA, Rai R, Loprete DM, Cunningham T, Svetlov V, Cooper TG. Cross regulation of four GATA factors that control nitrogen catabolic gene expression in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1997;179(11):3416–29.	doi:10.1128/jb.179.11.3416-3429.1997 PMID 9171383 PMC179131	–	21	5	15	–	21	4	Northern blot - wt vs TF mutant
280	Gross 2000 Gross C, Kelleher M, Iyer VR, Brown PO, Winge DR. Identification of the copper regulon in <i>Saccharomyces cerevisiae</i> by DNA microarrays. <i>J Biol Chem</i> 2000;275(41):32310–6.	doi:10.1074/jbc.M005946200 PMID 10922376	no GEO	21	3	21	6	13	4	Microarray analysis - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
281	Keller 2001 Keller G, Ray E, Brown PO, Winge DR. Haa1, a protein homologous to the copper-regulated transcription factor Ace1, is a novel transcriptional activator. <i>J Biol Chem</i> 2001;276(42):38697–702.	doi:10.1074/jbc.M107131200 PMID 11504737	no GEO	21	1	9	12	9	5	DNA footprinting
282	Ratnakumar 2010 Ratnakumar S, Young ET. Snf1 dependence of peroxisomal gene expression is mediated by Adr1. <i>J Biol Chem</i> 2010;285(14):10703–14.	doi:10.1074/jbc.M109.079848 PMID 20139423 PMC2856278	–	21	3	21	7	14	3	RT-PCR - wt vs TF mutant
283	McIsaac 2011 McIsaac RS, Silverman SJ, McClean MN, et al., Botstein D. Fast-acting and nearly gratuitous induction of gene expression and protein depletion in <i>Saccharomyces cerevisiae</i> . <i>Mol Biol Cell</i> 2011;22(22):4447–59.	doi:10.1091/mbc.E11-05-0466 PMID 21965290 PMC3216669	–	21	1	21	–	21	1	Microarray analysis - wt vs TF mutant
284	Platt 2013 Platt JM, Ryvkin P, Wanat JJ, et al., Johnson FB. Rap1 relocalization contributes to the chromatin-mediated gene expression profile and pace of cell senescence. <i>Genes Dev</i> 2013;27(12):1406–20.	doi:10.1101/gad.218776.113 PMID 23756653 PMC3701195	–	21	1	21	19	2	3	ChIP
285	Becker 2015 Becker E, Liu Y, Lardenois A, et al., Primig M. Integrated RNA- and protein profiling of fermentation and respiration in diploid budding yeast provides insight into nutrient control of cell growth and development. <i>J Proteomics</i> 2015;119:30–44.	doi:10.1016/j.jprot.2015.01.015 PMID 25662576 PMC5469100	–	21	1	21	2	19	2	Microarray analysis - wt vs TF mutant
286	Ghavidel 2018 Ghavidel A, Baxi K, Prusinkiewicz M, et al., Harkness TA. Rapid Nuclear Exclusion of Hcm1 in Aging <i>Saccharomyces cerevisiae</i> Leads to Vacuolar Alkalization and Replicative Senescence. <i>G3 (Bethesda)</i> 2018;8(5):1579–1592.	doi:10.1534/g3.118.200161 PMID 29519938 PMC5940150	–	21	3	19	4	17	3	RT-PCR - wt vs TF mutant
287	Rottensteiner 2002 Rottensteiner H, Palmieri L, Hartig A, et al., Gurvitz A. The peroxisomal transporter gene ANT1 is regulated by a deviant oleate response element (ORE): characterization of the signal for fatty acid induction. <i>Biochem J</i> 2002;365(Pt 1):109–17.	doi:10.1042/BJ20011495 PMID 12071844 PMC1222661	–	20	2	6	4	16	6	Western blot - wt vs TF mutant
288	Ho 2006 Ho SW, Jona G, Chen CT, Johnston M, Snyder M. Linking DNA-binding proteins to their recognition sequences by using protein microarrays. <i>Proc Natl Acad Sci U S A</i> 2006;103(26):9940–5.	doi:10.1073/pnas.0509185103 PMID 16785442 PMC1502558	–	20	1	20	20	–	2	ChIP
289	Tachibana 2007 Tachibana C, Biddick R, Law GL, Young ET. A poised initiation complex is activated by SNF1. <i>J Biol Chem</i> 2007;282(52):37308–15.	doi:10.1074/jbc.M707363200 PMID 17974563	–	20	2	20	20	–	2	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
290	Foster 2013 Foster HA, Cui M, Naveenathayalan A, Unden H, Schwanbeck R, Höfken T. The zinc cluster protein Sut1 contributes to filamentation in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2013;12(2):244–53.	doi:10.1128/EC.00214-12 PMID 23223039 PMC3571311	–	20	3	20	–	20	2	lacZ - wt vs TF overexpression
291	Mehlgarten 2015 Mehlgarten C, Krijger JJ, Lemnian I, et al., Breunig KD. Divergent Evolution of the Transcriptional Network Controlled by Snf1-Interacting Protein Sip4 in Budding Yeasts. <i>PLoS One</i> 2015;10(10):e0139464.	doi:10.1371/journal.pone.0139464 PMID 26440109 PMC4634231	–	20	2	20	–	20	1	RT-PCR - wt vs TF mutant
292	Mostovoy 2016 Mostovoy Y, Thiemicke A, Hsu TY, Brem RB. The Role of Transcription Factors at Antisense-Expressing Gene Pairs in Yeast. <i>Genome Biol Evol</i> 2016;8(6):1748–61.	doi:10.1093/gbe/evw104 PMID 27190003 PMC4943177	–	20	2	20	1	19	3	RT-PCR - wt vs TF mutant
293	Zhang 2017 Zhang T, Bu P, Zeng J, Vancura A. Increased heme synthesis in yeast induces a metabolic switch from fermentation to respiration even under conditions of glucose repression. <i>J Biol Chem</i> 2017;292(41):16942–16954.	doi:10.1074/jbc.M117.790923 PMID 28830930 PMC5641876	–	20	4	16	10	10	4	ChIP
294	Iacovella 2018 Iacovella MG, Bremang M, Basha O, et al., De Wulf P. Integrating Rio1 activities discloses its nutrient-activated network in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Res</i> 2018;46(15):7586–7611.	doi:10.1093/nar/gky618 PMID 30011030 PMC6125641	–	20	1	10	–	20	4	RT-PCR - wt vs TF mutant
295	Le Crom 2002 Le Crom S, Devaux F, Marc P, Zhang X, Moye-Rowley WS, Jacq C. New insights into the pleiotropic drug resistance network from genome-wide characterization of the YRR1 transcription factor regulation system. <i>Mol Cell Biol</i> 2002;22(8):2642–9.	doi:10.1128/MCB.22.8.2642-2649.2002 PMID 11909958 PMC133742	–	19	1	19	–	19	2	Microarray analysis - wt vs TF mutant
296	Tuttle 2003 Tuttle MS, Radisky D, Li L, Kaplan J. A dominant allele of PDR1 alters transition metal resistance in yeast. <i>J Biol Chem</i> 2003;278(2):1273–80.	doi:10.1074/jbc.M209631200 PMID 12411429	–	19	1	19	–	19	3	Microarray analysis - wt vs TF mutant
297	Conesa 2005 Conesa C, Ruotolo R, Soularue P, et al., Dieci G. Modulation of yeast genome expression in response to defective RNA polymerase III-dependent transcription. <i>Mol Cell Biol</i> 2005;25(19):8631–42.	doi:10.1128/MCB.25.19.8631-8642.2005 PMID 16166643 PMC1265737	–	19	1	19	–	19	1	Microarray analysis - wt vs TF mutant
298	Chen 2010 Chen L, Lopes JM. Multiple bHLH proteins regulate CIT2 expression in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 2010;27(6):345–59.	doi:10.1002/yea.1757 PMID 20162531	–	18	8	11	–	18	15	lacZ - wt vs TF mutant
299	Joo 2011 Joo YJ, Kim JH, Kang UB, Yu MH, Kim J. Gcn4p-mediated transcriptional repression of ribosomal protein genes under amino-acid starvation. <i>EMBO J</i> 2011;30(5):859–72.	doi:10.1038/emboj.2010.332 PMID 21183953 PMC3049204	–	18	3	18	13	5	2	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
300	Fei 2011 Fei W, Zhong L, Ta MT, Shui G, Wenk MR, Yang H. The size and phospholipid composition of lipid droplets can influence their proteome. <i>Biochem Biophys Res Commun</i> 2011;415(3):455–62.	doi:10.1016/j.bbrc.2011.10.091 PMID 22057011	–	18	1	18	–	18	1	Proteomics - wt vs TF mutant
301	Parua 2014 Parua PK, Dombek KM, Young ET. Yeast 14-3-3 protein functions as a comodulator of transcription by inhibiting coactivator functions. <i>J Biol Chem</i> 2014;289(51):35542–60.	doi:10.1074/jbc.M114.592287 PMID 25355315 PMC4271238	–	18	3	18	–	18	2	RT-PCR - wt vs TF mutant
302	Solís 2016 Solís EJ, Pandey JP, Zheng X, et al., Denic V. Defining the Essential Function of Yeast Hsf1 Reveals a Compact Transcriptional Program for Maintaining Eukaryotic Proteostasis. <i>Mol Cell</i> 2016;63(1):60–71.	doi:10.1016/j.molcel.2016.05.014 PMID 27320198 PMC4938784	–	18	1	18	–	18	1	Net-seq
303	Wang 2018 Wang SQ, Wang T, Liu JF, Deng L, Wang F. Overexpression of Ecm22 improves ergosterol biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>Lett Appl Microbiol</i> 2018;67(5):484–490.	doi:10.1111/lam.13061 PMID 30098030	–	18	1	18	–	18	1	RNA-seq - wt vs TF mutant
304	Kuo 1994 Kuo MH, Grayhack E. A library of yeast genomic MCM1 binding sites contains genes involved in cell cycle control, cell wall and membrane structure, and metabolism. <i>Mol Cell Biol</i> 1994;14(1):348–59.	doi:10.1128/mcb.14.1.348-359.1994 PMID 8264602 PMC358384	–	17	1	17	17	–	2	Library of genomic binding sites
305	Daugherty 1993 Daugherty JR, Rai R, el Berry HM, Cooper TG. Regulatory circuit for responses of nitrogen catabolic gene expression to the GLN3 and DAL80 proteins and nitrogen catabolite repression in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1993;175(1):64–73.	doi:10.1128/jb.175.1.64-73.1993 PMID 8416910 PMC196097	–	17	2	17	–	17	1	Northern blot - wt vs TF mutant
306	Hikkel 2003 Hikkel I, Lucau-Danila A, Delaveau T, Marc P, Devaux F, Jacq C. A general strategy to uncover transcription factor properties identifies a new regulator of drug resistance in yeast. <i>J Biol Chem</i> 2003;278(13):11427–32.	doi:10.1074/jbc.M208549200 PMID 12529331	–	17	1	17	1	16	2	Microarray analysis - wt vs TF mutant
307	Rubio-Teixeira 2007 Rubio-Teixeira M. Urmylation controls Nilp and Gln3p-dependent expression of nitrogen-catabolite repressed genes in <i>Saccharomyces cerevisiae</i> . <i>FEBS Lett</i> 2007;581(3):541–50.	doi:10.1016/j.febslet.2007.01.019 PMID 17254574	–	17	3	6	–	17	3	lacZ - wt vs TF mutant
308	Iliina 2008 Iliina Y, Sloma E, Maciaszczyk-Dziubinska E, et al., Tamás MJ. Characterization of the DNA-binding motif of the arsenic-responsive transcription factor Yap8p. <i>Biochem J</i> 2008;415(3):467–75.	doi:10.1042/BJ20080713 PMID 18593383	–	17	2	17	3	14	3	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
309	Munding 2013 Munding EM, Shiue L, Katzman S, Donohue JP, Ares M Jr. Competition between pre-mRNAs for the splicing machinery drives global regulation of splicing. <i>Mol Cell</i> 2013;51(3):338–48.	doi:10.1016/j.molcel.2013.06.012 PMID 23891561 PMC3771316	–	17	2	17	–	17	3	RNA-seq - wt vs TF mutant
310	Khakhina 2015 Khakhina S, Johnson SS, Manoharlal R, et al., Moye-Rowley WS. Control of Plasma Membrane Permeability by ABC Transporters. <i>Eukaryot Cell</i> 2015;14(5):442–53.	doi:10.1128/EC.00021-15 PMID 25724885 PMC4421010	GSE65264	17	1	17	–	17	1	Microarray analysis - wt vs TF mutant
311	Rajvanshi 2017 Rajvanshi PK, Arya M, Rajasekharan R. The stress-regulatory transcription factors Msn2 and Msn4 regulate fatty acid oxidation in budding yeast. <i>J Biol Chem</i> 2017;292(45):18628–18643.	doi:10.1074/jbc.M117.801704 PMID 28924051 PMC5682970	–	17	2	12	7	10	1	RT-PCR - wt vs TF mutant
312	Lundin 1994 Lundin M, Nehlin JO, Ronne H. Importance of a flanking AT-rich region in target site recognition by the GC box-binding zinc finger protein MIG1. <i>Mol Cell Biol</i> 1994;14(3):1979–85.	doi:10.1128/mcb.14.3.1979-1985.1994 PMID 8114729 PMC358557	–	16	1	13	16	–	2	EMSA
313	DeRisi 1997 DeRisi JL, Iyer VR, Brown PO. Exploring the metabolic and genetic control of gene expression on a genomic scale. <i>Science</i> 1997;278(5338):680–6.	doi:10.1126/science.278.5338.680 PMID 9381177	no GEO	16	1	16	–	16	1	Microarray analysis - wt vs TF overexpression
314	Rep 1999 Rep M, Reiser V, Gartner U, et al., Ruis H. Osmotic stress-induced gene expression in <i>Saccharomyces cerevisiae</i> requires Msn1p and the novel nuclear factor Hot1p. <i>Mol Cell Biol</i> 1999;19(8):5474–85.	doi:10.1128/MCB.19.8.5474 PMID 10409737 PMC84389	–	16	4	16	–	16	5	Northern blot - wt vs TF mutant
315	Puig 2000 Puig S, Pérez-Ortín JE. Stress response and expression patterns in wine fermentations of yeast genes induced at the diauxic shift. <i>Yeast</i> 2000;16(2):139–48.	doi:10.1002/(SICI)1097-0061(20000130)16:2<139::AID-YEA512>3.0.CO;2-J PMID 10641036	–	16	2	16	–	16	1	Northern blot - wt vs TF mutant
316	Klinkenberg 2006 Klinkenberg LG, Webb T, Zitomer RS. Synergy among differentially regulated repressors of the ribonucleotide diphosphate reductase genes of <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2006;5(7):1007–17.	doi:10.1128/EC.00045-06 PMID 16835445 PMC1489293	–	16	3	9	–	16	2	lacZ - wt vs TF mutant
317	Teixeira 2008 Teixeira MC, Dias PJ, Simões T, Sá-Correia I. Yeast adaptation to mancozeb involves the up-regulation of FLR1 under the coordinate control of Yap1, Rpn4, Pdr3, and Yrr1. <i>Biochem Biophys Res Commun</i> 2008;367(2):249–55.	doi:10.1016/j.bbrc.2007.12.056 PMID 18086556	–	16	5	16	–	16	2	RT-PCR - wt vs TF mutant
318	Ferrezuelo 2010 Ferrezuelo F, Colomina N, Futcher B, Aldea M. The transcriptional network activated by Cln3 cyclin at the G1-to-S transition of the yeast cell cycle. <i>Genome Biol</i> 2010;11(6):R67.	doi:10.1186/gb-2010-11-6-r67 PMID 20573214 PMC2911115	–	16	2	8	16	–	2	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
319	van der Felden 2014 van der Felden J, Weisser S, Brückner S, Lenz P, Mösch HU. The transcription factors Tec1 and Ste12 interact with coregulators Msa1 and Msa2 to activate adhesion and multicellular development. <i>Mol Cell Biol</i> 2014;34(12):2283–93.	doi:10.1128/MCB.01599-13 PMID 24732795 PMC4054286	–	16	4	16	5	11	1	Microarray analysis - wt vs TF mutant
320	Jin 2014 Jin M, He D, Backues SK, et al., Klionsky DJ. Transcriptional regulation by Pho23 modulates the frequency of autophagosome formation. <i>Curr Biol</i> 2014;24(12):1314–1322.	doi:10.1016/j.cub.2014.04.048 PMID 24881874 PMC4169046	–	16	2	16	–	16	3	RNA-seq - wt vs TF mutant
321	Vido 2001 Vido K, Spector D, Lagniel G, Lopez S, Toledano MB, Labarre J. A proteome analysis of the cadmium response in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2001;276(11):8469–74.	doi:10.1074/jbc.M008708200 PMID 11078740	–	15	2	15	–	15	1	Proteomics - wt vs TF mutant
322	Hall 2006 Hall DB, Wade JT, Struhl K. An HMG protein, Hmo1, associates with promoters of many ribosomal protein genes and throughout the rRNA gene locus in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2006;26(9):3672–9.	doi:10.1128/MCB.26.9.3672-3679.2006 PMID 16612005 PMC1447432	–	15	3	15	15	–	2	ChIP
323	Schlecht 2008 Schlecht U, Erb I, Demougin P, et al., Primig M. Genome-wide expression profiling, in vivo DNA binding analysis, and probabilistic motif prediction reveal novel Abf1 target genes during fermentation, respiration, and sporulation in yeast. <i>Mol Biol Cell</i> 2008;19(5):2193–207.	doi:10.1091/mbc.e07-12-1242 PMID 18305101 PMC2366881	–	15	1	15	–	15	1	Microarray analysis - wt vs TF mutant
324	Thepnok 2014 Thepnok P, Ratanakhanokchai K, Soontorngun N. The novel zinc cluster regulator Tog1 plays important roles in oleate utilization and oxidative stress response in <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2014;450(4):1276–82.	doi:10.1016/j.bbrc.2014.06.128 PMID 24998441	–	15	1	8	7	8	1	RT-PCR - wt vs TF mutant
325	Serratore 2018 Serratore ND, Baker KM, Macadlo LA, et al., Briggs SD. A Novel Sterol-Signaling Pathway Governs Azole Antifungal Drug Resistance and Hypoxic Gene Repression in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2018;208(3):1037–1055.	doi:10.1534/genetics.117.300554 PMID 29263028 PMC5844321	–	15	3	6	2	13	11	RT-PCR - wt vs TF mutant
326	Cha 2021 Cha S, Hong CP, Kang HA, Hahn JS. Differential activation mechanisms of two isoforms of Gcr1 transcription factor generated from spliced and un-spliced transcripts in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Res</i> 2021;49(2):745–759.	doi:10.1093/nar/gkaa1221 PMID 33367825 PMC7826247	–	15	2	11	8	7	3	ChIP
327	Sato 2021 Sato A, Kimura T, Hondo K, Kawano-Kawada M, Sekito T. The vacuolar amino acid transport system is a novel, direct target of GATA transcription factors. <i>Biosci Biotechnol Biochem</i> 2021;85(3):587–599.	doi:10.1093/bbb/zbaa041 PMID 33624780	–	15	4	12	1	14	3	RT-PCR - wt vs. TF double mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
328	Lucau-Danila 2003 Lucau-Danila A, Delaveau T, Lelandais G, Devaux F, Jacq C. Competitive promoter occupancy by two yeast paralogous transcription factors controlling the multidrug resistance phenomenon. <i>J Biol Chem</i> 2003;278(52):52641–50.	doi:10.1074/jbc.M309580200 PMID 14512416	–	14	1	14	–	14	1	Microarray analysis - wt vs TF mutant
329	Aow 2013 Aow JS, Xue X, Run JQ, Lim GF, Goh WS, Clarke ND. Differential binding of the related transcription factors Pho4 and Cbf1 can tune the sensitivity of promoters to different levels of an induction signal. <i>Nucleic Acids Res</i> 2013;41(9):4877–87.	doi:10.1093/nar/gkt210 PMID 23558744 PMC3643608	GSE26770	14	2	14	14	–	2	ChIP
330	Bernard 2015 Bernard A, Jin M, González-Rodríguez P, et al., Klionsky DJ. Rph1/KDM4 mediates nutrient-limitation signaling that leads to the transcriptional induction of autophagy. <i>Curr Biol</i> 2015;25(5):546–55.	doi:10.1016/j.cub.2014.12.049 PMID 25660547 PMC4348152	–	14	2	14	–	14	1	RT-PCR - wt vs TF mutant
331	Ling 2015 Ling H, Pratomo Juwono NK, Teo WS, Liu R, Leong SS, Chang MW. Engineering transcription factors to improve tolerance against alkane biofuels in <i>Saccharomyces cerevisiae</i> . <i>Biotechnol Biofuels</i> 2015;8:231.	doi:10.1186/s13068-015-0411-z PMID 26719765 PMC4696261	–	14	2	14	–	14	2	RT-PCR - wt vs TF mutant
332	Zhao 2016 Zhao X, Zou H, Chen J, Du G, Zhou J. The modification of Gat1p in nitrogen catabolite repression to enhance non-preferred nitrogen utilization in <i>Saccharomyces cerevisiae</i> . <i>Sci Rep</i> 2016;6:21603.	doi:10.1038/srep21603 PMID 26899143 PMC4761935	–	14	1	14	–	14	1	RT-PCR - wt vs TF mutant
333	Fermi 2016 Fermi B, Bosio MC, Dieci G. Promoter architecture and transcriptional regulation of Abf1-dependent ribosomal protein genes in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Res</i> 2016;44(13):6113–26.	doi:10.1093/nar/gkw194 PMID 27016735 PMC5291244	–	14	2	14	14	–	3	RT-PCR - wt vs mutant promoter sequence
334	Kim 2016 Kim MS, Hahn JS. Role of CK2-dependent phosphorylation of Ifh1 and Crf1 in transcriptional regulation of ribosomal protein genes in <i>Saccharomyces cerevisiae</i> . <i>Biochim Biophys Acta</i> 2016;1859(8):1004–13.	doi:10.1016/j.bbagr.2016.06.003 PMID 27321754	–	14	2	14	7	7	3	ChIP
335	Shi 2016 Shi BJ. Decoding common and divergent cellular functions of the domains of forkhead transcription factors Fkh1 and Fkh2. <i>Biochem J</i> 2016;473(21):3855–3869.	doi:10.1042/BCJ20160609 PMID 27555611	–	14	3	14	5	9	4	Northern blot - wt vs TF mutant
336	Sunwoo 2021 Sunwoo IY, Sukwong P, Park YR, et al., Kim SK. Enhancement of Galactose Uptake from <i>Kappaphycus alvarezii</i> Using <i>Saccharomyces cerevisiae</i> through Deletion of Negative Regulators of GAL Genes. <i>Appl Biochem Biotechnol</i> 2021;193(2):577–588.	doi:10.1007/s12010-020-03434-3 PMID 33043399	–	14	4	14	–	14	1	RT-PCR - wt vs. TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
337	Jordá 2021 Jordá T, Rozès N, Puig S. Sterol Composition Modulates the Response of <i>Saccharomyces cerevisiae</i> to Iron Deficiency. <i>J Fungi (Basel)</i> 2021;7(11).	doi:10.3390/jof7110901 PMID 34829190 PMC8620032	–	14	3	9	–	14	3	RT-PCR - wt vs TF mutant
338	Huang 2002 Huang D, Moffat J, Andrews B. Dissection of a complex phenotype by functional genomics reveals roles for the yeast cyclin-dependent protein kinase Pho85 in stress adaptation and cell integrity. <i>Mol Cell Biol</i> 2002;22(14):5076–88.	doi:10.1128/MCB.22.14.5076-5088.2002 PMID 12077337 PMC139770	no GEO	13	1	13	–	13	1	Microarray analysis - wt vs TF mutant
339	Oliveira 2003 Oliveira EM, Martins AS, Carvajal E, Bon EP. The role of the GATA factors Gln3p, Nil1p, Dal80p and the Ure2p on ASP3 regulation in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 2003;20(1):31–7.	doi:10.1002/yea.930 PMID 12489124	–	13	3	3	–	13	4	Northern blot - wt vs TF mutant
340	Galgoczy 2004 Galgoczy DJ, Cassidy-Stone A, Llinás M, et al., Johnson AD. Genomic dissection of the cell-type-specification circuit in <i>Saccharomyces cerevisiae</i> . <i>Proc Natl Acad Sci U S A</i> 2004;101(52):18069–74.	doi:10.1073/pnas.0407611102 PMID 15604142 PMC535907	–	13	1	13	13	–	1	Chip-on-chip
341	Sasaki 2005 Sasaki H, Kishimoto T, Mizuno T, Shinzato T, Uemura H. Expression of GCR1, the transcriptional activator of glycolytic enzyme genes in the yeast <i>Saccharomyces cerevisiae</i> , is positively autoregulated by Gcr1p. <i>Yeast</i> 2005;22(4):305–19.	doi:10.1002/yea.1212 PMID 15789351	–	13	2	7	–	13	3	lacZ - wt vs TF mutant
342	Manzanares-Estreded 2017 Manzanares-Estreded S, Espí-Bardisa J, Alarcón B, Pascual-Ahuir A, Proft M. Multilayered control of peroxisomal activity upon salt stress in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2017;104(5):851–868.	doi:10.1111/mmi.13669 PMID 28321934	–	13	3	9	4	9	1	RT-PCR - wt vs TF mutant
343	Rajakumar 2020 Rajakumar S, Suriyagandhi V, Nachiappan V. Impairment of MET transcriptional activators, MET4 and MET31 induced lipid accumulation in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Res</i> 2020;20(5).	doi:10.1093/femsyr/foaa039 PMID 32648914	–	13	3	9	–	13	2	RT-PCR - wt vs. TF mutant
344	Chidambaram 2022 Chidambaram R, Ramachandran G, Rajasekharan R, Nachiappan V. Impairment of transcription factor Gcr1p binding motif perturbs OPI3 transcription in <i>Saccharomyces cerevisiae</i> . <i>J Cell Biochem</i> 2022;123(6):1032–1052.	doi:10.1002/jcb.30245 PMID 35416329	–	13	4	5	3	10	4	lacZ - wt vs TF mutant
345	Lu 2022 Lu Z, Wu Y, Chen Y, et al., Huang R. Role of spt23 in <i>Saccharomyces cerevisiae</i> thermal tolerance. <i>Appl Microbiol Biotechnol</i> 2022;106(9–10):3691–3705.	doi:10.1007/s00253-022-11920-3 PMID 35476152 PMC9151549	–	13	1	7	–	13	2	RT-PCR - wt vs. TF mutant
346	Dang 1994 Dang VD, Valens M, Bolotin-Fukuhara M, Daignan-Fornier B. A genetic screen to isolate genes regulated by the yeast CCAAT-box binding protein Hap2p. <i>Yeast</i> 1994;10(10):1273–83.	doi:10.1002/yea.320101004 PMID 7900416	–	12	3	9	–	12	2	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
347	Vik 2001 Vik A, Rine J. Upc2p and Ecm22p, dual regulators of sterol biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2001; 21(19):6395–405.	doi:10.1128/MCB.21.19.6395-6405.2001 PMID 11533229 PMC99787	–	12	2	4	2	10	5	lacZ - wt vs TF mutant
348	Kandror 2004 Kandror O, Bretschneider N, Kreydin E, Cavalieri D, Goldberg AL. Yeast adapt to near-freezing temperatures by STRE/Msn2,4-dependent induction of trehalose synthesis and certain molecular chaperones. <i>Mol Cell</i> 2004;13(6):771–81.	doi:10.1016/s1097-2765(04)00148-0 PMID 15053871	–	12	2	12	–	12	2	Northern blot - wt vs TF mutant
349	Nosaka 2005 Nosaka K, Onozuka M, Konno H, et al., Akaji K. Genetic regulation mediated by thiamin pyrophosphate-binding motif in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2005;58(2):467–79.	doi:10.1111/j.1365-2958.2005.04835.x PMID 16194233	–	12	2	9	–	12	3	Microarray analysis - wt vs TF mutant
350	Som 2005 Som I, Mitsch RN, Urbanowski JL, Rolfes RJ. DNA-bound Bas1 recruits Pho2 to activate ADE genes in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2005;4(10):1725–35.	doi:10.1128/EC.4.10.1725-1735.2005 PMID 16215179 PMC1265903	–	12	2	12	12	–	2	ChIP
351	Abraham 2005 Abraham DS, Vershon AK. N-terminal arm of Mcm1 is required for transcription of a subset of genes involved in maintenance of the cell wall. <i>Eukaryot Cell</i> 2005;4(11):1808–19.	doi:10.1128/EC.4.11.1808-1819.2005 PMID 16278448 PMC1287865	–	12	1	7	–	12	2	Northern blot - wt vs TF mutant
352	Ruiz 2008 Ruiz A, Serrano R, Ariño J. Direct regulation of genes involved in glucose utilization by the calcium/calcineurin pathway. <i>J Biol Chem</i> 2008;283(20):13923–33.	doi:10.1074/jbc.M708683200 PMID 18362157	–	12	1	12	9	3	1	ChIP
353	Gurvitz 2009 Gurvitz A, Suomi F, Rottensteiner H, Hiltunen JK, Dawes IW. Avoiding unscheduled transcription in shared promoters: <i>Saccharomyces cerevisiae</i> Sum1p represses the divergent gene pair SPS18-SPS19 through a midsporulation element (MSE). <i>FEMS Yeast Res</i> 2009;9(6):821–31.	doi:10.1111/j.1567-1364.2009.00527.x PMID 19583587 PMC2784042	–	12	3	9	–	12	1	RT-PCR - wt vs TF mutant
354	Pastor 2009 Pastor MM, Proft M, Pascual-Ahuir A. Mitochondrial function is an inducible determinant of osmotic stress adaptation in yeast. <i>J Biol Chem</i> 2009;284(44):30307–17.	doi:10.1074/jbc.M109.050682 PMID 19720830 PMC2781586	–	12	4	12	–	12	1	Northern blot - wt vs TF mutant
355	Darieva 2010 Darieva Z, Clancy A, Bulmer R, et al., Sharrocks AD. A competitive transcription factor binding mechanism determines the timing of late cell cycle-dependent gene expression. <i>Mol Cell</i> 2010;38(1):29–40.	doi:10.1016/j.molcel.2010.02.030 PMID 20385087 PMC3566586	–	12	4	12	12	–	3	ChIP
356	Shetty 2010 Shetty A, Lopes JM. Derepression of INO1 transcription requires cooperation between the Ino2p-Ino4p heterodimer and Cbf1p and recruitment of the ISW2 chromatin-remodeling complex. <i>Eukaryot Cell</i> 2010;9(12):1845–55.	doi:10.1128/EC.00144-10 PMID 20935143 PMC3008276	–	12	7	8	4	8	3	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
357	Munding 2010 Munding EM, Igel AH, Shiue L, Dorigi KM, Treviño LR, Ares M Jr. Integration of a splicing regulatory network within the meiotic gene expression program of <i>Saccharomyces cerevisiae</i> . <i>Genes Dev</i> 2010;24(23):2693–704.	doi:10.1101/gad.1977410 PMID 21123654 PMC2994042	GSE24675	12	1	12	1	11	2	RT-PCR - wt vs TF mutant
358	Hernández 2011 Hernández H, Aranda C, Riego L, González A. Gln3-Gcn4 hybrid transcriptional activator determines catabolic and biosynthetic gene expression in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2011;404(3):859–64.	doi:10.1016/j.bbrc.2010.12.075 PMID 21184740	–	12	2	12	12	–	1	ChIP
359	Elsztein 2011 Elsztein C, de Lucena RM, de Morais MA Jr. The resistance of the yeast <i>Saccharomyces cerevisiae</i> to the biocide polyhexamethylene biguanide: involvement of cell wall integrity pathway and emerging role for YAP1. <i>BMC Mol Biol</i> 2011;12:38.	doi:10.1186/1471-2199-12-38 PMID 21854579 PMC3175164	–	12	1	12	–	12	2	RT-PCR - wt vs TF mutant
360	Landry 2014 Landry BD, Mapa CE, Arsenault HE, Poti KE, Benanti JA. Regulation of a transcription factor network by Cdk1 coordinates late cell cycle gene expression. <i>EMBO J</i> 2014;33(9):1044–60.	doi:10.1002/embj.201386877 PMID 24714560 PMC4193936	GSE55121	12	3	12	12	–	2	ChIP
361	Yan 2015 Yan C, Zhang D, Raygoza Garay JA, Mwangi MM, Bai L. Decoupling of divergent gene regulation by sequence-specific DNA binding factors. <i>Nucleic Acids Res</i> 2015;43(15):7292–305.	doi:10.1093/nar/gkv618 PMID 26082499 PMC4551913	–	12	7	12	10	2	2	ChIP
362	Nakazawa 2018 Nakazawa N, Yanata H, Ito N, Kaneta E, Takahashi K. Oxidative stress tolerance of a spore clone isolated from Shirakami kodama yeast depends on altered regulation of Msn2 leading to enhanced expression of ROS-degrading enzymes. <i>J Gen Appl Microbiol</i> 2018;64(4):149–157.	doi:10.2323/jgam.2017.11.002 PMID 29607878	–	12	4	4	–	12	5	Northern blot - wt vs TF mutant
363	Kim 2019 Kim MS, Cho KH, Park KH, Jang J, Hahn JS. Activation of Haa1 and War1 transcription factors by differential binding of weak acid anions in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Res</i> 2019;47(3):1211–1224.	doi:10.1093/nar/gky1188 PMID 30476185 PMC6379682	–	12	2	5	2	10	7	RT-PCR - wt vs TF mutant
364	Yang 2019 Yang L, Zheng C, Chen Y, Shi X, Ying Z, Ying H. Nitric oxide increases biofilm formation in <i>Saccharomyces cerevisiae</i> by activating the transcriptional factor Mac1p and thereby regulating the transmembrane protein Ctr1. <i>Biotechnol Biofuels</i> 2019;12:30.	doi:10.1186/s13068-019-1359-1 PMID 30809273 PMC6375214	–	12	1	6	–	12	1	RT-PCR - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
365	Cheng 2021 Cheng Y, Zhu H, Du Z, et al., He X. Eukaryotic translation factor eIF5A contributes to acetic acid tolerance in <i>Saccharomyces cerevisiae</i> via transcriptional factor Ume6p. <i>Biotechnol Biofuels</i> 2021;14(1):38.	doi:10.1186/s13068-021-01885-2 PMID 33557922 PMC7869214	–	12	1	3	–	12	2	RT-PCR - wt vs TF mutant
366	Mizuno 2021 Mizuno T, Irie K. Msn2/4 transcription factors positively regulate expression of Atg39 ER-phagy receptor. <i>Sci Rep</i> 2021;11(1):11919.	doi:10.1038/s41598-021-91480-0 PMID 34099851 PMC8184937	–	12	2	4	–	12	5	RT-PCR - wt vs TF mutant and Western blot - WT vs TF Mutant
367	Daignan-Fornier 1994 Daignan-Fornier B, Valens M, Lemire BD, Bolotin-Fukuhara M. Structure and regulation of SDH3, the yeast gene encoding the cytochrome b560 subunit of respiratory complex II. <i>J Biol Chem</i> 1994;269(22):15469–72.	PMID 8195189	–	11	3	10	–	11	2	lacZ - wt vs TF mutant
368	Grishin 1998 Grishin AV, Rothenberg M, Downs MA, Blumer KJ. Mot3, a Zn finger transcription factor that modulates gene expression and attenuates mating pheromone signaling in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 1998;149(2):879–92.	doi:10.1093/genetics/149.2.879 PMID 9611199 PMC1460180	–	11	1	11	–	11	1	lacZ - wt vs TF mutant
369	Chu 1998 Chu S, Herskowitz I. Gametogenesis in yeast is regulated by a transcriptional cascade dependent on Ndt80. <i>Mol Cell</i> 1998;1(5):685–96.	doi:10.1016/s1097-2765(00)80068-4 PMID 9660952	–	11	2	11	–	11	1	Northern blot - wt vs TF mutant
370	Zaragoza 2001 Zaragoza O, Vincent O, Gancedo JM. Regulatory elements in the FBP1 promoter respond differently to glucose-dependent signals in <i>Saccharomyces cerevisiae</i> . <i>Biochem J</i> 2001;359(Pt 1):193–201.	doi:10.1042/0264-6021:3590193 PMID 11563983 PMC1222135	–	11	5	9	2	9	8	lacZ - wt vs TF mutant
371	Akache 2002 Akache B, Turcotte B. New regulators of drug sensitivity in the family of yeast zinc cluster proteins. <i>J Biol Chem</i> 2002;277(24):21254–60.	doi:10.1074/jbc.M202566200 PMID 11943786	–	11	7	11	–	11	1	Northern blot - wt vs TF mutant
372	Wysocki 2004 Wysocki R, Fortier PK, Maciaszczyk E, et al., Tamás MJ. Transcriptional activation of metalloid tolerance genes in <i>Saccharomyces cerevisiae</i> requires the AP-1-like proteins Yap1p and Yap8p. <i>Mol Biol Cell</i> 2004;15(5):2049–60.	doi:10.1091/mbc.e03-04-0236 PMID 14978214 PMC404003	–	11	2	11	2	9	3	Northern blot - wt vs TF mutant
373	Georis 2009 Georis I, Feller A, Vierendeels F, Dubois E. The yeast GATA factor Gat1 occupies a central position in nitrogen catabolite repression-sensitive gene activation. <i>Mol Cell Biol</i> 2009;29(13):3803–15.	doi:10.1128/MCB.00399-09 PMID 19380492 PMC2698774	–	11	4	11	4	7	4	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
374	Castro-Prego 2009 Castro-Prego R, Lamas-Maceiras M, Soengas P, Carneiro I, González-Siso I, Cerdán ME. Regulatory factors controlling transcription of <i>Saccharomyces cerevisiae</i> IXR1 by oxygen levels: a model of transcriptional adaptation from aerobiosis to hypoxia implicating ROX1 and IXR1 cross-regulation. <i>Biochem J</i> 2009;425(1):235–43.	doi:10.1042/BJ20091500 PMID 19807692	–	11	5	9	5	6	3	Northern blot - wt vs TF mutant
375	Wan 2013 Wan Y, Arens CE, Wang S, et al., Liu H. Role of the repressor Oaf3p in the recruitment of transcription factors and chromatin dynamics during the oleate response. <i>Biochem J</i> 2013;449(2):507–17.	doi:10.1042/BJ20121029 PMID 23088601	–	11	4	11	8	3	1	ChIP
376	Mazzola 2015 Mazzola D, Pimentel C, Caetano S, et al., Rodrigues-Pousada C. Inhibition of Yap2 activity by MAPKAP kinase Rck1 affects yeast tolerance to cadmium. <i>FEBS Lett</i> 2015;589(19 Pt B):2841–9.	doi:10.1016/j.febslet.2015.07.049 PMID 26296316	–	11	2	11	–	11	2	RT-PCR - wt vs TF mutant
377	Höfken 2017 Höfken T. Ecm22 and Upc2 regulate yeast mating through control of expression of the mating genes PRM1 and PRM4. <i>Biochem Biophys Res Commun</i> 2017;493(4):1485–1490.	doi:10.1016/j.bbrc.2017.10.005 PMID 28986257	–	11	3	5	–	11	7	lacZ - wt vs TF mutant
378	Musa 2018 Musa M, Perić M, Bou Dib P, et al., Kriško A. Heat-induced longevity in budding yeast requires respiratory metabolism and glutathione recycling. <i>Aging (Albany NY)</i> 2018;10(9):2407–2427.	doi:10.18632/aging.101560 PMID 30227387 PMC6188503	–	11	1	11	–	11	1	RT-PCR - wt vs TF mutant
379	Karpov 2019 Karpov DS, Spasskaya DS, Nadolinskaia NI, Tutyaeva VV, Lysov YP, Karpov VL. Deregulation of the 19S proteasome complex increases yeast resistance to 4-NQO and oxidative stress via upregulation of Rpn4- and proteasome-dependent stress responsive genes. <i>FEMS Yeast Res</i> 2019;19(2).	doi:10.1093/femsyr/foz002 PMID 30629175	–	11	1	10	–	11	3	RT-PCR - wt vs TF mutant
380	Yan 2020 Yan H, Fang T, Xu H, Jiang L. The pH-sensing Rim101 pathway positively regulates the transcriptional expression of the calcium pump gene PMR1 to affect calcium sensitivity in budding yeast. <i>Biochem Biophys Res Commun</i> 2020;532(3):453–458.	doi:10.1016/j.bbrc.2020.08.083 PMID 32891431	–	11	2	4	–	11	2	LacZ - WT vs. TF double mutant
381	Lupo 2021 Lupo O, Krieger G, Jonas F, Barkai N. Accumulation of cis- and trans-regulatory variations is associated with phenotypic divergence of a complex trait between yeast species. <i>G3 (Bethesda)</i> 2021;11(2).	doi:10.1093/g3journal/jkab016 PMID 33609368 PMC8022985	–	11	2	11	11	–	1	ChEC-seq

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
382	Dzanaeva 2021 Dzanaeva L, Kruk B, Ruchala J, Sibirny A, Dmytruk K. The impact of transcription factors Znf1, Sip4, Adr1, Tup1, and Hap4 on xylose alcoholic fermentation in the engineered yeast <i>Saccharomyces cerevisiae</i> . <i>Antonie Van Leeuwenhoek</i> 2021; 114(9):1373–1385.	doi:10.1007/s10482-021-01607-6 PMID 34170419	–	11	4	11	–	11	1	RT-PCR - wt vs. TF mutant
383	Buchman 1988 Buchman AR, Lue NF, Kornberg RD. Connections between transcriptional activators, silencers, and telomeres as revealed by functional analysis of a yeast DNA-binding protein. <i>Mol Cell Biol</i> 1988;8(12):5086–99.	doi:10.1128/mcb.8.12.5086-5099.1988 PMID 3072472 PMC365610	–	10	1	10	7	–	2	EMSA
384	Martínez-Pastor 1996 Martínez-Pastor MT, Marchler G, Schüller C, Marchler-Bauer A, Ruis H, Estruch F. The <i>Saccharomyces cerevisiae</i> zinc finger proteins Msn2p and Msn4p are required for transcriptional induction through the stress response element (STRE). <i>EMBO J</i> 1996;15(9):2227–35.	PMID 8641288 PMC450147	–	10	2	10	–	10	2	Northern blot - wt vs TF overexpression
385	Ramil 2000 Ramil E, Agrimonti C, Shechter E, Gervais M, Guiard B. Regulation of the CYB2 gene expression: transcriptional co-ordination by the Hap1p, Hap2/3/4/5p and Adr1p transcription factors. <i>Mol Microbiol</i> 2000;37(5):1116–32.	doi:10.1046/j.1365-2958.2000.02065.x PMID 10972830	–	10	6	10	1	9	2	lacZ - wt vs TF mutant
386	Goldmark 2000 Goldmark JP, Fazzio TG, Estep PW, Church GM, Tsukiyama T. The Isw2 chromatin remodeling complex represses early meiotic genes upon recruitment by Ume6p. <i>Cell</i> 2000;103(3):423–33.	doi:10.1016/s0092-8674(00)00134-3 PMID 11081629	–	10	1	9	1	9	1	Northern blot - wt vs TF mutant
387	Blaiseau 2001 Blaiseau PL, Lesuisse E, Camadro JM. Aft2p, a novel iron-regulated transcription activator that modulates, with Aft1p, intracellular iron use and resistance to oxidative stress in yeast. <i>J Biol Chem</i> 2001;276(36):34221–6.	doi:10.1074/jbc.M104987200 PMID 11448968	–	10	2	3	–	10	2	Northern blot - wt vs TF mutant
388	Serrano 2002 Serrano R, Ruiz A, Bernal D, Chambers JR, Ariño J. The transcriptional response to alkaline pH in <i>Saccharomyces cerevisiae</i> : evidence for calcium-mediated signalling. <i>Mol Microbiol</i> 2002;46(5):1319–33.	doi:10.1046/j.1365-2958.2002.03246.x PMID 12453218	–	10	5	10	–	10	1	lacZ - wt vs TF mutant
389	Wong 2003 Wong CM, Ching YP, Zhou Y, Kung HF, Jin DY. Transcriptional regulation of yeast peroxiredoxin gene TSA2 through Hap1p, Rox1p, and Hap2/3/5p. <i>Free Radic Biol Med</i> 2003;34(5):585–97.	doi:10.1016/s0891-5849(02)01354-0 PMID 12614847	–	10	10	10	–	10	3	RT-PCR - wt vs TF mutant
390	Kolaczekowski 2004 Kolaczekowski M, Kolaczekowska A, Gaigg B, Schneiter R, Moye-Rowley WS. Differential regulation of ceramide synthase components LAC1 and LAG1 in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2004;3(4):880–92.	doi:10.1128/EC.3.4.880-892.2004 PMID 15302821 PMC500886	–	10	4	8	1	9	4	Western blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
391	van Dyk 2005 van Dyk D, Pretorius IS, Bauer FF. Mss11p is a central element of the regulatory network that controls FLO11 expression and invasive growth in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2005;169(1):91–106.	doi:10.1534/genetics.104.033704 PMID 15466424 PMC1448879	–	10	10	10	–	10	1	Northern blot - wt vs TF mutant
392	Zaim 2005 Zaim J, Speina E, Kierzek AM. Identification of new genes regulated by the Crt1 transcription factor, an effector of the DNA damage checkpoint pathway in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2005;280(1):28–37.	doi:10.1074/jbc.M404669200 PMID 15494396	–	10	1	6	–	10	2	RT-PCR - wt vs TF mutant
393	Fernandes 2005 Fernandes AR, Mira NP, Vargas RC, Canelhas I, Sá-Correia I. <i>Saccharomyces cerevisiae</i> adaptation to weak acids involves the transcription factor Haa1p and Haa1p-regulated genes. <i>Biochem Biophys Res Commun</i> 2005;337(1):95–103.	doi:10.1016/j.bbrc.2005.09.010 PMID 16176797	–	10	1	8	–	10	3	Western blot - wt vs TF mutant
394	Platara 2006 Platara M, Ruiz A, Serrano R, Palomino A, Moreno F, Ariño J. The transcriptional response of the yeast Na(+)-ATPase ENA1 gene to alkaline stress involves three main signaling pathways. <i>J Biol Chem</i> 2006;281(48):36632–42.	doi:10.1074/jbc.M606483200 PMID 17023428	–	10	5	5	–	10	2	lacZ - wt vs TF mutant
395	Fichtner 2007 Fichtner L, Schulze F, Braus GH. Differential Flo8p-dependent regulation of FLO1 and FLO11 for cell-cell and cell-substrate adherence of <i>S. cerevisiae</i> S288c. <i>Mol Microbiol</i> 2007;66(5):1276–89.	doi:10.1111/j.1365-2958.2007.06014.x PMID 18001350 PMC2780560	–	10	5	10	–	10	1	RT-PCR - wt vs TF mutant
396	Kolaczowska 2008 Kolaczowska A, Kolaczowski M, Goffeau A, Moye-Rowley WS. Compensatory activation of the multidrug transporters Pdr5p, Snq2p, and Yor1p by Pdr1p in <i>Saccharomyces cerevisiae</i> . <i>FEBS Lett</i> 2008;582(6):977–83.	doi:10.1016/j.febslet.2008.02.045 PMID 18307995 PMC2288637	–	10	2	10	–	10	1	lacZ - wt vs TF mutant
397	Shi 2013 Shi L, Tu BP. Acetyl-CoA induces transcription of the key G1 cyclin CLN3 to promote entry into the cell division cycle in <i>Saccharomyces cerevisiae</i> . <i>Proc Natl Acad Sci U S A</i> 2013;110(18):7318–23.	doi:10.1073/pnas.1302490110 PMID 23589851 PMC3645525	–	10	4	10	10	–	3	ChIP
398	Maoz 2015 Maoz N, Gabay O, Waldman Ben-Asher H, Cohen HY. The yeast forkhead HCM1 controls life span independent of calorie restriction. <i>J Gerontol A Biol Sci Med Sci</i> 2015;70(4):444–53.	doi:10.1093/gerona/glu059 PMID 24835838	–	10	1	10	–	10	1	RT-PCR - wt vs TF overexpression
399	Parnell 2014 Parnell EJ, Yu Y, Lucena R, et al., Stillman DJ. The Rts1 regulatory subunit of PP2A phosphatase controls expression of the HO endonuclease via localization of the Ace2 transcription factor. <i>J Biol Chem</i> 2014;289(51):35431–7.	doi:10.1074/jbc.M114.611715 PMID 25352596 PMC4271228	–	10	3	10	1	9	2	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
400	Tsang 2015 Tsang F, James C, Kato M, et al., Lin SJ. Reduced Ssy1-Ptr3-Ssy5 (SPS) signaling extends replicative life span by enhancing NAD+ homeostasis in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2015;290(20):12753–64.	doi:10.1074/jbc.M115.644534 PMID 25825491 PMC4432292	–	10	3	10	2	8	1	RT-PCR - wt vs TF mutant
401	Gomar-Alba 2015 Gomar-Alba M, Amaral C, Artacho A, et al., lí del Olmo M. The C-terminal region of the Hot1 transcription factor binds GGGACAAA-related sequences in the promoter of its target genes. <i>Biochim Biophys Acta</i> 2015;1849(12):1385–97.	doi:10.1016/j.bbagr.2015.10.007 PMID 26470684	–	10	1	10	10	–	1	ChIP
402	Romero 2019 Romero AM, Ramos-Alonso L, Montellá-Manuel S, et al., Puig S. A genome-wide transcriptional study reveals that iron deficiency inhibits the yeast TORC1 pathway. <i>Biochim Biophys Acta Gene Regul Mech</i> 2019;1862(9):194414.	doi:10.1016/j.bbagr.2019.194414 PMID 31394264	GSE127875	10	2	6	4	6	2	RT-PCR - wt vs TF mutant
403	Wang 2020 Wang YP, Wei XQ, Guo XW, Xiao DG. Effect of the Deletion of Genes Related to Amino Acid Metabolism on the Production of Higher Alcohols by <i>Saccharomyces cerevisiae</i> . <i>Biomed Res Int</i> 2020;2020:6802512.	doi:10.1155/2020/6802512 PMID 33204707 PMC7665916	–	10	2	10	–	10	1	RT-PCR - wt vs. TF mutant
404	Barba-Aliaga 2020 Barba-Aliaga M, Villarroel-Vicente C, Stanciu A, Corman A, Martínez-Pastor MT, Alepuz P. Yeast Translation Elongation Factor eIF5A Expression Is Regulated by Nutrient Availability through Different Signalling Pathways. <i>Int J Mol Sci</i> 2020; 22(1).	doi:10.3390/ijms22010219 PMID 33379337 PMC7794953	–	10	2	5	–	10	5	RT-PCR - wt vs. TF mutant
405	Yamada 2022 Yamada Y. UME6 Is Involved in the Suppression of Basal Transcription of ABC Transporters and Drug Resistance in the $\rho(+)$ Cells of <i>Saccharomyces cerevisiae</i> . <i>Microorganisms</i> 2022;10(3).	doi:10.3390/microorganisms10030601 PMID 35336175 PMC8953597	–	10	2	9	–	10	2	RT-PCR - wt vs. TF mutant
406	Kinney 1989 Kinney DM, Lusty CJ. Arginine restriction induced by delta-N-(phosphonacetyl)-L-ornithine signals increased expression of HIS3, TRP5, CPA1, and CPA2 in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1989;9(11):4882–8.	doi:10.1128/mcb.9.11.4882-4888.1989 PMID 2689869 PMC363638	–	9	1	4	–	9	3	lacZ - wt vs TF overexpression
407	Ozcan 1995 Ozcan S, Johnston M. Three different regulatory mechanisms enable yeast hexose transporter (HXT) genes to be induced by different levels of glucose. <i>Mol Cell Biol</i> 1995;15(3):1564–72.	doi:10.1128/MCB.15.3.1564 PMID 7862149 PMC230380	–	9	2	9	–	9	2	lacZ - wt vs TF mutant
408	Talibi 1995 Talibi D, Grenson M, André B. Cis- and trans-acting elements determining induction of the genes of the gamma-aminobutyrate (GABA) utilization pathway in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Res</i> 1995;23(4):550–7.	doi:10.1093/nar/23.4.550 PMID 7899074 PMC306719	–	9	4	9	–	9	2	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
409	Jackson 1996 Jackson JC, Lopes JM. The yeast UME6 gene is required for both negative and positive transcriptional regulation of phospholipid biosynthetic gene expression. <i>Nucleic Acids Res</i> 1996;24(7):1322–9.	doi:10.1093/nar/24.7.1322 PMID 8614637 PMC145798	–	9	2	9	–	9	3	Northern blot - wt vs TF mutant
410	Soussi-Boudekou 1997 Soussi-Boudekou S, Vissers S, Urrestarazu A, Jauniaux JC, André B. Gzf3p, a fourth GATA factor involved in nitrogen-regulated transcription in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 1997;23(6):1157–68.	doi:10.1046/j.1365-2958.1997.3021665.x PMID 9106207	–	9	4	5	–	9	3	lacZ - wt vs TF mutant
411	Kuras 1997 Kuras L, Barbey R, Thomas D. Assembly of a bZIP-bHLH transcription activation complex: formation of the yeast Cbfl-Met4-Met28 complex is regulated through Met28 stimulation of Cbfl DNA binding. <i>EMBO J</i> 1997;16(9):2441–51.	doi:10.1093/emboj/16.9.2441 PMID 9171357 PMC1169844	–	9	3	9	3	6	3	lacZ - wt vs TF mutant
412	Zhou 2001 Zhou H, Winston F. NRG1 is required for glucose repression of the SUC2 and GAL genes of <i>Saccharomyces cerevisiae</i> . <i>BMC Genet</i> 2001;2:5.	doi:10.1186/1471-2156-2-5 PMID 11281938 PMC31344	–	9	3	9	–	9	2	Northern blot - wt vs TF mutant
413	Deminoff 2001 Deminoff SJ, Santangelo GM. Rap1p requires Gcr1p and Gcr2p homodimers to activate ribosomal protein and glycolytic genes, respectively. <i>Genetics</i> 2001;158(1):133–43.	doi:10.1093/genetics/158.1.133 PMID 11333224 PMC1461654	–	9	2	9	–	9	1	Primer extension - wt vs TF mutant
414	Yamamoto 2007 Yamamoto A, Ueda J, Yamamoto N, Hashikawa N, Sakurai H. Role of heat shock transcription factor in <i>Saccharomyces cerevisiae</i> oxidative stress response. <i>Eukaryot Cell</i> 2007;6(8):1373–9.	doi:10.1128/EC.00098-07 PMID 17586717 PMC1951129	–	9	3	9	1	6	4	RT-PCR - wt vs TF mutant
415	Bandhakavi 2008 Bandhakavi S, Xie H, O’Callaghan B, Sakurai H, Kim DH, Griffin TJ. Hsf1 activation inhibits rapamycin resistance and TOR signaling in yeast revealed by combined proteomic and genetic analysis. <i>PLoS One</i> 2008;3(2):e1598.	doi:10.1371/journal.pone.0001598 PMID 18270585 PMC2225505	–	9	3	9	–	9	2	RT-PCR - wt vs TF mutant
416	Georis 2009 Georis I, Feller A, Tate JJ, Cooper TG, Dubois E. Nitrogen catabolite repression-sensitive transcription as a readout of Tor pathway regulation: the genetic background, reporter gene and GATA factor assayed determine the outcomes. <i>Genetics</i> 2009;181(3):861–74.	doi:10.1534/genetics.108.099051 PMID 19104072 PMC2651060	–	9	5	7	4	5	1	RT-PCR - wt vs TF mutant
417	Yukawa 2009 Yukawa M, Yo K, Hasegawa H, Ueno M, Tsuchiya E. The Rpd3/HDAC complex is present at the URS1 cis-element with hyperacetylated histone H3. <i>Biosci Biotechnol Biochem</i> 2009;73(2):378–84.	doi:10.1271/bbb.80621 PMID 19202282	–	9	1	8	–	9	2	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
418	Neklesa 2009 Neklesa TK, Davis RW. A genome-wide screen for regulators of TORC1 in response to amino acid starvation reveals a conserved Npr2/3 complex. <i>PLoS Genet</i> 2009;5(6):e1000515.	doi:10.1371/journal.pgen.1000515 PMID 19521502 PMC2686269	–	9	6	6	–	9	3	GFP - wt vs TF mutant
419	Ju 2010 Ju D, Wang X, Ha SW, Fu J, Xie Y. Inhibition of proteasomal degradation of rpn4 impairs nonhomologous end-joining repair of DNA double-strand breaks. <i>PLoS One</i> 2010;5(4):e9877.	doi:10.1371/journal.pone.0009877 PMID 20376190 PMC2848573	–	9	1	9	–	9	1	RT-PCR - wt vs TF mutant
420	Cardillo 2012 Cardillo SB, Levi CE, Bermúdez Moretti M, Correa García S. Interplay between the transcription factors acting on the GATA- and GABA-responsive elements of <i>Saccharomyces cerevisiae</i> UGA promoters. <i>Microbiology (Reading)</i> 2012; 158(Pt 4):925–935.	doi:10.1099/mic.0.051235-0 PMID 22282516	–	9	4	9	9	–	1	ChIP
421	Liu 2013 Liu J, Barrientos A. Transcriptional regulation of yeast oxidative phosphorylation hypoxic genes by oxidative stress. <i>Antioxid Redox Signal</i> 2013;19(16):1916–27.	doi:10.1089/ars.2012.4589 PMID 22703342 PMC3852346	–	9	3	6	5	4	6	ChIP
422	Lee 2013 Lee K, Hahn JS. Interplay of Aro80 and GATA activators in regulation of genes for catabolism of aromatic amino acids in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2013;88(6):1120–34.	doi:10.1111/mmi.12246 PMID 23651256	–	9	3	9	–	9	5	RT-PCR - wt vs TF mutant
423	Kahana-Edwin 2013 Kahana-Edwin S, Stark M, Kassir Y. Multiple MAPK cascades regulate the transcription of IME1, the master transcriptional activator of meiosis in <i>Saccharomyces cerevisiae</i> . <i>PLoS One</i> 2013;8(11):e78920.	doi:10.1371/journal.pone.0078920 PMID 24236068 PMC3827324	–	9	9	9	4	5	6	ChIP
424	Low 2014 Low YS, Bircham PW, Maass DR, Atkinson PH. Kinetochore genes are required to fully activate secretory pathway expansion in <i>S. cerevisiae</i> under induced ER stress. <i>Mol Biosyst</i> 2014;10(7):1790–802.	doi:10.1039/c3mb70414a PMID 24722431	–	9	5	9	–	9	1	GFP - wt vs TF mutant
425	Scherz 2014 Scherz K, Andersen, Bojsen R, et al., Regenber B. Genetic basis for <i>Saccharomyces cerevisiae</i> biofilm in liquid medium. <i>G3 (Bethesda)</i> 2014;4(9):1671–80.	doi:10.1534/g3.114.010892 PMID 25009170 PMC4169159	–	9	9	9	–	9	9	Northern blot - wt vs TF mutant
426	Knight 2014 Knight B, Kubik S, Ghosh B, et al., Shore D. Two distinct promoter architectures centered on dynamic nucleosomes control ribosomal protein gene transcription. <i>Genes Dev</i> 2014; 28(15):1695–709.	doi:10.1101/gad.244434.114 PMID 25085421 PMC4117944	–	9	3	9	9	–	1	ChIP
427	Pérez-Sampietro 2014 Pérez-Sampietro M, Herrero E. The PacC-family protein Rim101 prevents selenite toxicity in <i>Saccharomyces cerevisiae</i> by controlling vacuolar acidification. <i>Fungal Genet Biol</i> 2014; 71:76–85.	doi:10.1016/j.fgb.2014.09.001 PMID 25239548	–	9	3	9	–	9	5	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
428	Paul 2015 Paul E, Tirosh I, Lai W, Buck MJ, Palumbo MJ, Morse RH. Chromatin mediation of a transcriptional memory effect in yeast. <i>G3 (Bethesda)</i> 2015;5(5):829–38.	doi:10.1534/g3.115.017418 PMID 25748434 PMC4426369	–	9	1	9	9	–	1	ChIP
429	Singh 2016 Singh N, Yadav KK, Rajasekharan R. ZAP1-mediated modulation of triacylglycerol levels in yeast by transcriptional control of mitochondrial fatty acid biosynthesis. <i>Mol Microbiol</i> 2016;100(1):55–75.	doi:10.1111/mmi.13298 PMID 26711224	–	9	1	9	–	9	2	RT-PCR - wt vs TF mutant
430	Gaspar-Cordeiro 2018 Gaspar-Cordeiro A, Marques Caetano S, Amaral C, Rodrigues-Pousada C, Pimentel C. Ace1 prevents intracellular copper accumulation by regulating Fet3 expression and thereby restricting Aft1 activity. <i>FEBS J</i> 2018;285(10):1861–1872.	doi:10.1111/febs.14450 PMID 29604179	–	9	2	4	–	9	3	RT-PCR - wt vs TF mutant
431	Li 2020 Li L, Bertram S, Kaplan J, Jia X, Ward DM. The mitochondrial iron exporter genes MMT1 and MMT2 in yeast are transcriptionally regulated by Aft1 and Yap1. <i>J Biol Chem</i> 2020;295(6):1716–1726.	doi:10.1074/jbc.RA119.011154 PMID 31896574 PMC7008362	–	9	3	4	–	9	3	lacZ - wt vs TF mutant
432	Daignan-Fornier 1992 Daignan-Fornier B, Fink GR. Coregulation of purine and histidine biosynthesis by the transcriptional activators BAS1 and BAS2. <i>Proc Natl Acad Sci U S A</i> 1992;89(15):6746–50.	doi:10.1073/pnas.89.15.6746 PMID 1495962 PMC49580	–	8	2	8	4	4	3	lacZ - wt vs TF mutant
433	Messenguy 1991 Messenguy F, Dubois E, Boonchird C. Determination of the DNA-binding sequences of ARG1 proteins to arginine anabolic and catabolic promoters. <i>Mol Cell Biol</i> 1991;11(5):2852–63.	doi:10.1128/mcb.11.5.2852-2863.1991 PMID 2017180 PMC360071	–	8	2	8	8	–	1	EMSA
434	Althoefer 1995 Althoefer H, Schleiffer A, Wassmann K, Nordheim A, Ammerer G. Mcm1 is required to coordinate G2-specific transcription in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1995;15(11):5917–28.	doi:10.1128/MCB.15.11.5917 PMID 7565744 PMC230843	–	8	3	8	–	4	2	–
435	Nishi 1995 Nishi K, Park CS, Pepper AE, Eichinger G, Innis MA, Holland MJ. The GCR1 requirement for yeast glycolytic gene expression is suppressed by dominant mutations in the SGC1 gene, which encodes a novel basic-helix-loop-helix protein. <i>Mol Cell Biol</i> 1995;15(5):2646–53.	doi:10.1128/MCB.15.5.2646 PMID 7739544 PMC230494	–	8	2	8	–	8	1	Northern blot - wt vs TF mutant
436	Hu 1995 Hu Z, Nehlin JO, Ronne H, Michels CA. MIG1-dependent and MIG1-independent glucose regulation of MAL gene expression in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 1995;28(3):258–66.	doi:10.1007/BF00309785 PMID 8529272	–	8	1	5	1	4	4	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
437	Coffman 1996 Coffman JA, Rai R, Cunningham T, Svetlov V, Cooper TG. Gat1p, a GATA family protein whose production is sensitive to nitrogen catabolite repression, participates in transcriptional activation of nitrogen-catabolic genes in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1996;16(3):847–58.	doi:10.1128/MCB.16.3.847 PMID 8622686 PMC231065	–	8	3	8	–	8	2	Northern blot - wt vs TF mutant
438	Schmitt 1996 Schmitt AP, McEntee K. Msn2p, a zinc finger DNA-binding protein, is the transcriptional activator of the multistress response in <i>Saccharomyces cerevisiae</i> . <i>Proc Natl Acad Sci U S A</i> 1996;93(12):5777–82.	doi:10.1073/pnas.93.12.5777 PMID 8650168 PMC39137	–	8	2	8	–	8	2	Northern blot - wt vs TF overexpression
439	Lutfiyya 1996 Lutfiyya LL, Johnston M. Two zinc-finger-containing repressors are responsible for glucose repression of SUC2 expression. <i>Mol Cell Biol</i> 1996;16(9):4790–7.	doi:10.1128/MCB.16.9.4790 PMID 8756637 PMC231480	–	8	3	8	1	7	2	Invertase - wt vs TF mutant
440	McInerny 1997 McInerny CJ, Partridge JF, Mikesell GE, Creemer DP, Breeden LL. A novel Mcm1-dependent element in the SWI4, CLN3, CDC6, and CDC47 promoters activates M/G1-specific transcription. <i>Genes Dev</i> 1997;11(10):1277–88.	doi:10.1101/gad.11.10.1277 PMID 9171372	–	8	1	8	2	3	2	–
441	Coffman 1997 Coffman JA, Cooper TG. Nitrogen GATA factors participate in transcriptional regulation of vacuolar protease genes in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1997;179(17):5609–13.	doi:10.1128/jb.179.17.5609-5613.1997 PMID 9287023 PMC179439	–	8	4	6	–	8	3	Northern blot - wt vs TF mutant
442	Martins 1998 Martins LJ, Jensen LT, Simon JR, Keller GL, Winge DR. Metalloregulation of FRE1 and FRE2 homologs in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 1998;273(37):23716–21.	doi:10.1074/jbc.273.37.23716 PMID 9726978	–	8	2	8	8	–	3	DNA footprinting
443	Kennedy 1999 Kennedy MA, Barbuch R, Bard M. Transcriptional regulation of the squalene synthase gene (ERG9) in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochim Biophys Acta</i> 1999;1445(1):110–22.	doi:10.1016/s0167-4781(99)00035-4 PMID 10209263	–	8	7	7	–	8	3	lacZ - wt vs TF mutant
444	Kumar 2000 Kumar R, Reynolds DM, Shevchenko A, Shevchenko A, Goldstone SD, Dalton S. Forkhead transcription factors, Fkh1p and Fkh2p, collaborate with Mcm1p to control transcription required for M-phase. <i>Curr Biol</i> 2000;10(15):896–906.	doi:10.1016/s0960-9822(00)00618-7 PMID 10959837	–	8	3	8	–	6	2	–
445	Waters 2002 Waters BM, Eide DJ. Combinatorial control of yeast FET4 gene expression by iron, zinc, and oxygen. <i>J Biol Chem</i> 2002;277(37):33749–57.	doi:10.1074/jbc.M206214200 PMID 12095998	–	8	4	5	–	8	7	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
446	Hongay 2002 Hongay C, Jia N, Bard M, Winston F. Mot3 is a transcriptional repressor of ergosterol biosynthetic genes and is required for normal vacuolar function in <i>Saccharomyces cerevisiae</i> . <i>EMBO J</i> 2002;21(15):4114–24.	doi:10.1093/emboj/cdf415 PMID 12145211 PMC126159	–	8	1	4	2	6	3	Northern blot - wt vs TF mutant
447	Hagerman 2002 Hagerman RA, Willis RA. The yeast gene COQ5 is differentially regulated by Mig1p, Rtg3p and Hap2p. <i>Biochim Biophys Acta</i> 2002;1578(1–3):51–8.	doi:10.1016/s0167-4781(02)00496-7 PMID 12393187	–	8	4	4	–	8	4	Northern blot - wt vs TF mutant
448	Aranda 2003 Aranda A, del Olmo MI. Response to acetaldehyde stress in the yeast <i>Saccharomyces cerevisiae</i> involves a strain-dependent regulation of several ALD genes and is mediated by the general stress response pathway. <i>Yeast</i> 2003;20(8):747–59.	doi:10.1002/yea.991 PMID 12794936	–	8	2	8	–	8	1	Northern blot - wt vs TF mutant
449	Chen 2005 Chen XJ, Wang X, Kaufman BA, Butow RA. Aconitase couples metabolic regulation to mitochondrial DNA maintenance. <i>Science</i> 2005;307(5710):714–7.	doi:10.1126/science.1106391 PMID 15692048	–	8	3	8	–	8	3	Microarray analysis - wt vs TF mutant
450	Martens 2005 Martens JA, Wu PY, Winston F. Regulation of an intergenic transcript controls adjacent gene transcription in <i>Saccharomyces cerevisiae</i> . <i>Genes Dev</i> 2005;19(22):2695–704.	doi:10.1101/gad.1367605 PMID 16291644 PMC1283962	–	8	6	8	–	8	2	Northern blot - wt vs TF mutant
451	Pramila 2006 Pramila T, Wu W, Miles S, Noble WS, Breeden LL. The Forkhead transcription factor Hcm1 regulates chromosome segregation genes and fills the S-phase gap in the transcriptional circuitry of the cell cycle. <i>Genes Dev</i> 2006;20(16):2266–78.	doi:10.1101/gad.1450606 PMID 16912276 PMC1553209	GSE4987	8	1	8	6	1	2	DNA footprinting
452	Su 2008 Su NY, Ouni I, Papagiannis CV, Kaiser P. A dominant suppressor mutation of the met30 cell cycle defect suggests regulation of the <i>Saccharomyces cerevisiae</i> Met4-Cbf1 transcription complex by Met32. <i>J Biol Chem</i> 2008;283(17):11615–24.	doi:10.1074/jbc.M708230200 PMID 18308733 PMC2431082	–	8	2	7	–	8	4	RT-PCR - wt vs TF mutant
453	Melamed 2008 Melamed D, Pnueli L, Arava Y. Yeast translational response to high salinity: global analysis reveals regulation at multiple levels. <i>RNA</i> 2008;14(7):1337–51.	doi:10.1261/rna.864908 PMID 18495938 PMC2441982	–	8	2	8	–	8	2	Northern blot - wt vs TF mutant
454	Kupchak 2008 Kupchak BR, Villa NY, Kulemina LV, Lyons TJ. Dissecting the regulation of yeast genes by the osmotin receptor. <i>Biochem Biophys Res Commun</i> 2008;374(2):210–3.	doi:10.1016/j.bbrc.2008.07.002 PMID 18625204 PMC2537462	–	8	4	8	–	8	2	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
455	Hlynialuk 2008 Hlynialuk C, Schierholtz R, Vernooy A, van der Merwe G. Nsf1/Ypl230w participates in transcriptional activation during non-fermentative growth and in response to salt stress in <i>Saccharomyces cerevisiae</i> . <i>Microbiology (Reading)</i> 2008;154(Pt 8):2482–2491.	doi:10.1099/mic.0.2008/019976-0 PMID 18667581	–	8	3	8	–	8	3	Northern blot - wt vs TF mutant
456	Tsoi 2009 Tsoi BM, Beckhouse AG, Gelling CL, et al., Dawes IW. Essential role of one-carbon metabolism and Gcn4p and Bas1p transcriptional regulators during adaptation to anaerobic growth of <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2009; 284(17):11205–15.	doi:10.1074/jbc.M809225200 PMID 19224916 PMC2670125	–	8	2	3	–	8	4	lacZ - wt vs TF mutant
457	Pinson 2009 Pinson B, Vaur S, Sagot I, Couplier F, Lemoine S, Daignan-Fornier B. Metabolic intermediates selectively stimulate transcription factor interaction and modulate phosphate and purine pathways. <i>Genes Dev</i> 2009;23(12): 1399–407.	doi:10.1101/gad.521809 PMID 19528318 PMC2701576	–	8	2	8	8	–	1	ChIP
458	Wielemans 2010 Wielemans K, Jean C, Vissers S, André B. Amino acid signaling in yeast: post-genome duplication divergence of the Stp1 and Stp2 transcription factors. <i>J Biol Chem</i> 2010;285(2): 855–65.	doi:10.1074/jbc.M109.015263 PMID 19906648 PMC2801287	–	8	3	8	2	6	6	lacZ - wt vs TF mutant
459	Ihrig 2010 Ihrig J, Hausmann A, Hain A, et al., Mühlenhoff U. Iron regulation through the back door: iron-dependent metabolite levels contribute to transcriptional adaptation to iron deprivation in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2010; 9(3):460–71.	doi:10.1128/EC.00213-09 PMID 20008079 PMC2837980	–	8	7	8	–	8	2	Luciferase - wt vs mutant promoter sequence
460	Li 2010 Li Y, Chen G, Liu W. Alterations in the interaction between GAL4 and GAL80 effect regulation of the yeast GAL regulon mediated by the F box protein Dsg1. <i>Curr Microbiol</i> 2010; 61(3):210–6.	doi:10.1007/s00284-010-9598-1 PMID 20135317	–	8	2	2	–	8	7	lacZ - wt vs TF mutant
461	Cardillo 2010 Cardillo SB, Bermúdez Moretti M, Correa García S. Uga3 and Uga35/Dal81 transcription factors regulate UGA4 transcription in response to gamma-aminobutyric acid and leucine. <i>Eukaryot Cell</i> 2010;9(8):1262–71.	doi:10.1128/EC.00117-10 PMID 20581295 PMC2918931	–	8	5	8	3	5	1	lacZ - wt vs TF mutant
462	Kahana 2010 Kahana S, Pnueli L, Kainth P, Sassi HE, Andrews B, Kassir Y. Functional dissection of IME1 transcription using quantitative promoter-reporter screening. <i>Genetics</i> 2010;186(3):829–41.	doi:10.1534/genetics.110.122200 PMID 20739709 PMC2975282	–	8	8	8	5	3	4	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
463	Wawrzycka 2010 Wawrzycka D, Sobczak I, Bartosz G, Bocer T, Ulaszewski S, Goffeau A. Vmr 1p is a novel vacuolar multidrug resistance ABC transporter in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Res</i> 2010;10(7):828–38.	doi:10.1111/j.1567-1364.2010.00673.x PMID 20846144	–	8	6	8	–	8	2	lacZ - wt vs TF mutant
464	Braun 2013 Braun KA, Parua PK, Dombek KM, Miner GE, Young ET. 14-3-3 (Bmh) proteins regulate combinatorial transcription following RNA polymerase II recruitment by binding at Adr1-dependent promoters in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2013;33(4):712–24.	doi:10.1128/MCB.01226-12 PMID 23207903 PMC3571347	GSE40116	8	2	8	5	3	3	ChIP
465	Haynes 2013 Haynes BC, Maier EJ, Kramer MH, Wang PI, Brown H, Brent MR. Mapping functional transcription factor networks from gene expression data. <i>Genome Res</i> 2013;23(8):1319–28.	doi:10.1101/gr.150904.112 PMID 23636944 PMC3730105	–	8	2	8	–	8	1	Quantigene Plex Assay WT vs TF mutant
466	Zhang 2013 Zhang N, Quan Z, Rash B, Oliver SG. Synergistic effects of TOR and proteasome pathways on the yeast transcriptome and cell growth. <i>Open Biol</i> 2013;3(5):120137.	doi:10.1098/rsob.120137 PMID 23697803 PMC3866871	–	8	4	8	–	8	5	Northern blot - wt vs TF mutant
467	Rietzschel 2015 Rietzschel N, Pierik AJ, Bill E, Lill R, Mühlenhoff U. The basic leucine zipper stress response regulator Yap5 senses high-iron conditions by coordination of [2Fe-2S] clusters. <i>Mol Cell Biol</i> 2015;35(2):370–8.	doi:10.1128/MCB.01033-14 PMID 25368382 PMC4272429	–	8	8	8	–	8	3	Luciferase - wt vs TF mutant
468	Meurer 2017 Meurer M, Chevyreva V, Cerulus B, Knop M. The regulatable MAL32 promoter in <i>Saccharomyces cerevisiae</i> : characteristics and tools to facilitate its use. <i>Yeast</i> 2017;34(1):39–49.	doi:10.1002/yea.3214 PMID 27714848	–	8	1	4	–	8	2	GFP - wt vs TF mutant
469	Wang 2017 Wang Z, Bai X, Guo X, He X. Regulation of crucial enzymes and transcription factors on 2-phenylethanol biosynthesis via Ehrlich pathway in <i>Saccharomyces cerevisiae</i> . <i>J Ind Microbiol Biotechnol</i> 2017;44(1):129–139.	doi:10.1007/s10295-016-1852-5 PMID 27770224	–	8	2	5	2	6	2	RT-PCR - wt vs TF mutant
470	Li 2017 Li F, Zheng LD, Chen X, Zhao X, Briggs SD, Du HN. Gcn5-mediated Rph1 acetylation regulates its autophagic degradation under DNA damage stress. <i>Nucleic Acids Res</i> 2017;45(9):5183–5197.	doi:10.1093/nar/gkx129 PMID 28334815 PMC5435933	–	8	1	5	3	5	2	RT-PCR - wt vs TF overexpression
471	Wierman 2017 Wierman MB, Maqani N, Strickler E, Li M, Smith JS. Caloric Restriction Extends Yeast Chronological Life Span by Optimizing the Snf1 (AMPK) Signaling Pathway. <i>Mol Cell Biol</i> 2017;37(13).	doi:10.1128/MCB.00562-16 PMID 28373292 PMC5472825	–	8	2	4	–	8	2	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
472	Wang 2017 Wang H, Ouyang Y, Zhou C, et al., Ma M. YKL071W from <i>Saccharomyces cerevisiae</i> encodes a novel aldehyde reductase for detoxification of glycolaldehyde and furfural derived from lignocellulose. <i>Appl Microbiol Biotechnol</i> 2017;101(23–24): 8405–8418.	doi:10.1007/s00253-017-8567-z PMID 29034432	–	8	4	4	–	8	2	RT-PCR - wt vs TF mutant
473	Vamvakas 2019 Vamvakas SS, Kapolos J, Farmakis L, et al., Demoli D. Specific serine residues of Msn2/4 are responsible for regulation of alcohol fermentation rates and ethanol resistance. <i>Biotechnol Prog</i> 2019;35(2):e2759.	doi:10.1002/btpr.2759 PMID 30507007	–	8	2	8	–	8	1	RT-PCR - wt vs TF mutant
474	Pinson 2019 Pinson B, Ceschin J, Saint-Marc C, Daignan-Fornier B. Dual control of NAD(+) synthesis by purine metabolites in yeast. <i>Elife</i> 2019;8.	doi:10.7554/eLife.43808 PMID 30860478 PMC6430606	–	8	2	8	–	8	1	Northern blot - wt vs TF mutant
475	Yu 2019 Yu R, Sun L, Sun Y, Han X, Qin L, Dang W. Cellular response to moderate chromatin architectural defects promotes longevity. <i>Sci Adv</i> 2019;5(7):eaav1165.	doi:10.1126/sciadv.aav1165 PMID 31309140 PMC6620092	–	8	2	8	–	8	1	RT-PCR - wt vs. TF double mutant
476	Chen 2020 Chen LB, Ma S, Jiang TX, Qiu XB. Transcriptional upregulation of proteasome activator Blm10 antagonizes cellular aging. <i>Biochem Biophys Res Commun</i> 2020;532(2): 211–218.	doi:10.1016/j.bbrc.2020.07.003 PMID 32861419	–	8	2	2	–	8	6	RT-PCR - wt vs TF mutant
477	Spasskaya 2020 Spasskaya DS, Nadolinskaia NI, Tutyaeva VV, Lysov YP, Karpov VL, Karpov DS. Yeast Rpn4 Links the Proteasome and DNA Repair via RAD52 Regulation. <i>Int J Mol Sci</i> 2020; 21(21).	doi:10.3390/ijms21218097 PMID 33143019 PMC7672625	–	8	1	4	–	8	3	RT-PCR - wt vs TF mutant
478	Matos 2021 Matos GS, Madeira JB, Fernandes CM, et al., Montero-Lomeli M. Regulation of sphingolipid synthesis by the G1/S transcription factor Swi4. <i>Biochim Biophys Acta Mol Cell Biol Lipids</i> 2021;1866(9):158983.	doi:10.1016/j.bbalip.2021.158983 PMID 34062255 PMC8512607	–	8	1	8	–	8	4	RT-PCR - wt vs. TF mutant
479	Chasman 1990 Chasman DI, Lue NF, Buchman AR, LaPointe JW, Lorch Y, Kornberg RD. A yeast protein that influences the chromatin structure of UASG and functions as a powerful auxiliary gene activator. <i>Genes Dev</i> 1990;4(4):503–14.	doi:10.1101/gad.4.4.503 PMID 2361590	–	7	1	7	7	–	1	EMSA
480	Halladay 1995 Halladay JT, Craig EA. A heat shock transcription factor with reduced activity suppresses a yeast HSP70 mutant. <i>Mol Cell Biol</i> 1995;15(9):4890–7.	doi:10.1128/MCB.15.9.4890 PMID 7651408 PMC230735	–	7	1	7	2	5	4	Western blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
481	Schüller 1994 Schüller HJ, Schütz A, Knab S, Hoffmann B, Schweizer E. Importance of general regulatory factors Rap1p, Abf1p and Reb1p for the activation of yeast fatty acid synthase genes FAS1 and FAS2. <i>Eur J Biochem</i> 1994;225(1):213–22.	doi:10.1111/j.1432-1033.1994.00213.x PMID 7925441	–	7	4	7	4	2	2	lacZ - wt vs mutant promoter sequence
482	Vashee 1993 Vashee S, Xu H, Johnston SA, Kodadek T. How do "Zn2 cys6" proteins distinguish between similar upstream activation sites? Comparison of the DNA-binding specificity of the GAL4 protein in vitro and in vivo. <i>J Biol Chem</i> 1993;268(33):24699–706.	PMID 8227030	–	7	1	7	1	–	1	–
483	Steber 1995 Steber CM, Esposito RE. UME6 is a central component of a developmental regulatory switch controlling meiosis-specific gene expression. <i>Proc Natl Acad Sci U S A</i> 1995;92(26):12490–4.	doi:10.1073/pnas.92.26.12490 PMID 8618927 PMC40383	–	7	1	7	2	5	6	Northern blot - wt vs TF mutant
484	Yamaguchi-Iwai 1996 Yamaguchi-Iwai Y, Stearman R, Dancis A, Klausner RD. Iron-regulated DNA binding by the AFT1 protein controls the iron regulon in yeast. <i>EMBO J</i> 1996;15(13):3377–84.	PMID 8670839 PMC451901	–	7	1	6	3	4	3	Northern blot - wt vs TF overexpression
485	Ozcan 1996 Ozcan S, Johnston M. Two different repressors collaborate to restrict expression of the yeast glucose transporter genes HXT2 and HXT4 to low levels of glucose. <i>Mol Cell Biol</i> 1996;16(10):5536–45.	doi:10.1128/MCB.16.10.5536 PMID 8816466 PMC231553	–	7	3	7	2	4	3	lacZ - wt vs TF mutant
486	Dang 1996 Dang VD, Valens M, Bolotin-Fukuhara M, Daignan-Fornier B. Cloning of the ASN1 and ASN2 genes encoding asparagine synthetases in <i>Saccharomyces cerevisiae</i> : differential regulation by the CCAAT-box-binding factor. <i>Mol Microbiol</i> 1996;22(4):681–92.	doi:10.1046/j.1365-2958.1996.d01-1715.x PMID 8951815	–	7	5	7	1	5	3	lacZ - wt vs TF mutant
487	Dubois 1997 Dubois E, Messenguy F. Integration of the multiple controls regulating the expression of the arginase gene CAR1 of <i>Saccharomyces cerevisiae</i> in response to different nitrogen signals: role of Gln3p, ArgRp-Mcm1p, and Ume6p. <i>Mol Gen Genet</i> 1997;253(5):568–80.	doi:10.1007/s004380050359 PMID 9065690	–	7	7	7	–	7	3	Enzyme activity - wt vs mutant promoter sequence
488	Kwast 1997 Kwast KE, Burke PV, Brown K, Poyton RO. REO1 and ROX1 are alleles of the same gene which encodes a transcriptional repressor of hypoxic genes in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 1997;32(6):377–83.	doi:10.1007/s002940050291 PMID 9388292	–	7	1	7	–	7	1	Northern blot - wt vs TF mutant
489	Hepworth 1998 Hepworth SR, Friesen H, Segall J. NDT80 and the meiotic recombination checkpoint regulate expression of middle sporulation-specific genes in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1998;18(10):5750–61.	doi:10.1128/MCB.18.10.5750 PMID 9742092 PMC109161	–	7	2	7	–	7	3	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
490	Denis 1998 Denis V, Daignan-Fornier B. Synthesis of glutamine, glycine and 10-formyl tetrahydrofolate is coregulated with purine biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1998;259(3):246–55.	doi:10.1007/s004380050810 PMID 9749667	–	7	2	7	–	7	2	lacZ - wt vs TF mutant
491	Treger 1998 Treger JM, Schmitt AP, Simon JR, McEntee K. Transcriptional factor mutations reveal regulatory complexities of heat shock and newly identified stress genes in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 1998;273(41):26875–9.	doi:10.1074/jbc.273.41.26875 PMID 9756934	–	7	2	7	–	7	1	Northern blot - wt vs TF mutant
492	Mai 2000 Mai B, Breeden L. CLN1 and its repression by Xbp1 are important for efficient sporulation in budding yeast. <i>Mol Cell Biol</i> 2000;20(2):478–87.	doi:10.1128/MCB.20.2.478-487.2000 PMID 10611226 PMC85107	–	7	1	7	–	7	2	Northern blot - wt vs TF mutant
493	Bojunga 1999 Bojunga N, Entian KD. Cat8p, the activator of gluconeogenic genes in <i>Saccharomyces cerevisiae</i> , regulates carbon source-dependent expression of NADP-dependent cytosolic isocitrate dehydrogenase (Idp2p) and lactate permease (Jen1p). <i>Mol Gen Genet</i> 1999;262(4–5):869–75.	doi:10.1007/s004380051152 PMID 10628872	–	7	2	3	–	7	2	Northern blot - wt vs TF mutant
494	Zähringer 2000 Zähringer H, Thevelein JM, Nwaka S. Induction of neutral trehalase Nth1 by heat and osmotic stress is controlled by STRE elements and Msn2/Msn4 transcription factors: variations of PKA effect during stress and growth. <i>Mol Microbiol</i> 2000;35(2):397–406.	doi:10.1046/j.1365-2958.2000.01706.x PMID 10652100	–	7	2	2	–	7	3	lacZ - wt vs TF mutant
495	Blom 2000 Blom J, De Mattos MJ, Grivell LA. Redirection of the respiro-fermentative flux distribution in <i>Saccharomyces cerevisiae</i> by overexpression of the transcription factor Hap4p. <i>Appl Environ Microbiol</i> 2000;66(5):1970–3.	doi:10.1128/AEM.66.5.1970-1973.2000 PMID 10788368 PMC101441	–	7	4	7	–	4	2	Northern blot - wt vs TF overexpression
496	Koranda 2000 Koranda M, Schleiffer A, Endler L, Ammerer G. Forkhead-like transcription factors recruit Ndd1 to the chromatin of G2/M-specific promoters. <i>Nature</i> 2000;406(6791):94–8.	doi:10.1038/35017589 PMID 10894549	–	7	3	7	1	–	1	–
497	Pic 2000 Pic A, Lim FL, Ross SJ, et al., Morgan BA. The forkhead protein Fkh2 is a component of the yeast cell cycle transcription factor SFF. <i>EMBO J</i> 2000;19(14):3750–61.	doi:10.1093/emboj/19.14.3750 PMID 10899128 PMC313965	–	7	2	7	3	3	4	EMSA
498	Hallstrom 2000 Hallstrom TC, Moye-Rowley WS. Multiple signals from dysfunctional mitochondria activate the pleiotropic drug resistance pathway in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2000;275(48):37347–56.	doi:10.1074/jbc.M007338200 PMID 10980204	–	7	4	7	–	6	2	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
499	Gurvitz 2000 Gurvitz A, Wabnegger L, Rottensteiner H, et al., Hamilton B. Adr1p-dependent regulation of the oleic acid-inducible yeast gene SPS19 encoding the peroxisomal beta-oxidation auxiliary enzyme 2,4-dienoyl-CoA reductase. <i>Mol Cell Biol Res Commun</i> 2000;4(2):81–9.	doi:10.1006/mcbr.2000.0261 PMID 11170837	–	7	3	7	–	7	1	Northern blot - wt vs TF mutant
500	do Valle Matta 2001 do Valle Matta MA, Jonniaux JL, Balzi E, Goffeau A, van den Hazel B. Novel target genes of the yeast regulator Pdr1p: a contribution of the TPO1 gene in resistance to quinidine and other drugs. <i>Gene</i> 2001;272(1–2):111–9.	doi:10.1016/s0378-1119(01)00558-3 PMID 11470516	–	7	1	7	–	7	1	Northern blot - wt vs TF overexpression
501	Lodi 2002 Lodi T, Fontanesi F, Guiard B. Co-ordinate regulation of lactate metabolism genes in yeast: the role of the lactate permease gene JEN1. <i>Mol Genet Genomics</i> 2002;266(5):838–47.	doi:10.1007/s00438-001-0604-y PMID 11810259	–	7	7	7	1	3	3	–
502	Spode 2002 Spode I, Maiwald D, Hollenberg CP, Suckow M. ATF/CREB sites present in sub-telomeric regions of <i>Saccharomyces cerevisiae</i> chromosomes are part of promoters and act as UAS/URS of highly conserved COS genes. <i>J Mol Biol</i> 2002;319(2):407–20.	doi:10.1016/S0022-2836(02)00322-4 PMID 12051917	–	7	4	7	–	3	1	–
503	Guaragnella 2003 Guaragnella N, Butow RA. ATO3 encoding a putative outward ammonium transporter is an RTG-independent retrograde responsive gene regulated by GCN4 and the Ssy1-Ptr3-Ssy5 amino acid sensor system. <i>J Biol Chem</i> 2003;278(46):45882–7.	doi:10.1074/jbc.M309301200 PMID 12966084	–	7	3	7	–	7	2	Northern blot - wt vs TF mutant
504	Nevitt 2004 Nevitt T, Pereira J, Azevedo D, Guerreiro P, Rodrigues-Pousada C. Expression of YAP4 in <i>Saccharomyces cerevisiae</i> under osmotic stress. <i>Biochem J</i> 2004;379(Pt 2):367–74.	doi:10.1042/BJ20031127 PMID 14680476 PMC1224068	–	7	2	4	–	7	1	Northern blot - wt vs TF mutant
505	Barbey 2005 Barbey R, Baudouin-Cornu P, Lee TA, et al., Thomas D. Inducible dissociation of SCF(Met30) ubiquitin ligase mediates a rapid transcriptional response to cadmium. <i>EMBO J</i> 2005;24(3):521–32.	doi:10.1038/sj.emboj.7600556 PMID 15660125 PMC548661	–	7	1	7	3	4	1	Northern blot - wt vs TF mutant
506	Gardocki 2005 Gardocki ME, Bakewell M, Kamath D, Robinson K, Borovicka K, Lopes JM. Genomic analysis of PIS1 gene expression. <i>Eukaryot Cell</i> 2005;4(3):604–14.	doi:10.1128/EC.4.3.604-614.2005 PMID 15755922 PMC1087795	–	7	7	7	–	7	2	lacZ - wt vs TF mutant
507	Davies 2005 Davies BS, Wang HS, Rine J. Dual activators of the sterol biosynthetic pathway of <i>Saccharomyces cerevisiae</i> : similar activation/regulatory domains but different response mechanisms. <i>Mol Cell Biol</i> 2005;25(16):7375–85.	doi:10.1128/MCB.25.16.7375-7385.2005 PMID 16055745 PMC1190251	–	7	2	6	–	7	1	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
508	Ashe 2008 Ashe M, de Bruin RA, Kalashnikova T, McDonald WH, Yates JR 3rd, Wittenberg C. The SBF- and MBF-associated protein Msa1 is required for proper timing of G1-specific transcription in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2008;283(10):6040–9.	doi:10.1074/jbc.M708248200 PMID 18160399	–	7	2	7	7	–	1	ChIP
509	Barrales 2008 Barrales RR, Jimenez J, Ibeas JI. Identification of novel activation mechanisms for FLO11 regulation in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2008;178(1):145–56.	doi:10.1534/genetics.107.081315 PMID 18202364 PMC2206066	–	7	7	7	–	7	3	Northern blot - wt vs TF mutant
510	Castro-Prego 2010 Castro-Prego R, Lamas-Maceiras M, Soengas P, et al., Cerdán ME. Ixr1p regulates oxygen-dependent HEM13 transcription. <i>FEMS Yeast Res</i> 2010;10(3):309–21.	doi:10.1111/j.1567-1364.2010.00616.x PMID 20345897	–	7	1	2	3	4	6	ChIP
511	Georis 2011 Georis I, Tate JJ, Cooper TG, Dubois E. Nitrogen-responsive regulation of GATA protein family activators Gln3 and Gat1 occurs by two distinct pathways, one inhibited by rapamycin and the other by methionine sulfoximine. <i>J Biol Chem</i> 2011;286(52):44897–912.	doi:10.1074/jbc.M111.290577 PMID 22039046 PMC3248002	–	7	2	7	1	6	2	RT-PCR - wt vs TF mutant
512	Rodríguez-Colman 2013 Rodríguez-Colman MJ, Sorolla MA, Vall-Llaura N, Tamarit J, Ros J, Cabisco E. The FOX transcription factor Hcm1 regulates oxidative metabolism in response to early nutrient limitation in yeast. Role of Snf1 and Tor1/Sch9 kinases. <i>Biochim Biophys Acta</i> 2013;1833(8):2004–15.	doi:10.1016/j.bbamcr.2013.02.015 PMID 23481038	–	7	1	7	–	7	1	RT-PCR - wt vs TF mutant
513	Serra-Cardona 2014 Serra-Cardona A, Petrežsélyová S, Canadell D, Ramos J, Ariño J. Coregulated expression of the Na ⁺ /phosphate Pho89 transporter and Ena1 Na ⁺ -ATPase allows their functional coupling under high-pH stress. <i>Mol Cell Biol</i> 2014;34(24):4420–35.	doi:10.1128/MCB.01089-14 PMID 25266663 PMC4248728	–	7	6	7	–	7	4	lacZ - wt vs TF mutant
514	Nishizawa 2015 Nishizawa M. The regulators of yeast PHO system participate in the transcriptional regulation of G1 cyclin under alkaline stress conditions. <i>Yeast</i> 2015;32(3):367–78.	doi:10.1002/yea.3064 PMID 25582350	–	7	3	7	–	7	3	Northern blot - wt vs TF mutant
515	Caetano 2015 Caetano SM, Menezes R, Amaral C, Rodrigues-Pousada C, Pimentel C. Repression of the Low Affinity Iron Transporter Gene FET4: A NOVEL MECHANISM AGAINST CADMIUM TOXICITY ORCHESTRATED BY YAP1 VIA ROX1. <i>J Biol Chem</i> 2015;290(30):18584–95.	doi:10.1074/jbc.M114.600742 PMID 26063801 PMC4513117	–	7	2	7	1	6	1	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
516	Gutin 2015 Gutin J, Sadeh A, Rahat A, Aharoni A, Friedman N. Condition-specific genetic interaction maps reveal crosstalk between the cAMP/PKA and the HOG MAPK pathways in the activation of the general stress response. <i>Mol Syst Biol</i> 2015;11(10):829.	doi:10.15252/msb.20156451 PMID 26446933 PMC4631200	–	7	7	7	–	7	2	GFP - wt vs TF mutant
517	Takahashi 2017 Takahashi T, Ohara Y, Sawatari M, Sueno K. Isolation and characterization of sake yeast mutants with enhanced isoamyl acetate productivity. <i>J Biosci Bioeng</i> 2017;123(1):71–77.	doi:10.1016/j.jbiosc.2016.07.002 PMID 27475923	–	7	1	2	–	7	2	RT-PCR - wt vs TF mutant
518	Eapen 2017 Eapen VV, Waterman DP, Bernard A, et al., Haber JE. A pathway of targeted autophagy is induced by DNA damage in budding yeast. <i>Proc Natl Acad Sci U S A</i> 2017;114(7):E1158–E1167.	doi:10.1073/pnas.1614364114 PMID 28154131 PMC5320992	–	7	1	6	–	7	1	RT-PCR - wt vs TF overexpression
519	Chen 2017 Chen J, Tresenrider A, Chia M, et al., Ünal E. Kinetochores inactivation by expression of a repressive mRNA. <i>Elife</i> 2017;6.	doi:10.7554/eLife.27417 PMID 28906249 PMC5655150	–	7	3	5	6	1	5	ChIP
520	Zhao 2018 Zhao H, Wang Q, Liu C, et al., Li W. A Role for the Respiratory Chain in Regulating Meiosis Initiation in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2018;208(3):1181–1194.	doi:10.1534/genetics.118.300689 PMID 29301906 PMC5844330	–	7	2	3	3	4	5	ChIP
521	Romero 2018 Romero AM, Jordá T, Rozès N, Martínez-Pastor MT, Puig S. Regulation of yeast fatty acid desaturase in response to iron deficiency. <i>Biochim Biophys Acta Mol Cell Biol Lipids</i> 2018;1863(6):657–668.	doi:10.1016/j.bbalip.2018.03.008 PMID 29627385	–	7	2	3	–	7	5	RT-PCR - wt vs TF mutant
522	Parnell 2019 Parnell EJ, Stillman DJ. Multiple Negative Regulators Restrict Recruitment of the SWI/SNF Chromatin Remodeler to the HO Promoter in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2019;212(4):1181–1204.	doi:10.1534/genetics.119.302359 PMID 31167839 PMC6707452	–	7	6	6	3	4	2	RT-PCR - wt vs. TF mutant
523	Raithatha 2021 Raithatha SA, Vaza S, Islam MT, Greenwood B, Stuart DT. Ume6 Acts as a Stable Platform To Coordinate Repression and Activation of Early Meiosis-Specific Genes in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2021;41(7):e0037820.	doi:10.1128/MCB.00378-20 PMID 33941619 PMC8224235	–	7	1	2	6	1	4	ChIP
524	Keng 1992 Keng T, Richard C, Larocque R. Structure and regulation of yeast HEM3, the gene for porphobilinogen deaminase. <i>Mol Gen Genet</i> 1992;234(2):233–43.	doi:10.1007/BF00283844 PMID 1508149	–	6	4	6	2	4	2	lacZ - wt vs TF mutant
525	Cunningham 1991 Cunningham TS, Cooper TG. Expression of the DAL80 gene, whose product is homologous to the GATA factors and is a negative regulator of multiple nitrogen catabolic genes in <i>Saccharomyces cerevisiae</i> , is sensitive to nitrogen catabolite repression. <i>Mol Cell Biol</i> 1991;11(12):6205–15.	doi:10.1128/mcb.11.12.6205-6215.1991 PMID 1944286 PMC361806	–	6	1	6	–	6	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
526	Smith 1990 Smith HE, Su SS, Neigeborn L, Driscoll SE, Mitchell AP. Role of IME1 expression in regulation of meiosis in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1990;10(12):6103–13.	doi:10.1128/mcb.10.12.6103-6113.1990 PMID 2247050 PMC362885	–	6	1	6	–	6	2	Northern blot - wt vs TF mutant
527	Yoo 1985 Yoo HS, Genbauffe FS, Cooper TG. Identification of the ureidoglycolate hydrolase gene in the DAL gene cluster of <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1985;5(9):2279–88.	doi:10.1128/mcb.5.9.2279-2288.1985 PMID 3915539 PMC366954	–	6	2	6	–	6	1	Northern blot - wt vs TF mutant
528	Oehlen 1996 Oehlen LJ, McKinney JD, Cross FR. Ste12 and Mcm1 regulate cell cycle-dependent transcription of FAR1. <i>Mol Cell Biol</i> 1996;16(6):2830–7.	doi:10.1128/MCB.16.6.2830 PMID 8649392 PMC231275	–	6	2	6	–	6	3	Northern blot - wt vs TF mutant
529	Cunningham 1996 Cunningham TS, Svetlov VV, Rai R, Smart W, Cooper TG. Gln3p is capable of binding to UAS(NTR) elements and activating transcription in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1996;178(12):3470–9.	doi:10.1128/jb.178.12.3470-3479.1996 PMID 8655543 PMC178115	–	6	1	6	6	–	1	EMSA
530	Toone 1995 Toone WM, Johnson AL, Banks GR, et al., Johnston LH. Rme1, a negative regulator of meiosis, is also a positive activator of G1 cyclin gene expression. <i>EMBO J</i> 1995;14(23):5824–32.	doi:10.1002/j.1460-2075.1995.tb00270.x PMID 8846775 PMC394700	–	6	4	6	–	6	2	Northern blot - wt vs TF mutant
531	Karpichev 1997 Karpichev IV, Luo Y, Mariani RC, Small GM. A complex containing two transcription factors regulates peroxisome proliferation and the coordinate induction of beta-oxidation enzymes in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1997;17(1):69–80.	doi:10.1128/MCB.17.1.69 PMID 8972187 PMC231731	–	6	2	6	1	5	3	Northern blot - wt vs TF mutant
532	Blaiseau 1997 Blaiseau PL, Isnard AD, Surdin-Kerjan Y, Thomas D. Met31p and Met32p, two related zinc finger proteins, are involved in transcriptional regulation of yeast sulfur amino acid metabolism. <i>Mol Cell Biol</i> 1997;17(7):3640–8.	doi:10.1128/MCB.17.7.3640 PMID 9199298 PMC232216	–	6	2	6	–	6	2	Northern blot - wt vs TF mutant
533	Marini 1997 Marini AM, Soussi-Boudekou S, Vissers S, Andre B. A family of ammonium transporters in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1997;17(8):4282–93.	doi:10.1128/MCB.17.8.4282 PMID 9234685 PMC232281	–	6	2	6	–	6	3	lacZ - wt vs TF mutant
534	Kawahara 1997 Kawahara T, Yanagi H, Yura T, Mori K. Endoplasmic reticulum stress-induced mRNA splicing permits synthesis of transcription factor Hac1p/Ern4p that activates the unfolded protein response. <i>Mol Biol Cell</i> 1997;8(10):1845–62.	doi:10.1091/mbc.8.10.1845 PMID 9348528 PMC25627	–	6	1	6	–	6	1	Northern blot - wt vs TF mutant
535	Matheos 1997 Matheos DP, Kingsbury TJ, Ahsan US, Cunningham KW. Tcn1p/Crz1p, a calcineurin-dependent transcription factor that differentially regulates gene expression in <i>Saccharomyces cerevisiae</i> . <i>Genes Dev</i> 1997;11(24):3445–58.	doi:10.1101/gad.11.24.3445 PMID 9407036 PMC316804	–	6	1	6	–	6	4	lacZ - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
536	Smith 1998 Smith A, Ward MP, Garrett S. Yeast PKA represses Msn2p/Msn4p-dependent gene expression to regulate growth, stress response and glycogen accumulation. <i>EMBO J</i> 1998; 17(13):3556–64.	doi:10.1093/emboj/17.13.3556 PMID 9649426 PMC1170692	–	6	2	6	–	6	1	Northern blot - wt vs TF mutant
537	McBride 1999 McBride HJ, Yu Y, Stillman DJ. Distinct regions of the Swi5 and Ace2 transcription factors are required for specific gene activation. <i>J Biol Chem</i> 1999;274(30):21029–36.	doi:10.1074/jbc.274.30.21029 PMID 10409653	–	6	2	6	6	–	1	DNA footprinting
538	Türkel 1999 Türkel S, Bisson LF. Transcription of the HXT4 gene is regulated by Gcr1p and Gcr2p in the yeast <i>S. cerevisiae</i> . <i>Yeast</i> 1999;15(11):1045–57.	<a href="https://doi.org/10.1002/(SICI)1097-0061(199908)15:11<1045::AID-YEA433>3.0.CO;2-Y">doi:10.1002/(SICI)1097-0061(199908)15:11<1045::AID-YEA433>3.0.CO;2-Y PMID 10455229	–	6	2	3	–	6	2	lacZ - wt vs TF mutant
539	Gagiano 1999 Gagiano M, Van Dyk D, Bauer FF, Lambrechts MG, Pretorius IS. Divergent regulation of the evolutionarily closely related promoters of the <i>Saccharomyces cerevisiae</i> STA2 and MUC1 genes. <i>J Bacteriol</i> 1999;181(20):6497–508.	doi:10.1128/JB.181.20.6497-6508.1999 PMID 10515942 PMC103787	–	6	3	6	–	6	2	lacZ - wt vs TF overexpression
540	Grauslund 1999 Grauslund M, Lopes JM, Rønnow B. Expression of GUT1, which encodes glycerol kinase in <i>Saccharomyces cerevisiae</i> , is controlled by the positive regulators Adr1p, Ino2p and Ino4p and the negative regulator Opi1p in a carbon source-dependent fashion. <i>Nucleic Acids Res</i> 1999;27(22):4391–8.	doi:10.1093/nar/27.22.4391 PMID 10536147 PMC148721	–	6	6	6	5	1	2	lacZ - wt vs mutant promoter sequence
541	Pedruzzi 2000 Pedruzzi I, Bürckert N, Egger P, De Virgilio C. <i>Saccharomyces cerevisiae</i> Ras/cAMP pathway controls post-diauxic shift element-dependent transcription through the zinc finger protein Gis1. <i>EMBO J</i> 2000;19(11):2569–79.	doi:10.1093/emboj/19.11.2569 PMID 10835355 PMC212766	–	6	1	3	–	6	2	Northern blot - wt vs TF mutant
542	Raitt 2000 Raitt DC, Johnson AL, Erkin AM, et al., Johnston LH. The Skn7 response regulator of <i>Saccharomyces cerevisiae</i> interacts with Hsf1 in vivo and is required for the induction of heat shock genes by oxidative stress. <i>Mol Biol Cell</i> 2000;11(7):2335–47.	doi:10.1091/mbc.11.7.2335 PMID 10888672 PMC14923	–	6	1	4	1	5	4	Northern blot - wt vs TF mutant
543	Dormer 2000 Dormer UH, Westwater J, McLaren NF, Kent NA, Mellor J, Jamieson DJ. Cadmium-inducible expression of the yeast GSH1 gene requires a functional sulfur-amino acid regulatory network. <i>J Biol Chem</i> 2000;275(42):32611–6.	doi:10.1074/jbc.M004167200 PMID 10921921	–	6	6	6	1	4	3	lacZ - wt vs TF mutant
544	Gasch 2000 Gasch AP, Spellman PT, Kao CM, et al., Brown PO. Genomic expression programs in the response of yeast cells to environmental changes. <i>Mol Biol Cell</i> 2000;11(12):4241–57.	doi:10.1091/mbc.11.12.4241 PMID 11102521 PMC15070	no GEO	6	3	6	–	6	2	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
545	Vyas 2001 Vyas VK, Kuchin S, Carlson M. Interaction of the repressors Nrg1 and Nrg2 with the Snf1 protein kinase in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2001;158(2):563–72.	doi:10.1093/genetics/158.2.563 PMID 11404322 PMC1461687	–	6	2	4	–	6	3	lacZ - wt vs TF mutant
546	Hahn 2002 Hahn JS, Thiele DJ. Regulation of the <i>Saccharomyces cerevisiae</i> Slt2 kinase pathway by the stress-inducible Sdp1 dual specificity phosphatase. <i>J Biol Chem</i> 2002;277(24):21278–84.	doi:10.1074/jbc.M202557200 PMID 11923319	–	6	1	2	–	6	3	Northern blot - wt vs TF mutant
547	Riego 2002 Riego L, Avendaño A, DeLuna A, Rodríguez E, González A. GDH1 expression is regulated by GLN3, GCN4, and HAP4 under respiratory growth. <i>Biochem Biophys Res Commun</i> 2002;293(1):79–85.	doi:10.1016/S0006-291X(02)00174-2 PMID 12054566	–	6	6	6	–	6	2	lacZ - wt vs TF mutant
548	Kaniak 2004 Kaniak A, Xue Z, Macool D, Kim JH, Johnston M. Regulatory network connecting two glucose signal transduction pathways in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2004;3(1):221–31.	doi:10.1128/EC.3.1.221-231.2004 PMID 14871952 PMC329515	–	6	2	5	–	6	3	lacZ - wt vs TF mutant
549	Menezes 2004 Menezes RA, Amaral C, Delaunay A, Toledano M, Rodrigues-Pousada C. Yap8p activation in <i>Saccharomyces cerevisiae</i> under arsenic conditions. <i>FEBS Lett</i> 2004;566(1–3):141–6.	doi:10.1016/j.febslet.2004.04.019 PMID 15147884	–	6	2	6	–	6	1	Northern blot - wt vs TF mutant
550	Pnueli 2004 Pnueli L, Edry I, Cohen M, Kassir Y. Glucose and nitrogen regulate the switch from histone deacetylation to acetylation for expression of early meiosis-specific genes in budding yeast. <i>Mol Cell Biol</i> 2004;24(12):5197–208.	doi:10.1128/MCB.24.12.5197-5208.2004 PMID 15169885 PMC419861	–	6	2	6	3	2	2	ChIP
551	Zhu 2004 Zhu Y, Xiao W. Pdr3 is required for DNA damage induction of MAG1 and DDI1 via a bi-directional promoter element. <i>Nucleic Acids Res</i> 2004;32(17):5066–75.	doi:10.1093/nar/gkh838 PMID 15452273 PMC521653	–	6	3	6	2	4	1	lacZ - wt vs TF mutant
552	Kim 2004 Kim TS, Kim HY, Yoon JH, Kang HS. Recruitment of the Swi/Snf complex by Ste12-Tec1 promotes Flo8-Mss11-mediated activation of STA1 expression. <i>Mol Cell Biol</i> 2004;24(21):9542–56.	doi:10.1128/MCB.24.21.9542-9556.2004 PMID 15485921 PMC522284	–	6	6	6	2	4	2	lacZ - wt vs TF mutant
553	Pic-Taylor 2004 Pic-Taylor A, Darieva Z, Morgan BA, Sharrocks AD. Regulation of cell cycle-specific gene expression through cyclin-dependent kinase-mediated phosphorylation of the forkhead transcription factor Fkh2p. <i>Mol Cell Biol</i> 2004;24(22):10036–46.	doi:10.1128/MCB.24.22.10036-10046.2004 PMID 15509804 PMC525469	–	6	2	6	4	2	2	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
554	Harkness 2004 Harkness TA, Shea KA, Legrand C, Brahmania M, Davies GF. A functional analysis reveals dependence on the anaphase-promoting complex for prolonged life span in yeast. <i>Genetics</i> 2004;168(2):759–74.	doi:10.1534/genetics.104.027771 PMID 15514051 PMC1448841	–	6	2	6	–	6	3	Northern blot - wt vs TF mutant
555	Raithatha 2005 Raithatha SA, Stuart DT. Meiosis-specific regulation of the <i>Saccharomyces cerevisiae</i> S-phase cyclin CLB5 is dependent on MluI cell cycle box (MCB) elements in its promoter but is independent of MCB-binding factor activity. <i>Genetics</i> 2005;169(3):1329–42.	doi:10.1534/genetics.104.036103 PMID 15654101 PMC1449548	–	6	4	6	–	5	3	Northern blot - wt vs TF mutant
556	Imazu 2005 Imazu H, Sakurai H. <i>Saccharomyces cerevisiae</i> heat shock transcription factor regulates cell wall remodeling in response to heat shock. <i>Eukaryot Cell</i> 2005;4(6):1050–6.	doi:10.1128/EC.4.6.1050-1056.2005 PMID 15947197 PMC1151985	–	6	1	6	–	6	1	RT-PCR - wt vs TF mutant
557	Keller 2005 Keller G, Bird A, Winge DR. Independent metalloreulation of Ace1 and Mac1 in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2005;4(11):1863–71.	doi:10.1128/EC.4.11.1863-1871.2005 PMID 16278453 PMC1287855	–	6	2	6	5	–	2	ChIP
558	Ishida 2006 Ishida C, Aranda C, Valenzuela L, et al., González A. The UGA3-GLT1 intergenic region constitutes a promoter whose bidirectional nature is determined by chromatin organization in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2006;59(6):1790–806.	doi:10.1111/j.1365-2958.2006.05055.x PMID 16553884	–	6	3	6	2	4	3	Northern blot - wt vs TF mutant
559	Wu 2007 Wu CY, Bird AJ, Winge DR, Eide DJ. Regulation of the yeast TSA1 peroxiredoxin by ZAP1 is an adaptive response to the oxidative stress of zinc deficiency. <i>J Biol Chem</i> 2007;282(4):2184–95.	doi:10.1074/jbc.M606639200 PMID 17121842	–	6	1	6	1	5	1	Microarray analysis - wt vs TF mutant
560	Godard 2007 Godard P, Urrestarazu A, Vissers S, et al., André B. Effect of 21 different nitrogen sources on global gene expression in the yeast <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2007;27(8):3065–86.	doi:10.1128/MCB.01084-06 PMID 17308034 PMC1899933	–	6	2	6	3	3	2	RT-PCR - wt vs TF mutant
561	Galeote 2007 Galeote VA, Alexandre H, Bach B, Delobel P, Dequin S, Blondin B. Sfl1p acts as an activator of the HSP30 gene in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 2007;52(2):55–63.	doi:10.1007/s00294-007-0136-z PMID 17594096	–	6	1	6	–	6	1	Microarray analysis - wt vs TF mutant
562	Lenssen 2007 Lenssen E, Azzouz N, Michel A, Landrieux E, Collart MA. The Ccr4-not complex regulates Skn7 through Srb10 kinase. <i>Eukaryot Cell</i> 2007;6(12):2251–9.	doi:10.1128/EC.00327-06 PMID 17965252 PMC2168244	–	6	2	6	6	–	3	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
563	Nishizawa 2008 Nishizawa M, Komai T, Morohashi N, Shimizu M, Toh-e A. Transcriptional repression by the Pho4 transcription factor controls the timing of SNZ1 expression. <i>Eukaryot Cell</i> 2008;7(6):949–57.	doi:10.1128/EC.00366-07 PMID 18408055 PMC2446651	–	6	3	6	2	4	3	Northern blot - wt vs TF mutant
564	Menezes 2008 Menezes RA, Amaral C, Batista-Nascimento L, et al., Rodrigues-Pousada C. Contribution of Yap1 towards <i>Saccharomyces cerevisiae</i> adaptation to arsenic-mediated oxidative stress. <i>Biochem J</i> 2008;414(2):301–11.	doi:10.1042/BJ20071537 PMID 18439143	–	6	2	6	–	6	1	RT-PCR - wt vs TF mutant
565	Lotito 2009 Lotito L, Russo A, Bueno S, Chillemi G, Fogli MV, Capranico G. A specific transcriptional response of yeast cells to camptothecin dependent on the Swi4 and Mbp1 factors. <i>Eur J Pharmacol</i> 2009;603(1–3):29–36.	doi:10.1016/j.ejphar.2008.12.002 PMID 19094980	–	6	2	6	–	6	2	RT-PCR - wt vs TF mutant
566	He 2009 He XJ, Mulford KE, Fassler JS. Oxidative stress function of the <i>Saccharomyces cerevisiae</i> Skn7 receiver domain. <i>Eukaryot Cell</i> 2009;8(5):768–78.	doi:10.1128/EC.00021-09 PMID 19304952 PMC2681607	–	6	2	6	3	3	4	EMSA
567	Joo 2009 Joo YJ, Kim JA, Baek JH, et al., Kim J. Cooperative regulation of ADE3 transcription by Gcn4p and Bas1p in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2009;8(8):1268–77.	doi:10.1128/EC.00116-09 PMID 19525417 PMC2725562	–	6	2	6	1	5	2	Northern blot - wt vs TF mutant
568	Pondugula 2009 Pondugula S, Neef DW, Voth WP, et al., Kladde MP. Coupling phosphate homeostasis to cell cycle-specific transcription: mitotic activation of <i>Saccharomyces cerevisiae</i> PHO5 by Mcm1 and Forkhead proteins. <i>Mol Cell Biol</i> 2009;29(18):4891–905.	doi:10.1128/MCB.00222-09 PMID 19596791 PMC2738290	–	6	3	6	6	–	1	ChIP
569	Nishizawa 2010 Nishizawa M, Tanigawa M, Hayashi M, et al., Toh-e A. Pho85 kinase, a cyclin-dependent kinase, regulates nuclear accumulation of the Rim101 transcription factor in the stress response of <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2010;9(6):943–51.	doi:10.1128/EC.00247-09 PMID 20382759 PMC2901641	–	6	1	6	1	5	1	Northern blot - wt vs TF mutant
570	Tsaponina 2011 Tsaponina O, Barsoum E, Aström SU, Chabes A. Ixr1 is required for the expression of the ribonucleotide reductase Rnr1 and maintenance of dNTP pools. <i>PLoS Genet</i> 2011;7(5):e1002061.	doi:10.1371/journal.pgen.1002061 PMID 21573136 PMC3088718	–	6	1	6	3	3	2	ChIP
571	Lavut 2012 Lavut A, Raveh D. Sequestration of highly expressed mRNAs in cytoplasmic granules, P-bodies, and stress granules enhances cell viability. <i>PLoS Genet</i> 2012;8(2):e1002527.	doi:10.1371/journal.pgen.1002527 PMID 22383896 PMC3285586	–	6	2	2	–	6	3	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
572	Nosaka 2012 Nosaka K, Esaki H, Onozuka M, Konno H, Hattori Y, Akaji K. Facilitated recruitment of Pdc2p, a yeast transcriptional activator, in response to thiamin starvation. <i>FEMS Microbiol Lett</i> 2012;330(2):140–7.	doi:10.1111/j.1574-6968.2012.02543.x PMID 22404710	–	6	2	6	6	–	2	ChIP
573	Ruiz-Roig 2012 Ruiz-Roig C, Noriega N, Duch A, Posas F, de Nadal E. The Hog1 SAPK controls the Rtg1/Rtg3 transcriptional complex activity by multiple regulatory mechanisms. <i>Mol Biol Cell</i> 2012;23(21):4286–96.	doi:10.1091/mbc.E12-04-0289 PMID 22956768 PMC3484105	–	6	2	6	2	4	1	Northern blot - wt vs TF mutant
574	Shetty 2013 Shetty A, Swaminathan A, Lopes JM. Transcription regulation of a yeast gene from a downstream location. <i>J Mol Biol</i> 2013;425(3):457–65.	doi:10.1016/j.jmb.2012.11.018 PMID 23174187	–	6	3	3	–	6	2	RT-PCR - wt vs TF mutant
575	Han 2013 Han BK, Emr SD. The phosphatidylinositol 3,5-bisphosphate (PI(3,5)P2)-dependent Tup1 conversion (PIPTC) regulates metabolic reprogramming from glycolysis to gluconeogenesis. <i>J Biol Chem</i> 2013;288(28):20633–45.	doi:10.1074/jbc.M113.452813 PMID 23733183 PMC3711327	–	6	2	6	3	3	2	ChIP
576	Gururaj 2013 Gururaj C, Federman RS, Chang A. Orm proteins integrate multiple signals to maintain sphingolipid homeostasis. <i>J Biol Chem</i> 2013;288(28):20453–63.	doi:10.1074/jbc.M113.472860 PMID 23737533 PMC3711311	–	6	4	6	–	6	3	RT-PCR - wt vs TF mutant
577	Blanda 2013 Blanda C, Höfken T. Regulation of mating in the budding yeast <i>Saccharomyces cerevisiae</i> by the zinc cluster proteins Sut1 and Sut2. <i>Biochem Biophys Res Commun</i> 2013;438(1):66–70.	doi:10.1016/j.bbrc.2013.07.027 PMID 23872066	–	6	2	6	–	6	1	lacZ - wt vs TF mutant
578	Gallo-Ebert 2013 Gallo-Ebert C, Donigan M, Liu HY, et al., Nickels JT Jr. The yeast anaerobic response element AR1b regulates aerobic antifungal drug-dependent sterol gene expression. <i>J Biol Chem</i> 2013;288(49):35466–77.	doi:10.1074/jbc.M113.526087 PMID 24163365 PMC3853293	–	6	2	6	–	6	3	Northern blot - wt vs TF mutant
579	Yelamanchi 2014 Yelamanchi SK, Veis J, Anrather D, Klug H, Ammerer G. Genotoxic stress prevents Ndd1-dependent transcriptional activation of G2/M-specific genes in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2014;34(4):711–24.	doi:10.1128/MCB.01090-13 PMID 24324010 PMC3911480	–	6	2	6	6	–	1	ChIP
580	Fayyadkazan 2014 Fayyadkazan M, Tate JJ, Vierendeels F, Cooper TG, Dubois E, Georis I. Components of Golgi-to-vacuole trafficking are required for nitrogen- and TORC1-responsive regulation of the yeast GATA factors. <i>Microbiologyopen</i> 2014;3(3):271–87.	doi:10.1002/mbo3.168 PMID 24644271 PMC4082702	–	6	2	6	–	6	1	RT-PCR - wt vs TF mutant
581	Mayhew 2014 Mayhew D, Mitra RD. Transcription factor regulation and chromosome dynamics during pseudohyphal growth. <i>Mol Biol Cell</i> 2014;25(17):2669–76.	doi:10.1091/mbc.E14-04-0871 PMID 25009286 PMC4148256	–	6	5	6	6	–	1	Multiplex calling card analysis

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
582	Imamura 2015 Imamura Y, Yu F, Nakamura M, et al., Tsuchiya E. RSC Chromatin-Remodeling Complex Is Important for Mitochondrial Function in <i>Saccharomyces cerevisiae</i> . <i>PLoS One</i> 2015;10(6):e0130397.	doi:10.1371/journal.pone.0130397 PMID 26086550 PMC4472808	–	6	1	6	–	6	1	RT-PCR - wt vs TF mutant
583	Palavecino 2015 Palavecino MD, Correa-García SR, Bermúdez-Moretti M. Genes of Different Catabolic Pathways Are Coordinately Regulated by Dal81 in <i>Saccharomyces cerevisiae</i> . <i>J Amino Acids</i> 2015;2015:484702.	doi:10.1155/2015/484702 PMID 26457198 PMC4589623	–	6	2	6	2	4	4	ChIP
584	Milias-Argeitis 2016 Milias-Argeitis A, Oliveira AP, Gerosa L, Falter L, Sauer U, Lygeros J. Elucidation of Genetic Interactions in the Yeast GATA-Factor Network Using Bayesian Model Selection. <i>PLoS Comput Biol</i> 2016;12(3):e1004784.	doi:10.1371/journal.pcbi.1004784 PMID 26967983 PMC4788432	–	6	4	6	–	6	2	GFP - wt vs TF mutant
585	Gil 2017 Gil FN, Bellí G, Viegas CA. The <i>Saccharomyces cerevisiae</i> response to stress caused by the herbicidal active substance alachlor requires the iron regulon transcription factor Aft1p. <i>Environ Microbiol</i> 2017;19(2):485–499.	doi:10.1111/1462-2920.13439 PMID 27376881	–	6	1	3	–	6	1	RT-PCR - wt vs TF mutant
586	Cho 2017 Cho BR, Hahn JS. CK2-dependent phosphorylation positively regulates stress-induced activation of Msn2 in <i>Saccharomyces cerevisiae</i> . <i>Biochim Biophys Acta Gene Regul Mech</i> 2017; 1860(6):695–704.	doi:10.1016/j.bbagr.2017.03.003 PMID 28330760	–	6	1	2	–	6	3	RT-PCR - wt vs TF mutant
587	Henriques 2017 Henriques SF, Mira NP, Sá-Correia I. Genome-wide search for candidate genes for yeast robustness improvement against formic acid reveals novel susceptibility (Trk1 and positive regulators) and resistance (Haa1-regulon) determinants. <i>Biotechnol Biofuels</i> 2017;10:96.	doi:10.1186/s13068-017-0781-5 PMID 28428821 PMC5395885	–	6	1	5	–	6	2	RT-PCR - wt vs TF mutant
588	Chang 2018 Chang N, Yao S, Chen D, Zhang L, Huang J, Zhang L. The Hog1 positive regulated YCT1 gene expression under cadmium tolerance of budding yeast. <i>FEMS Microbiol Lett</i> 2018;365(17).	doi:10.1093/femsle/fny170 PMID 29982432	–	6	4	4	–	6	3	lacZ - wt vs TF mutant
589	Shi 2018 Shi X, Zou Y, Chen Y, Ying H. Overexpression of THI4 and HAP4 Improves Glucose Metabolism and Ethanol Production in <i>Saccharomyces cerevisiae</i> . <i>Front Microbiol</i> 2018;9:1444.	doi:10.3389/fmicb.2018.01444 PMID 29997610 PMC6030257	–	6	1	6	–	6	1	RT-PCR - wt vs TF overexpression
590	DeMille 2019 DeMille D, Pape JA, Bikman BT, Ghassemian M, Grose JH. The Regulation of Cbfl by PAS Kinase Is a Pivotal Control Point for Lipogenesis vs. Respiration in <i>Saccharomyces cerevisiae</i> . <i>G3 (Bethesda)</i> 2019;9(1):33–46.	doi:10.1534/g3.118.200663 PMID 30381292 PMC6325914	–	6	1	4	3	3	1	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
591	Simpson-Lavy 2019 Simpson-Lavy K, Kupiec M. Carbon Catabolite Repression in Yeast is Not Limited to Glucose. <i>Sci Rep</i> 2019;9(1):6491.	doi:10.1038/s41598-019-43032-w PMID 31019232 PMC6482301	–	6	3	4	–	6	2	lacZ - wt vs TF mutant
592	Vanacloig-Pedros 2019 Vanacloig-Pedros E, Lozano-Pérez C, Alarcón B, Pascual-Ahuir A, Proft M. Live-cell assays reveal selectivity and sensitivity of the multidrug response in budding yeast. <i>J Biol Chem</i> 2019;294(35):12933–12946.	doi:10.1074/jbc.RA119.009291 PMID 31296662 PMC6721934	–	6	2	6	–	6	1	Luciferase - wt vs TF mutant
593	Kasahara 2020 Kasahara K, Nakayama R, Shiwa Y, et al., Kokubo T. Fpr1, a primary target of rapamycin, functions as a transcription factor for ribosomal protein genes cooperatively with Hmo1 in <i>Saccharomyces cerevisiae</i> . <i>PLoS Genet</i> 2020;16(6):e1008865.	doi:10.1371/journal.pgen.1008865 PMID 32603360 PMC7357790	–	6	1	6	6	–	1	ChIP-seq
594	Pujol-Carrion 2021 Pujol-Carrion N, Pavón-Vergés M, Arroyo J, de la Torre-Ruiz MA. The MAPK Slt2/Mpk1 plays a role in iron homeostasis through direct regulation of the transcription factor Aft1. <i>Biochim Biophys Acta Mol Cell Res</i> 2021;1868(5):118974.	doi:10.1016/j.bbamcr.2021.118974 PMID 33549702	–	6	2	2	–	6	3	lacZ - wt vs TF mutant
595	Trawick 1992 Trawick JD, Kraut N, Simon FR, Poyton RO. Regulation of yeast COX6 by the general transcription factor ABF1 and separate HAP2- and heme-responsive elements. <i>Mol Cell Biol</i> 1992;12(5):2302–14.	doi:10.1128/mcb.12.5.2302-2314.1992 PMID 1314953 PMC364402	–	5	5	5	4	1	2	lacZ - wt vs mutant promoter sequence
596	Oechsner 1992 Oechsner U, Hermann H, Zollner A, Haid A, Bandlow W. Expression of yeast cytochrome c1 is controlled at the transcriptional level by glucose, oxygen and haem. <i>Mol Gen Genet</i> 1992;232(3):447–59.	doi:10.1007/BF00266250 PMID 1316998	–	5	5	5	3	2	2	lacZ - wt vs mutant promoter sequence
597	Dirick 1992 Dirick L, Möll T, Auer H, Nasmyth K. A central role for SWI6 in modulating cell cycle Start-specific transcription in yeast. <i>Nature</i> 1992;357(6378):508–13.	doi:10.1038/357508a0 PMID 1608451	–	5	2	5	1	–	1	–
598	Nehlin 1991 Nehlin JO, Carlberg M, Ronne H. Control of yeast GAL genes by MIG1 repressor: a transcriptional cascade in the glucose response. <i>EMBO J</i> 1991;10(11):3373–7.	doi:10.1002/j.1460-2075.1991.tb04901.x PMID 1915298 PMC453065	–	5	2	5	2	3	3	Northern blot - wt vs TF mutant
599	Cooper 1990 Cooper TG, Ferguson D, Rai R, Bysani N. The GLN3 gene product is required for transcriptional activation of allantoin system gene expression in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1990;172(2):1014–8.	doi:10.1128/jb.172.2.1014-1018.1990 PMID 2153652 PMC208530	–	5	1	5	–	5	1	Northern blot - wt vs TF mutant
600	Dorsman 1990 Dorsman JC, Grivell LA. Expression of the gene encoding subunit II of yeast QH2: cytochrome c oxidoreductase is regulated by multiple factors. <i>Curr Genet</i> 1990;17(6):459–64.	doi:10.1007/BF00313072 PMID 2167771	–	5	5	5	4	1	2	lacZ - wt vs mutant promoter sequence

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
601	Mitchell 1990 Mitchell AP, Driscoll SE, Smith HE. Positive control of sporulation-specific genes by the IME1 and IME2 products in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1990;10(5):2104–10.	doi:10.1128/mcb.10.5.2104-2110.1990 PMID 2183020 PMC360558	–	5	1	5	–	5	1	Northern blot - wt vs TF mutant
602	Trueblood 1988 Trueblood CE, Wright RM, Poyton RO. Differential regulation of the two genes encoding <i>Saccharomyces cerevisiae</i> cytochrome c oxidase subunit V by heme and the HAP2 and REO1 genes. <i>Mol Cell Biol</i> 1988;8(10):4537–40.	doi:10.1128/mcb.8.10.4537-4540.1988 PMID 2847035 PMC365531	–	5	5	5	–	2	2	–
603	Fields 1988 Fields S, Chaleff DT, Sprague GF Jr. Yeast STE7, STE11, and STE12 genes are required for expression of cell-type-specific genes. <i>Mol Cell Biol</i> 1988;8(2):551–6.	doi:10.1128/mcb.8.2.551-556.1988 PMID 3280969 PMC363179	–	5	1	5	–	5	1	Northern blot - wt vs TF mutant
604	Holland 1987 Holland MJ, Yokoi T, Holland JP, Myambo K, Innis MA. The GCR1 gene encodes a positive transcriptional regulator of the enolase and glyceraldehyde-3-phosphate dehydrogenase gene families in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1987;7(2):813–20.	doi:10.1128/mcb.7.2.813-820.1987 PMID 3547083 PMC365139	–	5	1	5	–	5	1	Western blot - wt vs TF mutant
605	Stanbrough 1995 Stanbrough M, Rowen DW, Magasanik B. Role of the GATA factors Gln3p and Nil1p of <i>Saccharomyces cerevisiae</i> in the expression of nitrogen-regulated genes. <i>Proc Natl Acad Sci U S A</i> 1995;92(21):9450–4.	doi:10.1073/pnas.92.21.9450 PMID 7568152 PMC40819	–	5	2	5	–	5	3	lacZ - wt vs TF mutant
606	Betina 1995 Betina S, Gavurníková G, Havierník P, Sabová L, Kolarov J. Expression of the AAC2 gene encoding the major mitochondrial ADP/ATP carrier in <i>Saccharomyces cerevisiae</i> is controlled at the transcriptional level by oxygen, heme and HAP2 factor. <i>Eur J Biochem</i> 1995;229(3):651–7.	doi:10.1111/j.1432-1033.1995.tb20510.x PMID 7758459	–	5	5	5	5	–	2	lacZ - wt vs mutant promoter sequence
607	O’Connell 1995 O’Connell KF, Surdin-Kerjan Y, Baker RE. Role of the <i>Saccharomyces cerevisiae</i> general regulatory factor CP1 in methionine biosynthetic gene transcription. <i>Mol Cell Biol</i> 1995;15(4):1879–88.	doi:10.1128/MCB.15.4.1879 PMID 7891681 PMC230413	–	5	3	5	–	5	2	Northern blot - wt vs TF mutant
608	Crawford 1995 Crawford MJ, Sherman DR, Goldberg DE. Regulation of <i>Saccharomyces cerevisiae</i> flavohemoglobin gene expression. <i>J Biol Chem</i> 1995;270(12):6991–6.	doi:10.1074/jbc.270.12.6991 PMID 7896850	–	5	5	5	–	2	2	–
609	Hohmann 1993 Hohmann S. Characterisation of PDC2, a gene necessary for high level expression of pyruvate decarboxylase structural genes in <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1993;241(5–6):657–66.	doi:10.1007/BF00279908 PMID 8264540	–	5	1	2	–	5	2	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
610	Rolfes 1993 Rolfes RJ, Hinnebusch AG. Translation of the yeast transcriptional activator GCN4 is stimulated by purine limitation: implications for activation of the protein kinase GCN2. <i>Mol Cell Biol</i> 1993;13(8):5099–111.	doi:10.1128/mcb.13.8.5099-5111.1993 PMID 8336737 PMC360163	–	5	1	5	–	5	2	Northern blot - wt vs TF mutant
611	Cunningham 1993 Cunningham TS, Cooper TG. The <i>Saccharomyces cerevisiae</i> DAL80 repressor protein binds to multiple copies of GATAA-containing sequences (URSGATA). <i>J Bacteriol</i> 1993;175(18):5851–61.	doi:10.1128/jb.175.18.5851-5861.1993 PMID 8376332 PMC206664	–	5	1	5	5	–	2	EMSA
612	Rahner 1996 Rahner A, Schöler A, Martens E, Gollwitzer B, Schüller HJ. Dual influence of the yeast Cat1p (Snf1p) protein kinase on carbon source-dependent transcriptional activation of gluconeogenic genes by the regulatory gene CAT8. <i>Nucleic Acids Res</i> 1996;24(12):2331–7.	doi:10.1093/nar/24.12.2331 PMID 8710504 PMC145921	–	5	3	5	–	5	2	lacZ - wt vs TF mutant
613	Watt 1997 Watt R, Piper PW. UBI4, the polyubiquitin gene of <i>Saccharomyces cerevisiae</i> , is a heat shock gene that is also subject to catabolite derepression control. <i>Mol Gen Genet</i> 1997;253(4):439–47.	doi:10.1007/s004380050341 PMID 9037103	–	5	5	5	4	1	2	lacZ - wt vs mutant promoter sequence
614	Zhong 1997 Zhong H, Vershon AK. The yeast homeodomain protein MATA1alpha2 shows extended DNA binding specificity in complex with Mcm1. <i>J Biol Chem</i> 1997;272(13):8402–9.	doi:10.1074/jbc.272.13.8402 PMID 9079665	–	5	1	5	1	4	2	lacZ - wt vs TF mutant
615	Pinkham 1997 Pinkham JL, Wang Z, Alsina J. Heme regulates SOD2 transcription by activation and repression in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 1997;31(4):281–91.	doi:10.1007/s002940050207 PMID 9108135	–	5	5	5	5	–	1	lacZ - wt vs mutant promoter sequence
616	Sweet 1997 Sweet DH, Jang YK, Sancar GB. Role of UME6 in transcriptional regulation of a DNA repair gene in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1997;17(11):6223–35.	doi:10.1128/MCB.17.11.6223 PMID 9343383 PMC232473	–	5	1	5	4	1	2	EMSA
617	Schroeder 1998 Schroeder SC, Weil PA. Genetic tests of the role of Abf1p in driving transcription of the yeast TATA box binding protein-encoding gene, SPT15. <i>J Biol Chem</i> 1998;273(31):19884–91.	doi:10.1074/jbc.273.31.19884 PMID 9677425	–	5	1	5	5	–	1	DNA footprinting
618	Huang 1998 Huang M, Zhou Z, Elledge SJ. The DNA replication and damage checkpoint pathways induce transcription by inhibition of the Crt1 repressor. <i>Cell</i> 1998;94(5):595–605.	doi:10.1016/s0092-8674(00)81601-3 PMID 9741624	–	5	1	5	5	–	3	ChIP
619	Vincent 1998 Vincent O, Carlson M. Sip4, a Snf1 kinase-dependent transcriptional activator, binds to the carbon source-responsive element of gluconeogenic genes. <i>EMBO J</i> 1998;17(23):7002–8.	doi:10.1093/emboj/17.23.7002 PMID 9843506 PMC1171048	–	5	2	5	1	4	3	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
620	Xie 1999 Xie J, Pierce M, Gailus-Durner V, Wagner M, Winter E, Vershon AK. Sum1 and Hst1 repress middle sporulation-specific gene expression during mitosis in <i>Saccharomyces cerevisiae</i> . <i>EMBO J</i> 1999;18(22):6448–54.	doi:10.1093/emboj/18.22.6448 PMID 10562556 PMC1171707	–	5	2	5	–	4	2	Northern blot - wt vs TF mutant
621	Lodi 1999 Lodi T, Alberti A, Guiard B, Ferrero I. Regulation of the <i>Saccharomyces cerevisiae</i> DLD1 gene encoding the mitochondrial protein D-lactate ferricytochrome c oxidoreductase by HAP1 and HAP2/3/4/5. <i>Mol Gen Genet</i> 1999;262(4–5):623–32.	doi:10.1007/s004380051125 PMID 10628845	–	5	5	5	5	–	1	lacZ - wt vs mutant promoter sequence
622	de Boer 2000 de Boer M, Nielsen PS, Bebelman JP, Heerikhuizen H, Andersen HA, Planta RJ. Stp1p, Stp2p and Abf1p are involved in regulation of expression of the amino acid transporter gene BAP3 of <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Res</i> 2000;28(4):974–81.	doi:10.1093/nar/28.4.974 PMID 10648791 PMC102570	–	5	3	5	2	3	1	Northern blot - wt vs TF mutant
623	Llorente 2000 Llorente B, Dujon B. Transcriptional regulation of the <i>Saccharomyces cerevisiae</i> DAL5 gene family and identification of the high affinity nicotinic acid permease TNA1 (YGR260w). <i>FEBS Lett</i> 2000;475(3):237–41.	doi:10.1016/s0014-5793(00)01698-7 PMID 10869563	–	5	1	5	–	2	2	–
624	Yuan 2000 Yuan DS. Zinc-regulated genes in <i>Saccharomyces cerevisiae</i> revealed by transposon tagging. <i>Genetics</i> 2000;156(1):45–58.	doi:10.1093/genetics/156.1.45 PMID 10978274 PMC1461233	–	5	1	5	–	5	1	lacZ - wt vs TF mutant
625	Lamb 2001 Lamb TM, Xu W, Diamond A, Mitchell AP. Alkaline response genes of <i>Saccharomyces cerevisiae</i> and their relationship to the RIM101 pathway. <i>J Biol Chem</i> 2001;276(3):1850–6.	doi:10.1074/jbc.M008381200 PMID 11050096	–	5	1	5	–	5	1	Northern blot - wt vs TF mutant
626	Jensen-Pergakes 2001 Jensen-Pergakes K, Guo Z, Giattina M, Sturley SL, Bard M. Transcriptional regulation of the two sterol esterification genes in the yeast <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 2001;183(17):4950–7.	doi:10.1128/JB.183.17.4950-4957.2001 PMID 11489845 PMC95368	–	5	2	3	–	5	2	lacZ - wt vs TF mutant
627	Mai 2002 Mai B, Miles S, Breeden LL. Characterization of the ECB binding complex responsible for the M/G(1)-specific transcription of CLN3 and SWI4. <i>Mol Cell Biol</i> 2002;22(2):430–41.	doi:10.1128/MCB.22.2.430-441.2002 PMID 11756540 PMC139728	–	5	1	5	5	–	2	ChIP
628	Hellauer 2002 Hellauer K, Akache B, MacPherson S, Sirard E, Turcotte B. Zinc cluster protein Rdr1p is a transcriptional repressor of the PDR5 gene encoding a multidrug transporter. <i>J Biol Chem</i> 2002;277(20):17671–6.	doi:10.1074/jbc.M201637200 PMID 11882665	–	5	1	5	–	5	1	Microarray analysis - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	Tfs	pairs	direct	indirect	cond.	top assay
629	Turner 2002 Turner SD, Ricci AR, Petropoulos H, Genereaux J, Skerjanc IS, Brandl CJ. The E2 ubiquitin conjugase Rad6 is required for the ArgR/Mcm1 repression of ARG1 transcription. <i>Mol Cell Biol</i> 2002;22(12):4011–9.	doi:10.1128/MCB.22.12.4011-4019.2002 PMID 12024015 PMC133851	–	5	5	5	3	2	1	lacZ - wt vs mutant promoter sequence
630	Köhler 2002 Köhler T, Wesche S, Taheri N, Braus GH, Mösch HU. Dual role of the <i>Saccharomyces cerevisiae</i> TEA/ATTS family transcription factor Tec1p in regulation of gene expression and cellular development. <i>Eukaryot Cell</i> 2002;1(5):673–86.	doi:10.1128/EC.1.5.673-686.2002 PMID 12455687 PMC126755	–	5	2	5	–	5	3	lacZ - wt vs TF mutant
631	De Freitas 2004 De Freitas JM, Kim JH, Poynton H, et al., Vulpe C. Exploratory and confirmatory gene expression profiling of mac1Delta. <i>J Biol Chem</i> 2004;279(6):4450–8.	doi:10.1074/jbc.M212308200 PMID 14534306	–	5	1	5	–	5	1	RT-PCR - wt vs TF mutant
632	Alberti 2003 Alberti A, Lodi T, Ferrero I, Donnini C. MIG1-dependent and MIG1-independent regulation of GAL gene expression in <i>Saccharomyces cerevisiae</i> : role of Imp2p. <i>Yeast</i> 2003;20(13):1085–96.	doi:10.1002/yea.1025 PMID 14558142	–	5	1	5	–	5	2	Northern blot - wt vs TF mutant
633	Yarragudi 2004 Yarragudi A, Miyake T, Li R, Morse RH. Comparison of ABF1 and RAP1 in chromatin opening and transactivator potentiation in the budding yeast <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2004;24(20):9152–64.	doi:10.1128/MCB.24.20.9152-9164.2004 PMID 15456886 PMC517901	–	5	2	5	5	–	1	ALPHA-galactosidase WT vs target promoter mutant
634	Avendaño 2005 Avendaño A, Riego L, DeLuna A, et al., González A. Swi/SNF-GCN5-dependent chromatin remodelling determines induced expression of GDH3, one of the paralogous genes responsible for ammonium assimilation and glutamate biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2005; 57(1):291–305.	doi:10.1111/j.1365-2958.2005.04689.x PMID 15948967	–	5	5	5	–	5	2	lacZ - wt vs TF mutant
635	Sarver 2005 Sarver A, DeRisi J. Fzflp regulates an inducible response to nitrosative stress in <i>Saccharomyces cerevisiae</i> . <i>Mol Biol Cell</i> 2005;16(10):4781–91.	doi:10.1091/mbc.e05-05-0436 PMID 16014606 PMC1237083	–	5	1	5	–	5	2	lacZ - wt vs TF overexpression
636	Chen 2006 Chen H, Fink GR. Feedback control of morphogenesis in fungi by aromatic alcohols. <i>Genes Dev</i> 2006;20(9):1150–61.	doi:10.1101/gad.1411806 PMID 16618799 PMC1472474	–	5	4	5	–	5	2	RT-PCR - wt vs TF mutant
637	Koehler 2007 Koehler RN, Rachfall N, Rolfes RJ. Activation of the ADE genes requires the chromatin remodeling complexes SAGA and SWI/SNF. <i>Eukaryot Cell</i> 2007;6(8):1474–85.	doi:10.1128/EC.00068-07 PMID 17573544 PMC1951130	–	5	2	5	2	3	3	ChIP
638	Hirasaki 2008 Hirasaki M, Kaneko Y, Harashima S. Protein phosphatase Siw14 controls intracellular localization of Gln3 in cooperation with Npr1 kinase in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> 2008; 409(1–2):34–43.	doi:10.1016/j.gene.2007.11.005 PMID 18166280	–	5	1	5	–	5	1	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
639	Morohashi 2008 Morohashi N, Nakajima K, Kuwana S, Tachiwana H, Kurumizaka H, Shimizu M. In vivo and in vitro footprinting of nucleosomes and transcriptional activators using an infrared-fluorescence DNA sequencer. <i>Biol Pharm Bull</i> 2008; 31(2):187–92.	doi:10.1248/bpb.31.187 PMID 18239271	–	5	5	5	5	–	1	In vivo UV-photofootprinting
640	Ikeda 2008 Ikeda M, Kihara A, Denpoh A, Igarashi Y. The Rim101 pathway is involved in Rsb1 expression induced by altered lipid asymmetry. <i>Mol Biol Cell</i> 2008;19(5):1922–31.	doi:10.1091/mbc.e07-08-0806 PMID 18287536 PMC2366876	–	5	4	4	–	5	3	Western blot - wt vs TF mutant
641	Izquierdo 2008 Izquierdo A, Casas C, Mühlenhoff U, Lillig CH, Herrero E. Saccharomyces cerevisiae Grx6 and Grx7 are monothiol glutaredoxins associated with the early secretory pathway. <i>Eukaryot Cell</i> 2008;7(8):1415–26.	doi:10.1128/EC.00133-08 PMID 18503006 PMC2519769	–	5	3	5	–	5	3	Northern blot - wt vs TF mutant
642	Pitoniak 2009 Pitoniak A, Birkaya B, Dionne HM, Vadaie N, Cullen PJ. The signaling mucins Msb2 and Hkr1 differentially regulate the filamentation mitogen-activated protein kinase pathway and contribute to a multimodal response. <i>Mol Biol Cell</i> 2009; 20(13):3101–14.	doi:10.1091/mbc.e08-07-0760 PMID 19439450 PMC2704161	–	5	1	5	–	5	1	lacZ - wt vs TF mutant
643	Furukawa 2009 Furukawa K, Sidoux-Walter F, Hohmann S. Expression of the yeast aquaporin Aqp2 affects cell surface properties under the control of osmoregulatory and morphogenic signalling pathways. <i>Mol Microbiol</i> 2009;74(5):1272–86.	doi:10.1111/j.1365-2958.2009.06933.x PMID 19889095	–	5	5	5	–	5	2	lacZ - wt vs TF mutant
644	Gurevich 2010 Gurevich V, Kassir Y. A switch from a gradient to a threshold mode in the regulation of a transcriptional cascade promotes robust execution of meiosis in budding yeast. <i>PLoS One</i> 2010; 5(6):e11005.	doi:10.1371/journal.pone.0011005 PMID 20543984 PMC2882377	–	5	1	5	5	–	1	RT-PCR - wt vs mutant promoter sequence
645	Dutoit 2010 Dutoit R, Dubois E, Jacobs E. Selection systems based on dominant-negative transcription factors for precise genetic engineering. <i>Nucleic Acids Res</i> 2010;38(19):e183.	doi:10.1093/nar/gkq708 PMID 20702421 PMC2965260	–	5	2	5	5	–	2	RT-PCR - wt vs mutant promoter sequence
646	Arroyo 2011 Arroyo J, Hutzler J, Bermejo C, et al., Strahl S. Functional and genomic analyses of blocked protein O-mannosylation in baker's yeast. <i>Mol Microbiol</i> 2011;79(6):1529–46.	doi:10.1111/j.1365-2958.2011.07537.x PMID 21231968	–	5	1	5	–	5	1	RT-PCR - wt vs TF mutant
647	Raab 2011 Raab AM, Hlavacek V, Bolotina N, Lang C. Shifting the fermentative/oxidative balance in <i>Saccharomyces cerevisiae</i> by transcriptional deregulation of Snf1 via overexpression of the upstream activating kinase Sak1p. <i>Appl Environ Microbiol</i> 2011;77(6):1981–9.	doi:10.1128/AEM.02219-10 PMID 21257817 PMC3067313	–	5	1	5	5	–	2	RT-PCR - wt vs mutant promoter sequence

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
648	Montañés 2011 Montañés FM, Pascual-Ahuir A, Proft M. Repression of ergosterol biosynthesis is essential for stress resistance and is mediated by the Hog1 MAP kinase and the Mot3 and Rox1 transcription factors. <i>Mol Microbiol</i> 2011;79(4):1008–23.	doi:10.1111/j.1365-2958.2010.07502.x PMID 21299653	–	5	3	5	1	4	2	Northern blot - wt vs TF mutant
649	Takahata 2011 Takahata S, Yu Y, Stillman DJ. Repressive chromatin affects factor binding at yeast HO (homothallic switching) promoter. <i>J Biol Chem</i> 2011;286(40):34809–19.	doi:10.1074/jbc.M111.281626 PMID 21840992 PMC3186355	–	5	3	5	5	–	1	ChIP
650	Li 2011 Li L, Jia X, Ward DM, Kaplan J. Yap5 protein-regulated transcription of the TYW1 gene protects yeast from high iron toxicity. <i>J Biol Chem</i> 2011;286(44):38488–38497.	doi:10.1074/jbc.M111.286666 PMID 21917924 PMC3207407	–	5	1	5	2	3	1	Microarray analysis - wt vs TF mutant
651	He 2012 He Y, Swaminathan A, Lopes JM. Transcription regulation of the <i>Saccharomyces cerevisiae</i> PHO5 gene by the Ino2p and Ino4p basic helix-loop-helix proteins. <i>Mol Microbiol</i> 2012; 83(2):395–407.	doi:10.1111/j.1365-2958.2011.07941.x PMID 22182244	–	5	5	5	3	2	2	ChIP
652	Brion 2013 Brion C, Ambroset C, Sanchez I, Legras JL, Blondin B. Differential adaptation to multi-stressed conditions of wine fermentation revealed by variations in yeast regulatory networks. <i>BMC Genomics</i> 2013;14:681.	doi:10.1186/1471-2164-14-681 PMID 24094006 PMC3870980	GSE41025	5	1	5	–	5	1	Microarray analysis - wt vs TF mutant
653	Pascual 2013 Pascual F, Soto-Cardalda A, Carman GM. PAH1-encoded phosphatidate phosphatase plays a role in the growth phase- and inositol-mediated regulation of lipid synthesis in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2013;288(50): 35781–92.	doi:10.1074/jbc.M113.525766 PMID 24196957 PMC3861629	–	5	5	5	–	5	1	lacZ - wt vs TF mutant
654	Kim 2014 Kim HY, Lee SB, Kang HS, Oh GT, Kim T. Two distinct domains of Flo8 activator mediates its role in transcriptional activation and the physical interaction with Mss11. <i>Biochem Biophys Res Commun</i> 2014;449(2):202–7.	doi:10.1016/j.bbrc.2014.04.161 PMID 24813990	–	5	2	5	1	4	1	lacZ - wt vs TF mutant
655	Burns 2014 Burns LT, Wentz SR. Casein kinase II regulation of the Hot1 transcription factor promotes stochastic gene expression. <i>J Biol Chem</i> 2014;289(25):17668–79.	doi:10.1074/jbc.M114.561217 PMID 24817120 PMC4067201	–	5	1	5	5	–	1	ChIP
656	Pougach 2014 Pougach K, Voet A, Kondrashov FA, et al., Verstrepen KJ. Duplication of a promiscuous transcription factor drives the emergence of a new regulatory network. <i>Nat Commun</i> 2014;5: 4868.	doi:10.1038/ncomms5868 PMID 25204769 PMC4172970	–	5	1	5	5	–	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
657	Andreadis 2014 Andreadis C, Nikolaou C, Fragiadakis GS, Tsiliki G, Alexandraki D. Rad9 interacts with Aft1 to facilitate genome surveillance in fragile genomic sites under non-DNA damage-inducing conditions in <i>S. cerevisiae</i> . <i>Nucleic Acids Res</i> 2014;42(20):12650–67.	doi:10.1093/nar/gku915 PMID 25300486 PMC4227768	–	5	1	5	5	–	1	ChIP
658	Laskar 2015 Laskar S, K S, Bhattacharyya MK, Nair AS, Dhar P, Bhattacharyya S. Heat stress-induced Cup9-dependent transcriptional regulation of SIR2. <i>Mol Cell Biol</i> 2015;35(2):437–50.	doi:10.1128/MCB.01046-14 PMID 25384977 PMC4272432	–	5	4	5	–	5	3	Western blot - wt vs TF mutant
659	Cahyani 2015 Cahyani I, Cridge AG, Engelke DR, Ganley AR, O'Sullivan JM. A sequence-specific interaction between the <i>Saccharomyces cerevisiae</i> rRNA gene repeats and a locus encoding an RNA polymerase I subunit affects ribosomal DNA stability. <i>Mol Cell Biol</i> 2015;35(3):544–54.	doi:10.1128/MCB.01249-14 PMID 25421713 PMC4285424	–	5	5	5	–	5	1	RT-PCR - wt vs TF mutant
660	Nishida-Aoki 2015 Nishida-Aoki N, Mori H, Kuroda K, Ueda M. Activation of the mitochondrial signaling pathway in response to organic solvent stress in yeast. <i>Curr Genet</i> 2015;61(2):153–64.	doi:10.1007/s00294-014-0463-9 PMID 25487302	–	5	5	5	–	5	1	RT-PCR - wt vs TF mutant
661	Kim 2015 Kim D, Song JY, Hahn JS. Improvement of glucose uptake rate and production of target chemicals by overexpressing hexose transporters and transcriptional activator Gcr1 in <i>Saccharomyces cerevisiae</i> . <i>Appl Environ Microbiol</i> 2015;81(24):8392–401.	doi:10.1128/AEM.02056-15 PMID 26431967 PMC4644637	–	5	1	5	5	–	1	RT-PCR - wt vs mutant promoter sequence
662	Li 2016 Li T, De Clercq N, Medina DA, et al., Alepuz P. The mRNA cap-binding protein Cbc1 is required for high and timely expression of genes by promoting the accumulation of gene-specific activators at promoters. <i>Biochim Biophys Acta</i> 2016;1859(2):405–19.	doi:10.1016/j.bbagr.2016.01.002 PMID 26775127	–	5	2	5	5	–	2	ChIP
663	Prugar 2017 Prugar E, Burnett C, Chen X, Hollingsworth NM. Coordination of Double Strand Break Repair and Meiotic Progression in Yeast by a Mek1-Ndt80 Negative Feedback Loop. <i>Genetics</i> 2017;206(1):497–512.	doi:10.1534/genetics.117.199703 PMID 28249986 PMC5419491	–	5	1	5	–	5	1	Western blot - wt vs TF mutant
664	Ye 2017 Ye C, Sutter BM, Wang Y, Kuang Z, Tu BP. A Metabolic Function for Phospholipid and Histone Methylation. <i>Mol Cell</i> 2017;66(2):180–193.e8.	doi:10.1016/j.molcel.2017.02.026 PMID 28366644 PMC5482412	–	5	1	5	–	5	1	RT-PCR - wt vs TF mutant
665	Chang 2017 Chang YL, Tseng SF, Huang YC, et al., Teng SC. Yeast Cip1 is activated by environmental stress to inhibit Cdk1-G1 cyclins via Mcm1 and Msn2/4. <i>Nat Commun</i> 2017;8(1):56.	doi:10.1038/s41467-017-00080-y PMID 28676626 PMC5496861	–	5	2	2	5	–	5	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
666	Wei 2018 Wei S, Liu Y, Wu M, et al., Bao X. Disruption of the transcription factors Thi2p and Nrm1p alleviates the post-glucose effect on xylose utilization in <i>Saccharomyces cerevisiae</i> . <i>Biotechnol Biofuels</i> 2018;11:112.	doi:10.1186/s13068-018-1112-1 PMID 29686730 PMC5901872	–	5	1	5	–	5	1	RT-PCR - wt vs TF mutant
667	Miller 2018 Miller D, Brandt N, Gresham D. Systematic identification of factors mediating accelerated mRNA degradation in response to changes in environmental nitrogen. <i>PLoS Genet</i> 2018;14(5): e1007406.	doi:10.1371/journal.pgen.1007406 PMID 29782489 PMC5983874	–	5	3	3	–	5	2	Barseq after FACS after mRNA FISH (BFF)
668	Hijazi 2020 Hijazi I, Knupp J, Chang A. Retrograde signaling mediates an adaptive survival response to endoplasmic reticulum stress in <i>Saccharomyces cerevisiae</i> . <i>J Cell Sci</i> 2020;133(6).	doi:10.1242/jcs.241539 PMID 32005698 PMC7132770	–	5	1	2	–	5	4	lacZ - wt vs TF mutant
669	Ros-Carrero 2020 Ros-Carrero C, Ramos-Alonso L, Romero AM, Bañó MC, Martínez-Pastor MT, Puig S. The yeast Aft1 transcription factor activates ribonucleotide reductase catalytic subunit RNR1 in response to iron deficiency. <i>Biochim Biophys Acta Gene Regul Mech</i> 2020;1863(7):194522.	doi:10.1016/j.bbagr.2020.194522 PMID 32147528	–	5	1	2	–	5	1	RT-PCR - wt vs TF mutant
670	Laidlaw 2021 Laidlaw KME, Bisinski DD, Shashkova S, et al., MacDonald C. A glucose-starvation response governs endocytic trafficking and eisosomal retention of surface cargoes in budding yeast. <i>J Cell Sci</i> 2021;134(2).	doi:10.1242/jcs.257733 PMID 33443082 PMC7860119	–	5	3	5	–	5	1	RT-PCR - wt vs TF mutant
671	Fang 2022 Fang W, Zhu Y, Yang S, Tong X, Ye C. Reciprocal regulation of phosphatidylcholine synthesis and H3K36 methylation programs metabolic adaptation. <i>Cell Rep</i> 2022;39(2):110672.	doi:10.1016/j.celrep.2022.110672 PMID 35417718	–	5	1	5	–	5	1	RT-PCR - wt vs TF mutant
672	Keng 1992 Keng T. HAP1 and ROX1 form a regulatory pathway in the repression of HEM13 transcription in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1992;12(6):2616–23.	doi:10.1128/mcb.12.6.2616-2623.1992 PMID 1588959 PMC364455	–	4	2	4	–	4	3	Northern blot - wt vs TF mutant
673	Huie 1992 Huie MA, Scott EW, Drazinic CM, et al., Baker HV. Characterization of the DNA-binding activity of GCR1: in vivo evidence for two GCR1-binding sites in the upstream activating sequence of TPI of <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1992;12(6):2690–700.	doi:10.1128/mcb.12.6.2690-2700.1992 PMID 1588965 PMC364463	–	4	1	4	4	–	2	EMSA
674	Ogas 1991 Ogas J, Andrews BJ, Herskowitz I. Transcriptional activation of CLN1, CLN2, and a putative new G1 cyclin (HCS26) by SWI4, a positive regulator of G1-specific transcription. <i>Cell</i> 1991;66(5):1015–26.	doi:10.1016/0092-8674(91)90445-5 PMID 1832336	–	4	1	4	–	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
675	Verma 1991 Verma R, Patapoutian A, Gordon CB, Campbell JL. Identification and purification of a factor that binds to the Mlu I cell cycle box of yeast DNA replication genes. <i>Proc Natl Acad Sci U S A</i> 1991;88(16):7155–9.	doi:10.1073/pnas.88.16.7155 PMID 1871128 PMC52252	–	4	2	4	1	–	1	–
676	Coschigano 1991 Coschigano PW, Magasanik B. The URE2 gene product of <i>Saccharomyces cerevisiae</i> plays an important role in the cellular response to the nitrogen source and has homology to glutathione s-transferases. <i>Mol Cell Biol</i> 1991;11(2):822–32.	doi:10.1128/mcb.11.2.822-832.1991 PMID 1990286 PMC359734	–	4	2	4	–	4	2	Enzyme activity - wt vs TF mutant
677	Repetto 1990 Repetto B, Tzagoloff A. Structure and regulation of KGD2, the structural gene for yeast dihydrolipoyl transsuccinylase. <i>Mol Cell Biol</i> 1990;10(8):4221–32.	doi:10.1128/mcb.10.8.4221-4232.1990 PMID 2115121 PMC360958	–	4	4	4	2	2	1	lacZ - wt vs TF mutant
678	Wang 1990 Wang H, Nicholson PR, Stillman DJ. Identification of a <i>Saccharomyces cerevisiae</i> DNA-binding protein involved in transcriptional regulation. <i>Mol Cell Biol</i> 1990;10(4):1743–53.	doi:10.1128/mcb.10.4.1743-1753.1990 PMID 2181283 PMC362280	–	4	1	4	1	–	1	–
679	Lowry 1990 Lowry CV, Cerdán ME, Zitomer RS. A hypoxic consensus operator and a constitutive activation region regulate the ANB1 gene of <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1990;10(11):5921–6.	doi:10.1128/mcb.10.11.5921-5926.1990 PMID 2233724 PMC361388	–	4	1	4	3	1	1	lacZ - wt vs mutant promoter sequence
680	Repetto 1989 Repetto B, Tzagoloff A. Structure and regulation of KGD1, the structural gene for yeast alpha-ketoglutarate dehydrogenase. <i>Mol Cell Biol</i> 1989;9(6):2695–705.	doi:10.1128/mcb.9.6.2695-2705.1989 PMID 2503710 PMC362342	–	4	4	4	–	4	1	lacZ - wt vs TF mutant
681	Trawick 1989 Trawick JD, Wright RM, Poyton RO. Transcription of yeast COX6, the gene for cytochrome c oxidase subunit VI, is dependent on heme and on the HAP2 gene. <i>J Biol Chem</i> 1989;264(12):7005–8.	PMID 2540169	–	4	4	4	–	–	1	–
682	Forsburg 1989 Forsburg SL, Guarente L. Communication between mitochondria and the nucleus in regulation of cytochrome genes in the yeast <i>Saccharomyces cerevisiae</i> . <i>Annu Rev Cell Biol</i> 1989;5:153–80.	doi:10.1146/annurev.cb.05.110189.001101 PMID 2557058	–	4	4	4	–	–	1	–
683	Passmore 1989 Passmore S, Elble R, Tye BK. A protein involved in minichromosome maintenance in yeast binds a transcriptional enhancer conserved in eukaryotes. <i>Genes Dev</i> 1989;3(7):921–35.	doi:10.1101/gad.3.7.921 PMID 2673922	–	4	1	4	4	–	2	EMSA
684	Olesen 1987 Olesen J, Hahn S, Guarente L. Yeast HAP2 and HAP3 activators both bind to the CYC1 upstream activation site, UAS2, in an interdependent manner. <i>Cell</i> 1987;51(6):953–61.	doi:10.1016/0092-8674(87)90582-4 PMID 2826015	–	4	4	4	4	–	1	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
685	Friden 1988 Friden P, Schimmel P. LEU3 of <i>Saccharomyces cerevisiae</i> activates multiple genes for branched-chain amino acid biosynthesis by binding to a common decanucleotide core sequence. <i>Mol Cell Biol</i> 1988;8(7):2690–7.	doi:10.1128/mcb.8.7.2690-2697.1988 PMID 3043190 PMC363479	–	4	1	4	–	4	1	lacZ - wt vs TF mutant
686	Shore 1987 Shore D, Nasmyth K. Purification and cloning of a DNA binding protein from yeast that binds to both silencer and activator elements. <i>Cell</i> 1987;51(5):721–32.	doi:10.1016/0092-8674(87)90095-x PMID 3315231	–	4	1	4	–	–	1	–
687	Keng 1987 Keng T, Guarente L. Constitutive expression of the yeast HEM1 gene is actually a composite of activation and repression. <i>Proc Natl Acad Sci U S A</i> 1987;84(24):9113–7.	doi:10.1073/pnas.84.24.9113 PMID 3321068 PMC299702	–	4	4	4	4	–	1	lacZ - wt vs mutant promoter sequence
688	Arndt 1986 Arndt K, Fink GR. GCN4 protein, a positive transcription factor in yeast, binds general control promoters at all 5' TGACTC 3' sequences. <i>Proc Natl Acad Sci U S A</i> 1986;83(22):8516–20.	doi:10.1073/pnas.83.22.8516 PMID 3464968 PMC386961	–	4	1	4	4	–	1	DNA footprinting
689	Simon 1995 Simon MM, Pavlik P, Hartig A, et al., Schanz B. A C-terminal region of the <i>Saccharomyces cerevisiae</i> transcription factor ADR1 plays an important role in the regulation of peroxisome proliferation by fatty acids. <i>Mol Gen Genet</i> 1995;249(3):289–96.	doi:10.1007/BF00290529 PMID 7500953	–	4	1	4	–	4	2	Northern blot - wt vs TF overexpression
690	Kuras 1995 Kuras L, Thomas D. Identification of the yeast methionine biosynthetic genes that require the centromere binding factor 1 for their transcriptional activation. <i>FEBS Lett</i> 1995;367(1):15–8.	doi:10.1016/0014-5793(95)00528-h PMID 7601277	–	4	1	4	–	4	1	Northern blot - wt vs TF mutant
691	Rosenkrantz 1994 Rosenkrantz M, Kell CS, Pennell EA, Devenish LJ. The HAP2,3,4 transcriptional activator is required for derepression of the yeast citrate synthase gene, CIT1. <i>Mol Microbiol</i> 1994;13(1):119–31.	doi:10.1111/j.1365-2958.1994.tb00407.x PMID 7984086	–	4	4	4	4	–	1	lacZ - wt vs mutant promoter sequence
692	Liao 1993 Liao X, Butow RA. RTG1 and RTG2: two yeast genes required for a novel path of communication from mitochondria to the nucleus. <i>Cell</i> 1993;72(1):61–71.	doi:10.1016/0092-8674(93)90050-z PMID 8422683	–	4	2	4	1	–	1	–
693	Messenguy 1993 Messenguy F, Dubois E. Genetic evidence for a role for MCM1 in the regulation of arginine metabolism in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1993;13(4):2586–92.	doi:10.1128/mcb.13.4.2586-2592.1993 PMID 8455631 PMC359592	–	4	2	4	–	4	1	Enzyme activity - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
694	Katzmann 1995 Katzmann DJ, Hallstrom TC, Voet M, et al., Moye-Rowley WS. Expression of an ATP-binding cassette transporter-encoding gene (YOR1) is required for oligomycin resistance in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1995; 15(12):6875–83.	doi:10.1128/MCB.15.12.6875 PMID 8524254 PMC230942	–	4	2	4	–	4	2	lacZ - wt vs TF mutant
695	Dang 1996 Dang VD, Bohn C, Bolotin-Fukuhara M, Daignan-Fornier B. The CCAAT box-binding factor stimulates ammonium assimilation in <i>Saccharomyces cerevisiae</i> , defining a new cross-pathway regulation between nitrogen and carbon metabolisms. <i>J Bacteriol</i> 1996;178(7):1842–9.	doi:10.1128/jb.178.7.1842-1849.1996 PMID 8606156 PMC177877	–	4	4	4	2	2	2	lacZ - wt vs TF mutant
696	Dohrmann 1996 Dohrmann PR, Voth WP, Stillman DJ. Role of negative regulation in promoter specificity of the homologous transcriptional activators Ace2p and Swi5p. <i>Mol Cell Biol</i> 1996;16(4):1746–58.	doi:10.1128/MCB.16.4.1746 PMID 8657150 PMC231161	–	4	2	4	2	2	2	DNA footprinting
697	Knapp 1996 Knapp D, Bhoite L, Stillman DJ, Nasmyth K. The transcription factor Swi5 regulates expression of the cyclin kinase inhibitor p40SIC1. <i>Mol Cell Biol</i> 1996;16(10):5701–7.	doi:10.1128/MCB.16.10.5701 PMID 8816483 PMC231570	–	4	2	4	–	2	3	–
698	Jansma 1996 Jansma DB, Archambault J, Mostachfi O, Friesen JD. Similar upstream regulatory elements of genes that encode the two largest subunits of RNA polymerase II in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Res</i> 1996;24(22):4543–51.	doi:10.1093/nar/24.22.4543 PMID 8948647 PMC146278	–	4	2	4	–	4	1	lacZ - wt vs TF mutant
699	Wang 1997 Wang J, Sirenko O, Needleman R. Genomic footprinting of Mig1p in the MAL62 promoter. Binding is dependent upon carbon source and competitive with the Mal63p activator. <i>J Biol Chem</i> 1997;272(7):4613–22.	doi:10.1074/jbc.272.7.4613 PMID 9020190	–	4	2	3	2	1	3	DNA footprinting
700	Flattery-O'Brien 1997 Flattery-O'Brien JA, Grant CM, Dawes IW. Stationary-phase regulation of the <i>Saccharomyces cerevisiae</i> SOD2 gene is dependent on additive effects of HAP2/3/4/5- and STRE-binding elements. <i>Mol Microbiol</i> 1997;23(2):303–12.	doi:10.1046/j.1365-2958.1997.2121581.x PMID 9044264	–	4	4	4	4	–	1	lacZ - wt vs mutant promoter sequence
701	Morgan 1997 Morgan BA, Banks GR, Toone WM, Raitt D, Kuge S, Johnston LH. The Skn7 response regulator controls gene expression in the oxidative stress response of the budding yeast <i>Saccharomyces cerevisiae</i> . <i>EMBO J</i> 1997;16(5):1035–44.	doi:10.1093/emboj/16.5.1035 PMID 9118942 PMC1169703	–	4	2	4	–	4	1	Northern blot - wt vs TF mutant
702	Bon 1997 Bon EP, Carvajal E, Stanbrough M, Rowen D, Magasanik B. Asparaginase II of <i>Saccharomyces cerevisiae</i> . GLN3/URE2 regulation of a periplasmic enzyme. <i>Appl Biochem Biotechnol</i> 1997;63–65:203–12.	doi:10.1007/978-1-4612-2312-2_19 PMID 9170245	–	4	1	4	–	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
703	Parrou 1997 Parrou JL, Teste MA, François J. Effects of various types of stress on the metabolism of reserve carbohydrates in <i>Saccharomyces cerevisiae</i> : genetic evidence for a stress-induced recycling of glycogen and trehalose. <i>Microbiology (Reading)</i> 1997;143 (Pt 6):1891–1900.	doi:10.1099/00221287-143-6-1891 PMID 9202465	–	4	2	4	–	4	1	lacZ - wt vs TF mutant
704	Nourani 1997 Nourani A, Wesolowski-Louvel M, Delaveau T, Jacq C, Delahodde A. Multiple-drug-resistance phenomenon in the yeast <i>Saccharomyces cerevisiae</i> : involvement of two hexose transporters. <i>Mol Cell Biol</i> 1997;17(9):5453–60.	doi:10.1128/MCB.17.9.5453 PMID 9271421 PMC232394	–	4	2	4	–	–	1	–
705	Kratzer 1997 Kratzer S, Schüller HJ. Transcriptional control of the yeast acetyl-CoA synthetase gene, ACS1, by the positive regulators CAT8 and ADR1 and the pleiotropic repressor UME6. <i>Mol Microbiol</i> 1997;26(4):631–41.	doi:10.1046/j.1365-2958.1997.5611937.x PMID 9427394	–	4	4	4	2	2	2	lacZ - wt vs TF mutant
706	Wolfger 1997 Wolfger H, Mahé Y, Parle-McDermott A, Delahodde A, Kuchler K. The yeast ATP binding cassette (ABC) protein genes PDR10 and PDR15 are novel targets for the Pdr1 and Pdr3 transcriptional regulators. <i>FEBS Lett</i> 1997;418(3):269–74.	doi:10.1016/s0014-5793(97)01382-3 PMID 9428726	–	4	2	4	–	–	1	–
707	Mori 1998 Mori K, Ogawa N, Kawahara T, Yanagi H, Yura T. Palindrome with spacer of one nucleotide is characteristic of the cis-acting unfolded protein response element in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 1998;273(16):9912–20.	doi:10.1074/jbc.273.16.9912 PMID 9545334	–	4	1	4	3	–	1	EMSA
708	Cui 1998 Cui Z, Shiraki T, Hirata D, Miyakawa T. Yeast gene YRR1, which is required for resistance to 4-nitroquinoline N-oxide, mediates transcriptional activation of the multidrug resistance transporter gene SNQ2. <i>Mol Microbiol</i> 1998;29(5):1307–15.	doi:10.1046/j.1365-2958.1998.01027.x PMID 9767597	–	4	3	4	–	4	1	Northern blot - wt vs TF mutant
709	Mannhaupt 1999 Mannhaupt G, Schnall R, Karpov V, Vetter I, Feldmann H. Rpn4p acts as a transcription factor by binding to PACE, a nonamer box found upstream of 26S proteasomal and other genes in yeast. <i>FEBS Lett</i> 1999;450(1–2):27–34.	doi:10.1016/s0014-5793(99)00467-6 PMID 10350051	–	4	1	4	2	2	2	CAT - wt vs TF mutant
710	Navarro-Aviño 1999 Navarro-Aviño JP, Prasad R, Miralles VJ, Benito RM, Serrano R. A proposal for nomenclature of aldehyde dehydrogenases in <i>Saccharomyces cerevisiae</i> and characterization of the stress-inducible ALD2 and ALD3 genes. <i>Yeast</i> 1999;15(10A):829–42.	<a href="https://doi.org/10.1002/(SICI)1097-0061(199907)15:10A<829::AID-YEA423>3.0.CO;2-9">doi:10.1002/(SICI)1097-0061(199907)15:10A<829::AID-YEA423>3.0.CO;2-9 PMID 10407263	–	4	2	4	–	4	1	lacZ - wt vs TF mutant
711	Garay-Arroyo 1999 Garay-Arroyo A, Covarrubias AA. Three genes whose expression is induced by stress in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 1999;15(10A):879–92.	<a href="https://doi.org/10.1002/(SICI)1097-0061(199907)15:10A<879::AID-YEA428>3.0.CO;2-Q">doi:10.1002/(SICI)1097-0061(199907)15:10A<879::AID-YEA428>3.0.CO;2-Q PMID 10407268	–	4	2	4	–	4	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
712	Liu 1999 Liu XD, Morano KA, Thiele DJ. The yeast Hsp110 family member, Sse1, is an Hsp90 cochaperone. <i>J Biol Chem</i> 1999; 274(38):26654–60.	doi:10.1074/jbc.274.38.26654 PMID 10480867	–	4	1	4	–	4	2	Northern blot - wt vs TF mutant
713	Garcia 2000 Garcia SC, Moretti MB, Batlle A. Constitutive expression of the UGA4 gene in <i>Saccharomyces cerevisiae</i> depends on two positive-acting proteins, Uga3p and Uga35p. <i>FEMS Microbiol Lett</i> 2000;184(2):219–24.	doi:10.1111/j.1574-6968.2000.tb09017.x PMID 10713424	–	4	4	4	–	4	1	lacZ - wt vs TF mutant
714	Yun 2000 Yun CW, Ferea T, Rashford J, et al., Philpott CC. Desferrioxamine-mediated iron uptake in <i>Saccharomyces cerevisiae</i> . Evidence for two pathways of iron uptake. <i>J Biol Chem</i> 2000;275(14):10709–15.	doi:10.1074/jbc.275.14.10709 PMID 10744769	no GEO	4	1	4	–	4	1	Northern blot - wt vs TF overexpression
715	Ross 2000 Ross SJ, Findlay VJ, Malakasi P, Morgan BA. Thioredoxin peroxidase is required for the transcriptional response to oxidative stress in budding yeast. <i>Mol Biol Cell</i> 2000;11(8):2631–42.	doi:10.1091/mbc.11.8.2631 PMID 10930459 PMC14945	–	4	2	4	–	4	1	Northern blot - wt vs TF mutant
716	Terashima 2000 Terashima H, Yabuki N, Arisawa M, Hamada K, Kitada K. Up-regulation of genes encoding glycosylphosphatidylinositol (GPI)-attached proteins in response to cell wall damage caused by disruption of FKS1 in <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 2000;264(1–2):64–74.	doi:10.1007/s004380000285 PMID 11016834	–	4	1	3	3	1	1	lacZ - wt vs mutant promoter sequence
717	Pan 2000 Pan X, Heitman J. Sok2 regulates yeast pseudohyphal differentiation via a transcription factor cascade that regulates cell-cell adhesion. <i>Mol Cell Biol</i> 2000;20(22):8364–72.	doi:10.1128/MCB.20.22.8364-8372.2000 PMID 11046133 PMC102143	–	4	3	4	–	4	1	Northern blot - wt vs TF mutant
718	Bourdineaud 2000 Bourdineaud JP, De Sampaio G, Lauquin GJ. A Rox1-independent hypoxic pathway in yeast. Antagonistic action of the repressor Ord1 and activator Yap1 for hypoxic expression of the SRP1/TIR1 gene. <i>Mol Microbiol</i> 2000;38(4):879–90.	doi:10.1046/j.1365-2958.2000.02188.x PMID 11115121	–	4	2	4	2	2	2	EMSA
719	Grauslund 2000 Grauslund M, Rønnow B. Carbon source-dependent transcriptional regulation of the mitochondrial glycerol-3-phosphate dehydrogenase gene, GUT2, from <i>Saccharomyces cerevisiae</i> . <i>Can J Microbiol</i> 2000;46(12):1096–100.	doi:10.1139/w00-105 PMID 11142398	–	4	4	4	–	4	1	Northern blot - wt vs TF mutant
720	Abramova 2001 Abramova N, Sertil O, Mehta S, Lowry CV. Reciprocal regulation of anaerobic and aerobic cell wall mannoprotein gene expression in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 2001; 183(9):2881–7.	doi:10.1128/JB.183.9.2881-2887.2001 PMID 11292809 PMC99506	–	4	2	4	–	4	2	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
721	Hallstrom 2001 Hallstrom TC, Lambert L, Schorling S, Balzi E, Goffeau A, Moye-Rowley WS. Coordinate control of sphingolipid biosynthesis and multidrug resistance in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2001;276(26):23674–80.	doi:10.1074/jbc.M101568200 PMID 11323424	–	4	2	4	2	2	2	DNA footprinting
722	Alepuz 2001 Alepuz PM, Jovanovic A, Reiser V, Ammerer G. Stress-induced map kinase Hog1 is part of transcription activation complexes. <i>Mol Cell</i> 2001;7(4):767–77.	doi:10.1016/s1097-2765(01)00221-0 PMID 11336700	–	4	1	4	4	–	2	ChIP
723	Bernard 2001 Bernard F, André B. Genetic analysis of the signalling pathway activated by external amino acids in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2001;41(2):489–502.	doi:10.1046/j.1365-2958.2001.02538.x PMID 11489133	–	4	2	4	–	4	1	lacZ - wt vs TF mutant
724	Wenz 2001 Wenz P, Schwank S, Hoja U, Schüller HJ. A downstream regulatory element located within the coding sequence mediates autoregulated expression of the yeast fatty acid synthase gene FAS2 by the FAS1 gene product. <i>Nucleic Acids Res</i> 2001;29(22):4625–32.	doi:10.1093/nar/29.22.4625 PMID 11713312 PMC92567	–	4	2	4	–	–	1	–
725	Frenz 2001 Frenz LM, Johnson AL, Johnston LH. Rme1, which controls CLN2 expression in <i>Saccharomyces cerevisiae</i> , is a nuclear protein that is cell cycle regulated. <i>Mol Genet Genomics</i> 2001;266(3):374–84.	doi:10.1007/s004380100515 PMID 11713667	–	4	2	4	–	4	3	Northern blot - wt vs TF mutant
726	Jiao 2002 Jiao K, Nau JJ, Cool M, Gray WM, Fassler JS, Malone RE. Phylogenetic footprinting reveals multiple regulatory elements involved in control of the meiotic recombination gene, REC102. <i>Yeast</i> 2002;19(2):99–114.	doi:10.1002/yea.800 PMID 11788965	–	4	4	4	–	4	1	lacZ - wt vs TF mutant
727	Hong 2002 Hong SK, Cha MK, Choi YS, Kim WC, Kim IH. Msn2p/Msn4p act as a key transcriptional activator of yeast cytoplasmic thiol peroxidase II. <i>J Biol Chem</i> 2002;277(14):12109–17.	doi:10.1074/jbc.M111341200 PMID 11821410	–	4	4	4	–	4	3	RT-PCR - wt vs TF mutant
728	Teixeira 2002 Teixeira MC, Sá-Correia I. <i>Saccharomyces cerevisiae</i> resistance to chlorinated phenoxyacetic acid herbicides involves Pdr1p-mediated transcriptional activation of TPO1 and PDR5 genes. <i>Biochem Biophys Res Commun</i> 2002;292(2):530–7.	doi:10.1006/bbrc.2002.6691 PMID 11906193	–	4	2	4	–	4	1	Northern blot - wt vs TF mutant
729	Walsh 2002 Walsh L, Schmuckli-Maurer J, Billinton N, Barker MG, Heyer WD, Walmsley RM. DNA-damage induction of RAD54 can be regulated independently of the RAD9- and DDC1-dependent checkpoints that regulate RNR2. <i>Curr Genet</i> 2002;41(4):232–40.	doi:10.1007/s00294-002-0302-2 PMID 12172964	–	4	2	3	–	4	4	GFP - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
730	Pak 2002 Pak J, Segall J. Regulation of the premiddle and middle phases of expression of the NDT80 gene during sporulation of <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2002;22(18):6417–29.	doi:10.1128/MCB.22.18.6417-6429.2002 PMID 12192041 PMC135636	–	4	2	4	–	4	2	lacZ - wt vs TF mutant
731	Unnikrishnan 2003 Unnikrishnan I, Miller S, Meinke M, LaPorte DC. Multiple positive and negative elements involved in the regulation of expression of GSY1 in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2003;278(29):26450–7.	doi:10.1074/jbc.M211808200 PMID 12697770	–	4	4	4	2	2	1	lacZ - wt vs TF mutant
732	Sertil 2003 Sertil O, Kapoor R, Cohen BD, Abramova N, Lowry CV. Synergistic repression of anaerobic genes by Mot3 and Rox1 in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Res</i> 2003;31(20):5831–7.	doi:10.1093/nar/gkg792 PMID 14530431 PMC219473	–	4	3	4	–	4	2	Northern blot - wt vs TF mutant
733	Schüller 2004 Schüller C, Mamnun YM, Mollapour M, et al., Kuchler K. Global phenotypic analysis and transcriptional profiling defines the weak acid stress response regulon in <i>Saccharomyces cerevisiae</i> . <i>Mol Biol Cell</i> 2004;15(2):706–20.	doi:10.1091/mbc.e03-05-0322 PMID 14617816 PMC338074	–	4	2	4	–	4	1	Microarray analysis - wt vs TF mutant
734	Reinoso-Martín 2003 Reinoso-Martín C, Schüller C, Schuetzer-Muehlbauer M, Kuchler K. The yeast protein kinase C cell integrity pathway mediates tolerance to the antifungal drug caspofungin through activation of Slt2p mitogen-activated protein kinase signaling. <i>Eukaryot Cell</i> 2003;2(6):1200–10.	doi:10.1128/EC.2.6.1200-1210.2003 PMID 14665455 PMC326656	–	4	2	4	–	4	1	Northern blot - wt vs TF mutant
735	Mennella 2003 Mennella TA, Klinkenberg LG, Zitomer RS. Recruitment of Tup1-Ssn6 by yeast hypoxic genes and chromatin-independent exclusion of TATA binding protein. <i>Eukaryot Cell</i> 2003;2(6):1288–303.	doi:10.1128/EC.2.6.1288-1303.2003 PMID 14665463 PMC326644	–	4	2	4	2	2	1	ChIP
736	Wehrschütz-Sigl 2004 Wehrschütz-Sigl E, Jungwirth H, Bergler H, Högenauer G. The transporters Pdr5p and Snq2p mediate diazaborine resistance and are under the control of the gain-of-function allele PDR1-12. <i>Eur J Biochem</i> 2004;271(6):1145–52.	doi:10.1111/j.1432-1033.2004.04018.x PMID 15009193	–	4	2	4	–	–	1	–
737	Abdel-Sater 2004 Abdel-Sater F, Iraqui I, Urrestarazu A, André B. The external amino acid signaling pathway promotes activation of Stp1 and Uga35/Dal81 transcription factors for induction of the AGP1 gene in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2004;166(4):1727–39.	doi:10.1534/genetics.166.4.1727 PMID 15126393 PMC1470804	–	4	4	4	–	4	3	Western blot - wt vs TF mutant
738	Almaguer 2004 Almaguer C, Cheng W, Nolder C, Patton-Vogt J. Glycerophosphoinositol, a novel phosphate source whose transport is regulated by multiple factors in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2004;279(30):31937–42.	doi:10.1074/jbc.M403648200 PMID 15145930	–	4	4	4	–	4	3	CAT - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
739	Enjalbert 2004 Enjalbert B, Parrou JL, Teste MA, François J. Combinatorial control by the protein kinases PKA, PHO85 and SNF1 of transcriptional induction of the <i>Saccharomyces cerevisiae</i> GSY2 gene at the diauxic shift. <i>Mol Genet Genomics</i> 2004; 271(6):697–708.	doi:10.1007/s00438-004-1014-8 PMID 15221454	–	4	4	4	–	4	3	lacZ - wt vs TF mutant
740	Yoon 2004 Yoon S, Govind CK, Qiu H, Kim SJ, Dong J, Hinnebusch AG. Recruitment of the ArgR/Mcm1p repressor is stimulated by the activator Gcn4p: a self-checking activation mechanism. <i>Proc Natl Acad Sci U S A</i> 2004;101(32):11713–8.	doi:10.1073/pnas.0404652101 PMID 15289616 PMC511042	–	4	4	4	3	1	1	ChIP
741	Bernstein 2004 Bernstein BE, Liu CL, Humphrey EL, Perlstein EO, Schreiber SL. Global nucleosome occupancy in yeast. <i>Genome Biol</i> 2004;5(9):R62.	doi:10.1186/gb-2004-5-9-r62 PMID 15345046 PMC522869	–	4	1	4	–	–	1	–
742	Nevitt 2004 Nevitt T, Pereira J, Rodrigues-Pousada C. YAP4 gene expression is induced in response to several forms of stress in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 2004;21(16):1365–74.	doi:10.1002/yea.1188 PMID 15565582	–	4	3	4	2	2	2	Northern blot - wt vs TF mutant
743	Klinkenberg 2005 Klinkenberg LG, Mennella TA, Luetkenhaus K, Zitomer RS. Combinatorial repression of the hypoxic genes of <i>Saccharomyces cerevisiae</i> by DNA binding proteins Rox1 and Mot3. <i>Eukaryot Cell</i> 2005;4(4):649–60.	doi:10.1128/EC.4.4.649-660.2005 PMID 15821125 PMC1087805	–	4	2	4	3	1	2	EMSA
744	Bourges 2005 Bourges I, Horan S, Meunier B. Effect of inhibition of the bcl complex on gene expression profile in yeast. <i>J Biol Chem</i> 2005;280(33):29743–9.	doi:10.1074/jbc.M505915200 PMID 15967791	–	4	3	4	–	4	1	Microarray analysis - wt vs TF overexpression
745	Mai 2006 Mai B, Breeden LL. Identification of target genes of a yeast transcriptional repressor. <i>Methods Mol Biol</i> 2006;317:267–77.	doi:10.1385/1-59259-968-0:267 PMID 16264235	–	4	1	4	–	3	1	Northern blot - wt vs TF mutant
746	Carrozza 2005 Carrozza MJ, Florens L, Swanson SK, et al., Workman JL. Stable incorporation of sequence specific repressors Ash1 and Ume6 into the Rpd3L complex. <i>Biochim Biophys Acta</i> 2005; 1731(2):77–87;discussion 75–6.	doi:10.1016/j.bbaexp.2005.09.005 PMID 16314178	–	4	2	4	2	2	1	ChIP
747	Kim 2005 Kim SJ, Swanson MJ, Qiu H, Govind CK, Hinnebusch AG. Activator Gcn4p and Cyc8p/Tup1p are interdependent for promoter occupancy at ARG1 in vivo. <i>Mol Cell Biol</i> 2005; 25(24):11171–83.	doi:10.1128/MCB.25.24.11171-11183.2005 PMID 16314536 PMC1316967	–	4	1	4	2	2	1	ChIP
748	Bester 2006 Bester MC, Pretorius IS, Bauer FF. The regulation of <i>Saccharomyces cerevisiae</i> FLO gene expression and Ca ²⁺ -dependent flocculation by Flo8p and Mss11p. <i>Curr Genet</i> 2006;49(6):375–83.	doi:10.1007/s00294-006-0068-z PMID 16568252	–	4	2	4	–	4	1	Northern blot - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
749	Kupchak 2007 Kupchak BR, Garitaonandia I, Villa NY, et al., Lyons TJ. Probing the mechanism of FET3 repression by Izh2p overexpression. <i>Biochim Biophys Acta</i> 2007;1773(7):1124–32.	doi:10.1016/j.bbamcr.2007.04.003 PMID 17553578 PMC1994572	–	4	4	4	–	4	2	Enzyme activity - wt vs TF mutant
750	Jin 2008 Jin R, Dobry CJ, McCown PJ, Kumar A. Large-scale analysis of yeast filamentous growth by systematic gene disruption and overexpression. <i>Mol Biol Cell</i> 2008;19(1):284–96.	doi:10.1091/mbc.e07-05-0519 PMID 17989363 PMC2174193	–	4	4	4	–	4	1	GFP - wt vs TF mutant
751	Erkina 2008 Erkina TY, Tschetter PA, Erkin AM. Different requirements of the SWI/SNF complex for robust nucleosome displacement at promoters of heat shock factor and Msn2- and Msn4-regulated heat shock genes. <i>Mol Cell Biol</i> 2008;28(4):1207–17.	doi:10.1128/MCB.01069-07 PMID 18070923 PMC2258741	–	4	2	4	4	–	2	ChIP
752	Coleman 2008 Coleman CB, Allen PL, Valles JM, Hammond TG. Transcriptional regulation of changes in growth, cell cycle, and gene expression of <i>Saccharomyces cerevisiae</i> due to changes in buoyancy. <i>Biotechnol Bioeng</i> 2008;100(2):334–43.	doi:10.1002/bit.21748 PMID 18078295	–	4	2	4	–	4	2	GFP - wt vs TF mutant
753	Soto 2008 Soto A, Carman GM. Regulation of the <i>Saccharomyces cerevisiae</i> CKI1-encoded choline kinase by zinc depletion. <i>J Biol Chem</i> 2008;283(15):10079–88.	doi:10.1074/jbc.M800502200 PMID 18276583 PMC2367329	–	4	4	4	–	4	3	lacZ - wt vs TF mutant
754	Tomar 2008 Tomar RS, Zheng S, Brunke-Reese D, Wolcott HN, Reese JC. Yeast Rap1 contributes to genomic integrity by activating DNA damage repair genes. <i>EMBO J</i> 2008;27(11):1575–84.	doi:10.1038/emboj.2008.93 PMID 18480842 PMC2426730	–	4	1	4	–	4	1	Northern blot - wt vs TF mutant
755	Gong 2009 Gong J, Siede W. SBF transcription factor complex positively regulates UV mutagenesis in <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2009;379(4):1009–14.	doi:10.1016/j.bbrc.2009.01.012 PMID 19150335 PMC2674285	–	4	2	2	–	4	1	Northern blot - wt vs TF mutant
756	Zhu 2009 Zhu C, Byers KJ, McCord RP, et al., Bulyk ML. High-resolution DNA-binding specificity analysis of yeast transcription factors. <i>Genome Res</i> 2009;19(4):556–66.	doi:10.1101/gr.090233.108 PMID 19158363 PMC2665775	GSE13684	4	1	4	4	–	2	ChIP
757	Yoon 2009 Yoon S, Hinnebusch AG. Mcm1p binding sites in ARG1 positively regulate Gcn4p binding and SWI/SNF recruitment. <i>Biochem Biophys Res Commun</i> 2009;381(1):123–8.	doi:10.1016/j.bbrc.2009.02.045 PMID 19233144 PMC2683423	–	4	2	4	4	–	1	ChIP
758	García 2009 García R, Rodríguez-Peña JM, Bermejo C, Nombela C, Arroyo J. The high osmotic response and cell wall integrity pathways cooperate to regulate transcriptional responses to zymolyase-induced cell wall stress in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2009;284(16):10901–11.	doi:10.1074/jbc.M808693200 PMID 19234305 PMC2667776	GSE12684	4	1	4	–	4	1	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
759	Yazawa 2009 Yazawa H, Iwahashi H, Kamisaka Y, Kimura K, Uemura H. Production of polyunsaturated fatty acids in yeast <i>Saccharomyces cerevisiae</i> and its relation to alkaline pH tolerance. <i>Yeast</i> 2009;26(3):167–84.	doi:10.1002/yea.1659 PMID 19243079	–	4	1	4	–	4	1	RT-PCR - wt vs TF mutant
760	Joo 2009 Joo YJ, Kim JH, Baek JH, Seong KM, Lee JY, Kim J. Determination of the core promoter regions of the <i>Saccharomyces cerevisiae</i> RPS3 gene. <i>Biochim Biophys Acta</i> 2009;1789(11–12):741–50.	doi:10.1016/j.bbagr.2009.10.002 PMID 19853675	–	4	4	4	3	1	2	lacZ - wt vs mutant promoter sequence
761	Archer 2010 Archer CT, Kodadek T. The hydrophobic patch of ubiquitin is required to protect transactivator-promoter complexes from destabilization by the proteasomal ATPases. <i>Nucleic Acids Res</i> 2010;38(3):789–96.	doi:10.1093/nar/gkp1066 PMID 19939937 PMC2817475	–	4	1	4	3	1	2	ChIP
762	Takatsume 2010 Takatsume Y, Ohdate T, Maeta K, Nomura W, Izawa S, Inoue Y. Calcineurin/Crz1 destabilizes Msn2 and Msn4 in the nucleus in response to Ca(2+) in <i>Saccharomyces cerevisiae</i> . <i>Biochem J</i> 2010;427(2):275–87.	doi:10.1042/BJ20091334 PMID 20121702	–	4	3	4	–	4	2	Enzyme activity - wt vs TF mutant
763	Wang 2010 Wang S, Zhao X, Suran R, Vogt VM, Lis JT, Shi H. Knocking down gene function with an RNA aptamer expressed as part of an intron. <i>Nucleic Acids Res</i> 2010;38(15):e154.	doi:10.1093/nar/gkq529 PMID 20542918 PMC2926621	–	4	1	4	–	4	1	RT-PCR - wt vs TF overexpression
764	Kuttykrishnan 2010 Kuttykrishnan S, Sabina J, Langton LL, Johnston M, Brent MR. A quantitative model of glucose signaling in yeast reveals an incoherent feed forward loop leading to a specific, transient pulse of transcription. <i>Proc Natl Acad Sci U S A</i> 2010;107(38):16743–8.	doi:10.1073/pnas.0912483107 PMID 20810924 PMC2944743	–	4	2	4	–	4	3	RT-PCR - wt vs TF mutant
765	Huang 2010 Huang YC, Chen HT, Teng SC. Intragenic transcription of a noncoding RNA modulates expression of ASP3 in budding yeast. <i>RNA</i> 2010;16(11):2085–93.	doi:10.1261/rna.2177410 PMID 20817754 PMC2957049	–	4	2	2	–	4	2	Northern blot - wt vs TF mutant
766	Dueñas-Sánchez 2010 Dueñas-Sánchez R, Codón AC, Rincón AM, Benítez T. Increased biomass production of industrial bakers' yeasts by overexpression of Hap4 gene. <i>Int J Food Microbiol</i> 2010;143(3):150–60.	doi:10.1016/j.ijfoodmicro.2010.08.009 PMID 20832886	–	4	1	4	4	–	1	RT-PCR - wt vs mutant promoter sequence
767	Teixeira 2010 Teixeira MC, Dias PJ, Monteiro PT, et al., Sá-Correia I. Refining current knowledge on the yeast FLR1 regulatory network by combined experimental and computational approaches. <i>Mol Biosyst</i> 2010;6(12):2471–81.	doi:10.1039/c004881j PMID 20938527	–	4	3	4	4	–	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
768	Miao 2011 Miao R, Holmes-Hampton GP, Lindahl PA. Biophysical investigation of the iron in Aft1-1(up) and Gal-YAH1 Saccharomyces cerevisiae. <i>Biochemistry</i> 2011;50(13):2660–71.	doi:10.1021/bi102015s PMID 21361388 PMC3068216	–	4	1	2	–	4	2	RT-PCR - wt vs TF mutant
769	Lin 2011 Lin AP, Anderson SL, Minard KI, McAlister-Henn L. Effects of excess succinate and retrograde control of metabolite accumulation in yeast tricarboxylic cycle mutants. <i>J Biol Chem</i> 2011;286(39):33737–46.	doi:10.1074/jbc.M111.266890 PMID 21841001 PMC3190802	–	4	1	4	–	4	1	Western blot - wt vs TF mutant
770	Orumets 2012 Orumets K, Kevvai K, Nisamedtinov I, Tamm T, Paalme T. YAP1 over-expression in Saccharomyces cerevisiae enhances glutathione accumulation at its biosynthesis and substrate availability levels. <i>Biotechnol J</i> 2012;7(4):566–8.	doi:10.1002/biot.201100363 PMID 22009669	–	4	1	4	–	4	1	RT-PCR - wt vs TF mutant
771	Kołaczkowska 2012 Kołaczkowska A, Manente M, Kołaczkowski M, Laba J, Ghislain M, Wawrzycka D. The regulatory inputs controlling pleiotropic drug resistance and hypoxic response in yeast converge at the promoter of the aminocholesterol resistance gene RTA1. <i>FEMS Yeast Res</i> 2012;12(3):279–92.	doi:10.1111/j.1567-1364.2011.00768.x PMID 22129104	–	4	4	4	–	4	1	lacZ - wt vs TF mutant
772	Galdieri 2012 Galdieri L, Desai P, Vancura A. Facilitated assembly of the preinitiation complex by separated tail and head/middle modules of the mediator. <i>J Mol Biol</i> 2012;415(3):464–74.	doi:10.1016/j.jmb.2011.11.020 PMID 22137896 PMC3259248	–	4	4	4	1	3	1	RT-PCR - wt vs TF mutant
773	Crisucci 2012 Crisucci EM, Arndt KM. Paf1 restricts Gcn4 occupancy and antisense transcription at the ARG1 promoter. <i>Mol Cell Biol</i> 2012;32(6):1150–63.	doi:10.1128/MCB.06262-11 PMID 22252319 PMC3295010	–	4	2	4	3	1	1	ChIP
774	Lin 2012 Lin L, Chamberlain L, Zhu LJ, Green MR. Analysis of Gal4-directed transcription activation using Tra1 mutants selectively defective for interaction with Gal4. <i>Proc Natl Acad Sci U S A</i> 2012;109(6):1997–2002.	doi:10.1073/pnas.1116340109 PMID 22308403 PMC3277556	GSE31389	4	1	2	4	–	2	ChIP
775	Denby 2012 Denby CM, Im JH, Yu RC, Pesce CG, Brem RB. Negative feedback confers mutational robustness in yeast transcription factor regulation. <i>Proc Natl Acad Sci U S A</i> 2012;109(10):3874–8.	doi:10.1073/pnas.1116360109 PMID 22355134 PMC3309721	–	4	4	4	–	4	1	GFP - wt vs TF mutant
776	Levi 2012 Levi CE, Cardillo SB, Bertotti S, Ríos C, Correa García S, Moretti MB. GABA induction of the Saccharomyces cerevisiae UGA4 gene depends on the quality of the carbon source: role of the key transcription factors acting in this process. <i>Biochem Biophys Res Commun</i> 2012;421(3):572–7.	doi:10.1016/j.bbrc.2012.04.047 PMID 22525679	–	4	4	4	4	–	2	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
777	Cardona 2012 Cardona F, Del Olmo ML, Aranda A. Phylogenetic origin and transcriptional regulation at the post-diauxic phase of SPI1, in <i>Saccharomyces cerevisiae</i> . <i>Cell Mol Biol Lett</i> 2012;17(3):393–407.	doi:10.2478/s11658-012-0017-4 PMID 22610976 PMC6275683	–	4	3	3	–	4	3	Northern blot - wt vs TF mutant
778	Torbensen 2012 Torbensen R, Møller HD, Gresham D, et al., Regenber B. Amino acid transporter genes are essential for FLO11-dependent and FLO11-independent biofilm formation and invasive growth in <i>Saccharomyces cerevisiae</i> . <i>PLoS One</i> 2012;7(7):e41272.	doi:10.1371/journal.pone.0041272 PMID 22844449 PMC3406018	–	4	1	4	–	4	1	Microarray analysis - wt vs TF mutant
779	Tanaka 2012 Tanaka K, Ishii Y, Ogawa J, Shima J. Enhancement of acetic acid tolerance in <i>Saccharomyces cerevisiae</i> by overexpression of the HAA1 gene, encoding a transcriptional activator. <i>Appl Environ Microbiol</i> 2012;78(22):8161–3.	doi:10.1128/AEM.02356-12 PMID 22961896 PMC3485965	–	4	1	4	–	4	1	RT-PCR - wt vs TF overexpression (altered promoter sequence)
780	Holmes 2013 Holmes DL, Lancaster AK, Lindquist S, Halfmann R. Heritable remodeling of yeast multicellularity by an environmentally responsive prion. <i>Cell</i> 2013;153(1):153–65.	doi:10.1016/j.cell.2013.02.026 PMID 23540696 PMC3759520	–	4	1	4	–	4	1	RT-PCR - wt vs TF mutant
781	Kodo 2013 Kodo N, Matsuda T, Doi S, Munakata H. Salicylic acid resistance is conferred by a novel YRR1 mutation in <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2013;434(1):42–7.	doi:10.1016/j.bbrc.2013.03.069 PMID 23545261	–	4	1	4	–	4	1	RT-PCR - wt vs TF mutant
782	Herzog 2013 Herzog B, Popova B, Jakobshagen A, Shahpasandzadeh H, Braus GH. Mutual cross talk between the regulators Hac1 of the unfolded protein response and Gcn4 of the general amino acid control of <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2013;12(8):1142–54.	doi:10.1128/EC.00123-13 PMID 23794510 PMC3754535	–	4	2	4	–	4	2	RT-PCR - wt vs TF mutant
783	Downey 2013 Downey M, Knight B, Vashisht AA, et al., Toczyski DP. Gcn5 and sirTuins regulate acetylation of the ribosomal protein transcription factor Ifh1. <i>Curr Biol</i> 2013;23(17):1638–48.	doi:10.1016/j.cub.2013.06.050 PMID 23973296 PMC3982851	–	4	1	4	–	4	1	RT-PCR - wt vs TF mutant
784	Lee 2013 Lee SB, Kang HS, Kim T. Nrg1 functions as a global transcriptional repressor of glucose-repressed genes through its direct binding to the specific promoter regions. <i>Biochem Biophys Res Commun</i> 2013;439(4):501–5.	doi:10.1016/j.bbrc.2013.09.015 PMID 24025681	–	4	1	4	4	–	1	ChIP
785	Busnelli 2013 Busnelli S, Tripodi F, Nicastro R, et al., Coccetti P. Snf1/AMPK promotes SBF and MBF-dependent transcription in budding yeast. <i>Biochim Biophys Acta</i> 2013;1833(12):3254–3264.	doi:10.1016/j.bbamcr.2013.09.014 PMID 24084603	–	4	2	4	4	–	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
786	Edenberg 2014 Edenberg ER, Vashisht A, Benanti JA, Wohlschlegel J, Toczyski DP. Rad53 downregulates mitotic gene transcription by inhibiting the transcriptional activator Ndd1. <i>Mol Cell Biol</i> 2014;34(4):725–38.	doi:10.1128/MCB.01056-13 PMID 24324011 PMC3911491	–	4	2	4	4	–	2	ChIP
787	Nishida 2014 Nishida N, Jing D, Kuroda K, Ueda M. Activation of signaling pathways related to cell wall integrity and multidrug resistance by organic solvent in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 2014;60(3):149–62.	doi:10.1007/s00294-013-0419-5 PMID 24378717	–	4	4	4	–	4	1	RT-PCR - wt vs TF mutant
788	Pautasso 2014 Pautasso C, Rossi S. Transcriptional regulation of the protein kinase A subunits in <i>Saccharomyces cerevisiae</i> : autoregulatory role of the kinase A activity. <i>Biochim Biophys Acta</i> 2014; 1839(4):275–87.	doi:10.1016/j.bbagr.2014.02.005 PMID 24530423	–	4	4	4	4	–	1	lacZ - wt vs mutant promoter sequence
789	Sakihama 2015 Sakihama Y, Hasunuma T, Kondo A. Improved ethanol production from xylose in the presence of acetic acid by the overexpression of the HAA1 gene in <i>Saccharomyces cerevisiae</i> . <i>J Biosci Bioeng</i> 2015;119(3):297–302.	doi:10.1016/j.jbiosc.2014.09.004 PMID 25282639	–	4	1	4	4	–	1	RT-PCR - wt vs mutant promoter sequence
790	Nasuno 2014 Nasuno R, Aitoku M, Manago Y, Nishimura A, Sasano Y, Takagi H. Nitric oxide-mediated antioxidative mechanism in yeast through the activation of the transcription factor Mac1. <i>PLoS One</i> 2014;9(11):e113788.	doi:10.1371/journal.pone.0113788 PMID 25423296 PMC4244153	GSE61836	4	1	4	2	2	1	ChIP
791	Fayyad-Kazan 2016 Fayyad-Kazan M, Feller A, Bodo E, et al., Georis I. Yeast nitrogen catabolite repression is sustained by signals distinct from glutamine and glutamate reservoirs. <i>Mol Microbiol</i> 2016; 99(2):360–79.	doi:10.1111/mmi.13236 PMID 26419331	–	4	2	4	–	4	4	RT-PCR - wt vs TF mutant
792	Zhang 2015 Zhang MM, Zhao XQ, Cheng C, Bai FW. Improved growth and ethanol fermentation of <i>Saccharomyces cerevisiae</i> in the presence of acetic acid by overexpression of SET5 and PPR1. <i>Biotechnol J</i> 2015;10(12):1903–11.	doi:10.1002/biot.201500508 PMID 26479519	–	4	1	4	–	4	2	RT-PCR - wt vs TF overexpression
793	Qiu 2016 Qiu H, Chereji RV, Hu C, et al., Hinnebusch AG. Genome-wide cooperation by HAT Gcn5, remodeler SWI/SNF, and chaperone Ydj1 in promoter nucleosome eviction and transcriptional activation. <i>Genome Res</i> 2016;26(2):211–25.	doi:10.1101/gr.196337.115 PMID 26602697 PMC4728374	–	4	1	4	4	–	1	ChIP
794	Wu 2016 Wu Y, Du J, Xu G, Jiang L. The transcription factor Ace2 and its paralog Swi5 regulate ethanol production during static fermentation through their targets Cts1 and Rps4a in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Res</i> 2016;16(3).	doi:10.1093/femsyr/fow022 PMID 26975390	–	4	2	4	–	4	1	Gene overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
795	Podholová 2016 Podholová K, Plocek V, Rešetárová S, et al., Palková Z. Divergent branches of mitochondrial signaling regulate specific genes and the viability of specialized cell types of differentiated yeast colonies. <i>Oncotarget</i> 2016;7(13):15299–314.	doi:10.18632/oncotarget.8084 PMID 26992228 PMC4941242	–	4	2	4	–	4	3	Western blot - wt vs TF mutant
796	Yan 2016 Yan C, Wu S, Pocetti C, Bai L. Regulation of cell-to-cell variability in divergent gene expression. <i>Nat Commun</i> 2016;7: 11099.	doi:10.1038/ncomms11099 PMID 27010670 PMC4820839	–	4	4	4	–	4	2	GFP - wt vs TF mutant
797	Malo 2016 Malo ME, Postnikoff SD, Arnason TG, Harkness TA. Mitotic degradation of yeast Fkh1 by the Anaphase Promoting Complex is required for normal longevity, genomic stability and stress resistance. <i>Aging (Albany NY)</i> 2016;8(4):810–30.	doi:10.18632/aging.100949 PMID 27099939 PMC4925830	–	4	2	4	4	–	2	ChIP
798	Xie 2016 Xie B, Horecka J, Chu A, Davis RW, Becker E, Primig M. Ndt80 activates the meiotic ORC1 transcript isoform and SMA2 via a bi-directional middle sporulation element in <i>Saccharomyces cerevisiae</i> . <i>RNA Biol</i> 2016;13(9):772–82.	doi:10.1080/15476286.2016.1191738 PMID 27362276 PMC5013992	–	4	2	4	1	3	4	Western blot - wt vs TF mutant
799	Chakrabortee 2016 Chakrabortee S, Byers JS, Jones S, et al., Jarosz DF. Intrinsically Disordered Proteins Drive Emergence and Inheritance of Biological Traits. <i>Cell</i> 2016;167(2):369–381.e12.	doi:10.1016/j.cell.2016.09.017 PMID 27693355 PMC5066306	–	4	3	4	–	4	1	RT-PCR - wt vs TF mutant
800	Albert 2016 Albert B, Knight B, Merwin J, et al., Shore D. A Molecular Titration System Coordinates Ribosomal Protein Gene Transcription with Ribosomal RNA Synthesis. <i>Mol Cell</i> 2016; 64(4):720–733.	doi:10.1016/j.molcel.2016.10.003 PMID 27818142	–	4	2	4	4	–	2	ChIP
801	Karpov 2017 Karpov DS, Grineva EN, Leinsoo AT, et al., Karpov VL. Functional analysis of <i>Debaryomyces hansenii</i> Rpn4 on a genetic background of <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Res</i> 2017;17(1).	doi:10.1093/femsyr/fow098 PMID 27856503	–	4	1	4	–	4	1	RT-PCR - wt vs TF mutant
802	Zyrina 2017 Zyrina AN, Smirnova EA, Markova OV, Severin FF, Knorre DA. Mitochondrial Superoxide Dismutase and Yap1p Act as a Signaling Module Contributing to Ethanol Tolerance of the Yeast <i>Saccharomyces cerevisiae</i> . <i>Appl Environ Microbiol</i> 2017; 83(3).	doi:10.1128/AEM.02759-16 PMID 27864171 PMC5244310	–	4	1	2	–	4	4	GFP - wt vs TF mutant
803	Swinnen 2017 Swinnen S, Henriques SF, Shrestha R, Ho PW, Sá-Correia I, Nevoigt E. Improvement of yeast tolerance to acetic acid through Haa1 transcription factor engineering: towards the underlying mechanisms. <i>Microb Cell Fact</i> 2017;16(1):7.	doi:10.1186/s12934-016-0621-5 PMID 28068993 PMC5220606	–	4	1	4	–	4	1	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
804	Collins 2017 Collins ME, Black JJ, Liu Z. Casein Kinase I Isoform Hrr25 Is a Negative Regulator of Haa1 in the Weak Acid Stress Response Pathway in <i>Saccharomyces cerevisiae</i> . <i>Appl Environ Microbiol</i> 2017;83(13).	doi:10.1128/AEM.00672-17 PMID 28432100 PMC5478977	–	4	1	2	–	4	2	lacZ - wt vs TF mutant
805	Jaiswal 2017 Jaiswal D, Jezek M, Quijote J, et al., Green EM. Repression of Middle Sporulation Genes in <i>Saccharomyces cerevisiae</i> by the Sum1-Rfm1-Hst1 Complex Is Maintained by Set1 and H3K4 Methylation. <i>G3 (Bethesda)</i> 2017;7(12):3971–3982.	doi:10.1534/g3.117.300150 PMID 29066473 PMC5714494	–	4	1	4	–	4	1	RT-PCR - wt vs TF mutant
806	Zhang 2017 Zhang T, Galdieri L, Hasek J, Vancura A. Yeast phospholipase C is required for stability of casein kinase I Yck2p and expression of hexose transporters. <i>FEMS Microbiol Lett</i> 2017;364(22).	doi:10.1093/femsle/fnx227 PMID 29087456 PMC5827616	–	4	1	4	–	4	1	RT-PCR - wt vs TF mutant
807	Aymoz 2018 Aymoz D, Solé C, Pierre JJ, et al., Pelet S. Timing of gene expression in a cell-fate decision system. <i>Mol Syst Biol</i> 2018; 14(4):e8024.	doi:10.15252/msb.20178024 PMID 29695607 PMC5916086	–	4	2	4	4	–	1	ChIP
808	Chow 2018 Chow J, Notaro M, Prabhakar A, Free SJ, Cullen PJ. Impact of Fungal MAPK Pathway Targets on the Cell Wall. <i>J Fungi (Basel)</i> 2018;4(3).	doi:10.3390/jof4030093 PMID 30096860 PMC6162559	–	4	1	4	–	4	1	RT-PCR - wt vs TF mutant
809	Ramos-Alonso 2018 Ramos-Alonso L, Wittmaack N, Mulet I, et al., Puig S. Molecular strategies to increase yeast iron accumulation and resistance. <i>Metallomics</i> 2018;10(9):1245–1256.	doi:10.1039/c8mt00124c PMID 30137082	–	4	1	4	–	4	1	RT-PCR - wt vs TF overexpression
810	Chowdhary 2019 Chowdhary S, Kainth AS, Pincus D, Gross DS. Heat Shock Factor 1 Drives Intergenic Association of Its Target Gene Loci upon Heat Shock. <i>Cell Rep</i> 2019;26(1):18–28.e5.	doi:10.1016/j.celrep.2018.12.034 PMID 30605674 PMC6435272	–	4	1	3	4	–	3	ChIP
811	Voutsina 2019 Voutsina A, Fragiadakis GS, Gkouskou K, Alexandraki D. Synergy of Hir1, Ssn6, and Snf2 global regulators is the functional determinant of a Mac1 transcriptional switch in <i>S. cerevisiae</i> copper homeostasis. <i>Curr Genet</i> 2019;65(3): 799–816.	doi:10.1007/s00294-019-00935-5 PMID 30689022	–	4	1	1	3	1	4	ChIP
812	Jiang 2019 Jiang Y, Berg MD, Genereaux J, et al., Lajoie P. Sfp1 links TORC1 and cell growth regulation to the yeast SAGA-complex component Tra1 in response to polyQ proteotoxicity. <i>Traffic</i> 2019;20(4):267–283.	doi:10.1111/tra.12637 PMID 30740854	–	4	1	4	–	4	4	lacZ - wt vs TF mutant
813	Chow 2019 Chow J, Dionne HM, Prabhakar A, et al., Cullen PJ. Aggregate Filamentous Growth Responses in Yeast. <i>mSphere</i> 2019;4(2).	doi:10.1128/mSphere.00702-18 PMID 30842272 PMC6403458	–	4	4	4	–	4	1	RT-PCR - wt vs. TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
814	Aref 2021 Aref R, Sanad MNME, Schüller HJ. Forkhead transcription factor Fkh1: insights into functional regulatory domains crucial for recruitment of Sin3 histone deacetylase complex. <i>Curr Genet</i> 2021;67(3):487–499.	doi:10.1007/s00294-021-01158-3 PMID 33635403 PMC8139909	–	4	1	4	4	–	2	ChIP
815	Ottillie 2022 Ottillie S, Luth MR, Hellemann E, et al., Winzeler EA. Adaptive laboratory evolution in <i>S. cerevisiae</i> highlights role of transcription factors in fungal xenobiotic resistance. <i>Commun Biol</i> 2022;5(1):128.	doi:10.1038/s42003-022-03076-7 PMID 35149760 PMC8837787	–	4	1	4	–	4	1	RT-PCR - wt vs TF mutant
816	Schüller 1992 Schüller HJ, Schorr R, Hoffmann B, Schweizer E. Regulatory gene INO4 of yeast phospholipid biosynthesis is positively autoregulated and functions as a transactivator of fatty acid synthase genes FAS1 and FAS2 from <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Res</i> 1992;20(22):5955–61.	doi:10.1093/nar/20.22.5955 PMID 1461729 PMC334460	–	3	1	3	–	3	2	Northern blot - wt vs TF mutant
817	Dohrmann 1992 Dohrmann PR, Butler G, Tamai K, et al., Stillman DJ. Parallel pathways of gene regulation: homologous regulators SWI5 and ACE2 differentially control transcription of HO and chitinase. <i>Genes Dev</i> 1992;6(1):93–104.	doi:10.1101/gad.6.1.93 PMID 1730413	–	3	3	3	–	3	1	Northern blot - wt vs TF mutant
818	Turi 1992 Turi TG, Loper JC. Multiple regulatory elements control expression of the gene encoding the <i>Saccharomyces cerevisiae</i> cytochrome P450, lanosterol 14 alpha-demethylase (ERG11). <i>J Biol Chem</i> 1992;267(3):2046–56.	PMID 1730736	–	3	2	3	1	2	2	Northern blot - wt vs TF mutant
819	Mountain 1991 Mountain HA, Byström AS, Larsen JT, Korch C. Four major transcriptional responses in the methionine/threonine biosynthetic pathway of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 1991;7(8):781–803.	doi:10.1002/yea.320070804 PMID 1789001	–	3	1	3	–	3	1	Northern blot - wt vs TF mutant
820	Hwang-Shum 1991 Hwang-Shum JJ, Hagen DC, Jarvis EE, Westby CA, Sprague GF Jr. Relative contributions of MCM1 and STE12 to transcriptional activation of a- and alpha-specific genes from <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1991;227(2):197–204.	doi:10.1007/BF00259671 PMID 1905781	–	3	2	3	2	1	2	EMSA
821	Moehle 1991 Moehle CM, Hinnebusch AG. Association of RAP1 binding sites with stringent control of ribosomal protein gene transcription in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1991;11(5):2723–35.	doi:10.1128/mcb.11.5.2723-2735.1991 PMID 2017175 PMC360042	–	3	1	3	–	3	2	lacZ - wt vs TF mutant
822	Roy 1990 Roy A, Exinger F, Losson R. cis- and trans-acting regulatory elements of the yeast URA3 promoter. <i>Mol Cell Biol</i> 1990;10(10):5257–70.	doi:10.1128/mcb.10.10.5257-5270.1990 PMID 2204810 PMC361211	–	3	1	3	3	–	1	DNA footprinting

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
823	Santangelo 1990 Santangelo GM, Tornow J. Efficient transcription of the glycolytic gene ADH1 and three translational component genes requires the GCR1 product, which can act through TUF/GRF/RAP binding sites. <i>Mol Cell Biol</i> 1990;10(2): 859–62.	doi:10.1128/mcb.10.2.859-862.1990 PMID 2405258 PMC360891	–	3	1	3	3	–	1	DNA footprinting
824	Thorsness 1989 Thorsness M, Schafer W, D’Ari L, Rine J. Positive and negative transcriptional control by heme of genes encoding 3-hydroxy-3-methylglutaryl coenzyme A reductase in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1989;9(12):5702–12.	doi:10.1128/mcb.9.12.5702-5712.1989 PMID 2685574 PMC363742	–	3	2	3	–	3	1	lacZ - wt vs TF mutant
825	Friden 1987 Friden P, Schimmel P. LEU3 of <i>Saccharomyces cerevisiae</i> encodes a factor for control of RNA levels of a group of leucine-specific genes. <i>Mol Cell Biol</i> 1987;7(8):2708–17.	doi:10.1128/mcb.7.8.2708-2717.1987 PMID 2823102 PMC367887	–	3	1	3	1	2	2	Northern blot - wt vs TF mutant
826	Baker 1986 Baker HV. Glycolytic gene expression in <i>Saccharomyces cerevisiae</i> : nucleotide sequence of GCR1, null mutants, and evidence for expression. <i>Mol Cell Biol</i> 1986;6(11):3774–84.	doi:10.1128/mcb.6.11.3774-3784.1986 PMID 3025612 PMC367138	–	3	1	3	–	3	1	Enzyme activity - wt vs TF mutant
827	Bailis 1987 Bailis AM, Poole MA, Carman GM, Henry SA. The membrane-associated enzyme phosphatidylserine synthase is regulated at the level of mRNA abundance. <i>Mol Cell Biol</i> 1987;7(1):167–76.	doi:10.1128/mcb.7.1.167-176.1987 PMID 3031455 PMC365053	–	3	3	3	–	3	1	Northern blot - wt vs TF mutant
828	Hashimoto 1983 Hashimoto H, Kikuchi Y, Nogi Y, Fukasawa T. Regulation of expression of the galactose gene cluster in <i>Saccharomyces cerevisiae</i> . Isolation and characterization of the regulatory gene GAL4. <i>Mol Gen Genet</i> 1983;191(1):31–8.	doi:10.1007/BF00330886 PMID 6350827	–	3	1	3	–	–	1	–
829	Stephen 1995 Stephen DW, Rivers SL, Jamieson DJ. The role of the YAP1 and YAP2 genes in the regulation of the adaptive oxidative stress responses of <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 1995;16(3):415–23.	doi:10.1111/j.1365-2958.1995.tb02407.x PMID 7565103	–	3	1	3	–	3	1	lacZ - wt vs TF mutant
830	Deckert 1995 Deckert J, Rodriguez Torres AM, Simon JT, Zitomer RS. Mutational analysis of Rox1, a DNA-bending repressor of hypoxic genes in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1995;15(11):6109–17.	doi:10.1128/MCB.15.11.6109-6117.1995 PMID 7565763 PMC230862	–	3	1	3	–	2	1	Northern blot - wt vs TF mutant
831	Blinder 1995 Blinder D, Magasanik B. Recognition of nitrogen-responsive upstream activation sequences of <i>Saccharomyces cerevisiae</i> by the product of the GLN3 gene. <i>J Bacteriol</i> 1995;177(14): 4190–3.	doi:10.1128/jb.177.14.4190-4193.1995 PMID 7608102 PMC177161	–	3	1	3	1	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
832	Yamaguchi-Iwai 1995 Yamaguchi-Iwai Y, Dancis A, Klausner RD. AFT1: a mediator of iron regulated transcriptional control in <i>Saccharomyces cerevisiae</i> . <i>EMBO J</i> 1995;14(6):1231–9.	doi:10.1002/j.1460-2075.1995.tb07106.x PMID 7720713 PMC398200	–	3	1	3	–	3	1	lacZ - wt vs TF overexpression
833	Deckert 1995 Deckert J, Perini R, Balasubramanian B, Zitomer RS. Multiple elements and auto-repression regulate Rox1, a repressor of hypoxic genes in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 1995;139(3):1149–58.	doi:10.1093/genetics/139.3.1149 PMID 7768429 PMC1206446	–	3	2	3	–	3	1	lacZ - wt vs TF mutant
834	Hosaka 1994 Hosaka K, Nikawa J, Kodaki T, Yamashita S. Cloning and characterization of the SCS1 gene required for the expression of genes in yeast phospholipid synthesis. <i>J Biochem</i> 1994; 115(1):131–6.	doi:10.1093/oxfordjournals.jbchem.a124287 PMID 8188619	–	3	1	3	–	3	1	Northern blot - wt vs TF mutant
835	Martens 1994 Martens JA, Brandl CJ. GCN4p activation of the yeast TRP3 gene is enhanced by ABF1p and uses a suboptimal TATA element. <i>J Biol Chem</i> 1994;269(22):15661–7.	PMID 8195216	–	3	2	3	2	1	2	EMSA
836	Lai 1994 Lai K, McGraw P. Dual control of inositol transport in <i>Saccharomyces cerevisiae</i> by irreversible inactivation of permease and regulation of permease synthesis by INO2, INO4, and OPI1. <i>J Biol Chem</i> 1994;269(3):2245–51.	PMID 8294482	–	3	3	3	–	3	1	Northern blot - wt vs TF mutant
837	Schwob 1993 Schwob E, Nasmyth K. CLB5 and CLB6, a new pair of B cyclins involved in DNA replication in <i>Saccharomyces cerevisiae</i> . <i>Genes Dev</i> 1993;7(7A):1160–75.	doi:10.1101/gad.7.7a.1160 PMID 8319908	–	3	2	3	2	–	2	EMSA
838	Young 1993 Young MR, Craig EA. <i>Saccharomyces cerevisiae</i> HSP70 heat shock elements are functionally distinct. <i>Mol Cell Biol</i> 1993; 13(9):5637–46.	doi:10.1128/mcb.13.9.5637-5646.1993 PMID 8355706 PMC360292	–	3	1	3	3	–	2	EMSA
839	Dorrington 1993 Dorrington RA, Cooper TG. The DAL82 protein of <i>Saccharomyces cerevisiae</i> binds to the DAL upstream induction sequence (UIS). <i>Nucleic Acids Res</i> 1993;21(16): 3777–84.	doi:10.1093/nar/21.16.3777 PMID 8367295 PMC309890	–	3	1	3	–	–	1	–
840	Scott 1993 Scott EW, Baker HV. Concerted action of the transcriptional activators REB1, RAP1, and GCR1 in the high-level expression of the glycolytic gene TPI. <i>Mol Cell Biol</i> 1993; 13(1):543–50.	doi:10.1128/mcb.13.1.543-550.1993 PMID 8417350 PMC358933	–	3	3	3	3	–	1	lacZ - wt vs mutant promoter sequence
841	Willett 1993 Willett CE, Gelfman CM, Holland MJ. A complex regulatory element from the yeast gene ENO2 modulates GCR1-dependent transcriptional activation. <i>Mol Cell Biol</i> 1993;13(4):2623–33.	doi:10.1128/mcb.13.4.2623-2633.1993 PMID 8455635 PMC359601	–	3	3	3	3	–	1	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
842	Amillet 1995 Amillet JM, Buisson N, Labbe-Bois R. Positive and negative elements involved in the differential regulation by heme and oxygen of the HEM13 gene (coproporphyrinogen oxidase) in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 1995;28(6):503–11.	doi:10.1007/BF00518161 PMID 8593679	–	3	2	3	1	2	2	Enzyme activity - wt vs TF mutant
843	Bobola 1996 Bobola N, Jansen RP, Shin TH, Nasmyth K. Asymmetric accumulation of Ash1p in postanaphase nuclei depends on a myosin and restricts yeast mating-type switching to mother cells. <i>Cell</i> 1996;84(5):699–709.	doi:10.1016/s0092-8674(00)81048-x PMID 8625408	–	3	3	3	–	3	1	Northern blot - wt vs TF mutant
844	Kurihara 1996 Kurihara LJ, Stewart BG, Gammie AE, Rose MD. Kar4p, a karyogamy-specific component of the yeast pheromone response pathway. <i>Mol Cell Biol</i> 1996;16(8):3990–4002.	doi:10.1128/MCB.16.8.3990 PMID 8754797 PMC231395	–	3	2	2	–	3	1	Northern blot - wt vs TF mutant
845	Smart 1996 Smart WC, Coffman JA, Cooper TG. Combinatorial regulation of the <i>Saccharomyces cerevisiae</i> CAR1 (arginase) promoter in response to multiple environmental signals. <i>Mol Cell Biol</i> 1996;16(10):5876–87.	doi:10.1128/MCB.16.10.5876 PMID 8816501 PMC231589	–	3	3	3	–	3	2	Northern blot - wt vs TF mutant
846	Hellauer 1996 Hellauer K, Rochon MH, Turcotte B. A novel DNA binding motif for yeast zinc cluster proteins: the Leu3p and Pdr3p transcriptional activators recognize everted repeats. <i>Mol Cell Biol</i> 1996;16(11):6096–102.	doi:10.1128/MCB.16.11.6096 PMID 8887639 PMC231612	–	3	1	3	2	1	2	EMSA
847	Springer 1996 Springer C, Künzler M, Balmelli T, Braus GH. Amino acid and adenine cross-pathway regulation act through the same 5'-TGACTC-3' motif in the yeast HIS7 promoter. <i>J Biol Chem</i> 1996;271(47):29637–43.	doi:10.1074/jbc.271.47.29637 PMID 8939895	–	3	3	3	2	1	2	EMSA
848	Rolfes 1997 Rolfes RJ, Zhang F, Hinnebusch AG. The transcriptional activators BAS1, BAS2, and ABF1 bind positive regulatory sites as the critical elements for adenine regulation of ADE5,7. <i>J Biol Chem</i> 1997;272(20):13343–54.	doi:10.1074/jbc.272.20.13343 PMID 9148957	–	3	3	3	1	2	2	lacZ - wt vs TF mutant
849	Labbé 1997 Labbé S, Zhu Z, Thiele DJ. Copper-specific transcriptional repression of yeast genes encoding critical components in the copper transport pathway. <i>J Biol Chem</i> 1997;272(25):15951–8.	doi:10.1074/jbc.272.25.15951 PMID 9188496	–	3	1	3	1	–	1	–
850	Yamaguchi-Iwai 1997 Yamaguchi-Iwai Y, Serpe M, Haile D, et al., Dancis A. Homeostatic regulation of copper uptake in yeast via direct binding of MAC1 protein to upstream regulatory sequences of FRE1 and CTR1. <i>J Biol Chem</i> 1997;272(28):17711–8.	doi:10.1074/jbc.272.28.17711 PMID 9211922	–	3	1	3	–	2	2	–
851	Zhao 1997 Zhao H, Eide DJ. Zap1p, a metalloregulatory protein involved in zinc-responsive transcriptional regulation in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1997;17(9):5044–52.	doi:10.1128/MCB.17.9.5044 PMID 9271382 PMC232355	–	3	1	3	–	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
852	Hiesinger 1997 Hiesinger M, Wagner C, Schüller HJ. The acetyl-CoA synthetase gene ACS2 of the yeast <i>Saccharomyces cerevisiae</i> is coregulated with structural genes of fatty acid biosynthesis by the transcriptional activators Ino2p and Ino4p. <i>FEBS Lett</i> 1997;415(1):16–20.	doi:10.1016/s0014-5793(97)01085-5 PMID 9326360	–	3	3	3	1	2	2	lacZ - wt vs TF mutant
853	Mai 1997 Mai B, Breeden L. Xbp1, a stress-induced transcriptional repressor of the <i>Saccharomyces cerevisiae</i> Swi4/Mbp1 family. <i>Mol Cell Biol</i> 1997;17(11):6491–501.	doi:10.1128/MCB.17.11.6491 PMID 9343412 PMC232502	–	3	1	3	–	3	1	Northern blot - wt vs TF overexpression
854	Aerne 1998 Aerne BL, Johnson AL, Toyn JH, Johnston LH. Swi5 controls a novel wave of cyclin synthesis in late mitosis. <i>Mol Biol Cell</i> 1998;9(4):945–56.	doi:10.1091/mbc.9.4.945 PMID 9529390 PMC25320	–	3	1	3	–	3	1	Northern blot - wt vs TF mutant
855	Dzugasová 1998 Dzugasová V, Obernauerová M, Horváthová K, Vachová M, Záková M, Subík J. Phosphatidylglycerolphosphate synthase encoded by the PEL1/PGS1 gene in <i>Saccharomyces cerevisiae</i> is localized in mitochondria and its expression is regulated by phospholipid precursors. <i>Curr Genet</i> 1998;34(4):297–302.	doi:10.1007/s002940050399 PMID 9799363	–	3	3	3	–	3	2	lacZ - wt vs TF mutant
856	Proft 1999 Proft M, Serrano R. Repressors and upstream repressing sequences of the stress-regulated ENA1 gene in <i>Saccharomyces cerevisiae</i> : bZIP protein Sko1p confers HOG-dependent osmotic regulation. <i>Mol Cell Biol</i> 1999;19(1):537–46.	doi:10.1128/MCB.19.1.537 PMID 9858577 PMC83911	–	3	3	3	1	2	2	lacZ - wt vs TF mutant
857	Rupp 1999 Rupp S, Summers E, Lo HJ, Madhani H, Fink G. MAP kinase and cAMP filamentation signaling pathways converge on the unusually large promoter of the yeast FLO11 gene. <i>EMBO J</i> 1999;18(5):1257–69.	doi:10.1093/emboj/18.5.1257 PMID 10064592 PMC1171216	–	3	3	3	–	3	2	Northern blot - wt vs TF mutant
858	Iraqi 1999 Iraqi I, Vissers S, André B, Urrestarazu A. Transcriptional induction by aromatic amino acids in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1999;19(5):3360–71.	doi:10.1128/MCB.19.5.3360 PMID 10207060 PMC84129	–	3	2	3	–	3	1	Northern blot - wt vs TF mutant
859	Charizanis 1999 Charizanis C, Juhnke H, Krems B, Entian KD. The oxidative stress response mediated via Pos9/Skn7 is negatively regulated by the Ras/PKA pathway in <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1999;261(4–5):740–52.	doi:10.1007/s004380050017 PMID 10394911	–	3	2	3	–	–	1	–
860	Wagner 1999 Wagner C, Blank M, Strohmam B, Schüller HJ. Overproduction of the Opi1 repressor inhibits transcriptional activation of structural genes required for phospholipid biosynthesis in the yeast <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 1999;15(10A):843–54.	<a href="https://doi.org/10.1002/(SICI)1097-0061(199907)15:10A<843::AID-YEA424>3.0.CO;2-M">doi:10.1002/(SICI)1097-0061(199907)15:10A<843::AID-YEA424>3.0.CO;2-M PMID 10407264	–	3	3	3	–	3	1	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
861	Inoue 1999 Inoue Y, Matsuda T, Sugiyama K, Izawa S, Kimura A. Genetic analysis of glutathione peroxidase in oxidative stress response of <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 1999; 274(38):27002–9.	doi:10.1074/jbc.274.38.27002 PMID 10480913	–	3	1	3	–	2	3	–
862	Herrero 1999 Herrero P, Flores L, de la Cera T, Moreno F. Functional characterization of transcriptional regulatory elements in the upstream region of the yeast <i>GLK1</i> gene. <i>Biochem J</i> 1999;343 Pt 2(Pt 2):319–25.	PMID 10510295 PMC1220556	–	3	3	3	3	–	2	EMSA
863	Park 1999 Park HD, Scott S, Rai R, Dorrington R, Cooper TG. Synergistic operation of the <i>CAR2</i> (Ornithine transaminase) promoter elements in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1999;181(22):7052–64.	doi:10.1128/JB.181.22.7052-7064.1999 PMID 10559172 PMC94181	–	3	3	3	–	3	1	lacZ - wt vs TF mutant
864	Kobayashi 1999 Kobayashi O, Yoshimoto H, Sone H. Analysis of the genes activated by the <i>FLO8</i> gene in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 1999;36(5):256–61.	doi:10.1007/s002940050498 PMID 10591965	–	3	1	3	–	3	1	Microarray analysis - wt vs TF mutant
865	Rouillon 2000 Rouillon A, Barbey R, Patton EE, Tyers M, Thomas D. Feedback-regulated degradation of the transcriptional activator Met4 is triggered by the SCF(Met30) complex. <i>EMBO J</i> 2000;19(2):282–94.	doi:10.1093/emboj/19.2.282 PMID 10637232 PMC305562	–	3	3	3	3	–	1	EMSA
866	Tsujimoto 2000 Tsujimoto Y, Izawa S, Inoue Y. Cooperative regulation of DOG2, encoding 2-deoxyglucose-6-phosphate phosphatase, by Snf1 kinase and the high-osmolarity glycerol-mitogen-activated protein kinase cascade in stress responses of <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 2000;182(18):5121–6.	doi:10.1128/JB.182.18.5121-5126.2000 PMID 10960096 PMC94660	–	3	3	3	–	3	2	lacZ - wt vs TF mutant
867	Oshiro 2000 Oshiro J, Rangaswamy S, Chen X, Han GS, Quinn JE, Carman GM. Regulation of the DPP1-encoded diacylglycerol pyrophosphate (DGPP) phosphatase by inositol and growth phase. Inhibition of DGPP phosphatase activity by CDP-diacylglyceron and activation of phosphatidylserine synthase activity by DGPP. <i>J Biol Chem</i> 2000;275(52):40887–96.	doi:10.1074/jbc.M008144200 PMID 11016943	–	3	3	3	–	3	1	lacZ - wt vs TF mutant
868	Jelinsky 2000 Jelinsky SA, Estep P, Church GM, Samson LD. Regulatory networks revealed by transcriptional profiling of damaged <i>Saccharomyces cerevisiae</i> cells: Rpn4 links base excision repair with proteasomes. <i>Mol Cell Biol</i> 2000;20(21):8157–67.	doi:10.1128/MCB.20.21.8157-8167.2000 PMID 11027285 PMC86425	–	3	1	3	–	3	1	Northern blot - wt vs TF mutant
869	Foury 2001 Foury F, Talibi D. Mitochondrial control of iron homeostasis. A genome wide analysis of gene expression in a yeast frataxin-deficient strain. <i>J Biol Chem</i> 2001;276(11):7762–8.	doi:10.1074/jbc.M005804200 PMID 11112771	–	3	1	3	–	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
870	Pascual-Ahuir 2001 Pascual-Ahuir A, Serrano R, Proft M. The Sko1p repressor and Gcn4p activator antagonistically modulate stress-regulated transcription in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2001;21(1):16–25.	doi:10.1128/MCB.21.1.16-25.2001 PMID 11113177 PMC86564	–	3	2	3	–	3	1	lacZ - wt vs TF mutant
871	Ren 2000 Ren B, Robert F, Wyrick JJ, et al., Young RA. Genome-wide location and function of DNA binding proteins. <i>Science</i> 2000; 290(5500):2306–9.	doi:10.1126/science.290.5500.2306 PMID 11125145	–	3	1	3	–	–	1	–
872	Proft 2001 Proft M, Pascual-Ahuir A, de Nadal E, Ariño J, Serrano R, Posas F. Regulation of the Sko1 transcriptional repressor by the Hog1 MAP kinase in response to osmotic stress. <i>EMBO J</i> 2001;20(5):1123–33.	doi:10.1093/emboj/20.5.1123 PMID 11230135 PMC145460	–	3	1	3	–	3	1	Northern blot - wt vs TF mutant
873	Eiznhamer 2001 Eiznhamer DA, Ashburner BP, Jackson JC, Gardenour KR, Lopes JM. Expression of the INO2 regulatory gene of <i>Saccharomyces cerevisiae</i> is controlled by positive and negative promoter elements and an upstream open reading frame. <i>Mol Microbiol</i> 2001;39(5):1395–405.	PMID 11251853	–	3	3	3	–	3	3	Northern blot - wt vs TF mutant
874	Protchenko 2001 Protchenko O, Ferea T, Rashford J, et al., Philpott CC. Three cell wall mannoproteins facilitate the uptake of iron in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2001;276(52): 49244–50.	doi:10.1074/jbc.M109220200 PMID 11673473	no GEO	3	1	3	–	3	1	Northern blot - wt vs TF overexpression
875	Monteiro 2002 Monteiro G, Pereira GA, Netto LE. Regulation of mitochondrial thioredoxin peroxidase I expression by two different pathways: one dependent on cAMP and the other on heme. <i>Free Radic Biol Med</i> 2002;32(3):278–88.	doi:10.1016/s0891-5849(01)00801-2 PMID 11827753	–	3	3	3	–	3	2	Northern blot - wt vs TF mutant
876	Kuchin 2002 Kuchin S, Vyas VK, Carlson M. Snf1 protein kinase and the repressors Nrg1 and Nrg2 regulate FLO11, haploid invasive growth, and diploid pseudohyphal differentiation. <i>Mol Cell Biol</i> 2002;22(12):3994–4000.	doi:10.1128/MCB.22.12.3994-4000.2002 PMID 12024013 PMC133850	–	3	2	3	–	2	2	Northern blot - wt vs TF mutant
877	Jensen 2002 Jensen LT, Culotta VC. Regulation of <i>Saccharomyces cerevisiae</i> FET4 by oxygen and iron. <i>J Mol Biol</i> 2002;318(2): 251–60.	doi:10.1016/S0022-2836(02)00093-1 PMID 12051835	–	3	2	3	–	3	2	lacZ - wt vs TF mutant
878	Rodríguez-Navarro 2002 Rodríguez-Navarro S, Llorente B, Rodríguez-Manzanegue MT, et al., Pérez-Ortín JE. Functional analysis of yeast gene families involved in metabolism of vitamins B1 and B6. <i>Yeast</i> 2002;19(14):1261–76.	doi:10.1002/yea.916 PMID 12271461	–	3	1	3	–	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
879	Wheeler 2002 Wheeler GL, Quinn KA, Perrone G, Dawes IW, Grant CM. Glutathione regulates the expression of gamma-glutamylcysteine synthetase via the Met4 transcription factor. <i>Mol Microbiol</i> 2002;46(2):545–56.	doi:10.1046/j.1365-2958.2002.03174.x PMID 12406228	–	3	3	3	–	–	1	–
880	Saxena 2003 Saxena D, Kannan KB, Brandriss MC. Rapamycin treatment results in GATA factor-independent hyperphosphorylation of the proline utilization pathway activator in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2003;2(3):552–9.	doi:10.1128/EC.2.3.552-559.2003 PMID 12796300 PMC161436	–	3	3	3	–	3	1	lacZ - wt vs TF mutant
881	Kim 2003 Kim JH, Polish J, Johnston M. Specificity and regulation of DNA binding by the yeast glucose transporter gene repressor Rgt1. <i>Mol Cell Biol</i> 2003;23(15):5208–16.	doi:10.1128/MCB.23.15.5208-5216.2003 PMID 12861007 PMC165726	–	3	1	3	2	–	1	ChIP
882	Bedalov 2003 Bedalov A, Hirao M, Posakony J, Nelson M, Simon JA. NAD ⁺ -dependent deacetylase Hst1p controls biosynthesis and cellular NAD ⁺ levels in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2003;23(19):7044–54.	doi:10.1128/MCB.23.19.7044-7054.2003 PMID 12972620 PMC193940	–	3	1	3	2	–	2	ChIP
883	Wheeler 2003 Wheeler GL, Trotter EW, Dawes IW, Grant CM. Coupling of the transcriptional regulation of glutathione biosynthesis to the availability of glutathione and methionine via the Met4 and Yap1 transcription factors. <i>J Biol Chem</i> 2003;278(50):49920–8.	doi:10.1074/jbc.M310156200 PMID 14514673	–	3	3	3	–	3	1	lacZ - wt vs TF mutant
884	Willis 2003 Willis KA, Barbara KE, Menon BB, Moffat J, Andrews B, Santangelo GM. The global transcriptional activator of <i>Saccharomyces cerevisiae</i> , Gcr1p, mediates the response to glucose by stimulating protein synthesis and CLN-dependent cell cycle progression. <i>Genetics</i> 2003;165(3):1017–29.	doi:10.1093/genetics/165.3.1017 PMID 14668361 PMC1462843	–	3	1	3	–	3	1	Primer extension - wt vs TF mutant
885	Halme 2004 Halme A, Bumgarner S, Styles C, Fink GR. Genetic and epigenetic regulation of the FLO gene family generates cell-surface variation in yeast. <i>Cell</i> 2004;116(3):405–15.	doi:10.1016/s0092-8674(04)00118-7 PMID 15016375	–	3	3	3	–	3	1	Northern blot - wt vs TF mutant
886	Lyons 2004 Lyons TJ, Villa NY, Regalla LM, Kupchak BR, Vagstad A, Eide DJ. Metalloregulation of yeast membrane steroid receptor homologs. <i>Proc Natl Acad Sci U S A</i> 2004;101(15):5506–11.	doi:10.1073/pnas.0306324101 PMID 15060275 PMC397413	–	3	1	1	–	3	3	lacZ - wt vs TF mutant
887	London 2004 London MK, Keck BI, Ramos PC, Dohmen RJ. Regulatory mechanisms controlling biogenesis of ubiquitin and the proteasome. <i>FEBS Lett</i> 2004;567(2–3):259–64.	doi:10.1016/j.febslet.2004.04.078 PMID 15178333	–	3	1	3	–	–	1	–
888	Kersting 2004 Kersting MC, Choi HS, Carman GM. Regulation of the yeast EKI1-encoded ethanolamine kinase by inositol and choline. <i>J Biol Chem</i> 2004;279(34):35353–9.	doi:10.1074/jbc.M405704200 PMID 15201274	–	3	3	3	–	3	2	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
889	Srikanth 2005 Srikanth CV, Vats P, Bourbouloux A, Delrot S, Bachhawat AK. Multiple cis-regulatory elements and the yeast sulphur regulatory network are required for the regulation of the yeast glutathione transporter, Hgt1p. <i>Curr Genet</i> 2005;47(6):345–58.	doi:10.1007/s00294-005-0571-7 PMID 15821937	–	3	3	3	–	–	1	–
890	He 2005 He XJ, Fassler JS. Identification of novel Yap1p and Skn7p binding sites involved in the oxidative stress response of <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2005;58(5):1454–67.	doi:10.1111/j.1365-2958.2005.04917.x PMID 16313629 PMC2916641	–	3	1	3	1	–	1	–
891	Alenquer 2006 Alenquer M, Tenreiro S, Sá-Correia I. Adaptive response to the antimalarial drug artesunate in yeast involves Pdr1p/Pdr3p-mediated transcriptional activation of the resistance determinants TPO1 and PDR5. <i>FEMS Yeast Res</i> 2006;6(8):1130–9.	doi:10.1111/j.1567-1364.2006.00095.x PMID 17156010	–	3	2	3	–	–	1	–
892	Kaur 2007 Kaur J, Bachhawat AK. Yct1p, a novel, high-affinity, cysteine-specific transporter from the yeast <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2007;176(2):877–90.	doi:10.1534/genetics.107.070342 PMID 17435223 PMC1894615	–	3	3	3	–	3	1	lacZ - wt vs TF mutant
893	Rudra 2007 Rudra D, Mallick J, Zhao Y, Warner JR. Potential interface between ribosomal protein production and pre-rRNA processing. <i>Mol Cell Biol</i> 2007;27(13):4815–24.	doi:10.1128/MCB.02062-06 PMID 17452446 PMC1951472	–	3	1	3	3	–	1	ChIP
894	Veis 2007 Veis J, Klug H, Koranda M, Ammerer G. Activation of the G2/M-specific gene CLB2 requires multiple cell cycle signals. <i>Mol Cell Biol</i> 2007;27(23):8364–73.	doi:10.1128/MCB.01253-07 PMID 17908798 PMC2169163	–	3	2	3	3	–	1	ChIP
895	Casey 2008 Casey L, Patterson EE, Müller U, Fox CA. Conversion of a replication origin to a silencer through a pathway shared by a Forkhead transcription factor and an S phase cyclin. <i>Mol Biol Cell</i> 2008;19(2):608–22.	doi:10.1091/mbc.e07-04-0323 PMID 18045995 PMC2230585	–	3	2	3	–	3	2	Northern blot - wt vs TF mutant
896	Ahn 2009 Ahn SH, Keogh MC, Buratowski S. Ctk1 promotes dissociation of basal transcription factors from elongating RNA polymerase II. <i>EMBO J</i> 2009;28(3):205–12.	doi:10.1038/emboj.2008.280 PMID 19131970 PMC2632940	–	3	1	3	3	–	1	ChIP
897	Shock 2009 Shock TR, Thompson J, Yates JR 3rd, Madhani HD. Hog1 mitogen-activated protein kinase (MAPK) interrupts signal transduction between the Kss1 MAPK and the Tec1 transcription factor to maintain pathway specificity. <i>Eukaryot Cell</i> 2009;8(4):606–16.	doi:10.1128/EC.00005-09 PMID 19218425 PMC2669196	–	3	1	3	3	–	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
898	Birkaya 2009 Birkaya B, Maddi A, Joshi J, Free SJ, Cullen PJ. Role of the cell wall integrity and filamentous growth mitogen-activated protein kinase pathways in cell wall remodeling during filamentous growth. <i>Eukaryot Cell</i> 2009;8(8):1118–33.	doi:10.1128/EC.00006-09 PMID 19502582 PMC2725558	–	3	1	3	–	3	1	Northern blot - wt vs TF mutant
899	El Kaderi 2009 El Kaderi B, Medler S, Raghunayakula S, Ansari A. Gene looping is conferred by activator-dependent interaction of transcription initiation and termination machineries. <i>J Biol Chem</i> 2009;284(37):25015–25.	doi:10.1074/jbc.M109.007948 PMID 19602510 PMC2757206	–	3	3	3	3	–	1	ChIP
900	Di Talia 2009 Di Talia S, Wang H, Skotheim JM, Rosebrock AP, Fitcher B, Cross FR. Daughter-specific transcription factors regulate cell size control in budding yeast. <i>PLoS Biol</i> 2009;7(10):e1000221.	doi:10.1371/journal.pbio.1000221 PMID 19841732 PMC2756959	–	3	3	3	3	–	1	ChIP
901	Chen 2010 Chen S, Liu D, Finley RL Jr, Greenberg ML. Loss of mitochondrial DNA in the yeast cardiolipin synthase <i>crd1</i> mutant leads to up-regulation of the protein kinase Swelp that regulates the G2/M transition. <i>J Biol Chem</i> 2010;285(14):10397–407.	doi:10.1074/jbc.M110.100784 PMID 20086012 PMC2856246	–	3	1	3	–	3	1	RT-PCR - wt vs TF mutant
902	Herbig 2010 Herbig E, Warfield L, Fish L, et al., Hahn S. Mechanism of Mediator recruitment by tandem Gcn4 activation domains and three Gal11 activator-binding domains. <i>Mol Cell Biol</i> 2010;30(10):2376–90.	doi:10.1128/MCB.01046-09 PMID 20308326 PMC2863704	–	3	1	3	–	3	1	RT-PCR - wt vs TF mutant
903	Pérez 2010 Pérez J, Gómez A, Roncero C. Upregulation of the PRB1 gene in the <i>Saccharomyces cerevisiae</i> rim101Delta mutant produces proteolytic artefacts that differentially affect some proteins. <i>Yeast</i> 2010;27(8):575–81.	doi:10.1002/yea.1776 PMID 20641016	–	3	1	2	–	3	2	Western blot - wt vs TF mutant
904	Landstetter 2010 Landstetter N, Glaser W, Gregori C, Seipelt J, Kuchler K. Functional genomics of drug-induced ion homeostasis identifies a novel regulatory crosstalk of iron and zinc regulons in yeast. <i>OMICS</i> 2010;14(6):651–63.	doi:10.1089/omi.2010.0031 PMID 20695822	–	3	1	3	–	3	1	Northern blot - wt vs TF mutant
905	Yu 2010 Yu Y, Neiman AM, Sternglanz R. The JmjC domain of Gis1 is dispensable for transcriptional activation. <i>FEMS Yeast Res</i> 2010;10(7):793–801.	doi:10.1111/j.1567-1364.2010.00680.x PMID 20868382 PMC2970738	–	3	1	3	3	–	1	RT-PCR - wt vs mutant promoter sequence
906	Li 2010 Li Y, Chen G, Liu W. Multiple metabolic signals influence GAL gene activation by modulating the interaction of Gal80p with the transcriptional activator Gal4p. <i>Mol Microbiol</i> 2010;78(2):414–28.	doi:10.1111/j.1365-2958.2010.07343.x PMID 20979343	–	3	1	3	–	3	2	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
907	Bakhrat 2011 Bakhrat A, Eltzov E, Finkelstein Y, Marks RS, Raveh D. UV and arsenate toxicity: a specific and sensitive yeast bioluminescence assay. <i>Cell Biol Toxicol</i> 2011;27(3):227–36.	doi:10.1007/s10565-011-9184-8 PMID 21365285	–	3	3	3	–	3	1	Luciferase - wt vs TF mutant
908	Kim 2011 Kim HS, Kim NR, Yang J, Choi W. Identification of novel genes responsible for ethanol and/or thermotolerance by transposon mutagenesis in <i>Saccharomyces cerevisiae</i> . <i>Appl Microbiol Biotechnol</i> 2011;91(4):1159–72.	doi:10.1007/s00253-011-3298-z PMID 21556919	–	3	1	3	–	3	1	Northern blot - wt vs TF mutant
909	Cardillo 2011 Cardillo SB, Correa García S, Bermúdez Moretti M. Common features and differences in the expression of the three genes forming the UGA regulon in <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2011;410(4):885–9.	doi:10.1016/j.bbrc.2011.06.086 PMID 21708130	–	3	1	3	–	3	1	RT-PCR - wt vs TF mutant
910	Sadeh 2011 Sadeh A, Movshovich N, Volokh M, Gheber L, Aharoni A. Fine-tuning of the Msn2/4-mediated yeast stress responses as revealed by systematic deletion of Msn2/4 partners. <i>Mol Biol Cell</i> 2011;22(17):3127–38.	doi:10.1091/mbc.E10-12-1007 PMID 21757539 PMC3164460	–	3	2	3	–	3	1	GFP - wt vs TF mutant
911	Cabrito 2011 Cabrito TR, Teixeira MC, Singh A, Prasad R, Sá-Correia I. The yeast ABC transporter Pdr18 (ORF YNR070w) controls plasma membrane sterol composition, playing a role in multidrug resistance. <i>Biochem J</i> 2011;440(2):195–202.	doi:10.1042/BJ20110876 PMID 21831043 PMC3215286	–	3	3	3	–	3	1	RT-PCR - wt vs TF mutant
912	Siggers 2011 Siggers T, Duyzend MH, Reddy J, Khan S, Bulyk ML. Non-DNA-binding cofactors enhance DNA-binding specificity of a transcriptional regulatory complex. <i>Mol Syst Biol</i> 2011;7:555.	doi:10.1038/msb.2011.89 PMID 22146299 PMC3737730	–	3	1	3	3	–	2	RT-PCR - wt vs mutant promoter sequence
913	Sasano 2012 Sasano Y, Haitani Y, Hashida K, Ohtsu I, Shima J, Takagi H. Overexpression of the transcription activator Msn2 enhances the fermentation ability of industrial baker's yeast in frozen dough. <i>Biosci Biotechnol Biochem</i> 2012;76(3):624–7.	doi:10.1271/bbb.110959 PMID 22451415	–	3	1	3	3	–	1	RT-PCR - wt vs mutant promoter sequence
914	Escoté 2012 Escoté X, Miranda M, Menoyo S, et al., Vendrell J. Resveratrol induces antioxidant defence via transcription factor Yap1p. <i>Yeast</i> 2012;29(7):251–63.	doi:10.1002/yea.2903 PMID 22674736	–	3	1	3	–	3	1	RT-PCR - wt vs TF mutant
915	Ferreira 2012 Ferreira RT, Silva ARC, Pimentel C, Batista-Nascimento L, Rodrigues-Pousada C, Menezes RA. Arsenic stress elicits cytosolic Ca(2+) bursts and Crz1 activation in <i>Saccharomyces cerevisiae</i> . <i>Microbiology (Reading)</i> 2012;158(Pt 9):2293–2302.	doi:10.1099/mic.0.059170-0 PMID 22745270	–	3	2	3	–	3	1	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
916	Young 2012 Young ET, Zhang C, Shokat KM, Parua PK, Braun KA. The AMP-activated protein kinase Snf1 regulates transcription factor binding, RNA polymerase II activity, and mRNA stability of glucose-repressed genes in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2012;287(34):29021–34.	doi:10.1074/jbc.M112.380147 PMID 22761425 PMC3436521	–	3	1	3	3	–	1	ChIP
917	Bontron 2013 Bontron S, Jaquenoud M, Vaga S, et al., De Virgilio C. Yeast endosulfines control entry into quiescence and chronological life span by inhibiting protein phosphatase 2A. <i>Cell Rep</i> 2013; 3(1):16–22.	doi:10.1016/j.celrep.2012.11.025 PMID 23273919	–	3	3	3	–	3	1	RT-PCR - wt vs TF mutant
918	Baker 2013 Baker CR, Hanson-Smith V, Johnson AD. Following gene duplication, paralog interference constrains transcriptional circuit evolution. <i>Science</i> 2013;342(6154):104–8.	doi:10.1126/science.1240810 PMID 24092741 PMC3911913	–	3	1	3	–	3	1	NanoString wt vs TF mutant
919	Johnson 2014 Johnson C, Kweon HK, Sheidy D, et al., Kumar A. The yeast Sks1p kinase signaling network regulates pseudohyphal growth and glucose response. <i>PLoS Genet</i> 2014;10(3):e1004183.	doi:10.1371/journal.pgen.1004183 PMID 24603354 PMC3945295	–	3	3	3	–	3	3	RT-PCR - wt vs TF mutant
920	Sugiyama 2014 Sugiyama M, Akase SP, Nakanishi R, Horie H, Kaneko Y, Harashima S. Nuclear localization of Haa1, which is linked to its phosphorylation status, mediates lactic acid tolerance in <i>Saccharomyces cerevisiae</i> . <i>Appl Environ Microbiol</i> 2014;80(11):3488–95.	doi:10.1128/AEM.04241-13 PMID 24682296 PMC4018848	–	3	1	3	–	3	1	RT-PCR - wt vs TF overexpression
921	Sarkar 2014 Sarkar S, Dalgaard JZ, Millar JB, Arumugam P. The Rim15-endosulfine-PP2Acdc55 signalling module regulates entry into gametogenesis and quiescence via distinct mechanisms in budding yeast. <i>PLoS Genet</i> 2014;10(6):e1004456.	doi:10.1371/journal.pgen.1004456 PMID 24968058 PMC4072559	–	3	3	3	–	3	1	Western blot - wt vs TF mutant
922	Barajas 2014 Barajas D, Xu K, Sharma M, Wu CY, Nagy PD. Tombusviruses upregulate phospholipid biosynthesis via interaction between p33 replication protein and yeast lipid sensor proteins during virus replication in yeast. <i>Virology</i> 2014;471–473:72–80.	doi:10.1016/j.virol.2014.10.005 PMID 25461533 PMC4775091	–	3	1	3	–	3	1	Northern blot - wt vs TF mutant
923	Lardenois 2015 Lardenois A, Stuparevic I, Liu Y, et al., Primig M. The conserved histone deacetylase Rpd3 and its DNA binding subunit Ume6 control dynamic transcript architecture during mitotic growth and meiotic development. <i>Nucleic Acids Res</i> 2015;43(1):115–28.	doi:10.1093/nar/gku1185 PMID 25477386 PMC4288150	–	3	1	3	2	1	3	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
924	Yao 2015 Yao Y, Tsuchiyama S, Yang C, et al., Schmidt M. Proteasomes, Sir2, and Hxk2 form an interconnected aging network that impinges on the AMPK/Snf1-regulated transcriptional repressor Mig1. <i>PLoS Genet</i> 2015;11(1):e1004968.	doi:10.1371/journal.pgen.1004968 PMID 25629410 PMC4309596	–	3	1	3	–	3	1	RT-PCR - wt vs TF mutant
925	Merhej 2015 Merhej J, Delaveau T, Guitard J, et al., Devaux F. Yap7 is a transcriptional repressor of nitric oxide oxidase in yeasts, which arose from neofunctionalization after whole genome duplication. <i>Mol Microbiol</i> 2015;96(5):951–72.	doi:10.1111/mmi.12983 PMID 25732006	–	3	3	3	–	3	2	RT-PCR - wt vs TF mutant
926	Piccirillo 2015 Piccirillo S, Morales R, White MG, Smith K, Kapros T, Honigberg SM. Cell Differentiation and Spatial Organization in Yeast Colonies: Role of Cell-Wall Integrity Pathway. <i>Genetics</i> 2015;201(4):1427–38.	doi:10.1534/genetics.115.180919 PMID 26510787 PMC4676521	–	3	2	3	–	3	1	lacZ - wt vs TF mutant
927	Kingsbury 2016 Kingsbury JM, Shamaprasad N, Billmyre RB, Heitman J, Cardenas ME. Cancer-associated isocitrate dehydrogenase mutations induce mitochondrial DNA instability. <i>Hum Mol Genet</i> 2016;25(16):3524–3538.	doi:10.1093/hmg/ddw195 PMID 27427385 PMC5179947	–	3	2	3	–	3	3	RT-PCR - wt vs TF mutant
928	MacDiarmid 2016 MacDiarmid CW, Taggart J, Jeong J, Kerdsonboon K, Eide DJ. Activation of the Yeast UBI4 Polyubiquitin Gene by Zap1 Transcription Factor via an Intragenic Promoter Is Critical for Zinc-deficient Growth. <i>J Biol Chem</i> 2016;291(36):18880–96.	doi:10.1074/jbc.M116.743120 PMID 27432887 PMC5009263	–	3	3	3	2	1	2	RT-PCR - wt vs mutant promoter sequence
929	Rousseau 2016 Rousseau A, Bertolotti A. An evolutionarily conserved pathway controls proteasome homeostasis. <i>Nature</i> 2016; 536(7615):184–9.	doi:10.1038/nature18943 PMID 27462806 PMC4990136	–	3	2	3	–	3	3	Western blot - wt vs TF mutant
930	Kliewe 2017 Kliewe F, Kumme J, Grigat M, Hintze S, Schüller HJ. Opi1 mediates repression of phospholipid biosynthesis by phosphate limitation in the yeast <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 2017; 34(2):67–81.	doi:10.1002/yea.3215 PMID 27743455	–	3	2	3	3	–	1	ChIP
931	Fang 2017 Fang Z, Chen Z, Wang S, et al., Huang Z. Overexpression of OLE1 Enhances Cytoplasmic Membrane Stability and Confers Resistance to Cadmium in <i>Saccharomyces cerevisiae</i> . <i>Appl Environ Microbiol</i> 2017;83(1).	doi:10.1128/AEM.02319-16 PMID 27793829 PMC5165106	–	3	2	2	–	3	2	RT-PCR - wt vs TF mutant
932	Palma 2017 Palma M, Dias PJ, Roque FC, Luzia L, Guerreiro JF, Sá-Correia I. The <i>Zygosaccharomyces bailii</i> transcription factor Haa1 is required for acetic acid and copper stress responses suggesting subfunctionalization of the ancestral bifunctional protein Haa1/Cup2. <i>BMC Genomics</i> 2017;18(1): 75.	doi:10.1186/s12864-016-3443-2 PMID 28086780 PMC5234253	–	3	1	1	–	3	3	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
933	Shedlovskiy 2017 Shedlovskiy D, Zinskie JA, Gardner E, Pestov DG, Shcherbik N. Endonucleolytic cleavage in the expansion segment 7 of 25S rRNA is an early marker of low-level oxidative stress in yeast. <i>J Biol Chem</i> 2017;292(45):18469–18485.	doi:10.1074/jbc.M117.800003 PMID 28939771 PMC5682959	–	3	1	3	–	3	3	RT-PCR - wt vs TF mutant
934	Adamczyk 2017 Adamczyk M, Szatkowska R. Low RNA Polymerase III activity results in up regulation of HXT2 glucose transporter independently of glucose signaling and despite changing environment. <i>PLoS One</i> 2017;12(9):e0185516.	doi:10.1371/journal.pone.0185516 PMID 28961268 PMC5621690	–	3	3	3	3	–	1	ChIP
935	Yamaguchi 2018 Yamaguchi Y, Katsuki Y, Tanaka S, et al., Tani M. Protective role of the HOG pathway against the growth defect caused by impaired biosynthesis of complex sphingolipids in yeast <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2018;107(3):363–386.	doi:10.1111/mmi.13886 PMID 29215176	–	3	3	3	–	3	2	lacZ - wt vs TF mutant
936	Yang 2018 Yang L, Zheng C, Chen Y, Ying H. FLO Genes Family and Transcription Factor MIG1 Regulate <i>Saccharomyces cerevisiae</i> Biofilm Formation During Immobilized Fermentation. <i>Front Microbiol</i> 2018;9:1860.	doi:10.3389/fmicb.2018.01860 PMID 30210459 PMC6119776	–	3	1	3	–	3	1	RT-PCR - wt vs TF mutant
937	Mizuno 2020 Mizuno T, Muroi K, Irie K. Snf1 AMPK positively regulates ER-phagy via expression control of Atg39 autophagy receptor in yeast ER stress response. <i>PLoS Genet</i> 2020;16(9):e1009053.	doi:10.1371/journal.pgen.1009053 PMID 32986716 PMC7544123	–	3	3	3	–	3	1	RT-PCR - wt vs. TF double mutant
938	PMID 33402426 PubMed returns no record for this id.	PMID 33402426	–	3	2	2	1	2	1	RT-PCR - wt vs TF mutant
939	Wang 2021 Wang YP, Liu L, Wang XS, et al., Xiao DG. GAT1 Gene, the GATA Transcription Activator, Regulates the Production of Higher Alcohol during Wheat Beer Fermentation by <i>Saccharomyces cerevisiae</i> . <i>Bioengineering (Basel)</i> 2021;8(5).	doi:10.3390/bioengineering8050061 PMID 34066902 PMC8151594	–	3	1	3	–	3	1	RT-PCR - wt vs. TF mutant
940	Jethmalani 2021 Jethmalani Y, Tran K, Negesse MY, et al., Green EM. Set4 regulates stress response genes and coordinates histone deacetylases within yeast subtelomeres. <i>Life Sci Alliance</i> 2021;4(12).	doi:10.26508/lsa.202101126 PMID 34625508 PMC8507492	–	3	1	3	–	3	1	RT-PCR - wt vs TF mutant
941	Romero 2022 Romero AM, Maciaszczyk-Dziubinska E, Mombeinipour M, et al., Tamás MJ. Etp1 confers arsenite resistance by affecting ACR3 expression. <i>FEMS Yeast Res</i> 2022;22(1).	doi:10.1093/femsyr/foac018 PMID 35323907 PMC9041338	–	3	1	1	–	3	3	RT-PCR - wt vs TF mutant
942	de Winde 1992 de Winde JH, Grivell LA. Global regulation of mitochondrial biogenesis in <i>Saccharomyces cerevisiae</i> : ABF1 and CPF1 play opposite roles in regulating expression of the QCR8 gene, which encodes subunit VIII of the mitochondrial ubiquinol-cytochrome c oxidoreductase. <i>Mol Cell Biol</i> 1992;12(6):2872–83.	doi:10.1128/mcb.12.6.2872-2883.1992 PMID 1317009 PMC364482	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
943	Li 1992 Li W, Brandriss MC. Proline biosynthesis in <i>Saccharomyces cerevisiae</i> : molecular analysis of the PRO1 gene, which encodes gamma-glutamyl kinase. <i>J Bacteriol</i> 1992;174(12): 4148–56.	doi:10.1128/jb.174.12.4148-4156.1992 PMID 1350780 PMC206127	–	2	1	2	–	2	2	Northern blot - wt vs TF mutant
944	Nehlin 1992 Nehlin JO, Carlberg M, Ronne H. Yeast SKO1 gene encodes a bZIP protein that binds to the CRE motif and acts as a repressor of transcription. <i>Nucleic Acids Res</i> 1992;20(20): 5271–8.	doi:10.1093/nar/20.20.5271 PMID 1437546 PMC334331	–	2	2	2	–	2	2	Northern blot - wt vs TF mutant
945	Fantino 1992 Fantino E, Marguet D, Lauquin GJ. Downstream activating sequence within the coding region of a yeast gene: specific binding in vitro of RAP1 protein. <i>Mol Gen Genet</i> 1992;236(1): 65–75.	doi:10.1007/BF00279644 PMID 1494352	–	2	1	2	2	–	2	EMSA
946	Belova 1992 Belova IV, Sambuk EV, Padkina MV, Smirnov MN. [PHO2 and GCN4 transcription activators in the regulation of <i>Saccharomyces cerevisiae</i> acid phosphatase synthesis]. <i>Genetika</i> 1992;28(5):11–8.	PMID 1639254	–	2	1	2	–	2	1	Western blot - wt vs TF mutant
947	Nasmyth 1991 Nasmyth K, Dirick L. The role of SWI4 and SWI6 in the activity of G1 cyclins in yeast. <i>Cell</i> 1991;66(5):995–1013.	doi:10.1016/0092-8674(91)90444-4 PMID 1832338	–	2	1	2	–	–	1	–
948	Simon 1991 Simon M, Adam G, Rapatz W, Spevak W, Ruis H. The <i>Saccharomyces cerevisiae</i> ADR1 gene is a positive regulator of transcription of genes encoding peroxisomal proteins. <i>Mol Cell Biol</i> 1991;11(2):699–704.	doi:10.1128/mcb.11.2.699-704.1991 PMID 1899286 PMC359720	–	2	1	2	1	1	1	CAT - wt vs TF mutant
949	van Vuuren 1991 van Vuuren HJ, Daugherty JR, Rai R, Cooper TG. Upstream induction sequence, the cis-acting element required for response to the allantoin pathway inducer and enhancement of operation of the nitrogen-regulated upstream activation sequence in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1991; 173(22):7186–95.	doi:10.1128/jb.173.22.7186-7195.1991 PMID 1938916 PMC209224	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
950	Compagno 1991 Compagno C, Ranzi BM, Martegani E. The promoter of <i>Saccharomyces cerevisiae</i> FBA1 gene contains a single positive upstream regulatory element. <i>FEBS Lett</i> 1991;293(1–2): 97–100.	doi:10.1016/0014-5793(91)81160-a PMID 1959676	–	2	1	1	2	–	2	lacZ - wt vs mutant promoter sequence
951	Elble 1991 Elble R, Tye BK. Both activation and repression of a-mating-type-specific genes in yeast require transcription factor Mcm1. <i>Proc Natl Acad Sci U S A</i> 1991;88(23): 10966–70.	doi:10.1073/pnas.88.23.10966 PMID 1961765 PMC53053	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
952	Butler 1991 Butler G, Thiele DJ. ACE2, an activator of yeast metallothionein expression which is homologous to SWI5. <i>Mol Cell Biol</i> 1991;11(1):476–85.	doi:10.1128/mcb.11.1.476-485.1991 PMID 1986241 PMC359653	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
953	Axelrod 1991 Axelrod JD, Majors J, Brandriss MC. Proline-independent binding of PUT3 transcriptional activator protein detected by footprinting in vivo. <i>Mol Cell Biol</i> 1991;11(1):564–7.	doi:10.1128/mcb.11.1.564-567.1991 PMID 1986247 PMC359669	–	2	1	2	–	–	1	–
954	Silar 1991 Silar P, Butler G, Thiele DJ. Heat shock transcription factor activates transcription of the yeast metallothionein gene. <i>Mol Cell Biol</i> 1991;11(3):1232–8.	doi:10.1128/mcb.11.3.1232-1238.1991 PMID 1996089 PMC369394	–	2	1	2	–	–	1	–
955	Dubois 1991 Dubois E, Messenguy F. In vitro studies of the binding of the ARGR proteins to the ARG5,6 promoter. <i>Mol Cell Biol</i> 1991;11(4):2162–8.	doi:10.1128/mcb.11.4.2162-2168.1991 PMID 2005902 PMC359904	–	2	2	2	–	–	1	–
956	Kihara 1991 Kihara K, Nakamura M, Akada R, Yamashita I. Positive and negative elements upstream of the meiosis-specific glucoamylase gene in <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1991;226(3):383–92.	doi:10.1007/BF00260650 PMID 2038303	–	2	2	2	–	2	1	Enzyme activity - wt vs TF mutant
957	Wittenberg 1990 Wittenberg C, Sugimoto K, Reed SI. G1-specific cyclins of <i>S. cerevisiae</i> : cell cycle periodicity, regulation by mating pheromone, and association with the p34CDC28 protein kinase. <i>Cell</i> 1990;62(2):225–37.	doi:10.1016/0092-8674(90)90361-h PMID 2142620	–	2	1	2	–	–	1	–
958	Kawasaki 1990 Kawasaki Y, Nosaka K, Kaneko Y, Nishimura H, Iwashima A. Regulation of thiamine biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1990;172(10):6145–7.	doi:10.1128/jb.172.10.6145-6147.1990 PMID 2170344 PMC526944	–	2	1	2	–	2	1	Enzyme activity - wt vs TF mutant
959	Della Seta 1990 Della Seta F, Ciafré SA, Marck C, et al., Bozzoni I. The ABF1 factor is the transcriptional activator of the L2 ribosomal protein genes in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1990;10(5):2437–41.	doi:10.1128/mcb.10.5.2437-2441.1990 PMID 2183035 PMC360595	–	2	1	2	2	–	1	ChIP
960	Brisco 1990 Brisco PR, Kohlhaw GB. Regulation of yeast LEU2. Total deletion of regulatory gene LEU3 unmasks GCN4-dependent basal level expression of LEU2. <i>J Biol Chem</i> 1990;265(20):11667–75.	PMID 2195025	–	2	2	2	1	1	2	EMSA
961	Brindle 1990 Brindle PK, Holland JP, Willett CE, Innis MA, Holland MJ. Multiple factors bind the upstream activation sites of the yeast enolase genes ENO1 and ENO2: ABFI protein, like repressor activator protein RAP1, binds cis-acting sequences which modulate repression or activation of transcription. <i>Mol Cell Biol</i> 1990;10(9):4872–85.	doi:10.1128/mcb.10.9.4872-4885.1990 PMID 2201905 PMC361101	–	2	1	2	2	–	1	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
962	Kim 1989 Kim KS, Guarente L. Mutations that alter transcriptional activation but not DNA binding in the zinc finger of yeast activator HAPI. <i>Nature</i> 1989;342(6246):200–3.	doi:10.1038/342200a0 PMID 2509943	–	2	1	2	–	–	1	–
963	Paravicini 1989 Paravicini G, Mösch HU, Schmidheini T, Braus G. The general control activator protein GCN4 is essential for a basal level of ARO3 gene expression in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1989;9(1):144–51.	doi:10.1128/mcb.9.1.144-151.1989 PMID 2564634 PMC362155	–	2	1	2	–	1	2	–
964	Visser 1989 Visser S, Andre B, Muyldermans F, Grenson M. Positive and negative regulatory elements control the expression of the UGA4 gene coding for the inducible 4-aminobutyric-acid-specific permease in <i>Saccharomyces cerevisiae</i> . <i>Eur J Biochem</i> 1989;181(2):357–61.	doi:10.1111/j.1432-1033.1989.tb14732.x PMID 2653828	–	2	2	2	–	2	1	Enzyme activity - wt vs TF mutant
965	Braus 1989 Braus G, Mösch HU, Vogel K, Hinnen A, Hütter R. Interpathway regulation of the TRP4 gene of yeast. <i>EMBO J</i> 1989;8(3):939–45.	doi:10.1002/j.1460-2075.1989.tb03455.x PMID 2656261 PMC400894	–	2	2	2	2	–	1	DNA footprinting
966	Vogel 1989 Vogel K, Hörz W, Hinnen A. The two positively acting regulatory proteins PHO2 and PHO4 physically interact with PHO5 upstream activation regions. <i>Mol Cell Biol</i> 1989;9(5):2050–7.	doi:10.1128/mcb.9.5.2050-2057.1989 PMID 2664469 PMC362998	–	2	2	2	1	–	1	–
967	Smith 1989 Smith HE, Mitchell AP. A transcriptional cascade governs entry into meiosis in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1989;9(5):2142–52.	doi:10.1128/mcb.9.5.2142-2152.1989 PMID 2664470 PMC363008	–	2	1	1	–	2	2	Northern blot - wt vs TF mutant
968	Dolan 1989 Dolan JW, Kirkman C, Fields S. The yeast STE12 protein binds to the DNA sequence mediating pheromone induction. <i>Proc Natl Acad Sci U S A</i> 1989;86(15):5703–7.	doi:10.1073/pnas.86.15.5703 PMID 2668945 PMC297698	–	2	1	2	2	–	1	EMSA
969	Herruer 1989 Herruer MH, Mager WH, Doorenbosch TM, Wessels PL, Wassenaar TM, Planta RJ. The extended promoter of the gene encoding ribosomal protein S33 in yeast consists of multiple protein binding elements. <i>Nucleic Acids Res</i> 1989;17(18):7427–39.	doi:10.1093/nar/17.18.7427 PMID 2677998 PMC334821	–	2	1	2	1	–	1	–
970	Halfter 1989 Halfter H, Müller U, Winnacker EL, Gallwitz D. Isolation and DNA-binding characteristics of a protein involved in transcription activation of two divergently transcribed, essential yeast genes. <i>EMBO J</i> 1989;8(10):3029–37.	doi:10.1002/j.1460-2075.1989.tb08453.x PMID 2684633 PMC401380	–	2	1	2	2	–	1	DNA footprinting
971	Chambers 1989 Chambers A, Tsang JS, Stanway C, Kingsman AJ, Kingsman SM. Transcriptional control of the <i>Saccharomyces cerevisiae</i> PGK gene by RAP1. <i>Mol Cell Biol</i> 1989;9(12):5516–24.	doi:10.1128/mcb.9.12.5516-5524.1989 PMID 2685568 PMC363722	–	2	1	2	2	–	1	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
972	Siddiqui 1989 Siddiqui AH, Brandriss MC. The Saccharomyces cerevisiae PUT3 activator protein associates with proline-specific upstream activation sequences. <i>Mol Cell Biol</i> 1989;9(11):4706–12.	doi:10.1128/mcb.9.11.4706-4712.1989 PMID 2689862 PMC363617	–	2	1	2	–	–	1	–
973	Trueblood 1988 Trueblood CE, Poyton RO. Identification of REO1, a gene involved in negative regulation of COX5b and ANB1 in aerobically grown Saccharomyces cerevisiae. <i>Genetics</i> 1988;120(3):671–80.	doi:10.1093/genetics/120.3.671 PMID 2852136 PMC1203546	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
974	Hirsch 1986 Hirsch JP, Henry SA. Expression of the Saccharomyces cerevisiae inositol-1-phosphate synthase (INO1) gene is regulated by factors that affect phospholipid synthesis. <i>Mol Cell Biol</i> 1986;6(10):3320–8.	doi:10.1128/mcb.6.10.3320-3328.1986 PMID 3025587 PMC367077	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
975	Charron 1986 Charron MJ, Dubin RA, Michels CA. Structural and functional analysis of the MAL1 locus of Saccharomyces cerevisiae. <i>Mol Cell Biol</i> 1986;6(11):3891–9.	doi:10.1128/mcb.6.11.3891-3899.1986 PMID 3025617 PMC367152	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
976	Thiele 1988 Thiele DJ. ACE1 regulates expression of the Saccharomyces cerevisiae metallothionein gene. <i>Mol Cell Biol</i> 1988;8(7):2745–52.	doi:10.1128/mcb.8.7.2745-2752.1988 PMID 3043194 PMC363491	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
977	Chang 1988 Chang YS, Dubin RA, Perkins E, Forrest D, Michels CA, Needleman RB. MAL63 codes for a positive regulator of maltose fermentation in Saccharomyces cerevisiae. <i>Curr Genet</i> 1988;14(3):201–9.	doi:10.1007/BF00376740 PMID 3058330	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
978	Lowry 1988 Lowry CV, Zitomer RS. ROX1 encodes a heme-induced repression factor regulating ANB1 and CYC7 of Saccharomyces cerevisiae. <i>Mol Cell Biol</i> 1988;8(11):4651–8.	doi:10.1128/mcb.8.11.4651-4658.1988 PMID 3062365 PMC365554	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
979	Brandriss 1987 Brandriss MC. Evidence for positive regulation of the proline utilization pathway in Saccharomyces cerevisiae. <i>Genetics</i> 1987;117(3):429–35.	doi:10.1093/genetics/117.3.429 PMID 3121434 PMC1203219	–	2	1	2	–	2	2	lacZ - wt vs TF mutant
980	Ramos 1988 Ramos F, Dubois E, Piérard A. Control of enzyme synthesis in the lysine biosynthetic pathway of Saccharomyces cerevisiae. Evidence for a regulatory role of gene LYS14. <i>Eur J Biochem</i> 1988;171(1–2):171–6.	doi:10.1111/j.1432-1033.1988.tb13773.x PMID 3123231	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
981	Arndt 1987 Arndt KT, Styles C, Fink GR. Multiple global regulators control HIS4 transcription in yeast. <i>Science</i> 1987;237(4817):874–80.	doi:10.1126/science.3303332 PMID 3303332	–	2	2	2	–	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
982	Brisco 1987 Brisco PR, Cunningham TS, Kohlhaw GB. Cloning, disruption and chromosomal mapping of yeast LEU3, a putative regulatory gene. <i>Genetics</i> 1987;115(1):91–9.	doi:10.1093/genetics/115.1.91 PMID 3549446 PMC1203067	–	2	1	2	–	2	1	Enzyme activity - wt vs TF mutant
983	Giniger 1985 Giniger E, Varnum SM, Ptashne M. Specific DNA binding of GAL4, a positive regulatory protein of yeast. <i>Cell</i> 1985;40(4):767–74.	doi:10.1016/0092-8674(85)90336-8 PMID 3886158	–	2	1	2	–	–	1	–
984	Hinnebusch 1983 Hinnebusch AG, Fink GR. Positive regulation in the general amino acid control of <i>Saccharomyces cerevisiae</i> . <i>Proc Natl Acad Sci U S A</i> 1983;80(17):5374–8.	doi:10.1073/pnas.80.17.5374 PMID 6351059 PMC384258	–	2	1	2	–	–	1	–
985	West 1984 West RW Jr, Yocum RR, Ptashne M. <i>Saccharomyces cerevisiae</i> GAL1-GAL10 divergent promoter region: location and function of the upstream activating sequence UASG. <i>Mol Cell Biol</i> 1984;4(11):2467–78.	doi:10.1128/mcb.4.11.2467-2478.1984 PMID 6392852 PMC369078	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
986	Donahue 1981 Donahue TF, Henry SA. myo-Inositol-1-phosphate synthase. Characteristics of the enzyme and identification of its structural gene in yeast. <i>J Biol Chem</i> 1981;256(13):7077–85.	PMID 7016881	–	2	2	2	–	–	1	–
987	Sewell 1995 Sewell AK, Yokoya F, Yu W, Miyagawa T, Murayama T, Winge DR. Mutated yeast heat shock transcription factor exhibits elevated basal transcriptional activation and confers metal resistance. <i>J Biol Chem</i> 1995;270(42):25079–86.	doi:10.1074/jbc.270.42.25079 PMID 7559639	–	2	1	2	–	–	1	–
988	Delahodde 1995 Delahodde A, Delaveau T, Jacq C. Positive autoregulation of the yeast transcription factor Pdr3p, which is involved in control of drug resistance. <i>Mol Cell Biol</i> 1995;15(8):4043–51.	doi:10.1128/MCB.15.8.4043 PMID 7623800 PMC230643	–	2	2	2	–	2	2	Northern blot - wt vs TF mutant
989	Piatti 1995 Piatti S, Lengauer C, Nasmyth K. Cdc6 is an unstable protein whose de novo synthesis in G1 is important for the onset of S phase and for preventing a 'reductional' anaphase in the budding yeast <i>Saccharomyces cerevisiae</i> . <i>EMBO J</i> 1995;14(15):3788–99.	doi:10.1002/j.1460-2075.1995.tb00048.x PMID 7641697 PMC394453	–	2	2	2	–	1	2	–
990	Künzler 1995 Künzler M, Springer C, Braus GH. Activation and repression of the yeast ARO3 gene by global transcription factors. <i>Mol Microbiol</i> 1995;15(1):167–78.	doi:10.1111/j.1365-2958.1995.tb02231.x PMID 7752892	–	2	2	2	1	1	2	EMSA
991	Stanbrough 1995 Stanbrough M, Magasanik B. Transcriptional and posttranslational regulation of the general amino acid permease of <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1995;177(1):94–102.	doi:10.1128/jb.177.1.94-102.1995 PMID 7798155 PMC176561	–	2	2	2	–	2	2	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
992	André 1995 André B, Talibi D, Soussi Boudekou S, Hein C, Vissers S, Coornaert D. Two mutually exclusive regulatory systems inhibit UASGATA, a cluster of 5'-GAT(A/T)A-3' upstream from the UGA4 gene of <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Res</i> 1995;23(4):558–64.	doi:10.1093/nar/23.4.558 PMID 7899075 PMC306720	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
993	Graham 1994 Graham IR, Chambers A. A Reb1p-binding site is required for efficient activation of the yeast RAP1 gene, but multiple binding sites for Rap1p are not essential. <i>Mol Microbiol</i> 1994; 12(6):931–40.	doi:10.1111/j.1365-2958.1994.tb01081.x PMID 7934900	–	2	2	2	1	–	1	–
994	Tamai 1994 Tamai KT, Liu X, Silar P, Sosinowski T, Thiele DJ. Heat shock transcription factor activates yeast metallothionein gene expression in response to heat and glucose starvation via distinct signalling pathways. <i>Mol Cell Biol</i> 1994;14(12): 8155–65.	doi:10.1128/mcb.14.12.8155-8165.1994 PMID 7969152 PMC359354	–	2	1	2	–	–	1	–
995	Katzmann 1994 Katzmann DJ, Burnett PE, Golin J, Mahé Y, Moye-Rowley WS. Transcriptional control of the yeast PDR5 gene by the PDR3 gene product. <i>Mol Cell Biol</i> 1994;14(7):4653–61.	doi:10.1128/mcb.14.7.4653-4661.1994 PMID 8007969 PMC358838	–	2	2	2	1	1	2	DNA footprinting
996	Yagi 1994 Yagi S, Yagi K, Fukuoka J, Suzuki M. The UAS of the yeast GAPDH promoter consists of multiple general functional elements including RAP1 and GRF2 binding sites. <i>J Vet Med Sci</i> 1994;56(2):235–44.	doi:10.1292/jvms.56.235 PMID 8075210	–	2	2	2	1	–	1	–
997	Hasslacher 1993 Hasslacher M, Ivessa AS, Paltauf F, Kohlwein SD. Acetyl-CoA carboxylase from yeast is an essential enzyme and is regulated by factors that control phospholipid metabolism. <i>J Biol Chem</i> 1993;268(15):10946–52.	PMID 8098706	–	2	2	2	–	1	2	–
998	Nikoloff 1994 Nikoloff DM, Henry SA. Functional characterization of the INO2 gene of <i>Saccharomyces cerevisiae</i> . A positive regulator of phospholipid biosynthesis. <i>J Biol Chem</i> 1994;269(10):7402–11.	PMID 8125958	–	2	1	1	–	2	2	lacZ - wt vs TF mutant
999	Decottignies 1994 Decottignies A, Kolaczowski M, Balzi E, Goffeau A. Solubilization and characterization of the overexpressed PDR5 multidrug resistance nucleotide triphosphatase of yeast. <i>J Biol Chem</i> 1994;269(17):12797–803.	PMID 8175692	–	2	2	2	–	2	1	Western blot - wt vs TF overexpression
1000	Kovari 1993 Kovari LZ, Fourie M, Park HD, Kovari IA, Van Vuuren HJ, Cooper TG. Analysis of the inducer-responsive CAR1 upstream activation sequence (UAS1) and the factors required for its operation. <i>Yeast</i> 1993;9(8):835–45.	doi:10.1002/yea.320090804 PMID 8212891	–	2	2	2	2	–	1	lacZ - wt vs mutant promoter sequence

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1001	Jungmann 1993 Jungmann J, Reins HA, Lee J, et al., Jentsch S. MAC1, a nuclear regulatory protein related to Cu-dependent transcription factors is involved in Cu/Fe utilization and stress resistance in yeast. <i>EMBO J</i> 1993;12(13):5051–6.	doi:10.1002/j.1460-2075.1993.tb06198.x PMID 8262047 PMC413765	–	2	1	2	–	2	2	Northern blot - wt vs TF mutant
1002	Li 1993 Li Z, Brendel M. Co-regulation with genes of phospholipid biosynthesis of the CTR/HNM1-encoded choline/nitrogen mustard permease in <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1993;241(5–6):680–4.	doi:10.1007/BF00279911 PMID 8264542	–	2	2	2	1	–	1	–
1003	Stotz 1993 Stotz A, Müller PP, Linder P. Regulation of the ADE2 gene from <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 1993;24(6):472–80.	doi:10.1007/BF00351708 PMID 8299166	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1004	Estruch 1993 Estruch F, Carlson M. Two homologous zinc finger genes identified by multicopy suppression in a SNF1 protein kinase mutant of <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1993;13(7):3872–81.	doi:10.1128/mcb.13.7.3872-3881.1993 PMID 8321194 PMC359918	–	2	2	2	2	–	1	EMSA
1005	Su 1993 Su SS, Mitchell AP. Molecular characterization of the yeast meiotic regulatory gene RIM1. <i>Nucleic Acids Res</i> 1993;21(16):3789–97.	doi:10.1093/nar/21.16.3789 PMID 8367297 PMC309893	–	2	1	2	–	2	2	His3 - wt vs TF mutant
1006	Koch 1993 Koch C, Moll T, Neuberg M, Ahorn H, Nasmyth K. A role for the transcription factors Mbp1 and Swi4 in progression from G1 to S phase. <i>Science</i> 1993;261(5128):1551–7.	doi:10.1126/science.8372350 PMID 8372350	–	2	1	2	–	–	1	–
1007	André 1993 André B, Hein C, Grenson M, Jauniaux JC. Cloning and expression of the UGA4 gene coding for the inducible GABA-specific transport protein of <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1993;237(1–2):17–25.	doi:10.1007/BF00282779 PMID 8455553	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1008	Bowdish 1993 Bowdish KS, Mitchell AP. Bipartite structure of an early meiotic upstream activation sequence from <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1993;13(4):2172–81.	doi:10.1128/mcb.13.4.2172-2181.1993 PMID 8455605 PMC359538	–	2	2	2	2	–	2	lacZ - wt vs mutant promoter sequence
1009	Foster 1993 Foster R, Mikesell GE, Breeden L. Multiple SWI6-dependent cis-acting elements control SWI4 transcription through the cell cycle. <i>Mol Cell Biol</i> 1993;13(6):3792–801.	doi:10.1128/mcb.13.6.3792-3801.1993 PMID 8497280 PMC359864	–	2	2	2	1	–	2	–
1010	Ward 1995 Ward MP, Gimeno CJ, Fink GR, Garrett S. SOK2 may regulate cyclic AMP-dependent protein kinase-stimulated growth and pseudohyphal development by repressing transcription. <i>Mol Cell Biol</i> 1995;15(12):6854–63.	doi:10.1128/MCB.15.12.6854 PMID 8524252 PMC230940	–	2	1	2	–	2	2	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1011	Stanbrough 1996 Stanbrough M, Magasanik B. Two transcription factors, Gln3p and Ntl1p, use the same GATAAG sites to activate the expression of GAP1 of <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1996;178(8):2465–8.	doi:10.1128/jb.178.8.2465-2468.1996 PMID 8636059 PMC177966	–	2	2	2	–	2	2	lacZ - wt vs TF mutant
1012	Kovacech 1996 Kovacech B, Nasmyth K, Schuster T. EGT2 gene transcription is induced predominantly by Swi5 in early G1. <i>Mol Cell Biol</i> 1996;16(7):3264–74.	doi:10.1128/MCB.16.7.3264 PMID 8668141 PMC231320	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1013	Katzmann 1996 Katzmann DJ, Hallstrom TC, Mahé Y, Moye-Rowley WS. Multiple Pdr1p/Pdr3p binding sites are essential for normal expression of the ATP binding cassette transporter protein-encoding gene PDR5. <i>J Biol Chem</i> 1996;271(38):23049–54.	doi:10.1074/jbc.271.38.23049 PMID 8798494	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1014	Miyahara 1996 Miyahara K, Hirata D, Miyakawa T. yAP-1- and yAP-2-mediated, heat shock-induced transcriptional activation of the multidrug resistance ABC transporter genes in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 1996;29(2):103–5.	doi:10.1007/BF02221572 PMID 8821655	–	2	1	2	2	–	1	lacZ - wt vs mutant promoter sequence
1015	Mahé 1996 Mahé Y, Parle-McDermott A, Nourani A, Delahodde A, Lamprecht A, Kuchler K. The ATP-binding cassette multidrug transporter Snq2 of <i>Saccharomyces cerevisiae</i> : a novel target for the transcription factors Pdr1 and Pdr3. <i>Mol Microbiol</i> 1996;20(1):109–17.	doi:10.1111/j.1365-2958.1996.tb02493.x PMID 8861209	–	2	2	2	–	–	1	–
1016	Liesen 1996 Liesen T, Hollenberg CP, Heinisch JJ. ERA, a novel cis-acting element required for autoregulation and ethanol repression of PDC1 transcription in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 1996;21(3):621–32.	doi:10.1111/j.1365-2958.1996.tb02570.x PMID 8866484	–	2	2	2	1	1	2	DNA footprinting
1017	Cox 1996 Cox JS, Walter P. A novel mechanism for regulating activity of a transcription factor that controls the unfolded protein response. <i>Cell</i> 1996;87(3):391–404.	doi:10.1016/s0092-8674(00)81360-4 PMID 8898193	–	2	1	2	–	1	2	–
1018	Jia 1997 Jia Y, Rothermel B, Thornton J, Butow RA. A basic helix-loop-helix-leucine zipper transcription complex in yeast functions in a signaling pathway from mitochondria to the nucleus. <i>Mol Cell Biol</i> 1997;17(3):1110–7.	doi:10.1128/MCB.17.3.1110 PMID 9032238 PMC231836	–	2	2	2	1	1	2	EMSA
1019	Madhani 1997 Madhani HD, Fink GR. Combinatorial control required for the specificity of yeast MAPK signaling. <i>Science</i> 1997;275(5304):1314–7.	doi:10.1126/science.275.5304.1314 PMID 9036858	–	2	2	2	2	–	1	DNA footprinting

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1020	Naik 1997 Naik RR, Nebes V, Jones EW. Regulation of the proteinase B structural gene PRB1 in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1997;179(5):1469–74.	doi:10.1128/jb.179.5.1469-1474.1997 PMID 9045801 PMC178854	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1021	Jorgensen 1997 Jorgensen MU, Gjermansen C, Andersen HA, Kielland-Brandt MC. STP1, a gene involved in pre-tRNA processing in yeast, is important for amino-acid uptake and transcription of the permease gene BAP2. <i>Curr Genet</i> 1997;31(3):241–7.	doi:10.1007/s002940050201 PMID 9065387	–	2	1	2	–	2	2	lacZ - wt vs TF mutant
1022	Partridge 1997 Partridge JF, Mikesell GE, Breeden LL. Cell cycle-dependent transcription of CLN1 involves swi4 binding to MCB-like elements. <i>J Biol Chem</i> 1997;272(14):9071–7.	doi:10.1074/jbc.272.14.9071 PMID 9083033	–	2	1	2	1	1	1	EMSA
1023	Lin 1997 Lin SJ, Pufahl RA, Dancis A, O'Halloran TV, Culotta VC. A role for the <i>Saccharomyces cerevisiae</i> ATX1 gene in copper trafficking and iron transport. <i>J Biol Chem</i> 1997;272(14):9215–20.	PMID 9083054	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
1024	Randez-Gil 1997 Randez-Gil F, Bojunga N, Proft M, Entian KD. Glucose derepression of gluconeogenic enzymes in <i>Saccharomyces cerevisiae</i> correlates with phosphorylation of the gene activator Cat8p. <i>Mol Cell Biol</i> 1997;17(5):2502–10.	doi:10.1128/MCB.17.5.2502 PMID 9111319 PMC232099	–	2	2	2	1	1	2	Northern blot - wt vs TF mutant
1025	Coleman 1997 Coleman ST, Tseng E, Moye-Rowley WS. <i>Saccharomyces cerevisiae</i> basic region-leucine zipper protein regulatory networks converge at the ATR1 structural gene. <i>J Biol Chem</i> 1997;272(37):23224–30.	doi:10.1074/jbc.272.37.23224 PMID 9287330	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1026	Zarzov 1997 Zarzov P, Boucherie H, Mann C. A yeast heat shock transcription factor (Hsf1) mutant is defective in both Hsc82/Hsp82 synthesis and spindle pole body duplication. <i>J Cell Sci</i> 1997;110 (Pt 16):1879–91.	doi:10.1242/jcs.110.16.1879 PMID 9296388	–	2	1	2	–	2	1	Proteomics - wt vs TF mutant
1027	Dagkessamanskaya 1997 Dagkessamanskaya A, Ter-Avanesyan M, Mager WH. Transcriptional regulation of SUP35 and SUP45 in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 1997;13(13):1265–74.	<a href="https://doi.org/10.1002/(SICI)1097-0061(199710)13:13<1265::AID-YEA175>3.0.CO;2-#">doi:10.1002/(SICI)1097-0061(199710)13:13<1265::AID-YEA175>3.0.CO;2-# PMID 9364750	–	2	1	2	2	–	1	Northern blot - wt vs mutant promoter sequence
1028	Inoue 1998 Inoue Y, Tsujimoto Y, Kimura A. Expression of the glyoxalase I gene of <i>Saccharomyces cerevisiae</i> is regulated by high osmolarity glycerol mitogen-activated protein kinase pathway in osmotic stress response. <i>J Biol Chem</i> 1998;273(5):2977–83.	doi:10.1074/jbc.273.5.2977 PMID 9446611	–	2	2	2	–	2	1	CAT - wt vs TF mutant
1029	Hallstrom 1998 Hallstrom TC, Katzmman DJ, Torres RJ, Sharp WJ, Moye-Rowley WS. Regulation of transcription factor Pdr1p function by an Hsp70 protein in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1998;18(3):1147–55.	doi:10.1128/MCB.18.3.1147 PMID 9488429 PMC108827	–	2	1	2	–	1	2	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1030	Sagee 1998 Sagee S, Sherman A, Shenhar G, et al., Kassir Y. Multiple and distinct activation and repression sequences mediate the regulated transcription of IME1, a transcriptional activator of meiosis-specific genes in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1998;18(4):1985–95.	doi:10.1128/MCB.18.4.1985 PMID 9528770 PMC121428	–	2	2	2	2	–	1	EMSA
1031	Mendizabal 1998 Mendizabal I, Rios G, Mulet JM, Serrano R, de Larrinoa IF. Yeast putative transcription factors involved in salt tolerance. <i>FEBS Lett</i> 1998;425(2):323–8.	doi:10.1016/s0014-5793(98)00249-x PMID 9559673	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1032	Shen 1998 Shen H, Dowhan W. Regulation of phosphatidylglycerophosphate synthase levels in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 1998;273(19):11638–42.	doi:10.1074/jbc.273.19.11638 PMID 9565583	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1033	Barbaric 1998 Barbaric S, Münsterkötter M, Goding C, Hörz W. Cooperative Pho2-Pho4 interactions at the PHO5 promoter are critical for binding of Pho4 to UASp1 and for efficient transactivation by Pho4 at UASp2. <i>Mol Cell Biol</i> 1998;18(5):2629–39.	doi:10.1128/MCB.18.5.2629 PMID 9566882 PMC110642	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1034	Noël 1998 Noël J, Turcotte B. Zinc cluster proteins Leu3p and Uga3p recognize highly related but distinct DNA targets. <i>J Biol Chem</i> 1998;273(28):17463–8.	doi:10.1074/jbc.273.28.17463 PMID 9651335	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1035	Oehlen 1998 Oehlen L, Cross FR. The mating factor response pathway regulates transcription of TEC1, a gene involved in pseudohyphal differentiation of <i>Saccharomyces cerevisiae</i> . <i>FEBS Lett</i> 1998;429(1):83–8.	doi:10.1016/s0014-5793(98)00568-7 PMID 9657388	–	2	1	2	–	–	1	–
1036	Valenzuela 1998 Valenzuela L, Ballario P, Aranda C, Filetici P, González A. Regulation of expression of GLT1, the gene encoding glutamate synthase in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1998;180(14):3533–40.	doi:10.1128/JB.180.14.3533-3540.1998 PMID 9657994 PMC107319	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1037	Becker 1998 Becker B, Feller A, el Alami M, Dubois E, Piérard A. A nonameric core sequence is required upstream of the LYS genes of <i>Saccharomyces cerevisiae</i> for Lys14p-mediated activation and apparent repression by lysine. <i>Mol Microbiol</i> 1998;29(1):151–63.	doi:10.1046/j.1365-2958.1998.00916.x PMID 9701810	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
1038	Pierce 1998 Pierce M, Wagner M, Xie J, et al., Winter E. Transcriptional regulation of the SMK1 mitogen-activated protein kinase gene during meiotic development in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1998;18(10):5970–80.	doi:10.1128/MCB.18.10.5970 PMID 9742114 PMC109183	–	2	1	2	2	–	2	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1039	Reagan 1998 Reagan MS, Majors JE. The chromatin structure of the GAL1 promoter forms independently of Reb1p in <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1998;259(2):142–9.	doi:10.1007/s004380050799 PMID 9747705	–	2	1	2	2	–	1	DNA footprinting
1040	Blaiseau 1998 Blaiseau PL, Thomas D. Multiple transcriptional activation complexes tether the yeast activator Met4 to DNA. <i>EMBO J</i> 1998;17(21):6327–36.	doi:10.1093/emboj/17.21.6327 PMID 9799240 PMC1170957	–	2	2	2	2	–	1	EMSA
1041	De Boer 1998 De Boer M, Bebelman JP, Gonçalves PM, Maat J, Van Heerikhuizen H, Planta RJ. Regulation of expression of the amino acid transporter gene BAP3 in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 1998;30(3):603–13.	doi:10.1046/j.1365-2958.1998.01094.x PMID 9822825	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1042	Lutfiyya 1998 Lutfiyya LL, Iyer VR, DeRisi J, DeVit MJ, Brown PO, Johnston M. Characterization of three related glucose repressors and genes they regulate in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 1998;150(4):1377–91.	doi:10.1093/genetics/150.4.1377 PMID 9832517 PMC1460414	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1043	Bogengruber 1998 Bogengruber E, Eichberger T, Briza P, Dawes IW, Breitenbach M, Schriker R. Sporulation-specific expression of the yeast DIT1/DIT2 promoter is controlled by a newly identified repressor element and the short form of Rim101p. <i>Eur J Biochem</i> 1998;258(2):430–6.	doi:10.1046/j.1432-1327.1998.2580430.x PMID 9874208	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
1044	Zhang 1999 Zhang S, Skalsky Y, Garfinkel DJ. MGA2 or SPT23 is required for transcription of the delta9 fatty acid desaturase gene, OLE1, and nuclear membrane integrity in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 1999;151(2):473–83.	doi:10.1093/genetics/151.2.473 PMID 9927444 PMC1460504	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1045	Frolova 1999 Frolova E, Johnston M, Majors J. Binding of the glucose-dependent Mig1p repressor to the GAL1 and GAL4 promoters in vivo: regulation by glucose and chromatin structure. <i>Nucleic Acids Res</i> 1999;27(5):1350–8.	doi:10.1093/nar/27.5.1350 PMID 9973625 PMC148323	–	2	1	2	2	–	1	DNA footprinting
1046	Lee 1999 Lee J, Spector D, Godon C, Labarre J, Toledano MB. A new antioxidant with alkyl hydroperoxide defense properties in yeast. <i>J Biol Chem</i> 1999;274(8):4537–44.	doi:10.1074/jbc.274.8.4537 PMID 9988687	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1047	Simon 1999 Simon JR, Treger JM, McEntee K. Multiple independent regulatory pathways control UBI4 expression after heat shock in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 1999;31(3):823–32.	doi:10.1046/j.1365-2958.1999.01220.x PMID 10048026	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1048	Dudley 1999 Dudley AM, Gansheroff LJ, Winston F. Specific components of the SAGA complex are required for Gcn4- and Gcr1-mediated activation of the his4-912delta promoter in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 1999;151(4):1365–78.	doi:10.1093/genetics/151.4.1365 PMID 10101163 PMC1460567	–	2	2	2	2	–	1	lacZ - wt vs mutant promoter sequence

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1049	Cosma 1999 Cosma MP, Tanaka T, Nasmyth K. Ordered recruitment of transcription and chromatin remodeling factors to a cell cycle- and developmentally regulated promoter. <i>Cell</i> 1999;97(3):299–311.	doi:10.1016/S0092-8674(00)80740-0 PMID 10319811	–	2	1	2	2	–	1	ChIP
1050	Stroobants 1999 Stroobants AK, Hetteema EH, van den Berg M, Tabak HF. Enlargement of the endoplasmic reticulum membrane in <i>Saccharomyces cerevisiae</i> is not necessarily linked to the unfolded protein response via Ire1p. <i>FEBS Lett</i> 1999;453(1–2):210–4.	doi:10.1016/S0014-5793(99)00721-8 PMID 10403405	–	2	2	2	2	–	1	EMSA
1051	Fujiwara 1999 Fujiwara D, Kobayashi O, Yoshimoto H, Harashima S, Tamai Y. Molecular mechanism of the multiple regulation of the <i>Saccharomyces cerevisiae</i> ATF1 gene encoding alcohol acetyltransferase. <i>Yeast</i> 1999;15(12):1183–97.	<a href="https://doi.org/10.1002/(SICI)1097-0061(19990915)15:12<1183::AID-YEA444>3.0.CO;2-J">doi:10.1002/(SICI)1097-0061(19990915)15:12<1183::AID-YEA444>3.0.CO;2-J PMID 10487921	–	2	2	2	1	1	2	EMSA
1052	Rai 1999 Rai R, Daugherty JR, Cunningham TS, Cooper TG. Overlapping positive and negative GATA factor binding sites mediate inducible DAL7 gene expression in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 1999;274(39):28026–34.	doi:10.1074/jbc.274.39.28026 PMID 10488154	–	2	2	2	1	1	2	EMSA
1053	Chelstowska 1999 Chelstowska A, Liu Z, Jia Y, Amberg D, Butow RA. Signalling between mitochondria and the nucleus regulates the expression of a new D-lactate dehydrogenase activity in yeast. <i>Yeast</i> 1999;15(13):1377–91.	<a href="https://doi.org/10.1002/(SICI)1097-0061(19990930)15:13<1377::AID-YEA473>3.0.CO;2-0">doi:10.1002/(SICI)1097-0061(19990930)15:13<1377::AID-YEA473>3.0.CO;2-0 PMID 10509019	–	2	2	2	1	–	1	–
1054	Jang 1999 Jang YK, Wang L, Sancar GB. RPH1 and GIS1 are damage-responsive repressors of PHR1. <i>Mol Cell Biol</i> 1999;19(11):7630–8.	doi:10.1128/MCB.19.11.7630 PMID 10523651 PMC84794	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1055	Zhong 1999 Zhong H, McCord R, Vershon AK. Identification of target sites of the alpha2-Mcm1 repressor complex in the yeast genome. <i>Genome Res</i> 1999;9(11):1040–7.	doi:10.1101/gr.9.11.1040 PMID 10568744	–	2	1	2	2	–	1	EMSA
1056	Brôco 1999 Brôco N, Tenreiro S, Viegas CA, Sá-Correia I. FLR1 gene (ORF YBR008c) is required for benomyl and methotrexate resistance in <i>Saccharomyces cerevisiae</i> and its benomyl-induced expression is dependent on pdr3 transcriptional regulator. <i>Yeast</i> 1999;15(15):1595–608.	<a href="https://doi.org/10.1002/(SICI)1097-0061(199911)15:15<1595::AID-YEA484>3.0.CO;2-6">doi:10.1002/(SICI)1097-0061(199911)15:15<1595::AID-YEA484>3.0.CO;2-6 PMID 10572257	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1057	Delneri 1999 Delneri D, Gardner DC, Oliver SG. Analysis of the seven-member AAD gene set demonstrates that genetic redundancy in yeast may be more apparent than real. <i>Genetics</i> 1999;153(4):1591–600.	doi:10.1093/genetics/153.4.1591 PMID 10581269 PMC1460870	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1058	Garland 1999 Garland SA, Hoff K, Vickery LE, Culotta VC. Saccharomyces cerevisiae ISU1 and ISU2: members of a well-conserved gene family for iron-sulfur cluster assembly. <i>J Mol Biol</i> 1999;294(4): 897–907.	doi:10.1006/jmbi.1999.3294 PMID 10588895	–	2	1	2	–	2	2	Northern blot - wt vs TF overexpression
1059	Gurvitz 1999 Gurvitz A, Hamilton B, Hartig A, Ruis H, Dawes IW, Rottensteiner H. A novel element in the promoter of the Saccharomyces cerevisiae gene SPS19 enhances ORE-dependent up-regulation in oleic acid and is essential for de-repression. <i>Mol Gen Genet</i> 1999;262(3):481–92.	doi:10.1007/s004380051109 PMID 10589836	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1060	Huang 2000 Huang HL, Brandriss MC. The regulator of the yeast proline utilization pathway is differentially phosphorylated in response to the quality of the nitrogen source. <i>Mol Cell Biol</i> 2000;20(3):892–9.	doi:10.1128/MCB.20.3.892-899.2000 PMID 10629046 PMC85206	–	2	1	2	–	1	2	–
1061	Hollenhorst 2000 Hollenhorst PC, Bose ME, Mielke MR, Müller U, Fox CA. Forkhead genes in transcriptional silencing, cell morphology and the cell cycle. Overlapping and distinct functions for FKH1 and FKH2 in Saccharomyces cerevisiae. <i>Genetics</i> 2000;154(4):1533–48.	doi:10.1093/genetics/154.4.1533 PMID 10747051 PMC1461039	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1062	Munsterkötter 2000 Munsterkötter M, Barbaric S, Hörz W. Transcriptional regulation of the yeast PHO8 promoter in comparison to the coregulated PHO5 promoter. <i>J Biol Chem</i> 2000;275(30):22678–85.	doi:10.1074/jbc.M001409200 PMID 10801809	–	2	2	2	1	1	1	lacZ - wt vs TF mutant
1063	Messenguy 2000 Messenguy F, Vierendeels F, Scherens B, Dubois E. In Saccharomyces cerevisiae, expression of arginine catabolic genes CAR1 and CAR2 in response to exogenous nitrogen availability is mediated by the Ume6 (CargRI)-Sin3 (CargRII)-Rpd3 (CargRIII) complex. <i>J Bacteriol</i> 2000;182(11):3158–64.	doi:10.1128/JB.182.11.3158-3164.2000 PMID 10809695 PMC94502	–	2	1	2	–	2	1	Enzyme activity - wt vs TF mutant
1064	Sugiyama 2000 Sugiyama K, Izawa S, Inoue Y. The Yap1p-dependent induction of glutathione synthesis in heat shock response of Saccharomyces cerevisiae. <i>J Biol Chem</i> 2000;275(20):15535–40.	doi:10.1074/jbc.275.20.15535 PMID 10809786	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
1065	Garcia-Gimeno 2000 Garcia-Gimeno MA, Struhl K. Aca1 and Aca2, ATF/CREB activators in Saccharomyces cerevisiae, are important for carbon source utilization but not the response to stress. <i>Mol Cell Biol</i> 2000;20(12):4340–9.	doi:10.1128/MCB.20.12.4340-4349.2000 PMID 10825197 PMC85801	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1066	Jungwirth 2000 Jungwirth H, Wendler F, Platzer B, Bergler H, Högenauer G. Diazaborine resistance in yeast involves the efflux pumps Ycf1p and Flr1p and is enhanced by a gain-of-function allele of gene YAP1. <i>Eur J Biochem</i> 2000;267(15):4809–16.	doi:10.1046/j.1432-1327.2000.01537.x PMID 10903515	–	2	1	2	–	–	1	–
1067	Elkhaïmi 2000 Elkhaïmi M, Kaadige MR, Kamath D, Jackson JC, Biliran H Jr, Lopes JM. Combinatorial regulation of phospholipid biosynthetic gene expression by the UME6, SIN3 and RPD3 genes. <i>Nucleic Acids Res</i> 2000;28(16):3160–7.	doi:10.1093/nar/28.16.3160 PMID 10931932 PMC108424	–	2	1	2	–	2	2	Northern blot - wt vs TF mutant
1068	Kaiser 2000 Kaiser P, Flick K, Wittenberg C, Reed SI. Regulation of transcription by ubiquitination without proteolysis: Cdc34/SCF(Met30)-mediated inactivation of the transcription factor Met4. <i>Cell</i> 2000;102(3):303–14.	doi:10.1016/s0092-8674(00)00036-2 PMID 10975521	–	2	2	2	1	–	1	–
1069	Kastaniotis 2000 Kastaniotis AJ, Mennella TA, Konrad C, Torres AM, Zitomer RS. Roles of transcription factor Mot3 and chromatin in repression of the hypoxic gene ANB1 in yeast. <i>Mol Cell Biol</i> 2000;20(19):7088–98.	doi:10.1128/MCB.20.19.7088-7098.2000 PMID 10982825 PMC86251	–	2	2	2	1	–	2	–
1070	Delaunay 2000 Delaunay A, Isnard AD, Toledano MB. H2O2 sensing through oxidation of the Yap1 transcription factor. <i>EMBO J</i> 2000; 19(19):5157–66.	doi:10.1093/emboj/19.19.5157 PMID 11013218 PMC302088	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1071	Lindgren 2000 Lindgren A, Bungard D, Pierce M, Xie J, Vershon A, Winter E. The pachytene checkpoint in <i>Saccharomyces cerevisiae</i> requires the Sum1 transcriptional repressor. <i>EMBO J</i> 2000; 19(23):6489–97.	doi:10.1093/emboj/19.23.6489 PMID 11101521 PMC305847	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
1072	Sokolíková 2000 Sokolíková B, Sabová L, Kissová I, Kolarov J. A carbon-source-responsive element is required for regulation of the hypoxic ADP/ATP carrier (AAC3) isoform in <i>Saccharomyces cerevisiae</i> . <i>Biochem J</i> 2000;352 Pt 3(Pt 3): 893–8.	PMID 11104700 PMC1221531	–	2	2	2	1	1	2	EMSA
1073	Han 2001 Han GS, Johnston CN, Chen X, Athenstaedt K, Daum G, Carman GM. Regulation of the <i>Saccharomyces cerevisiae</i> DPP1-encoded diacylglycerol pyrophosphate phosphatase by zinc. <i>J Biol Chem</i> 2001;276(13):10126–33.	doi:10.1074/jbc.M011421200 PMID 11139591	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
1074	Moreno-Herrero 2001 Moreno-Herrero F, Herrero P, Colchero J, Baró AM, Moreno F. Imaging and mapping protein-binding sites on DNA regulatory regions with atomic force microscopy. <i>Biochem Biophys Res Commun</i> 2001;280(1):151–7.	doi:10.1006/bbrc.2000.4074 PMID 11162492	–	2	2	2	2	–	2	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1075	Tenreiro 2001 Tenreiro S, Fernandes AR, Sá-Correia I. Transcriptional activation of FLR1 gene during <i>Saccharomyces cerevisiae</i> adaptation to growth with benomyl: role of Yap1p and Pdr3p. <i>Biochem Biophys Res Commun</i> 2001;280(1):216–22.	doi:10.1006/bbrc.2000.4100 PMID 11162502	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1076	Roth 2001 Roth S, Schüller HJ. Cat8 and Sip4 mediate regulated transcriptional activation of the yeast malate dehydrogenase gene MDH2 by three carbon source-responsive promoter elements. <i>Yeast</i> 2001;18(2):151–62.	<a href="https://doi.org/10.1002/1097-0061(20010130)18:2<151::AID-YEA662>3.0.CO;2-Q">doi:10.1002/1097-0061(20010130)18:2<151::AID-YEA662>3.0.CO;2-Q PMID 11169757	–	2	2	2	1	–	1	–
1077	Simonics 2000 Simonics T, Kozovska Z, Michalkova-Papajova D, Delahodde A, Jacq C, Subik J. Isolation and molecular characterization of the carboxy-terminal pdr3 mutants in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 2000;38(5):248–55.	doi:10.1007/s002940000164 PMID 11191208	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
1078	Nielsen 2001 Nielsen PS, van den Hazel B, Didion T, et al., Andersen HA. Transcriptional regulation of the <i>Saccharomyces cerevisiae</i> amino acid permease gene BAP2. <i>Mol Gen Genet</i> 2001;264(5):613–22.	doi:10.1007/s004380000347 PMID 11212916	–	2	2	2	2	–	1	EMSA
1079	Valenzuela 2001 Valenzuela L, Aranda C, González A. TOR modulates GCN4-dependent expression of genes turned on by nitrogen limitation. <i>J Bacteriol</i> 2001;183(7):2331–4.	doi:10.1128/JB.183.7.2331-2334.2001 PMID 11244074 PMC95141	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
1080	Xie 2001 Xie Y, Varshavsky A. RPN4 is a ligand, substrate, and transcriptional regulator of the 26S proteasome: a negative feedback circuit. <i>Proc Natl Acad Sci U S A</i> 2001;98(6):3056–61.	doi:10.1073/pnas.071022298 PMID 11248031 PMC30606	–	2	1	2	–	2	1	Western blot - wt vs TF overexpression
1081	Régnacq 2001 Régnacq M, Alimardani P, El Moudni B, Bergès T. SUT1p interaction with Cyc8p(Ssn6p) relieves hypoxic genes from Cyc8p-Tup1p repression in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2001;40(5):1085–96.	doi:10.1046/j.1365-2958.2001.02450.x PMID 11401714	–	2	1	2	–	2	1	Northern blot - wt vs TF overexpression
1082	Gurvitz 2001 Gurvitz A, Hiltunen JK, Erdmann R, et al., Rottensteiner H. <i>Saccharomyces cerevisiae</i> Adr1p governs fatty acid beta-oxidation and peroxisome proliferation by regulating POX1 and PEX11. <i>J Biol Chem</i> 2001;276(34):31825–30.	doi:10.1074/jbc.M105989200 PMID 11431484	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
1083	Williams 2001 Williams KE, Cyert MS. The eukaryotic response regulator Skn7p regulates calcineurin signaling through stabilization of Crz1p. <i>EMBO J</i> 2001;20(13):3473–83.	doi:10.1093/emboj/20.13.3473 PMID 11432834 PMC125528	–	2	1	2	–	2	1	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1084	Crespo 2001 Crespo JL, Daicho K, Ushimaru T, Hall MN. The GATA transcription factors GLN3 and GAT1 link TOR to salt stress in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2001;276(37):34441–4.	doi:10.1074/jbc.M103601200 PMID 11457832	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1085	Walther 2001 Walther K, Schüller HJ. Adr1 and Cat8 synergistically activate the glucose-regulated alcohol dehydrogenase gene ADH2 of the yeast <i>Saccharomyces cerevisiae</i> . <i>Microbiology (Reading)</i> 2001;147(Pt 8):2037–2044.	doi:10.1099/00221287-147-8-2037 PMID 11495982	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1086	Chellappa 2001 Chellappa R, Kandasamy P, Oh CS, Jiang Y, Vemula M, Martin CE. The membrane proteins, Spt23p and Mga2p, play distinct roles in the activation of <i>Saccharomyces cerevisiae</i> OLE1 gene expression. Fatty acid-mediated regulation of Mga2p activity is independent of its proteolytic processing into a soluble transcription activator. <i>J Biol Chem</i> 2001;276(47):43548–56.	doi:10.1074/jbc.M107845200 PMID 11557770	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1087	Zhang 2001 Zhang X, Moye-Rowley WS. <i>Saccharomyces cerevisiae</i> multidrug resistance gene expression inversely correlates with the status of the F(0) component of the mitochondrial ATPase. <i>J Biol Chem</i> 2001;276(51):47844–52.	doi:10.1074/jbc.M106285200 PMID 11602584	–	2	2	2	–	–	1	–
1088	Wong 2002 Wong CM, Zhou Y, Ng RW, Kung Hf HF, Jin DY. Cooperation of yeast peroxiredoxins Tsa1p and Tsa2p in the cellular defense against oxidative and nitrosative stress. <i>J Biol Chem</i> 2002;277(7):5385–94.	doi:10.1074/jbc.M106846200 PMID 11741925	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
1089	Westwater 2002 Westwater J, McLaren NF, Dormer UH, Jamieson DJ. The adaptive response of <i>Saccharomyces cerevisiae</i> to mercury exposure. <i>Yeast</i> 2002;19(3):233–9.	doi:10.1002/yea.835 PMID 11816031	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1090	Portnoy 2002 Portnoy ME, Jensen LT, Culotta VC. The distinct methods by which manganese and iron regulate the Nramp transporters in yeast. <i>Biochem J</i> 2002;362(Pt 1):119–24.	doi:10.1042/0264-6021:3620119 PMID 11829747 PMC1222367	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1091	Newcomb 2002 Newcomb LL, Hall DD, Heideman W. AZF1 is a glucose-dependent positive regulator of CLN3 transcription in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2002;22(5):1607–14.	doi:10.1128/MCB.22.5.1607-1614.2002 PMID 11839825 PMC134705	–	2	1	2	–	2	2	Northern blot - wt vs TF mutant
1092	Owsianik 2002 Owsianik G, Balzi l L, Ghislain M. Control of 26S proteasome expression by transcription factors regulating multidrug resistance in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2002;43(5):1295–308.	doi:10.1046/j.1365-2958.2002.02823.x PMID 11918814	–	2	2	2	–	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1093	Kryndushkin 2002 Kryndushkin DS, Smirnov VN, Ter-Avanesyan MD, Kushnirov VV. Increased expression of Hsp40 chaperones, transcriptional factors, and ribosomal protein Rpp0 can cure yeast prions. <i>J Biol Chem</i> 2002;277(26):23702–8.	doi:10.1074/jbc.M111547200 PMID 11923285	–	2	1	2	–	2	1	lacZ - wt vs TF overexpression
1094	Cohen 2002 Cohen BA, Pilpel Y, Mitra RD, Church GM. Discrimination between paralogs using microarray analysis: application to the Yap1p and Yap2p transcriptional networks. <i>Mol Biol Cell</i> 2002;13(5):1608–14.	doi:10.1091/mbc.01-10-0472 PMID 12006656 PMC111130	no GEO	2	2	2	–	2	2	lacZ - wt vs TF mutant
1095	Terrell 2002 Terrell AR, Wongwisansri S, Pilon JL, Laybourn PJ. Reconstitution of nucleosome positioning, remodeling, histone acetylation, and transcriptional activation on the PHO5 promoter. <i>J Biol Chem</i> 2002;277(34):31038–47.	doi:10.1074/jbc.M204662200 PMID 12060664	–	2	2	2	2	–	1	EMSA
1096	Xia 1999 Xia ZX, Ao SZ. PHO4 and PHO2 Protein Interact with Upstream Sequence of PHO81 Gene. <i>Sheng Wu Hua Xue Yu Sheng Wu Li Xue Bao (Shanghai)</i> 1999;31(2):191–196.	PMID 12136214	–	2	2	2	2	–	1	EMSA
1097	Idicula 2002 Idicula AM, Blatch GL, Cooper TG, Dorrington RA. Binding and activation by the zinc cluster transcription factors of <i>Saccharomyces cerevisiae</i> . Redefining the UASGABA and its interaction with Uga3p. <i>J Biol Chem</i> 2002;277(48):45977–83.	doi:10.1074/jbc.M201789200 PMID 12235130 PMC4384467	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
1098	Sharma 2002 Sharma KG, Mason DL, Liu G, Rea PA, Bachhawat AK, Michaelis S. Localization, regulation, and substrate transport properties of Bpt1p, a <i>Saccharomyces cerevisiae</i> MRP-type ABC transporter. <i>Eukaryot Cell</i> 2002;1(3):391–400.	doi:10.1128/EC.1.3.391-400.2002 PMID 12455987 PMC118024	–	2	1	2	–	2	1	lacZ - wt vs TF overexpression
1099	de Nadal 2003 de Nadal E, Casadomé L, Posas F. Targeting the MEF2-like transcription factor Smp1 by the stress-activated Hog1 mitogen-activated protein kinase. <i>Mol Cell Biol</i> 2003;23(1):229–37.	doi:10.1128/MCB.23.1.229-237.2003 PMID 12482976 PMC140668	–	2	1	2	–	2	2	lacZ - wt vs TF mutant
1100	Gagiano 2003 Gagiano M, Bester M, van Dyk D, Franken J, Bauer FF, Pretorius IS. Mss11p is a transcription factor regulating pseudohyphal differentiation, invasive growth and starch metabolism in <i>Saccharomyces cerevisiae</i> in response to nutrient availability. <i>Mol Microbiol</i> 2003;47(1):119–34.	doi:10.1046/j.1365-2958.2003.03247.x PMID 12492858	–	2	1	2	–	–	1	–
1101	Angermayr 2003 Angermayr M, Bandlow W. Permanent nucleosome exclusion from the Gal4p-inducible yeast GCY1 promoter. <i>J Biol Chem</i> 2003;278(13):11026–31.	doi:10.1074/jbc.M210932200 PMID 12536147	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1102	Ter Linde 2003 Ter Linde JJ, Régnacq M, Steensma HY. Transcriptional regulation of YML083c under aerobic and anaerobic conditions. <i>Yeast</i> 2003;20(5):439–54.	doi:10.1002/yea.975 PMID 12673627	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1103	Rottensteiner 2003 Rottensteiner H, Hartig A, Hamilton B, Ruis H, Erdmann R, Gurvitz A. Saccharomyces cerevisiae Pip2p-Oaf1p regulates PEX25 transcription through an adenine-less ORE. <i>Eur J Biochem</i> 2003;270(9):2013–22.	doi:10.1046/j.1432-1033.2003.03575.x PMID 12709061	–	2	2	2	2	–	1	EMSA
1104	Rottensteiner 2003 Rottensteiner H, Wabnegger L, Erdmann R, et al., Gurvitz A. Saccharomyces cerevisiae PIP2 mediating oleic acid induction and peroxisome proliferation is regulated by Adr1p and Pip2p-Oaf1p. <i>J Biol Chem</i> 2003;278(30):27605–11.	doi:10.1074/jbc.M304097200 PMID 12748191	–	2	2	2	1	1	2	EMSA
1105	Neef 2003 Neef DW, Kladde MP. Polyphosphate loss promotes SNF/SWI- and Gcn5-dependent mitotic induction of PHO5. <i>Mol Cell Biol</i> 2003;23(11):3788–97.	doi:10.1128/MCB.23.11.3788-3797.2003 PMID 12748282 PMC155216	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1106	Oshiro 2003 Oshiro J, Han GS, Iwanyshyn WM, Conover K, Carman GM. Regulation of the yeast DPP1-encoded diacylglycerol pyrophosphate phosphatase by transcription factor Gis1p. <i>J Biol Chem</i> 2003;278(34):31495–503.	doi:10.1074/jbc.M305452200 PMID 12799368	–	2	2	2	–	2	2	Enzyme activity - wt vs TF mutant
1107	Shimizu 2003 Shimizu M, Takahashi K, Lamb TM, Shindo H, Mitchell AP. Yeast Ume6p repressor permits activator binding but restricts TBP binding at the HOP1 promoter. <i>Nucleic Acids Res</i> 2003;31(12):3033–7.	doi:10.1093/nar/gkg425 PMID 12799429 PMC162329	–	2	2	2	1	1	2	DNA footprinting
1108	Hahn 2004 Hahn JS, Thiele DJ. Activation of the Saccharomyces cerevisiae heat shock transcription factor under glucose starvation conditions by Snf1 protein kinase. <i>J Biol Chem</i> 2004;279(7):5169–76.	doi:10.1074/jbc.M311005200 PMID 14612437	–	2	1	2	–	2	2	Northern blot - wt vs TF mutant
1109	Springer 2003 Springer M, Wykoff DD, Miller N, O'Shea EK. Partially phosphorylated Pho4 activates transcription of a subset of phosphate-responsive genes. <i>PLoS Biol</i> 2003;1(2):E28.	doi:10.1371/journal.pbio.0000028 PMID 14624238 PMC261874	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
1110	Higgins 2003 Higgins VJ, Rogers PJ, Dawes IW. Application of genome-wide expression analysis to identify molecular markers useful in monitoring industrial fermentations. <i>Appl Environ Microbiol</i> 2003;69(12):7535–40.	doi:10.1128/AEM.69.12.7535-7540.2003 PMID 14660410 PMC310037	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1111	van Dyk 2003 van Dyk D, Hansson G, Pretorius IS, Bauer FF. Cellular differentiation in response to nutrient availability: The repressor of meiosis, Rme1p, positively regulates invasive growth in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2003;165(3):1045–58.	doi:10.1093/genetics/165.3.1045 PMID 14668363 PMC1462853	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
1112	Geng 2004 Geng F, Laurent BC. Roles of SWI/SNF and HATs throughout the dynamic transcription of a yeast glucose-repressible gene. <i>EMBO J</i> 2004;23(1):127–37.	doi:10.1038/sj.emboj.7600035 PMID 14685262 PMC1271673	–	2	2	2	–	2	1	RT-PCR - wt vs TF mutant
1113	Bird 2004 Bird AJ, Blankman E, Stillman DJ, Eide DJ, Winge DR. The Zap1 transcriptional activator also acts as a repressor by binding downstream of the TATA box in ZRT2. <i>EMBO J</i> 2004;23(5):1123–32.	doi:10.1038/sj.emboj.7600122 PMID 14976557 PMC380977	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
1114	Tomás-Cobos 2004 Tomás-Cobos L, Casadomé L, Mas G, Sanz P, Posas F. Expression of the HXT1 low affinity glucose transporter requires the coordinated activities of the HOG and glucose signalling pathways. <i>J Biol Chem</i> 2004;279(21):22010–9.	doi:10.1074/jbc.M400609200 PMID 15014083	–	2	2	2	–	–	1	–
1115	Seppä 2004 Seppä L, Hänninen AL, Makarow M. Upregulation of the Hsp104 chaperone at physiological temperature during recovery from thermal insult. <i>Mol Microbiol</i> 2004;52(1):217–25.	doi:10.1111/j.1365-2958.2003.03959.x PMID 15049822	–	2	2	2	–	–	1	–
1116	Tsuzi 2004 Tsuzi D, Maeta K, Takatsume Y, Izawa S, Inoue Y. Regulation of the yeast phospholipid hydroperoxide glutathione peroxidase GPX2 by oxidative stress is mediated by Yap1 and Skn7. <i>FEBS Lett</i> 2004;565(1–3):148–54.	doi:10.1016/j.febslet.2004.03.091 PMID 15135069	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1117	Kim 2004 Kim TS, Lee SB, Kang HS. Glucose repression of STA1 expression is mediated by the Nrg1 and Sfl1 repressors and the Srb8-11 complex. <i>Mol Cell Biol</i> 2004;24(17):7695–706.	doi:10.1128/MCB.24.17.7695-7706.2004 PMID 15314176 PMC507009	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1118	Monteiro 2004 Monteiro G, Netto LE. Glucose repression of PRX1 expression is mediated by Tor1p and Ras2p through inhibition of Msn2/4p in <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiol Lett</i> 2004;241(2):221–8.	doi:10.1016/j.femsle.2004.10.024 PMID 15598536	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1119	Martin 2004 Martin DE, Soulard A, Hall MN. TOR regulates ribosomal protein gene expression via PKA and the Forkhead transcription factor FHL1. <i>Cell</i> 2004;119(7):969–79.	doi:10.1016/j.cell.2004.11.047 PMID 15620355	–	2	2	2	–	2	2	Northern blot - wt vs TF mutant
1120	Oliveira 2005 Oliveira EM, Mansure JJ, Bon EP. Gln3p and Nil1p regulation of invertase activity and SUC2 expression in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Res</i> 2005;5(6–7):605–9.	doi:10.1016/j.femsyr.2004.11.011 PMID 15780659	–	2	2	2	–	2	1	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1121	Doniger 2005 Doniger SW, Huh J, Fay JC. Identification of functional transcription factor binding sites using closely related <i>Saccharomyces</i> species. <i>Genome Res</i> 2005;15(5):701–9.	doi:10.1101/gr.3578205 PMID 15837806 PMC1088298	–	2	2	2	1	1	2	Microarray analysis - wt vs TF mutant
1122	Gatti 2005 Gatti X, de Bettignies G, Claret S, Doignon F, Crouzet M, Thoraval D. RGD1, encoding a RhoGAP involved in low-pH survival, is an Msn2p/Msn4p regulated gene in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> 2005;351:159–69.	doi:10.1016/j.gene.2005.03.034 PMID 15922872	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1123	Rothfels 2005 Rothfels K, Tanny JC, Molnar E, Friesen H, Commisso C, Segall J. Components of the ESCRT pathway, DFG16, and YGR122w are required for Rim101 to act as a corepressor with Nrg1 at the negative regulatory element of the DIT1 gene of <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2005;25(15):6772–88.	doi:10.1128/MCB.25.15.6772-6788.2005 PMID 16024810 PMC1190364	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1124	Zhang 2005 Zhang Z, Reese JC. Molecular genetic analysis of the yeast repressor Rfx1/Crt1 reveals a novel two-step regulatory mechanism. <i>Mol Cell Biol</i> 2005;25(17):7399–411.	doi:10.1128/MCB.25.17.7399-7411.2005 PMID 16107689 PMC1190298	–	2	1	2	1	1	2	ChIP
1125	Takemori 2006 Takemori Y, Sakaguchi A, Matsuda S, Mizukami Y, Sakurai H. Stress-induced transcription of the endoplasmic reticulum oxidoreductin gene ERO1 in the yeast <i>Saccharomyces cerevisiae</i> . <i>Mol Genet Genomics</i> 2006;275(1):89–96.	doi:10.1007/s00438-005-0065-9 PMID 16292667	–	2	2	2	–	2	2	Northern blot - wt vs TF mutant
1126	Kim 2006 Kim JH, Brachet V, Moriya H, Johnston M. Integration of transcriptional and posttranslational regulation in a glucose signal transduction pathway in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2006;5(1):167–73.	doi:10.1128/EC.5.1.167-173.2006 PMID 16400179 PMC1360249	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
1127	Kleinschmidt 2006 Kleinschmidt M, Schulz R, Braus GH. The yeast CPC2/ASC1 gene is regulated by the transcription factors Fhl1p and Ifh1p. <i>Curr Genet</i> 2006;49(4):218–28.	doi:10.1007/s00294-005-0049-7 PMID 16402205	–	2	2	2	–	2	2	Northern blot - wt vs TF mutant
1128	Panwar 2006 Panwar SL, Moye-Rowley WS. Long chain base tolerance in <i>Saccharomyces cerevisiae</i> is induced by retrograde signals from the mitochondria. <i>J Biol Chem</i> 2006;281(10):6376–84.	doi:10.1074/jbc.M512115200 PMID 16407254	–	2	2	2	–	1	2	–
1129	Hahn 2006 Hahn JS, Neef DW, Thiele DJ. A stress regulatory network for co-ordinated activation of proteasome expression mediated by yeast heat shock transcription factor. <i>Mol Microbiol</i> 2006; 60(1):240–51.	doi:10.1111/j.1365-2958.2006.05097.x PMID 16556235	–	2	1	2	1	–	1	–
1130	Boban 2006 Boban M, Zargari A, Andréasson C, Heessen S, Thyberg J, Ljungdahl PO. Asi1 is an inner nuclear membrane protein that restricts promoter access of two latent transcription factors. <i>J Cell Biol</i> 2006;173(5):695–707.	doi:10.1083/jcb.200601011 PMID 16735580 PMC2063887	–	2	1	2	2	–	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1131	Jesch 2006 Jesch SA, Liu P, Zhao X, Wells MT, Henry SA. Multiple endoplasmic reticulum-to-nucleus signaling pathways coordinate phospholipid metabolism with gene expression by distinct mechanisms. <i>J Biol Chem</i> 2006;281(33):24070–83.	doi:10.1074/jbc.M604541200 PMID 16777852	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
1132	Jackevicius 2007 Jackevicius CA, Paterson JM, Naglie G. Concordance between discharge prescriptions and insurance claims in post-myocardial infarction patients. <i>Pharmacoepidemiol Drug Saf</i> 2007;16(2):207–15.	doi:10.1002/pds.1289 PMID 16862605	–	2	1	2	–	–	1	–
1133	Reynolds 2006 Reynolds TB. The Opilp transcription factor affects expression of FLO11, mat formation, and invasive growth in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2006;5(8):1266–75.	doi:10.1128/EC.00022-06 PMID 16896211 PMC1539139	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1134	Martin 2007 Martin CE, Oh CS, Jiang Y. Regulation of long chain unsaturated fatty acid synthesis in yeast. <i>Biochim Biophys Acta</i> 2007;1771(3):271–85.	doi:10.1016/j.bbailip.2006.06.010 PMID 16920014	–	2	2	2	–	–	1	–
1135	Simões 2006 Simões T, Mira NP, Fernandes AR, Sá-Correia I. The SPI1 gene, encoding a glycosylphosphatidylinositol-anchored cell wall protein, plays a prominent role in the development of yeast resistance to lipophilic weak-acid food preservatives. <i>Appl Environ Microbiol</i> 2006;72(11):7168–75.	doi:10.1128/AEM.01476-06 PMID 16980434 PMC1636168	–	2	1	1	–	2	2	Northern blot - wt vs TF mutant
1136	Wiles 2006 Wiles AM, Cai H, Naider F, Becker JM. Nutrient regulation of oligopeptide transport in <i>Saccharomyces cerevisiae</i> . <i>Microbiology (Reading)</i> 2006;152(Pt 10):3133–3145.	doi:10.1099/mic.0.29055-0 PMID 17005992	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1137	Paes de Faria 2006 Paes de Faria J, Fernandes L. Protection against oxidative stress through SUA7/TFIIB regulation in <i>Saccharomyces cerevisiae</i> . <i>Free Radic Biol Med</i> 2006;41(11):1684–93.	doi:10.1016/j.freeradbiomed.2006.09.003 PMID 17145557	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
1138	Tevzadze 2007 Tevzadze GG, Pierce JV, Esposito RE. Genetic evidence for a SPO1-dependent signaling pathway controlling meiotic progression in yeast. <i>Genetics</i> 2007;175(3):1213–27.	doi:10.1534/genetics.106.069252 PMID 17179081 PMC1840080	–	2	2	2	–	2	2	Northern blot - wt vs TF mutant
1139	Sherriff 2007 Sherriff JA, Kent NA, Mellor J. The Isw2 chromatin-remodeling ATPase cooperates with the Fkh2 transcription factor to repress transcription of the B-type cyclin gene CLB2. <i>Mol Cell Biol</i> 2007;27(8):2848–60.	doi:10.1128/MCB.01798-06 PMID 17283050 PMC1899929	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1140	Wiesenberger 2007 Wiesenberger G, Steinleitner K, Malli R, et al., Stadler JA. Mg2+ deprivation elicits rapid Ca2+ uptake and activates Ca2+/calcineurin signaling in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2007;6(4):592–9.	doi:10.1128/EC.00382-06 PMID 17337637 PMC1865649	GSE6687	2	1	2	–	2	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1141	Boban 2007 Boban M, Ljungdahl PO. Dal81 enhances Stp1- and Stp2-dependent transcription necessitating negative modulation by inner nuclear membrane protein Asil in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2007;176(4):2087–97.	doi:10.1534/genetics.107.075077 PMID 17603098 PMC1950616	–	2	1	2	2	–	1	ChIP
1142	Cai 2007 Cai H, Hauser M, Naider F, Becker JM. Differential regulation and substrate preferences in two peptide transporters of <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2007;6(10):1805–13.	doi:10.1128/EC.00257-06 PMID 17693598 PMC2043388	–	2	1	2	–	2	2	RT-PCR - wt vs TF mutant
1143	Chen 2007 Chen XJ, Wang X, Butow RA. Yeast aconitase binds and provides metabolically coupled protection to mitochondrial DNA. <i>Proc Natl Acad Sci U S A</i> 2007;104(34):13738–43.	doi:10.1073/pnas.0703078104 PMID 17698960 PMC1959452	–	2	2	2	–	2	1	Western blot - wt vs TF mutant
1144	Doyon 2007 Doyon JB, Liu DR. Identification of eukaryotic promoter regulatory elements using nonhomologous random recombination. <i>Nucleic Acids Res</i> 2007;35(17):5851–60.	doi:10.1093/nar/gkm634 PMID 17720707 PMC2034452	–	2	1	1	–	2	2	lacZ - wt vs TF mutant
1145	Han 2008 Han Q, Lu J, Duan J, et al., Huang B. Gcn5- and Elp3-induced histone H3 acetylation regulates hsp70 gene transcription in yeast. <i>Biochem J</i> 2008;409(3):779–88.	doi:10.1042/BJ20070578 PMID 17910533	–	2	1	2	2	–	1	ChIP
1146	Medvedik 2007 Medvedik O, Lamming DW, Kim KD, Sinclair DA. MSN2 and MSN4 link calorie restriction and TOR to sirtuin-mediated lifespan extension in <i>Saccharomyces cerevisiae</i> . <i>PLoS Biol</i> 2007;5(10):e261.	doi:10.1371/journal.pbio.0050261 PMID 17914901 PMC1994990	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1147	Luzzani 2007 Luzzani C, Cardillo SB, Bermúdez Moretti M, Correa García S. New insights into the regulation of the <i>Saccharomyces cerevisiae</i> UGA4 gene: two parallel pathways participate in carbon-regulated transcription. <i>Microbiology (Reading)</i> 2007;153(Pt 11):3677–3684.	doi:10.1099/mic.0.2007/010231-0 PMID 17975075	–	2	1	1	–	2	2	lacZ - wt vs TF mutant
1148	Bermejo 2008 Bermejo C, Rodríguez E, García R, et al., Arroyo J. The sequential activation of the yeast HOG and SLT2 pathways is required for cell survival to cell wall stress. <i>Mol Biol Cell</i> 2008;19(3):1113–24.	doi:10.1091/mbc.e07-08-0742 PMID 18184748 PMC2262984	–	2	1	1	–	2	2	lacZ - wt vs TF mutant
1149	Kobayashi 2008 Kobayashi Y, Inai T, Mizunuma M, et al., Miyakawa T. Identification of Tup1 and Cyc8 mutations defective in the responses to osmotic stress. <i>Biochem Biophys Res Commun</i> 2008;368(1):50–5.	doi:10.1016/j.bbrc.2008.01.033 PMID 18201562	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1150	Ng 2008 Ng CH, Tan SX, Perrone GG, Thorpe GW, Higgins VJ, Dawes IW. Adaptation to hydrogen peroxide in <i>Saccharomyces cerevisiae</i> : the role of NADPH-generating systems and the SKN7 transcription factor. <i>Free Radic Biol Med</i> 2008;44(6):1131–45.	doi:10.1016/j.freeradbiomed.2007.12.008 PMID 18206664	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1151	Karpova 2008 Karpova TS, Kim MJ, Spriet C, et al., McNally JG. Concurrent fast and slow cycling of a transcriptional activator at an endogenous promoter. <i>Science</i> 2008;319(5862):466–9.	doi:10.1126/science.1150559 PMID 18218898	–	2	1	2	2	–	1	FRAP
1152	Hausmann 2008 Hausmann A, Samans B, Lill R, Mühlenhoff U. Cellular and mitochondrial remodeling upon defects in iron-sulfur protein biogenesis. <i>J Biol Chem</i> 2008;283(13):8318–30.	doi:10.1074/jbc.M705570200 PMID 18227070	–	2	1	2	–	2	2	promoter assay
1153	Georis 2008 Georis I, Tate JJ, Cooper TG, Dubois E. Tor pathway control of the nitrogen-responsive DAL5 gene bifurcates at the level of Gln3 and Gat1 regulation in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2008;283(14):8919–29.	doi:10.1074/jbc.M708811200 PMID 18245087 PMC2276367	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1154	Yamamoto 2008 Yamamoto N, Maeda Y, Ikeda A, Sakurai H. Regulation of thermotolerance by stress-induced transcription factors in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2008;7(5):783–90.	doi:10.1128/EC.00029-08 PMID 18359875 PMC2394977	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1155	Shaner 2008 Shaner L, Gibney PA, Morano KA. The Hsp110 protein chaperone Sse1 is required for yeast cell wall integrity and morphogenesis. <i>Curr Genet</i> 2008;54(1):1–11.	doi:10.1007/s00294-008-0193-y PMID 18478233 PMC5635425	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1156	Weiss 2008 Weiss P, Huppert S, Kölling R. ESCRT-III protein Snf7 mediates high-level expression of the SUC2 gene via the Rim101 pathway. <i>Eukaryot Cell</i> 2008;7(11):1888–94.	doi:10.1128/EC.00194-08 PMID 18806212 PMC2583539	–	2	2	2	–	2	1	Enzyme activity - wt vs TF mutant
1157	Woo 2009 Woo DK, Phang TL, Trawick JD, Poyton RO. Multiple pathways of mitochondrial-nuclear communication in yeast: intergenomic signaling involves ABF1 and affects a different set of genes than retrograde regulation. <i>Biochim Biophys Acta</i> 2009;1789(2):135–45.	doi:10.1016/j.bbagr.2008.09.008 PMID 18977319	–	2	2	2	2	–	1	lacZ - wt vs mutant promoter sequence
1158	Wood 2009 Wood LK, Thiele DJ. Transcriptional activation in yeast in response to copper deficiency involves copper-zinc superoxide dismutase. <i>J Biol Chem</i> 2009;284(1):404–413.	doi:10.1074/jbc.M807027200 PMID 18977757 PMC2610518	–	2	1	2	1	1	2	ChIP
1159	Prasad 2008 Prasad V, Venkatesh KV. Stochastic analysis of the GAL genetic switch in <i>Saccharomyces cerevisiae</i> : modeling and experiments reveal hierarchy in glucose repression. <i>BMC Syst Biol</i> 2008;2:97.	doi:10.1186/1752-0509-2-97 PMID 19014615 PMC2614938	–	2	1	2	–	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1160	Jani 2008 Jani NM, Lopes JM. Transcription regulation of the <i>Saccharomyces cerevisiae</i> PIS1 gene by inositol and the pleiotropic regulator, Ume6p. <i>Mol Microbiol</i> 2008;70(6):1529–39.	doi:10.1111/j.1365-2958.2008.06506.x PMID 19019152	–	2	2	2	1	1	1	CAT - wt vs TF mutant
1161	Thakur 2009 Thakur JK, Arthanari H, Yang F, Chau KH, Wagner G, Näär AM. Mediator subunit Gal11p/MED15 is required for fatty acid-dependent gene activation by yeast transcription factor Oaf1p. <i>J Biol Chem</i> 2009;284(7):4422–8.	doi:10.1074/jbc.M808263200 PMID 19056732 PMC3837390	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1162	Sullivan 2009 Sullivan DP, Georgiev A, Menon AK. Tritium suicide selection identifies proteins involved in the uptake and intracellular transport of sterols in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2009;8(2):161–9.	doi:10.1128/EC.00135-08 PMID 19060182 PMC2643606	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1163	Hosiner 2009 Hosiner D, Lempiäinen H, Reiter W, et al., Schüller C. Arsenic toxicity to <i>Saccharomyces cerevisiae</i> is a consequence of inhibition of the TORC1 kinase combined with a chronic stress response. <i>Mol Biol Cell</i> 2009;20(3):1048–57.	doi:10.1091/mbc.e08-04-0438 PMID 19073887 PMC2633375	–	2	1	2	2	–	2	ChIP
1164	Van Mulders 2009 Van Mulders SE, Christianen E, Saerens SM, et al., Delvaux FR. Phenotypic diversity of Flo protein family-mediated adhesion in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Res</i> 2009;9(2):178–90.	doi:10.1111/j.1567-1364.2008.00462.x PMID 19087208	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1165	Lorenz 2009 Lorenz DR, Cantor CR, Collins JJ. A network biology approach to aging in yeast. <i>Proc Natl Acad Sci U S A</i> 2009;106(4):1145–50.	doi:10.1073/pnas.0812551106 PMID 19164565 PMC2629491	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
1166	Ahmed 2009 Ahmed NT, Bungard D, Shin ME, Moore M, Winter E. The Ime2 protein kinase enhances the disassociation of the Sum1 repressor from middle meiotic promoters. <i>Mol Cell Biol</i> 2009;29(16):4352–62.	doi:10.1128/MCB.00305-09 PMID 19528232 PMC2725727	–	2	2	2	2	–	1	ChIP
1167	Leverentz 2009 Leverentz MK, Campbell RN, Connolly Y, Whetton AD, Reece RJ. Mutation of a phosphorylatable residue in Put3p affects the magnitude of rapamycin-induced PUT1 activation in a Gat1p-dependent manner. <i>J Biol Chem</i> 2009;284(36):24115–22.	doi:10.1074/jbc.M109.030361 PMID 19574222 PMC2782005	–	2	2	2	2	–	1	ChIP
1168	Gurvitz 2009 Gurvitz A. A novel circuit overrides Adr1p control during expression of <i>Saccharomyces cerevisiae</i> 2-trans-enoyl-ACP reductase Etr1p of mitochondrial type 2 fatty acid synthase. <i>FEMS Microbiol Lett</i> 2009;297(2):255–60.	doi:10.1111/j.1574-6968.2009.01688.x PMID 19583790 PMC2784040	–	2	1	2	–	2	1	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1169	Truman 2009 Truman AW, Kim KY, Levin DE. Mechanism of Mpk1 mitogen-activated protein kinase binding to the Swi4 transcription factor and its regulation by a novel caffeine-induced phosphorylation. <i>Mol Cell Biol</i> 2009;29(24):6449–61.	doi:10.1128/MCB.00794-09 PMID 19805511 PMC2786871	–	2	1	2	–	2	2	lacZ - wt vs TF mutant
1170	Collins 2009 Collins GA, Lipford JR, Deshaies RJ, Tansey WP. Gal4 turnover and transcription activation. <i>Nature</i> 2009;461(7265):E7;discussion E8.	doi:10.1038/nature08406 PMID 19812621 PMC3072683	–	2	1	2	2	–	1	ChIP
1171	Louw 2010 Louw C, Young PR, van Rensburg P, Divol B. Regulation of endo-polygalacturonase activity in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Res</i> 2010;10(1):44–57.	doi:10.1111/j.1567-1364.2009.00584.x PMID 19840115	–	2	2	2	–	2	1	Overexpression of TF
1172	Ju 2010 Ju D, Xu H, Wang X, Xie Y. The transcription activation domain of Rpn4 is separate from its degrons. <i>Int J Biochem Cell Biol</i> 2010;42(2):282–6.	doi:10.1016/j.biocel.2009.11.003 PMID 19914394	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1173	Wang 2010 Wang X, Xu H, Ha SW, Ju D, Xie Y. Proteasomal degradation of Rpn4 in <i>Saccharomyces cerevisiae</i> is critical for cell viability under stressed conditions. <i>Genetics</i> 2010;184(2):335–42.	doi:10.1534/genetics.109.112227 PMID 19933873 PMC2828715	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1174	Bosis 2010 Bosis E, Salomon D, Ohayon O, Sivan G, Bar-Nun S, Rabinovich E. Ssz1 restores endoplasmic reticulum-associated protein degradation in cells expressing defective cdc48-ufd1-npl4 complex by upregulating cdc48. <i>Genetics</i> 2010;184(3):695–706.	doi:10.1534/genetics.109.111419 PMID 20038635 PMC2845338	–	2	1	2	–	2	1	Western blot - wt vs TF mutant
1175	Sekiya 2009 Sekiya M, Nogami S, Ohya Y. Transcription factors of M-phase cyclin CLB2 in the yeast cell wall integrity checkpoint. <i>Genes Genet Syst</i> 2009;84(4):269–76.	doi:10.1266/ggs.84.269 PMID 20057164	–	2	2	2	–	2	2	Northern blot - wt vs TF mutant
1176	Boysen 2010 Boysen JH, Subramanian S, Mitchell AP. Intervention of Bro1 in pH-responsive Rim20 localization in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2010;9(4):532–8.	doi:10.1128/EC.00027-10 PMID 20190076 PMC2863415	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1177	Desimone 2010 Desimone AM, Laney JD. Corepressor-directed preacetylation of histone H3 in promoter chromatin primes rapid transcriptional switching of cell-type-specific genes in yeast. <i>Mol Cell Biol</i> 2010;30(13):3342–56.	doi:10.1128/MCB.01450-09 PMID 20439496 PMC2897591	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1178	Hong 2011 Hong S, Yoon S. Mcm1p binding sites in the ARG1 promoter positively regulate ARG1 transcription and <i>S. cerevisiae</i> growth in the absence of arginine and Gcn4p. <i>Amino Acids</i> 2011;40(2):623–31.	doi:10.1007/s00726-010-0687-z PMID 20625780	–	2	2	2	2	–	1	Northern blot - wt vs mutant promoter sequence

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1179	Georis 2011 Georis I, Tate JJ, Feller A, Cooper TG, Dubois E. Intranuclear function for protein phosphatase 2A: Pph21 and Pph22 are required for rapamycin-induced GATA factor binding to the DAL5 promoter in yeast. <i>Mol Cell Biol</i> 2011;31(1):92–104.	doi:10.1128/MCB.00482-10 PMID 20974806 PMC3019842	–	2	2	2	2	–	1	ChIP
1180	Eriksson 2011 Eriksson PR, Ganguli D, Clark DJ. Spt10 and Swi4 control the timing of histone H2A/H2B gene activation in budding yeast. <i>Mol Cell Biol</i> 2011;31(3):557–72.	doi:10.1128/MCB.00909-10 PMID 21115727 PMC3028627	–	2	1	2	2	–	1	EMSA
1181	Teixeira 2011 Teixeira MC, Cabrito TR, Hanif ZM, Vargas RC, Tenreiro S, Sá-Correia I. Yeast response and tolerance to polyamine toxicity involving the drug : H ⁺ antiporter Qdr3 and the transcription factors Yap1 and Gcn4. <i>Microbiology (Reading)</i> 2011;157(Pt 4):945–956.	doi:10.1099/mic.0.043661-0 PMID 21148207	–	2	2	2	–	2	2	RT-PCR - wt vs TF mutant
1182	Knutson 2011 Knutson BA, Hahn S. Domains of Tra1 important for activator recruitment and transcription coactivator functions of SAGA and NuA4 complexes. <i>Mol Cell Biol</i> 2011;31(4):818–31.	doi:10.1128/MCB.00687-10 PMID 21149579 PMC3028633	–	2	2	2	2	–	1	ChIP
1183	Cocklin 2011 Cocklin R, Heyen J, Larry T, Tyers M, Goebel M. New insight into the role of the Cdc34 ubiquitin-conjugating enzyme in cell cycle regulation via Ace2 and Sic1. <i>Genetics</i> 2011;187(3):701–15.	doi:10.1534/genetics.110.125302 PMID 21196523 PMC3063666	GSE8453	2	2	2	–	2	1	Western blot - wt vs TF mutant
1184	Zhang 2011 Zhang M, Zhang C, Li J, et al., Xiao W. Inactivation of YAP1 enhances sensitivity of the yeast RNR3-lacZ genotoxicity testing system to a broad range of DNA-damaging agents. <i>Toxicol Sci</i> 2011;120(2):310–21.	doi:10.1093/toxsci/kfq391 PMID 21205635	–	2	1	1	–	2	2	lacZ - wt vs TF mutant
1185	Hirasaki 2011 Hirasaki M, Horiguchi M, Numamoto M, et al., Harashima S. Saccharomyces cerevisiae protein phosphatase Ppz1 and protein kinases Sat4 and Hal5 are involved in the control of subcellular localization of Gln3 by likely regulating its phosphorylation state. <i>J Biosci Bioeng</i> 2011;111(3):249–54.	doi:10.1016/j.jbiosc.2010.11.013 PMID 21237705	–	2	1	1	–	2	2	lacZ - wt vs TF mutant
1186	Lim 2011 Lim MK, Siew WL, Zhao J, Tay YC, Ang E, Lehming N. Galactose induction of the GAL1 gene requires conditional degradation of the Mig2 repressor. <i>Biochem J</i> 2011;435(3):641–9.	doi:10.1042/BJ20102034 PMID 21323640	–	2	2	2	2	–	2	ChIP
1187	Dhaoui 2011 Dhaoui M, Auchère F, Blaiseau PL, et al., Belgareh-Touzé N. Gex1 is a yeast glutathione exchanger that interferes with pH and redox homeostasis. <i>Mol Biol Cell</i> 2011;22(12):2054–67.	doi:10.1091/mbc.E10-11-0906 PMID 21490148 PMC3113770	–	2	2	2	–	2	1	Western blot - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1188	Bai 2011 Bai L, Ondracka A, Cross FR. Multiple sequence-specific factors generate the nucleosome-depleted region on CLN2 promoter. <i>Mol Cell</i> 2011;42(4):465–76.	doi:10.1016/j.molcel.2011.03.028 PMID 21596311 PMC3119483	–	2	2	2	2	–	1	ChIP
1189	Quan 2011 Quan Z, Oliver SG, Zhang N. JmjN interacts with JmjC to ensure selective proteolysis of Gis1 by the proteasome. <i>Microbiology (Reading)</i> 2011;157(Pt 9):2694–2701.	doi:10.1099/mic.0.048199-0 PMID 21680636	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
1190	Toesca 2011 Toesca I, Nery CR, Fernandez CF, Sayani S, Chanfreau GF. Cryptic transcription mediates repression of subtelomeric metal homeostasis genes. <i>PLoS Genet</i> 2011;7(6):e1002163.	doi:10.1371/journal.pgen.1002163 PMID 21738494 PMC3128112	–	2	2	2	1	1	1	ChIP
1191	Jouandot 2011 Jouandot D 2nd, Roy A, Kim JH. Functional dissection of the glucose signaling pathways that regulate the yeast glucose transporter gene (HXT) repressor Rgt1. <i>J Cell Biochem</i> 2011;112(11):3268–75.	doi:10.1002/jcb.23253 PMID 21748783 PMC3341738	–	2	1	2	2	–	1	ChIP
1192	Brückner 2011 Brückner S, Kern S, Birke R, Saugar I, Ulrich HD, Mösch HU. The TEA transcription factor Tec1 links TOR and MAPK pathways to coordinate yeast development. <i>Genetics</i> 2011;189(2):479–94.	doi:10.1534/genetics.111.133629 PMID 21840851 PMC3189810	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
1193	Spasskaia 2011 Spasskaia DS, Karpov DS, Karpov VL. [Escherichia coli Dam methylase as a molecular tool for mapping binding sites of the yeast transcription factor Rpn4]. <i>Mol Biol (Mosk)</i> 2011;45(4):642–51.	PMID 21954596	–	2	1	2	2	–	1	Dam-Rpn4 interaction with the target gene (measuring level of methylation of the target promoter)
1194	Dos Santos 2011 Dos Santos SC, Sá-Correia I. A genome-wide screen identifies yeast genes required for protection against or enhanced cytotoxicity of the antimalarial drug quinine. <i>Mol Genet Genomics</i> 2011;286(5–6):333–46.	doi:10.1007/s00438-011-0649-5 PMID 21960436	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1195	Baudouin-Cornu 2012 Baudouin-Cornu P, Lagniel G, Kumar C, Huang ME, Labarre J. Glutathione degradation is a key determinant of glutathione homeostasis. <i>J Biol Chem</i> 2012;287(7):4552–61.	doi:10.1074/jbc.M111.315705 PMID 22170048 PMC3281641	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1196	Xue 2011 Xue Y, Rushton MD, Maringe L. A novel checkpoint and RPA inhibitory pathway regulated by Rif1. <i>PLoS Genet</i> 2011;7(12):e1002417.	doi:10.1371/journal.pgen.1002417 PMID 22194703 PMC3240596	–	2	1	2	2	–	1	ChIP
1197	Brzovic 2011 Brzovic PS, Heikaus CC, Kisselev L, et al., Hahn S. The acidic transcription activator Gcn4 binds the mediator subunit Gal11/Med15 using a simple protein interface forming a fuzzy complex. <i>Mol Cell</i> 2011;44(6):942–53.	doi:10.1016/j.molcel.2011.11.008 PMID 22195967 PMC3246216	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1198	Malik 2012 Malik S, Chaurasia P, Lahudkar S, Uprety B, Bhaumik SR. Rad26p regulates the occupancy of histone H2A-H2B dimer at the active genes in vivo. <i>Nucleic Acids Res</i> 2012;40(8):3348–63.	doi:10.1093/nar/gkr1244 PMID 22199252 PMC3333851	–	2	1	1	2	–	2	ChIP
1199	Davidson 2012 Davidson MB, Katou Y, Keszthelyi A, et al., Brown GW. Endogenous DNA replication stress results in expansion of dNTP pools and a mutator phenotype. <i>EMBO J</i> 2012;31(4):895–907.	doi:10.1038/emboj.2011.485 PMID 22234187 PMC3280564	–	2	1	2	–	2	1	Western blot - wt vs TF mutant
1200	Abate 2012 Abate G, Bastonini E, Braun KA, Verdone L, Young ET, Caserta M. Snf1/AMPK regulates Gcn5 occupancy, H3 acetylation and chromatin remodelling at <i>S. cerevisiae</i> ADY2 promoter. <i>Biochim Biophys Acta</i> 2012;1819(5):419–27.	doi:10.1016/j.bbagr.2012.01.009 PMID 22306658 PMC3319277	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1201	Flom 2012 Flom GA, Langner E, Johnson JL. Identification of an Hsp90 mutation that selectively disrupts cAMP/PKA signaling in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 2012;58(3):149–63.	doi:10.1007/s00294-012-0373-7 PMID 22461145	GSE34757	2	2	2	–	2	1	Western blot - wt vs TF mutant
1202	Ohdate 2012 Ohdate T, Inoue Y. Involvement of glutathione peroxidase 1 in growth and peroxisome formation in <i>Saccharomyces cerevisiae</i> in oleic acid medium. <i>Biochim Biophys Acta</i> 2012;1821(9):1295–305.	doi:10.1016/j.bbailip.2012.05.004 PMID 22659048	–	2	2	2	–	2	1	Western blot - wt vs TF mutant
1203	Qi 2012 Qi J, Han A, Yang Z, Li C. Metal-sensing transcription factors Mac1p and Aft1p coordinately regulate vacuolar copper transporter CTR2 in <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2012;423(2):424–8.	doi:10.1016/j.bbrc.2012.05.150 PMID 22683637	–	2	2	2	–	2	2	RT-PCR - wt vs TF mutant
1204	Bartholomew 2012 Bartholomew CR, Suzuki T, Du Z, et al., Klionsky DJ. Ume6 transcription factor is part of a signaling cascade that regulates autophagy. <i>Proc Natl Acad Sci U S A</i> 2012;109(28):11206–10.	doi:10.1073/pnas.1200313109 PMID 22733735 PMC3396506	–	2	1	2	1	1	1	ChIP
1205	Hanscho 2012 Hanscho M, Ruckerbauer DE, Chauhan N, et al., Natter K. Nutritional requirements of the BY series of <i>Saccharomyces cerevisiae</i> strains for optimum growth. <i>FEMS Yeast Res</i> 2012;12(7):796–808.	doi:10.1111/j.1567-1364.2012.00830.x PMID 22780918	–	2	1	1	–	2	2	RT-PCR - wt vs TF mutant
1206	Li 2012 Li L, Miao R, Bertram S, Jia X, Ward DM, Kaplan J. A role for iron-sulfur clusters in the regulation of transcription factor Yap5-dependent high iron transcriptional responses in yeast. <i>J Biol Chem</i> 2012;287(42):35709–35721.	doi:10.1074/jbc.M112.395533 PMID 22915593 PMC3471690	–	2	1	1	–	2	2	lacZ - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1207	Darieva 2012 Darieva Z, Han N, Warwood S, Doris KS, Morgan BA, Sharrocks AD. Protein kinase C regulates late cell cycle-dependent gene expression. <i>Mol Cell Biol</i> 2012;32(22):4651–61.	doi:10.1128/MCB.06000-11 PMID 22966207 PMC3486177	–	2	1	2	2	–	1	ChIP
1208	Barreto 2012 Barreto L, Canadell D, Valverde-Saubí D, Casamayor A, Ariño J. The short-term response of yeast to potassium starvation. <i>Environ Microbiol</i> 2012;14(11):3026–42.	doi:10.1111/j.1462-2920.2012.02887.x PMID 23039231	GSE24712	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1209	Peñalosa-Ruiz 2012 Peñalosa-Ruiz G, Aranda C, Ongay-Larios L, Colon M, Quezada H, Gonzalez A. Paralogous ALT1 and ALT2 retention and diversification have generated catalytically active and inactive aminotransferases in <i>Saccharomyces cerevisiae</i> . <i>PLoS One</i> 2012;7(9):e45702.	doi:10.1371/journal.pone.0045702 PMID 23049841 PMC3458083	–	2	1	2	2	–	1	ChIP
1210	Walkey 2012 Walkey CJ, Luo Z, Madilao LL, van Vuuren HJ. The fermentation stress response protein Aaf1p/Yml081Wp regulates acetate production in <i>Saccharomyces cerevisiae</i> . <i>PLoS One</i> 2012;7(12):e51551.	doi:10.1371/journal.pone.0051551 PMID 23240040 PMC3519862	–	2	1	2	2	–	1	RT-PCR - wt vs mutant promoter sequence
1211	Tennen 2013 Tennen RI, Haye JE, Wijayatilake HD, Arlow T, Ponzio D, Gammie AE. Cell-cycle and DNA damage regulation of the DNA mismatch repair protein Msh2 occurs at the transcriptional and post-transcriptional level. <i>DNA Repair (Amst)</i> 2013;12(2):97–109.	doi:10.1016/j.dnarep.2012.11.002 PMID 23261051 PMC3749301	–	2	2	2	–	2	1	Luciferase - wt vs TF mutant
1212	Fujiwara 2013 Fujiwara H, Kawai S, Murata K. Significance of sulfiredoxin/peroxiredoxin and mitochondrial respiratory chain in response to and protection from 100% O(2) in <i>Saccharomyces cerevisiae</i> . <i>Mitochondrion</i> 2013;13(1):52–8.	doi:10.1016/j.mito.2012.12.007 PMID 23291433	–	2	2	2	–	2	1	RT-PCR - wt vs TF mutant
1213	Mirzaei 2013 Mirzaei H, Knijnenburg TA, Kim B, et al., Ranish J. Systematic measurement of transcription factor-DNA interactions by targeted mass spectrometry identifies candidate gene regulatory proteins. <i>Proc Natl Acad Sci U S A</i> 2013;110(9):3645–50.	doi:10.1073/pnas.1216918110 PMID 23388641 PMC3587231	–	2	1	2	2	–	1	TF Selected Reaction Monitoring (SRM) assay
1214	Vidal 2013 Vidal SE, Pincus D, Stewart-Ornstein J, El-Samad H. Formation of subnuclear foci is a unique spatial behavior of mating MAPKs during hyperosmotic stress. <i>Cell Rep</i> 2013;3(2):328–34.	doi:10.1016/j.celrep.2013.01.022 PMID 23416049 PMC3616637	–	2	1	2	2	–	1	ChIP
1215	Harris 2013 Harris MR, Lee D, Farmer S, Lowndes NF, de Bruin RA. Binding specificity of the G1/S transcriptional regulators in budding yeast. <i>PLoS One</i> 2013;8(4):e61059.	doi:10.1371/journal.pone.0061059 PMID 23593391 PMC3617184	–	2	2	2	2	–	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1216	Dionne 2013 Dionne I, Larose S, Dandjinou AT, Abou Elela S, Wellinger RJ. Cell cycle-dependent transcription factors control the expression of yeast telomerase RNA. <i>RNA</i> 2013;19(7):992–1002.	doi:10.1261/rna.037663.112 PMID 23690630 PMC3683933	–	2	2	2	2	–	1	ChIP
1217	González 2013 González A, Casado C, Ariño J, Casamayor A. Ptc6 is required for proper rapamycin-induced down-regulation of the genes coding for ribosomal and rRNA processing proteins in <i>S. cerevisiae</i> . <i>PLoS One</i> 2013;8(5):e64470.	doi:10.1371/journal.pone.0064470 PMID 23704987 PMC3660562	GSE38260	2	1	2	2	–	1	ChIP
1218	Li 2013 Li M, Valsakumar V, Poorey K, Bekiranov S, Smith JS. Genome-wide analysis of functional sirtuin chromatin targets in yeast. <i>Genome Biol</i> 2013;14(5):R48.	doi:10.1186/gb-2013-14-5-r48 PMID 23710766 PMC4053722	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1219	Kim 2014 Kim B, Cho BR, Hahn JS. Metabolic engineering of <i>Saccharomyces cerevisiae</i> for the production of 2-phenylethanol via Ehrlich pathway. <i>Biotechnol Bioeng</i> 2014;111(1):115–24.	doi:10.1002/bit.24993 PMID 23836015	–	2	1	2	2	–	1	RT-PCR - wt vs mutant promoter sequence
1220	Lee 2013 Lee JC, Tsoi A, Kornfeld GD, Dawes IW. Cellular responses to L-serine in <i>Saccharomyces cerevisiae</i> : roles of general amino acid control, compartmentalization, and aspartate synthesis. <i>FEMS Yeast Res</i> 2013;13(7):618–34.	doi:10.1111/1567-1364.12063 PMID 23837815	–	2	1	2	–	2	1	Northern blot - wt vs TF mutant
1221	Bleoanca 2013 Bleoanca I, Silva AR, Pimentel C, Rodrigues-Pousada C, Menezes Rde A. Relationship between ethanol and oxidative stress in laboratory and brewing yeast strains. <i>J Biosci Bioeng</i> 2013;116(6):697–705.	doi:10.1016/j.jbiosc.2013.05.037 PMID 23838012	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1222	Lee 2013 Lee K, Sung C, Kim BG, Hahn JS. Activation of Aro80 transcription factor by heat-induced aromatic amino acid influx in <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2013;438(1):43–7.	doi:10.1016/j.bbrc.2013.07.019 PMID 23860270	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1223	Linke 2013 Linke C, Klipp E, Lehrach H, Barberis M, Krobitch S. Fkh1 and Fkh2 associate with Sir2 to control CLB2 transcription under normal and oxidative stress conditions. <i>Front Physiol</i> 2013;4:173.	doi:10.3389/fphys.2013.00173 PMID 23874301 PMC3709100	–	2	2	2	–	2	2	RT-PCR - wt vs TF mutant
1224	Dejos 2013 Dejos C, Régnacq M, Bernard M, Voisin P, Bergès T. The MFS-type efflux pump Fli1 induced by Yap1 promotes canthin-6-one resistance in yeast. <i>FEBS Lett</i> 2013;587(18):3045–51.	doi:10.1016/j.febslet.2013.07.040 PMID 23912082	–	2	1	2	2	–	1	RT-PCR - wt vs mutant promoter sequence

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1225	Zhao 2013 Zhao Y, Du J, Xiong B, Xu H, Jiang L. ESCRT components regulate the expression of the ER/Golgi calcium pump gene PMR1 through the Rim101/Nrg1 pathway in budding yeast. <i>J Mol Cell Biol</i> 2013;5(5):336–44.	doi:10.1093/jmcb/mjt025 PMID 23933635	–	2	2	2	1	1	2	ChIP
1226	Lee 2013 Lee SK, Chen X, Huang L, Stargell LA. The head module of Mediator directs activation of preloaded RNAPII in vivo. <i>Nucleic Acids Res</i> 2013;41(22):10124–34.	doi:10.1093/nar/gkt796 PMID 24005039 PMC3905900	–	2	1	2	2	–	1	ChIP
1227	Tsaponina 2013 Tsaponina O, Chabes A. Pre-activation of the genome integrity checkpoint increases DNA damage tolerance. <i>Nucleic Acids Res</i> 2013;41(22):10371–8.	doi:10.1093/nar/gkt820 PMID 24049076 PMC3905891	–	2	1	2	–	2	1	Western blot - wt vs TF mutant
1228	Chang 2013 Chang J, Zhou Y, Hu X, et al., Fraser HB. The molecular mechanism of a cis-regulatory adaptation in yeast. <i>PLoS Genet</i> 2013;9(9):e1003813.	doi:10.1371/journal.pgen.1003813 PMID 24068973 PMC3778017	–	2	2	2	–	2	1	RT-PCR - wt vs TF mutant
1229	Zhao 2014 Zhao X, Zou H, Fu J, Zhou J, Du G, Chen J. Metabolic engineering of the regulators in nitrogen catabolite repression to reduce the production of ethyl carbamate in a model rice wine system. <i>Appl Environ Microbiol</i> 2014;80(1):392–8.	doi:10.1128/AEM.03055-13 PMID 24185848 PMC3910993	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1230	Watanabe 2013 Watanabe D, Hashimoto N, Mizuno M, Zhou Y, Akao T, Shimoi H. Accelerated alcoholic fermentation caused by defective gene expression related to glucose derepression in <i>Saccharomyces cerevisiae</i> . <i>Biosci Biotechnol Biochem</i> 2013;77(11):2255–62.	doi:10.1271/bbb.130519 PMID 24200791	–	2	2	2	–	2	1	RT-PCR - wt vs TF mutant
1231	Law 2014 Law MJ, Mallory MJ, Dunbrack RL Jr, Strich R. Acetylation of the transcriptional repressor Ume6p allows efficient promoter release and timely induction of the meiotic transient transcription program in yeast. <i>Mol Cell Biol</i> 2014;34(4):631–42.	doi:10.1128/MCB.00256-13 PMID 24298021 PMC3911482	–	2	1	2	1	1	1	ChIP
1232	Zapata 2014 Zapata J, Dephoure N, Macdonough T, et al., Kellogg DR. PP2A ^{Rts1} is a master regulator of pathways that control cell size. <i>J Cell Biol</i> 2014;204(3):359–76.	doi:10.1083/jcb.201309119 PMID 24493588 PMC3912523	–	2	2	2	2	–	1	ChIP
1233	Zhou 2014 Zhou Q, Wang W, He X, et al., Yan Q. Mechanistic study on the nuclear modifier gene MSS1 mutation suppressing neomycin sensitivity of the mitochondrial 15S rRNA C1477G mutation in <i>Saccharomyces cerevisiae</i> . <i>PLoS One</i> 2014;9(3):e90336.	doi:10.1371/journal.pone.0090336 PMID 24595024 PMC3940847	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1234	Ndoja 2014 Ndoja A, Cohen RE, Yao T. Ubiquitin signals proteolysis-independent stripping of transcription factors. <i>Mol Cell</i> 2014;53(6):893–903.	doi:10.1016/j.molcel.2014.02.002 PMID 24613342 PMC4005849	–	2	2	2	2	–	2	ChIP
1235	Aihara 2014 Aihara M, Jin X, Kurihara Y, et al., Kanki T. Tor and the Sin3-Rpd3 complex regulate expression of the mitophagy receptor protein Atg32 in yeast. <i>J Cell Sci</i> 2014;127(Pt 14): 3184–96.	doi:10.1242/jcs.153254 PMID 24838945	–	2	1	2	2	–	1	ChIP
1236	Kang 2014 Kang HJ, Chang M, Kang CM, et al., Yun CW. The expression of PHO92 is regulated by Gcr1, and Pho92 is involved in glucose metabolism in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 2014;60(4):247–53.	doi:10.1007/s00294-014-0430-5 PMID 24850134	–	2	1	2	–	2	2	RT-PCR - wt vs TF mutant
1237	Ha 2014 Ha CW, Kim K, Chang YJ, Kim B, Huh WK. The β -1,3-glucanosyltransferase Gas1 regulates Sir2-mediated rDNA stability in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Res</i> 2014;42(13):8486–99.	doi:10.1093/nar/gku570 PMID 24981510 PMC4117787	–	2	2	2	2	–	1	ChIP
1238	Zhao 2014 Zhao Y, Xiong B, Xu H, Jiang L. Expression of NYV1 encoding the negative regulator of Pmc1 is repressed by two transcriptional repressors, Nrg1 and Mig1. <i>FEBS Lett</i> 2014; 588(17):3195–201.	doi:10.1016/j.febslet.2014.06.062 PMID 25017437	–	2	2	2	–	2	1	RT-PCR - wt vs TF mutant
1239	Park 2014 Park SH, Kim S, Hahn JS. Metabolic engineering of <i>Saccharomyces cerevisiae</i> for the production of isobutanol and 3-methyl-1-butanol. <i>Appl Microbiol Biotechnol</i> 2014;98(21): 9139–47.	doi:10.1007/s00253-014-6081-0 PMID 25280745	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1240	Nguyen 2014 Nguyen T, Fischl H, Howe FS, et al., Mellor J. Transcription mediated insulation and interference direct gene cluster expression switches. <i>Elife</i> 2014;3:e03635.	doi:10.7554/eLife.03635 PMID 25407679 PMC4275577	–	2	2	2	–	2	1	Northern blot - wt vs TF mutant
1241	Numamoto 2015 Numamoto M, Tagami S, Ueda Y, et al., Harashima S. Nuclear localization domains of GATA activator Gln3 are required for transcription of target genes through dephosphorylation in <i>Saccharomyces cerevisiae</i> . <i>J Biosci Bioeng</i> 2015;120(2):121–7.	doi:10.1016/j.jbiosc.2014.12.017 PMID 25641578	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1242	Yang 2015 Yang H, Tong J, Lee CW, Ha S, Eom SH, Im YJ. Structural mechanism of ergosterol regulation by fungal sterol transcription factor Upc2. <i>Nat Commun</i> 2015;6:6129.	doi:10.1038/ncomms7129 PMID 25655993	–	2	1	2	–	2	2	RT-PCR - wt vs TF mutant
1243	Gancedo 2015 Gancedo JM, Flores CL, Gancedo C. The repressor Rgt1 and the cAMP-dependent protein kinases control the expression of the SUC2 gene in <i>Saccharomyces cerevisiae</i> . <i>Biochim Biophys Acta</i> 2015;1850(7):1362–7.	doi:10.1016/j.bbagen.2015.03.006 PMID 25810078	–	2	2	2	–	2	1	Enzymatic measurements

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1244	Marqués 2015 Marqués MC, Zamarbide-Forés S, Pedelini L, Llopis-Torregrosa V, Yenush L. A functional Rim101 complex is required for proper accumulation of the Ena1 Na ⁺ -ATPase protein in response to salt stress in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Res</i> 2015;15(4):fov017.	doi:10.1093/femsyr/fov017 PMID 25934176	–	2	2	2	–	2	2	Western blot - wt vs TF mutant
1245	Yi 2015 Yi DG, Huh WK. PKA, PHO and stress response pathways regulate the expression of UDP-glucose pyrophosphorylase through Msn2/4 in budding yeast. <i>FEBS Lett</i> 2015;589(18):2409–16.	doi:10.1016/j.febslet.2015.07.015 PMID 26188548	–	2	2	2	–	2	1	RT-PCR - wt vs TF mutant
1246	Georis 2015 Georis I, Tate JJ, Vierendeels F, Cooper TG, Dubois E. Premature termination of GAT1 transcription explains paradoxical negative correlation between nitrogen-responsive mRNA, but constitutive low-level protein production. <i>RNA Biol</i> 2015;12(8):824–37.	doi:10.1080/15476286.2015.1058476 PMID 26259534 PMC4615157	–	2	2	2	–	2	2	RT-PCR - wt vs TF mutant
1247	Rai 2015 Rai R, Tate JJ, Shanmuganatham K, Howe MM, Nelson D, Cooper TG. Nuclear Gln3 Import Is Regulated by Nitrogen Catabolite Repression Whereas Export Is Specifically Regulated by Glutamine. <i>Genetics</i> 2015;201(3):989–1016.	doi:10.1534/genetics.115.177725 PMID 26333687 PMC4649666	–	2	1	2	1	1	2	EMSA
1248	An 2015 An X, Zhang C, Sclafani RA, Seligman P, Huang M. The late-annotated small ORF LSO1 is a target gene of the iron regulon of <i>Saccharomyces cerevisiae</i> . <i>Microbiologyopen</i> 2015;4(6):941–51.	doi:10.1002/mbo3.303 PMID 26450372 PMC4694146	–	2	2	2	–	2	1	Western blot - wt vs TF mutant
1249	Lin 2015 Lin Y, Sohn CH, Dalal CK, Cai L, Elowitz MB. Combinatorial gene regulation by modulation of relative pulse timing. <i>Nature</i> 2015;527(7576):54–8.	doi:10.1038/nature15710 PMID 26466562 PMC4870307	–	2	2	2	–	2	1	RT-PCR - wt vs TF mutant
1250	Yao 2015 Yao Z, Delorme-Axford E, Backues SK, Klionsky DJ. Atg41/Icy2 regulates autophagosome formation. <i>Autophagy</i> 2015;11(12):2288–99.	doi:10.1080/15548627.2015.1107692 PMID 26565778 PMC4835205	–	2	1	2	1	1	2	ChIP
1251	Baek 2016 Baek SH, Kwon EY, Kim YH, Hahn JS. Metabolic engineering and adaptive evolution for efficient production of D-lactic acid in <i>Saccharomyces cerevisiae</i> . <i>Appl Microbiol Biotechnol</i> 2016;100(6):2737–48.	doi:10.1007/s00253-015-7174-0 PMID 26596574	–	2	1	2	2	–	1	RT-PCR - wt vs mutant promoter sequence
1252	Yarrington 2016 Yarrington RM, Goodrum JM, Stillman DJ. Nucleosomes Are Essential for Proper Regulation of a Multigated Promoter in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2016;202(2):551–63.	doi:10.1534/genetics.115.183715 PMID 26627840 PMC4788235	–	2	2	2	1	1	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1253	Kumar 2015 Kumar NV, Yang J, Pillai JK, et al., Tamás MJ. Arsenic Directly Binds to and Activates the Yeast AP-1-Like Transcription Factor Yap8. <i>Mol Cell Biol</i> 2015;36(6):913–22.	doi:10.1128/MCB.00842-15 PMID 26711267 PMC4810476	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1254	Vega 2016 Vega M, Riera A, Fernández-Cid A, Herrero P, Moreno F. Hexokinase 2 Is an Intracellular Glucose Sensor of Yeast Cells That Maintains the Structure and Activity of Mig1 Protein Repressor Complex. <i>J Biol Chem</i> 2016;291(14):7267–85.	doi:10.1074/jbc.M115.711408 PMID 26865637 PMC4817161	–	2	2	2	2	–	1	ChIP
1255	García 2016 García R, Sanz AB, Rodríguez-Peña JM, Nombela C, Arroyo J. Rlm1 mediates positive autoregulatory transcriptional feedback that is essential for Slt2-dependent gene expression. <i>J Cell Sci</i> 2016;129(8):1649–60.	doi:10.1242/jcs.180190 PMID 26933180	–	2	1	2	2	–	1	ChIP
1256	Umekawa 2016 Umekawa M, Ujihara M, Makishima K, Yamamoto S, Takematsu H, Wakayama M. The signaling pathways underlying starvation-induced upregulation of α -mannosidase Ams1 in <i>Saccharomyces cerevisiae</i> . <i>Biochim Biophys Acta</i> 2016;1860(6):1192–201.	doi:10.1016/j.bbagen.2016.02.018 PMID 26947009	–	2	2	2	–	2	1	RT-PCR - wt vs TF mutant
1257	Chen 2016 Chen Y, Luo W, Gong R, et al., Xiao D. Improved ethyl caproate production of Chinese liquor yeast by overexpressing fatty acid synthesis genes with OPI1 deletion. <i>J Ind Microbiol Biotechnol</i> 2016;43(9):1261–70.	doi:10.1007/s10295-016-1795-x PMID 27344573	–	2	1	2	–	2	1	RT-PCR - wt vs TF mutant
1258	Yadav 2016 Yadav PK, Rajasekharan R. Misregulation of a DDHD Domain-containing Lipase Causes Mitochondrial Dysfunction in Yeast. <i>J Biol Chem</i> 2016;291(35):18562–81.	doi:10.1074/jbc.M116.733378 PMID 27402848 PMC5000100	–	2	2	2	2	–	1	EMSA
1259	Lee 2017 Lee YM, Kim E, An J, et al., Kim W. Dissection of the HOG pathway activated by hydrogen peroxide in <i>Saccharomyces cerevisiae</i> . <i>Environ Microbiol</i> 2017;19(2):584–597.	doi:10.1111/1462-2920.13499 PMID 27554843	–	2	1	2	–	2	2	Western blot - wt vs TF mutant
1260	Chia 2016 Chia M, van Werven FJ. Temporal Expression of a Master Regulator Drives Synchronous Sporulation in Budding Yeast. <i>G3 (Bethesda)</i> 2016;6(11):3553–3560.	doi:10.1534/g3.116.034983 PMID 27605516 PMC5100854	–	2	2	2	1	1	2	ChIP
1261	Regenberg 2016 Regenberg B, Hanghøj KE, Andersen KS, Boomsma JJ. Clonal yeast biofilms can reap competitive advantages through cell differentiation without being obligatorily multicellular. <i>Proc Biol Sci</i> 2016;283(1842).	doi:10.1098/rspb.2016.1303 PMID 27807261 PMC5124089	–	2	2	2	–	2	2	RNA FISH
1262	Kliewe 2017 Kliewe F, Engelhardt M, Aref R, Schüller HJ. Promoter recruitment of corepressors Sin3 and Cyc8 by activator proteins of the yeast <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 2017;63(4):739–750.	doi:10.1007/s00294-017-0677-8 PMID 28175933	–	2	1	2	2	–	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1263	Godinho 2017 Godinho CP, Mira NP, Cabrito TR, et al., Sá-Correia I. Yeast response and tolerance to benzoic acid involves the Gcn4- and Stp1-regulated multidrug/multixenobiotic resistance transporter Tpo1. <i>Appl Microbiol Biotechnol</i> 2017;101(12): 5005–5018.	doi:10.1007/s00253-017-8277-6 PMID 28409382 PMC5486834	–	2	2	2	–	2	1	RT-PCR - wt vs TF mutant
1264	Kimura 2017 Kimura Y, Irie K, Mizuno T. Expression control of the AMPK regulatory subunit and its functional significance in yeast ER stress response. <i>Sci Rep</i> 2017;7:46713.	doi:10.1038/srep46713 PMID 28429799 PMC5399461	–	2	1	1	–	2	1	RT-PCR - wt vs TF mutant
1265	Hovsepian 2017 Hovsepian J, Defenouillère Q, Albanèse V, et al., Léon S. Multilevel regulation of an α -arrestin by glucose depletion controls hexose transporter endocytosis. <i>J Cell Biol</i> 2017; 216(6):1811–1831.	doi:10.1083/jcb.201610094 PMID 28468835 PMC5461024	–	2	2	2	–	2	1	lacZ - wt vs TF mutant
1266	Teunissen 2017 Teunissen JHM, Crooijmans ME, Teunisse PPP, van Heusden GPH. Lack of 14-3-3 proteins in <i>Saccharomyces cerevisiae</i> results in cell-to-cell heterogeneity in the expression of Pho4-regulated genes SPL2 and PHO84. <i>BMC Genomics</i> 2017;18(1):701.	doi:10.1186/s12864-017-4105-8 PMID 28877665 PMC5588707	–	2	1	1	–	2	1	GFP - wt vs TF mutant
1267	Gaspar 2017 Gaspar ML, Chang YF, Jesch SA, Aregullin M, Henry SA. Interaction between repressor Opi1p and ER membrane protein Scs2p facilitates transit of phosphatidic acid from the ER to mitochondria and is essential for INO1 gene expression in the presence of choline. <i>J Biol Chem</i> 2017;292(45): 18713–18728.	doi:10.1074/jbc.M117.809970 PMID 28924045 PMC5683188	–	2	1	1	–	2	2	RT-PCR - wt vs TF mutant
1268	Anton 2017 Anton C, Zanolari B, Arcones I, et al., Roncero C. Involvement of the exomer complex in the polarized transport of Ena1 required for <i>Saccharomyces cerevisiae</i> survival against toxic cations. <i>Mol Biol Cell</i> 2017;28(25):3672–3685.	doi:10.1091/mbc.E17-09-0549 PMID 29021337 PMC5706994	–	2	1	1	–	2	2	Western blot - wt vs TF mutant
1269	Krakowiak 2018 Krakowiak J, Zheng X, Patel N, et al., Pincus D. Hsf1 and Hsp70 constitute a two-component feedback loop that regulates the yeast heat shock response. <i>Elife</i> 2018;7.	doi:10.7554/eLife.31668 PMID 29393852 PMC5809143	–	2	1	2	–	2	1	Western blot - wt vs TF mutant
1270	Mizuno 2018 Mizuno T, Nakamura M, Irie K. Induction of Ptp2 and Cmp2 protein phosphatases is crucial for the adaptive response to ER stress in <i>Saccharomyces cerevisiae</i> . <i>Sci Rep</i> 2018;8(1):13078.	doi:10.1038/s41598-018-31413-6 PMID 30166606 PMC6117328	–	2	1	1	–	2	2	RT-PCR - wt vs TF mutant
1271	Sanz 2018 Sanz AB, García R, Rodríguez-Peña JM, Nombela C, Arroyo J. Slr2 MAPK association with chromatin is required for transcriptional activation of Rlm1 dependent genes upon cell wall stress. <i>Biochim Biophys Acta Gene Regul Mech</i> 2018; 1861(11):1029–1039.	doi:10.1016/j.bbaggm.2018.09.005 PMID 30343693	–	2	1	1	1	1	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1272	Xu 2019 Xu H, Fang T, Yan H, Jiang L. The protein kinase Cmk2 negatively regulates the calcium/calciueurin signalling pathway and expression of calcium pump genes PMR1 and PMC1 in budding yeast. <i>Cell Commun Signal</i> 2019;17(1):7.	doi:10.1186/s12964-019-0320-z PMID 30665402 PMC6341702	–	2	1	2	–	2	1	lacZ - wt vs TF mutant
1273	Donovan 2019 Donovan BT, Huynh A, Ball DA, et al., Lenstra TL. Live-cell imaging reveals the interplay between transcription factors, nucleosomes, and bursting. <i>EMBO J</i> 2019;38(12).	doi:10.15252/embj.2018100809 PMID 31101674 PMC6576174	–	2	1	1	2	–	1	3D Orbital Tracking Microscopy
1274	Tatip 2020 Tatip S, Taggart J, Wang Y, MacDiarmid CW, Eide DJ. Changes in transcription start sites of Zap1-regulated genes during zinc deficiency: Implications for HNT1 gene regulation. <i>Mol Microbiol</i> 2020;113(1):285–296.	doi:10.1111/mmi.14416 PMID 31692084 PMC7007328	–	2	1	2	–	2	1	RT-PCR - wt vs. TF mutant
1275	Steidle 2020 Steidle EA, Morrisette VA, Fujimaki K, et al., Rolfes RJ. The InsP(7) phosphatase Siw14 regulates inositol pyrophosphate levels to control localization of the general stress response transcription factor Msn2. <i>J Biol Chem</i> 2020;295(7):2043–2056.	doi:10.1074/jbc.RA119.012148 PMID 31848224 PMC7029108	–	2	1	1	–	2	2	RT-PCR - wt vs TF mutant
1276	Willis 2020 Willis SD, Hanley SE, Beishke T, Tati PD, Cooper KF. Ubiquitin-proteasome-mediated cyclin C degradation promotes cell survival following nitrogen starvation. <i>Mol Biol Cell</i> 2020;31(10):1015–1031.	doi:10.1091/mbc.E19-11-0622 PMID 32160104 PMC7346723	–	2	1	1	–	2	2	RT-PCR - wt vs TF mutant
1277	Kim 2021 Kim B, Lee Y, Choi H, Huh WK. The trehalose-6-phosphate phosphatase Tps2 regulates ATG8 transcription and autophagy in <i>Saccharomyces cerevisiae</i> . <i>Autophagy</i> 2021;17(4):1013–1027.	doi:10.1080/15548627.2020.1746592 PMID 32240040 PMC8078664	–	2	1	1	–	2	2	RT-PCR - wt vs TF mutant
1278	Yarrington 2020 Yarrington RM, Yu Y, Yan C, Bai L, Stillman DJ. A Role for Mediator Core in Limiting Coactivator Recruitment in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2020;215(2):407–420.	doi:10.1534/genetics.120.303254 PMID 32327563 PMC7268993	–	2	1	1	1	1	2	ChIP
1279	Galkina 2020 Galkina KV, Finkelberg JM, Markova OV, et al., Knorre DA. Protonophore FCCP provides fitness advantage to PDR-deficient yeast cells. <i>J Bioenerg Biomembr</i> 2020;52(5):383–395.	doi:10.1007/s10863-020-09849-1 PMID 32808242	–	2	2	2	–	2	1	GFP - wt vs TF double mutant
1280	Walden 2020 Walden EA, Fong RY, Pham TT, et al., Baetz K. Phenomic screen identifies a role for the yeast lysine acetyltransferase NuA4 in the control of Bcy1 subcellular localization, glycogen biosynthesis, and mitochondrial morphology. <i>PLoS Genet</i> 2020;16(11):e1009220.	doi:10.1371/journal.pgen.1009220 PMID 33253187 PMC7728387	–	2	2	2	–	2	1	RT-PCR - wt vs TF mutant and western blot

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1281	Stokdyk 2021 Stokdyk K, Berroyer A, Grami ZA, Kim N. The activity of yeast Apn2 AP endonuclease at uracil-derived AP sites is dependent on the major carbon source. <i>Curr Genet</i> 2021;67(2):283–294.	doi:10.1007/s00294-020-01141-4 PMID 33386486 PMC8035244	–	2	1	1	–	2	2	RT-PCR - wt vs. TF double mutant
1282	Swagatika 2021 Swagatika S, Tomar RS. ABC transporter Pdr5 is required for cantharidin resistance in <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2021;553:141–147.	doi:10.1016/j.bbrc.2021.03.074 PMID 33770579	–	2	2	2	–	2	1	RT-PCR - wt vs TF double mutant
1283	Guaragnella 2021 Guaragnella N, Ždravlević M, Palková Z, Giannattasio S. Analysis of Mitochondrial Retrograde Signaling in Yeast Model Systems. <i>Methods Mol Biol</i> 2021;2276:87–102.	doi:10.1007/978-1-0716-1266-8_6 PMID 34060034	–	2	2	2	–	2	1	Western blot - wt vs TF mutant
1284	Nasuno 2021 Nasuno R, Yoshioka N, Yoshikawa Y, Takagi H. Cysteine residues in the fourth zinc finger are important for activation of the nitric oxide-inducible transcription factor Fzfl in the yeast <i>Saccharomyces cerevisiae</i> . <i>Genes Cells</i> 2021;26(10):823–829.	doi:10.1111/gtc.12885 PMID 34245655	–	2	1	2	–	2	1	RT-PCR - wt vs. TF mutant
1285	Delago 2021 Delago A, Gregor R, Dubinsky L, et al., Meijler MM. A Bacterial Quorum Sensing Molecule Elicits a General Stress Response in <i>Saccharomyces cerevisiae</i> . <i>Front Microbiol</i> 2021;12:632658.	doi:10.3389/fmicb.2021.632658 PMID 34603220 PMC8481950	–	2	2	2	–	2	1	flow cytometry - GFP fluorescence
1286	Dokládál 2021 Dokládál L, Stumpe M, Hu Z, Jaquenoud M, Dengjel J, De Virgilio C. Phosphoproteomic responses of TORC1 target kinases reveal discrete and convergent mechanisms that orchestrate the quiescence program in yeast. <i>Cell Rep</i> 2021;37(13):110149.	doi:10.1016/j.celrep.2021.110149 PMID 34965436	–	2	1	1	–	2	1	BETA-galactosidase assay
1287	Li 2022 Li L, Wang J, Yang Z, et al., Lou H. Metabolic remodeling maintains a reducing environment for rapid activation of the yeast DNA replication checkpoint. <i>EMBO J</i> 2022;41(4):e108290.	doi:10.15252/embj.2021108290 PMID 35028974 PMC8844976	–	2	2	2	–	2	1	RT-PCR - wt vs. TF double mutant
1288	Ciriacy 1976 Ciriacy M. Cis-dominant regulatory mutations affecting the formation of glucose-repressible alcohol dehydrogenase (ADHII) in <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1976;145(3):327–33.	doi:10.1007/BF00325831 PMID 781520	–	1	1	1	–	1	1	Enzyme activity - wt vs TF mutant
1289	Bowman 1992 Bowman SB, Zaman Z, Collinson LP, Brown AJ, Dawes IW. Positive regulation of the LPD1 gene of <i>Saccharomyces cerevisiae</i> by the HAP2/HAP3/HAP4 activation system. <i>Mol Gen Genet</i> 1992;231(2):296–303.	doi:10.1007/BF00279803 PMID 1310523	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1290	Meyers 1992 Meyers S, Schauer W, Balzi E, Wagner M, Goffeau A, Golin J. Interaction of the yeast pleiotropic drug resistance genes PDR1 and PDR5. <i>Curr Genet</i> 1992;21(6):431–6.	doi:10.1007/BF00351651 PMID 1319843	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1291	Silve 1992 Silve S, Rhode PR, Coll B, Campbell J, Poyton RO. ABF1 is a phosphoprotein and plays a role in carbon source control of COX6 transcription in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1992;12(9):4197–208.	doi:10.1128/mcb.12.9.4197-4208.1992 PMID 1324416 PMC360325	–	1	1	1	–	–	1	–
1292	Remacle 1992 Remacle JE, Holmberg S. A REB1-binding site is required for GCN4-independent ILV1 basal level transcription and can be functionally replaced by an ABF1-binding site. <i>Mol Cell Biol</i> 1992;12(12):5516–26.	doi:10.1128/mcb.12.12.5516-5526.1992 PMID 1448083 PMC360489	–	1	1	1	1	–	1	EMSA
1293	Lanker 1992 Lanker S, Bushman JL, Hinnebusch AG, Trachsel H, Mueller PP. Autoregulation of the yeast lysyl-tRNA synthetase gene GCD5/KRS1 by translational and transcriptional control mechanisms. <i>Cell</i> 1992;70(4):647–57.	doi:10.1016/0092-8674(92)90433-d PMID 1505029	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1294	Pizzagalli 1992 Pizzagalli A, Piatti S, Derossi D, Gander I, Plevani P, Lucchini G. Positive cis-acting regulatory sequences mediate proper control of POL1 transcription in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 1992;21(3):183–9.	doi:10.1007/BF00336839 PMID 1563043	–	1	1	1	–	–	1	–
1295	Barbarić 1992 Barbarić S, Fascher KD, Hörz W. Activation of the weakly regulated PHO8 promoter in <i>S. cerevisiae</i> : chromatin transition and binding sites for the positive regulatory protein PHO4. <i>Nucleic Acids Res</i> 1992;20(5):1031–8.	doi:10.1093/nar/20.5.1031 PMID 1567507 PMC312087	–	1	1	1	1	–	1	EMSA
1296	Zitomer 1992 Zitomer RS, Lowry CV. Regulation of gene expression by oxygen in <i>Saccharomyces cerevisiae</i> . <i>Microbiol Rev</i> 1992;56(1):1–11.	doi:10.1128/mr.56.1.1-11.1992 PMID 1579104 PMC372851	–	1	1	1	–	–	1	–
1297	Park 1992 Park HD, Luche RM, Cooper TG. The yeast UME6 gene product is required for transcriptional repression mediated by the CAR1 URS1 repressor binding site. <i>Nucleic Acids Res</i> 1992;20(8):1909–15.	doi:10.1093/nar/20.8.1909 PMID 1579492 PMC312306	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1298	Schneider 1991 Schneider JC, Guarente L. Regulation of the yeast CYT1 gene encoding cytochrome c1 by HAP1 and HAP2/3/4. <i>Mol Cell Biol</i> 1991;11(10):4934–42.	doi:10.1128/mcb.11.10.4934-4942.1991 PMID 1656218 PMC361470	–	1	1	1	1	–	1	EMSA
1299	Mösch 1990 Mösch HU, Graf R, Schmidheini T, Braus G. Three GCN4 responsive elements act synergistically as upstream and as TATA-like elements in the yeast TRP4 promoter. <i>EMBO J</i> 1990;9(9):2951–7.	doi:10.1002/j.1460-2075.1990.tb07487.x PMID 1697266 PMC552012	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1300	Verdière 1991 Verdière J, Gaisne M, Labbe-Bois R. CYP1 (HAP1) is a determinant effector of alternative expression of heme-dependent transcribed genes in yeast [corrected]. <i>Mol Gen Genet</i> 1991;228(1-2):300-6.	doi:10.1007/BF00282480 PMID 1715975	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1301	Lydall 1991 Lydall D, Ammerer G, Nasmyth K. A new role for MCM1 in yeast: cell cycle regulation of SW15 transcription. <i>Genes Dev</i> 1991;5(12B):2405-19.	doi:10.1101/gad.5.12b.2405 PMID 1752436	–	1	1	1	1	–	1	DNA footprinting
1302	Bitter 1991 Bitter GA, Chang KK, Egan KM. A multi-component upstream activation sequence of the <i>Saccharomyces cerevisiae</i> glyceraldehyde-3-phosphate dehydrogenase gene promoter. <i>Mol Gen Genet</i> 1991;231(1):22-32.	doi:10.1007/BF00293817 PMID 1753943	–	1	1	1	1	–	1	EMSA
1303	Primig 1991 Primig M, Winkler H, Ammerer G. The DNA binding and oligomerization domain of MCM1 is sufficient for its interaction with other regulatory proteins. <i>EMBO J</i> 1991;10(13):4209-18.	doi:10.1002/j.1460-2075.1991.tb04999.x PMID 1756728 PMC453173	–	1	1	1	1	–	1	EMSA
1304	Jones 1991 Jones DG, Reusser U, Braus GH. Molecular cloning, characterization and analysis of the regulation of the ARO2 gene, encoding chorismate synthase, of <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 1991;5(9):2143-52.	doi:10.1111/j.1365-2958.1991.tb02144.x PMID 1837329	–	1	1	1	1	–	1	DNA footprinting
1305	Hagen 1991 Hagen DC, McCaffrey G, Sprague GF Jr. Pheromone response elements are necessary and sufficient for basal and pheromone-induced transcription of the FUS1 gene of <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1991;11(6):2952-61.	doi:10.1128/mcb.11.6.2952-2961.1991 PMID 1903837 PMC360123	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1306	Kovari 1991 Kovari LZ, Cooper TG. Participation of ABF-1 protein in expression of the <i>Saccharomyces cerevisiae</i> CAR1 gene. <i>J Bacteriol</i> 1991;173(20):6332-8.	doi:10.1128/jb.173.20.6332-6338.1991 PMID 1917865 PMC208964	–	1	1	1	1	–	1	EMSA
1307	Gralla 1991 Gralla EB, Thiele DJ, Silar P, Valentine JS. ACE1, a copper-dependent transcription factor, activates expression of the yeast copper, zinc superoxide dismutase gene. <i>Proc Natl Acad Sci U S A</i> 1991;88(19):8558-62.	doi:10.1073/pnas.88.19.8558 PMID 1924315 PMC52548	–	1	1	1	1	–	1	DNA footprinting
1308	Mösch 1991 Mösch HU, Scheier B, Lahti R, Mäntsälä P, Braus GH. Transcriptional activation of yeast nucleotide biosynthetic gene ADE4 by GCN4. <i>J Biol Chem</i> 1991;266(30):20453-6.	PMID 1939099	–	1	1	1	–	1	1	Enzyme activity - wt vs TF mutant
1309	Hayashi 1991 Hayashi N, Oshima Y. Specific cis-acting sequence for PHO8 expression interacts with PHO4 protein, a positive regulatory factor, in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1991;11(2):785-94.	doi:10.1128/mcb.11.2.785-794.1991 PMID 1990283 PMC359730	–	1	1	1	1	–	1	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1310	Carri 1991 Carri MT, Galiazzo F, Ciriolo MR, Rotilio G. Evidence for co-regulation of Cu,Zn superoxide dismutase and metallothionein gene expression in yeast through transcriptional control by copper via the ACE 1 factor. <i>FEBS Lett</i> 1991;278(2):263–6.	doi:10.1016/0014-5793(91)80131-1 PMID 1991520	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1311	Morris 1991 Morris ME, Jinks-Robertson S. Nucleotide sequence of the LYS2 gene of <i>Saccharomyces cerevisiae</i> : homology to <i>Bacillus brevis</i> tyrocidine synthetase 1. <i>Gene</i> 1991;98(1):141–5.	doi:10.1016/0378-1119(91)90117-t PMID 2013406	–	1	1	1	–	–	1	–
1312	Smith 1991 Smith BJ, Yaffe MP. A mutation in the yeast heat-shock factor gene causes temperature-sensitive defects in both mitochondrial protein import and the cell cycle. <i>Mol Cell Biol</i> 1991;11(5):2647–55.	doi:10.1128/mcb.11.5.2647-2655.1991 PMID 2017170 PMC360034	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1313	Lodi 1991 Lodi T, Guiard B. Complex transcriptional regulation of the <i>Saccharomyces cerevisiae</i> CYB2 gene encoding cytochrome b2: CYP1(HAP1) activator binds to the CYB2 upstream activation site UAS1-B2. <i>Mol Cell Biol</i> 1991;11(7):3762–72.	doi:10.1128/mcb.11.7.3762-3772.1991 PMID 2046677 PMC361146	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1314	Boorstein 1990 Boorstein WR, Craig EA. Structure and regulation of the SSA4 HSP70 gene of <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 1990;265(31):18912–21.	PMID 2121731	–	1	1	1	–	–	1	–
1315	Oh 1990 Oh D, Hopper JE. Transcription of a yeast phosphoglucomutase isozyme gene is galactose inducible and glucose repressible. <i>Mol Cell Biol</i> 1990;10(4):1415–22.	doi:10.1128/mcb.10.4.1415-1422.1990 PMID 2138705 PMC362244	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1316	Nehlin 1990 Nehlin JO, Ronne H. Yeast MIG1 repressor is related to the mammalian early growth response and Wilms' tumour finger proteins. <i>EMBO J</i> 1990;9(9):2891–8.	doi:10.1002/j.1460-2075.1990.tb07479.x PMID 2167835 PMC552004	–	1	1	1	1	–	1	EMSA
1317	Stout 1990 Stout VG, Iandolo JJ. Chromosomal gene transfer during conjugation by <i>Staphylococcus aureus</i> is mediated by transposon-facilitated mobilization. <i>J Bacteriol</i> 1990;172(10):6148–50.	doi:10.1128/jb.172.10.6148-6150.1990 PMID 2170345 PMC526945	–	1	1	1	–	1	1	Enzyme activity - wt vs TF mutant
1318	Chesley 1990 Chesley SF, Schatzki AD, DeUrrutia J, Greenblatt DJ, Shader RI, Miller LG. Cocaine augments peripheral benzodiazepine binding in humans. <i>J Clin Psychiatry</i> 1990;51(10):404–6.	PMID 2170346	–	1	1	1	–	1	1	Enzyme activity - wt vs TF mutant
1319	Engelbrecht 1990 Engelbrecht J, Roeder GS. MER1, a yeast gene required for chromosome pairing and genetic recombination, is induced in meiosis. <i>Mol Cell Biol</i> 1990;10(5):2379–89.	doi:10.1128/mcb.10.5.2379-2389.1990 PMID 2183032 PMC360586	–	1	1	1	–	1	1	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1320	Magdolen 1990 Magdolen V, Oechsner U, Trommler P, Bandlow W. Transcriptional control by galactose of a yeast gene encoding a protein homologous to mammalian aldo/keto reductases. <i>Gene</i> 1990;90(1):105–14.	doi:10.1016/0378-1119(90)90445-w PMID 2199324	–	1	1	1	–	–	1	–
1321	Thomas 1990 Thomas D, Barbey R, Surdin-Kerjan Y. Gene-enzyme relationship in the sulfate assimilation pathway of <i>Saccharomyces cerevisiae</i> . Study of the 3'-phosphoadenylylsulfate reductase structural gene. <i>J Biol Chem</i> 1990;265(26):15518–24.	PMID 2203779	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1322	Ju 1990 Ju QD, Morrow BE, Warner JR. REB1, a yeast DNA-binding protein with many targets, is essential for growth and bears some resemblance to the oncogene myb. <i>Mol Cell Biol</i> 1990;10(10):5226–34.	doi:10.1128/mcb.10.10.5226-5234.1990 PMID 2204808 PMC361205	–	1	1	1	1	–	1	DNA footprinting
1323	Uemura 1990 Uemura H, Fraenkel DG. gcr2, a new mutation affecting glycolytic gene expression in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1990;10(12):6389–96.	doi:10.1128/mcb.10.12.6389-6396.1990 PMID 2247062 PMC362915	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1324	Mellor 1990 Mellor J, Jiang W, Funk M, et al., Philippsen P. CPF1, a yeast protein which functions in centromeres and promoters. <i>EMBO J</i> 1990;9(12):4017–26.	doi:10.1002/j.1460-2075.1990.tb07623.x PMID 2249662 PMC552174	–	1	1	1	1	–	1	DNA footprinting
1325	Butler 1990 Butler G, Dawes IW, McConnell DJ. TUF factor binds to the upstream region of the pyruvate decarboxylase structural gene (PDC1) of <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1990;223(3):449–56.	doi:10.1007/BF00264453 PMID 2270085	–	1	1	1	1	–	1	EMSA
1326	Buchman 1990 Buchman AR, Kornberg RD. A yeast ARS-binding protein activates transcription synergistically in combination with other weak activating factors. <i>Mol Cell Biol</i> 1990;10(3):887–97.	doi:10.1128/mcb.10.3.887-897.1990 PMID 2406570 PMC360926	–	1	1	1	1	–	1	EMSA
1327	Capieaux 1989 Capieaux E, Vignais ML, Sentenac A, Goffeau A. The yeast H ⁺ -ATPase gene is controlled by the promoter binding factor TUF. <i>J Biol Chem</i> 1989;264(13):7437–46.	PMID 2523395	–	1	1	1	1	–	1	EMSA
1328	Hodge 1989 Hodge MR, Kim G, Singh K, Cumsy MG. Inverse regulation of the yeast COX5 genes by oxygen and heme. <i>Mol Cell Biol</i> 1989;9(5):1958–64.	doi:10.1128/mcb.9.5.1958-1964.1989 PMID 2546055 PMC362987	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1329	Errede 1989 Errede B, Ammerer G. STE12, a protein involved in cell-type-specific transcription and signal transduction in yeast, is part of protein-DNA complexes. <i>Genes Dev</i> 1989;3(9):1349–61.	doi:10.1101/gad.3.9.1349 PMID 2558054	–	1	1	1	1	–	1	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1330	Yoshida 1989 Yoshida K, Ogawa N, Oshima Y. Function of the PHO regulatory genes for repressible acid phosphatase synthesis in <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1989;217(1):40–6.	doi:10.1007/BF00330940 PMID 2671650	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1331	Tice-Baldwin 1989 Tice-Baldwin K, Fink GR, Arndt KT. BAS1 has a Myb motif and activates HIS4 transcription only in combination with BAS2. <i>Science</i> 1989;246(4932):931–5.	doi:10.1126/science.2683089 PMID 2683089	–	1	1	1	–	–	1	–
1332	Marczak 1989 Marczak JE, Brandriss MC. Isolation of constitutive mutations affecting the proline utilization pathway in <i>Saccharomyces cerevisiae</i> and molecular analysis of the PUT3 transcriptional activator. <i>Mol Cell Biol</i> 1989;9(11):4696–705.	doi:10.1128/mcb.9.11.4696-4705.1989 PMID 2689861 PMC363616	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1333	Cooper 1987 Cooper TG, Chisholm VT, Cho HJ, Yoo HS. Allantoin transport in <i>Saccharomyces cerevisiae</i> is regulated by two induction systems. <i>J Bacteriol</i> 1987;169(10):4660–7.	doi:10.1128/jb.169.10.4660-4667.1987 PMID 2820939 PMC213836	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1334	Prezant 1987 Prezant T, Pfeifer K, Guarente L. Organization of the regulatory region of the yeast CYC7 gene: multiple factors are involved in regulation. <i>Mol Cell Biol</i> 1987;7(9):3252–9.	doi:10.1128/mcb.7.9.3252-3259.1987 PMID 2823123 PMC367962	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1335	Winkler 1988 Winkler H, Adam G, Mattes E, Schanz M, Hartig A, Ruis H. Co-ordinate control of synthesis of mitochondrial and non-mitochondrial hemoproteins: a binding site for the HAP1 (CYP1) protein in the UAS region of the yeast catalase T gene (CTT1). <i>EMBO J</i> 1988;7(6):1799–804.	doi:10.1002/j.1460-2075.1988.tb03011.x PMID 2844525 PMC457171	–	1	1	1	1	–	1	Northern blot - wt vs mutant promoter sequence
1336	Genbauffe 1986 Genbauffe FS, Cooper TG. Induction and repression of the urea amidolyase gene in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1986;6(11):3954–64.	doi:10.1128/mcb.6.11.3954-3964.1986 PMID 3025621 PMC367159	–	1	1	1	–	1	1	Slot blot - wt vs TF mutant
1337	Pfeifer 1987 Pfeifer K, Prezant T, Guarente L. Yeast HAP1 activator binds to two upstream activation sites of different sequence. <i>Cell</i> 1987;49(1):19–27.	doi:10.1016/0092-8674(87)90751-3 PMID 3030565	–	1	1	1	1	–	1	EMSA
1338	Hamil 1988 Hamil KG, Nam HG, Fried HM. Constitutive transcription of yeast ribosomal protein gene TCM1 is promoted by uncommon cis- and trans-acting elements. <i>Mol Cell Biol</i> 1988; 8(10):4328–41.	doi:10.1128/mcb.8.10.4328-4341.1988 PMID 3054514 PMC365506	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1339	Xiao 1988 Xiao W, Rank GH. The yeast ILV2 gene is under general amino acid control. <i>Genome</i> 1988;30(6):984–6.	doi:10.1139/g88-156 PMID 3069583	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1340	Sengstag 1988 Sengstag C, Hinnen A. A 28-bp segment of the <i>Saccharomyces cerevisiae</i> PHO5 upstream activator sequence confers phosphate control to the CYC1-lacZ gene fusion. <i>Gene</i> 1988; 67(2):223–8.	doi:10.1016/0378-1119(88)90399-x PMID 3139496	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1341	Kassir 1988 Kassir Y, Granot D, Simchen G. IME1, a positive regulator gene of meiosis in <i>S. cerevisiae</i> . <i>Cell</i> 1988;52(6):853–62.	doi:10.1016/0092-8674(88)90427-8 PMID 3280136	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1342	Johnston 1987 Johnston LH, White JH, Johnson AL, Lucchini G, Plevani P. The yeast DNA polymerase I transcript is regulated in both the mitotic cell cycle and in meiosis and is also induced after DNA damage. <i>Nucleic Acids Res</i> 1987;15(13):5017–30.	doi:10.1093/nar/15.13.5017 PMID 3299263 PMC305944	–	1	1	1	–	–	1	–
1343	McCaffrey 1987 McCaffrey G, Clay FJ, Kelsay K, Sprague GF Jr. Identification and regulation of a gene required for cell fusion during mating of the yeast <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1987;7(8):2680–90.	doi:10.1128/mcb.7.8.2680-2690.1987 PMID 3313002 PMC367884	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1344	Breeden 1987 Breeden L, Nasmyth K. Cell cycle control of the yeast HO gene: cis- and trans-acting regulators. <i>Cell</i> 1987;48(3):389–97.	doi:10.1016/0092-8674(87)90190-5 PMID 3542227	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1345	Hope 1985 Hope IA, Struhl K. GCN4 protein, synthesized in vitro, binds HIS3 regulatory sequences: implications for general control of amino acid biosynthetic genes in yeast. <i>Cell</i> 1985;43(1):177–88.	doi:10.1016/0092-8674(85)90022-4 PMID 3907851	–	1	1	1	1	–	1	EMSA
1346	Shimada 1985 Shimada H, Fukasawa T. Controlled transcription of the yeast regulatory gene GAL80. <i>Gene</i> 1985;39(1):1–9.	doi:10.1016/0378-1119(85)90100-3 PMID 3908221	–	1	1	1	–	–	1	–
1347	Lucchini 1984 Lucchini G, Hinnebusch AG, Chen C, Fink GR. Positive regulatory interactions of the HIS4 gene of <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1984;4(7):1326–33.	doi:10.1128/mcb.4.7.1326-1333.1984 PMID 6095062 PMC368915	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1348	Post-Beittenmiller 1984 Post-Beittenmiller MA, Hamilton RW, Hopper JE. Regulation of basal and induced levels of the MEL1 transcript in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1984;4(7):1238–45.	doi:10.1128/mcb.4.7.1238-1245.1984 PMID 6209559 PMC368904	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1349	Guarente 1984 Guarente L, Lalonde B, Gifford P, Alani E. Distinctly regulated tandem upstream activation sites mediate catabolite repression of the CYC1 gene of <i>S. cerevisiae</i> . <i>Cell</i> 1984;36(2):503–11.	doi:10.1016/0092-8674(84)90243-5 PMID 6319028	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1350	Nasmyth 1983 Nasmyth K. Molecular analysis of a cell lineage. <i>Nature</i> 1983; 302(5910):670–6.	doi:10.1038/302670a0 PMID 6339953	–	1	1	1	–	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1351	Klig 1984 Klig LS, Henry SA. Isolation of the yeast INO1 gene: located on an autonomously replicating plasmid, the gene is fully regulated. <i>Proc Natl Acad Sci U S A</i> 1984;81(12):3816–20.	doi:10.1073/pnas.81.12.3816 PMID 6374665 PMC345311	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1352	Melcher 1995 Melcher K, Rose M, Künzler M, Braus GH, Entian KD. Molecular analysis of the yeast SER1 gene encoding 3-phosphoserine aminotransferase: regulation by general control and serine repression. <i>Curr Genet</i> 1995;27(6):501–8.	doi:10.1007/BF00314439 PMID 7553933	–	1	1	1	1	–	1	DNA footprinting
1353	De Winde 1995 De Winde JH, Grivell LA. Regulation of mitochondrial biogenesis in <i>Saccharomyces cerevisiae</i> . Intricate interplay between general and specific transcription factors in the promoter of the QCR8 gene. <i>Eur J Biochem</i> 1995;233(1):200–8.	doi:10.1111/j.1432-1033.1995.200_1.x PMID 7588747	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1354	Ozsarac 1995 Ozsarac N, Bhattacharyya M, Dawes IW, Clancy MJ. The SPR3 gene encodes a sporulation-specific homologue of the yeast CDC3/10/11/12 family of bud neck microfilaments and is regulated by ABFI. <i>Gene</i> 1995;164(1):157–62.	doi:10.1016/0378-1119(95)00438-c PMID 7590307	–	1	1	1	1	–	1	EMSA
1355	Johnston 1995 Johnston LH, Johnson AL. The DNA repair genes RAD54 and UNG1 are cell cycle regulated in budding yeast but MCB promoter elements have no essential role in the DNA damage response. <i>Nucleic Acids Res</i> 1995;23(12):2147–52.	doi:10.1093/nar/23.12.2147 PMID 7610042 PMC307001	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1356	Jung 1995 Jung SY, Yoo HY, Kim YH, Kim J, Rho HM. The glucose-dependent transactivation activity of ABF1 on the expression of the TDH3 gene in yeast. <i>Curr Genet</i> 1995;27(4):312–7.	doi:10.1007/BF00352099 PMID 7614553	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1357	Chelstowska 1995 Chelstowska A, Butow RA. RTG genes in yeast that function in communication between mitochondria and the nucleus are also required for expression of genes encoding peroxisomal proteins. <i>J Biol Chem</i> 1995;270(30):18141–6.	doi:10.1074/jbc.270.30.18141 PMID 7629125	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1358	Yoo 1995 Yoo HY, Jung SY, Kim YH, Kim J, Jung G, Rho HM. Transcriptional control of the <i>Saccharomyces cerevisiae</i> ADH1 gene by autonomously replicating sequence binding factor 1. <i>Curr Microbiol</i> 1995;31(3):163–8.	doi:10.1007/BF00293548 PMID 7663308	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1359	Giardina 1995 Giardina C, Lis JT. Dynamic protein-DNA architecture of a yeast heat shock promoter. <i>Mol Cell Biol</i> 1995;15(5):2737–44.	doi:10.1128/MCB.15.5.2737 PMID 7739554 PMC230504	–	1	1	1	1	–	1	DNA footprinting

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1360	Rai 1995 Rai R, Daugherty JR, Cooper TG. UASNTR functioning in combination with other UAS elements underlies exceptional patterns of nitrogen regulation in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 1995;11(3):247–60.	doi:10.1002/yea.320110307 PMID 7785325	–	1	1	1	–	–	1	–
1361	Wemmie 1994 Wemmie JA, Szczypka MS, Thiele DJ, Moye-Rowley WS. Cadmium tolerance mediated by the yeast AP-1 protein requires the presence of an ATP-binding cassette transporter-encoding gene, YCF1. <i>J Biol Chem</i> 1994;269(51):32592–7.	PMID 7798263	–	1	1	1	1	–	1	DNA footprinting
1362	Hu 1995 Hu Y, Cooper TG, Kohlhaw GB. The <i>Saccharomyces cerevisiae</i> Leu3 protein activates expression of GDH1, a key gene in nitrogen assimilation. <i>Mol Cell Biol</i> 1995;15(1):52–7.	doi:10.1128/MCB.15.1.52 PMID 7799961 PMC231907	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1363	Henry 1994 Henry YA, López MC, Gibbs JM, et al., Stanway CA. The yeast protein Gcr1p binds to the PGK UAS and contributes to the activation of transcription of the PGK gene. <i>Mol Gen Genet</i> 1994;245(4):506–11.	doi:10.1007/BF00302263 PMID 7808400	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1364	Ram 1995 Ram AF, Brekelmans SS, Oehlen LJ, Klis FM. Identification of two cell cycle regulated genes affecting the beta 1,3-glucan content of cell walls in <i>Saccharomyces cerevisiae</i> . <i>FEBS Lett</i> 1995;358(2):165–70.	doi:10.1016/0014-5793(94)01418-z PMID 7828729	–	1	1	1	–	–	1	–
1365	Covitz 1994 Covitz PA, Song W, Mitchell AP. Requirement for RGR1 and SIN4 in RME1-dependent repression in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 1994;138(3):577–86.	doi:10.1093/genetics/138.3.577 PMID 7851756 PMC1206209	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1366	Hedges 1995 Hedges D, Proft M, Entian KD. CAT8, a new zinc cluster-encoding gene necessary for derepression of gluconeogenic enzymes in the yeast <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1995;15(4):1915–22.	doi:10.1128/MCB.15.4.1915 PMID 7891685 PMC230417	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1367	Brazas 1993 Brazas RM, Stillman DJ. The Swi5 zinc-finger and Grf10 homeodomain proteins bind DNA cooperatively at the yeast HO promoter. <i>Proc Natl Acad Sci U S A</i> 1993;90(23):11237–41.	doi:10.1073/pnas.90.23.11237 PMID 7902583 PMC47957	–	1	1	1	1	–	1	EMSA
1368	Navarro 1994 Navarro B, Igual JC. ADR1 and SNF1 mediate different mechanisms in transcriptional regulation of yeast POT1 gene. <i>Biochem Biophys Res Commun</i> 1994;202(2):960–6.	doi:10.1006/bbrc.1994.2023 PMID 7914081	–	1	1	1	–	1	1	Luciferase - wt vs TF mutant
1369	Wu 1994 Wu AL, Moye-Rowley WS. GSH1, which encodes gamma-glutamylcysteine synthetase, is a target gene for yAP-1 transcriptional regulation. <i>Mol Cell Biol</i> 1994;14(9):5832–9.	doi:10.1128/mcb.14.9.5832-5839.1994 PMID 7915005 PMC359109	–	1	1	1	–	1	1	lacZ - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1370	Culotta 1994 Culotta VC, Howard WR, Liu XF. CRS5 encodes a metallothionein-like protein in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 1994;269(41):25295–302.	PMID 7929222	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1371	García-Arranz 1994 García-Arranz M, Maldonado AM, Mazón MJ, Portillo F. Transcriptional control of yeast plasma membrane H(+)-ATPase by glucose. Cloning and characterization of a new gene involved in this regulation. <i>J Biol Chem</i> 1994; 269(27):18076–82.	PMID 8027068	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1372	Lambert 1994 Lambert JR, Bilanchone VW, Cumsy MG. The ORD1 gene encodes a transcription factor involved in oxygen regulation and is identical to IXR1, a gene that confers cisplatin sensitivity to <i>Saccharomyces cerevisiae</i> . <i>Proc Natl Acad Sci U S A</i> 1994;91(15):7345–9.	doi:10.1073/pnas.91.15.7345 PMID 8041793 PMC44396	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1373	Cunningham 1994 Cunningham TS, Dorrington RA, Cooper TG. The UGA4 UASNTR site required for GLN3-dependent transcriptional activation also mediates DAL80-responsive regulation and DAL80 protein binding in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1994;176(15):4718–25.	doi:10.1128/jb.176.15.4718-4725.1994 PMID 8045902 PMC196294	–	1	1	1	1	–	1	EMSA
1374	Gounalaki 1994 Gounalaki N, Thireos G. Yap1p, a yeast transcriptional activator that mediates multidrug resistance, regulates the metabolic stress response. <i>EMBO J</i> 1994;13(17):4036–41.	doi:10.1002/j.1460-2075.1994.tb06720.x PMID 8076599 PMC395324	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1375	Murakami 1993 Murakami Y, Hurwitz J. Functional interactions between SV40 T antigen and other replication proteins at the replication fork. <i>J Biol Chem</i> 1993;268(15):11008–17.	PMID 8098707	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1376	Ahmed 1993 Ahmed M, Borsch CM, Neyfakh AA, Schuldiner S. Mutants of the <i>Bacillus subtilis</i> multidrug transporter Bmr with altered sensitivity to the antihypertensive alkaloid reserpine. <i>J Biol Chem</i> 1993;268(15):11086–9.	PMID 8098708	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1377	Osheroov 1993 Osheroov N, Gazit A, Gilon C, Levitzki A. Selective inhibition of the epidermal growth factor and HER2/neu receptors by tyrphostins. <i>J Biol Chem</i> 1993;268(15):11134–42.	PMID 8098709	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1378	Kuroda 1994 Kuroda S, Otaka S, Fujisawa Y. Fermentable and nonfermentable carbon sources sustain constitutive levels of expression of yeast triosephosphate dehydrogenase 3 gene from distinct promoter elements. <i>J Biol Chem</i> 1994;269(8):6153–62.	PMID 8119960	–	1	1	1	–	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1379	Carmen 1994 Carmen AA, Holland MJ. The upstream repression sequence from the yeast enolase gene ENO1 is a complex regulatory element that binds multiple trans-acting factors including REB1. <i>J Biol Chem</i> 1994;269(13):9790–7.	PMID 8144571	–	1	1	1	1	–	1	EMSA
1380	Ganter 1993 Ganter B, Tan S, Richmond TJ. Genomic footprinting of the promoter regions of STE2 and STE3 genes in the yeast <i>Saccharomyces cerevisiae</i> . <i>J Mol Biol</i> 1993;234(4):975–87.	doi:10.1006/jmbi.1993.1652 PMID 8263944	–	1	1	1	1	–	1	DNA footprinting
1381	Kuge 1994 Kuge S, Jones N. YAP1 dependent activation of TRX2 is essential for the response of <i>Saccharomyces cerevisiae</i> to oxidative stress by hydroperoxides. <i>EMBO J</i> 1994;13(3):655–64.	doi:10.1002/j.1460-2075.1994.tb06304.x PMID 8313910 PMC394856	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1382	ElBerry 1993 ElBerry HM, Majumdar ML, Cunningham TS, Sumrada RA, Cooper TG. Regulation of the urea active transporter gene (DUR3) in <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1993;175(15):4688–98.	doi:10.1128/jb.175.15.4688-4698.1993 PMID 8335627 PMC204920	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1383	Pavlik 1993 Pavlik P, Simon M, Schuster T, Ruis H. The glycerol kinase (GUT1) gene of <i>Saccharomyces cerevisiae</i> : cloning and characterization. <i>Curr Genet</i> 1993;24(1–2):21–5.	doi:10.1007/BF00324660 PMID 8358828	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1384	Kuenzler 1993 Kuenzler M, Balmelli T, Egli CM, Paravicini G, Braus GH. Cloning, primary structure, and regulation of the HIS7 gene encoding a bifunctional glutamine amidotransferase: cyclase from <i>Saccharomyces cerevisiae</i> . <i>J Bacteriol</i> 1993;175(17):5548–58.	doi:10.1128/jb.175.17.5548-5558.1993 PMID 8366040 PMC206611	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1385	Pittman 1993 Pittman D, Lu W, Malone RE. Genetic and molecular analysis of REC114, an early meiotic recombination gene in yeast. <i>Curr Genet</i> 1993;23(4):295–304.	doi:10.1007/BF00310890 PMID 8385581	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1386	Hagen 1993 Hagen DC, Bruhn L, Westby CA, Sprague GF Jr. Transcription of alpha-specific genes in <i>Saccharomyces cerevisiae</i> : DNA sequence requirements for activity of the coregulator alpha 1. <i>Mol Cell Biol</i> 1993;13(11):6866–75.	doi:10.1128/mcb.13.11.6866-6875.1993 PMID 8413280 PMC364749	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1387	Kovari 1993 Kovari LZ, Kovari I, Cooper TG. Participation of RAP1 protein in expression of the <i>Saccharomyces cerevisiae</i> arginase (CAR1) gene. <i>J Bacteriol</i> 1993;175(4):941–51.	doi:10.1128/jb.175.4.941-951.1993 PMID 8432717 PMC193005	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1388	Mountain 1993 Mountain HA, Byström AS, Korch C. The general amino acid control regulates MET4, which encodes a methionine-pathway-specific transcriptional activator of <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 1993;7(2):215–28.	doi:10.1111/j.1365-2958.1993.tb01113.x PMID 8446029	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1389	Rodicio 1993 Rodicio R, Heinisch JJ, Hollenberg CP. Transcriptional control of yeast phosphoglycerate mutase-encoding gene. <i>Gene</i> 1993;125(2):125–33.	doi:10.1016/0378-1119(93)90319-x PMID 8462867	–	1	1	1	1	–	1	EMSA
1390	Sabová 1993 Sabová L, Zeman I, Supek F, Kolarov J. Transcriptional control of AAC3 gene encoding mitochondrial ADP/ATP translocator in <i>Saccharomyces cerevisiae</i> by oxygen, heme and ROX1 factor. <i>Eur J Biochem</i> 1993;213(1):547–53.	doi:10.1111/j.1432-1033.1993.tb17793.x PMID 8477727	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1391	Kirkman-Correia 1993 Kirkman-Correia C, Stroke IL, Fields S. Functional domains of the yeast STE12 protein, a pheromone-responsive transcriptional activator. <i>Mol Cell Biol</i> 1993;13(6):3765–72.	doi:10.1128/mcb.13.6.3765-3772.1993 PMID 8497278 PMC359857	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1392	Morgan 1995 Morgan BA, Bouquin N, Merrill GF, Johnston LH. A yeast transcription factor bypassing the requirement for SBF and DSC1/MBF in budding yeast has homology to bacterial signal transduction proteins. <i>EMBO J</i> 1995;14(22):5679–89.	doi:10.1002/j.1460-2075.1995.tb00255.x PMID 8521825 PMC394683	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1393	Koch 1996 Koch C, Schleiffer A, Ammerer G, Nasmyth K. Switching transcription on and off during the yeast cell cycle: Cln/Cdc28 kinases activate bound transcription factor SBF (Swi4/Swi6) at start, whereas Clb/Cdc28 kinases displace it from the promoter in G2. <i>Genes Dev</i> 1996;10(2):129–41.	doi:10.1101/gad.10.2.129 PMID 8566747	–	1	1	1	1	–	1	DNA footprinting
1394	Packham 1996 Packham EA, Graham IR, Chambers A. The multifunctional transcription factors Abf1p, Rap1p and Reb1p are required for full transcriptional activation of the chromosomal PGK gene in <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1996;250(3):348–56.	doi:10.1007/BF02174393 PMID 8602150	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1395	Gailus-Durner 1996 Gailus-Durner V, Xie J, Chintamaneni C, Vershon AK. Participation of the yeast activator Abf1 in meiosis-specific expression of the HOP1 gene. <i>Mol Cell Biol</i> 1996;16(6):2777–86.	doi:10.1128/MCB.16.6.2777 PMID 8649386 PMC231269	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1396	Graham 1996 Graham IR, Chambers A. Rap1p is a negative regulator of the RAP1 gene. <i>Curr Genet</i> 1996;30(2):93–100.	doi:10.1007/s002940050106 PMID 8660466	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1397	Lambrechts 1996 Lambrechts MG, Sollitti P, Marmur J, Pretorius IS. A multicopy suppressor gene, MSS10, restores STA2 expression in <i>Saccharomyces cerevisiae</i> strains containing the STA10 repressor gene. <i>Curr Genet</i> 1996;29(6):523–9.	PMID 8662191	–	1	1	1	–	1	1	Northern blot - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1398	Luo 1996 Luo Y, Karpichev IV, Kohanski RA, Small GM. Purification, identification, and properties of a <i>Saccharomyces cerevisiae</i> oleate-activated upstream activating sequence-binding protein that is involved in the activation of POX1. <i>J Biol Chem</i> 1996; 271(20):12068–75.	doi:10.1074/jbc.271.20.12068 PMID 8662598	–	1	1	1	1	–	1	EMSA
1399	Muchardt 1996 Muchardt C, Reyes JC, Bourachot B, Leguoy E, Yaniv M. The hbrm and BRG-1 proteins, components of the human SNF/SWI complex, are phosphorylated and excluded from the condensed chromosomes during mitosis. <i>EMBO J</i> 1996;15(13): 3394–402.	PMID 8670841 PMC451903	–	1	1	1	1	–	1	In vivo footprinting WT vs TF mutant vs TF overexpressing mutant
1400	Oechsner 1996 Oechsner U, Bandlow W. Interactions of the yeast centromere and promoter factor, Cpflp, with the cytochrome c1 upstream region and functional implications on regulated gene expression. <i>Nucleic Acids Res</i> 1996;24(12):2395–403.	doi:10.1093/nar/24.12.2395 PMID 8710512 PMC145936	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1401	Lambrechts 1996 Lambrechts MG, Bauer FF, Marmur J, Pretorius IS. Muc1, a mucin-like protein that is regulated by Mss10, is critical for pseudohyphal differentiation in yeast. <i>Proc Natl Acad Sci U S A</i> 1996;93(16):8419–24.	doi:10.1073/pnas.93.16.8419 PMID 8710886 PMC38686	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1402	Kobayashi 1996 Kobayashi O, Suda H, Ohtani T, Sone H. Molecular cloning and analysis of the dominant flocculation gene FLO8 from <i>Saccharomyces cerevisiae</i> . <i>Mol Gen Genet</i> 1996;251(6):707–15.	doi:10.1007/BF02174120 PMID 8757402	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1403	Amillet 1996 Amillet JM, Buisson N, Labbe-Bois R. Characterization of an upstream activation sequence and two Rox1p-responsive sites controlling the induction of the yeast HEM13 gene by oxygen and heme deficiency. <i>J Biol Chem</i> 1996;271(40):24425–32.	doi:10.1074/jbc.271.40.24425 PMID 8798700	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1404	McLean 1995 McLean M, Hubberstey AV, Bouman DJ, Pece N, Mastrangelo P, Wildeman AG. Organization of the <i>Saccharomyces cerevisiae</i> actin gene UAS: functional significance of reiterated REB1 binding sites and AT-rich elements. <i>Mol Microbiol</i> 1995;18(4):605–14.	doi:10.1111/j.1365-2958.1995.mmi_18040605.x PMID 8817483	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1405	Grant 1996 Grant CM, Collinson LP, Roe JH, Dawes IW. Yeast glutathione reductase is required for protection against oxidative stress and is a target gene for yAP-1 transcriptional regulation. <i>Mol Microbiol</i> 1996;21(1):171–9.	doi:10.1046/j.1365-2958.1996.6351340.x PMID 8843443	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1406	Holmberg 1996 Holmberg S, Schjerling P. Cha4p of <i>Saccharomyces cerevisiae</i> activates transcription via serine/threonine response elements. <i>Genetics</i> 1996;144(2):467–78.	doi:10.1093/genetics/144.2.467 PMID 8889513 PMC1207543	–	1	1	1	1	–	1	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1407	Igual 1996 Igual JC, Johnson AL, Johnston LH. Coordinated regulation of gene expression by the cell cycle transcription factor Swi4 and the protein kinase C MAP kinase pathway for yeast cell integrity. <i>EMBO J</i> 1996;15(18):5001–13.	PMID 8890173 PMC452238	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1408	Nouraini 1996 Nouraini S, Hu J, McBroom LD, Friesen JD. Mutations in an Abflp binding site in the promoter of yeast RPO26 shift the transcription start sites and reduce the level of RPO26 mRNA. <i>Yeast</i> 1996;12(13):1339–50.	doi:10.1002/(SICI)1097-0061(199610)12:13%3C1339::AID-YEA31%3E3.0.CO;2-C PMID 8923739	–	1	1	1	–	–	1	–
1409	Nikawa 1996 Nikawa J, Akiyoshi M, Hirata S, Fukuda T. Saccharomyces cerevisiae IRE2/HAC1 is involved in IRE1-mediated KAR2 expression. <i>Nucleic Acids Res</i> 1996;24(21):4222–6.	doi:10.1093/nar/24.21.4222 PMID 8932376 PMC146251	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1410	Erkine 1996 Erkine AM, Adams CC, Diken T, Gross DS. Heat shock factor gains access to the yeast HSC82 promoter independently of other sequence-specific factors and antagonizes nucleosomal repression of basal and induced transcription. <i>Mol Cell Biol</i> 1996;16(12):7004–17.	doi:10.1128/MCB.16.12.7004 PMID 8943356 PMC231704	–	1	1	1	1	–	1	DNA footprinting
1411	Toyn 1997 Toyn JH, Johnson AL, Donovan JD, Toone WM, Johnston LH. The Swi5 transcription factor of Saccharomyces cerevisiae has a role in exit from mitosis through induction of the cdk-inhibitor Sic1 in telophase. <i>Genetics</i> 1997;145(1):85–96.	doi:10.1093/genetics/145.1.85 PMID 9017392 PMC1207787	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1412	Ozsarac 1997 Ozsarac N, Straffon MJ, Dalton HE, Dawes IW. Regulation of gene expression during meiosis in Saccharomyces cerevisiae: SPR3 is controlled by both ABFI and a new sporulation control element. <i>Mol Cell Biol</i> 1997;17(3):1152–9.	doi:10.1128/MCB.17.3.1152 PMID 9032242 PMC231840	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1413	Rowen 1997 Rowen DW, Esiobu N, Magasanik B. Role of GATA factor Nil2p in nitrogen regulation of gene expression in Saccharomyces cerevisiae. <i>J Bacteriol</i> 1997;179(11):3761–6.	doi:10.1128/jb.179.11.3761-3766.1997 PMID 9171427 PMC179175	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1414	Pemberton 1997 Pemberton LF, Blobel G. Characterization of the Wtm proteins, a novel family of Saccharomyces cerevisiae transcriptional modulators with roles in meiotic regulation and silencing. <i>Mol Cell Biol</i> 1997;17(8):4830–41.	doi:10.1128/MCB.17.8.4830 PMID 9234739 PMC232335	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1415	Alarco 1997 Alarco AM, Balan I, Talibi D, Mainville N, Raymond M. AP1-mediated multidrug resistance in Saccharomyces cerevisiae requires FLR1 encoding a transporter of the major facilitator superfamily. <i>J Biol Chem</i> 1997;272(31):19304–13.	doi:10.1074/jbc.272.31.19304 PMID 9235926	–	1	1	1	–	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1416	Szeto 1997 Szeto L, Fafalios MK, Zhong H, Vershon AK, Broach JR. Alpha2p controls donor preference during mating type interconversion in yeast by inactivating a recombinational enhancer of chromosome III. <i>Genes Dev</i> 1997;11(15):1899–911.	doi:10.1101/gad.11.15.1899 PMID 9271114 PMC316413	–	1	1	1	–	–	1	–
1417	Webber 1997 Webber AL, Lambrechts MG, Pretorius IS. MSS11, a novel yeast gene involved in the regulation of starch metabolism. <i>Curr Genet</i> 1997;32(4):260–6.	doi:10.1007/s002940050275 PMID 9342405	–	1	1	1	–	1	1	Enzyme activity - wt vs TF mutant
1418	Springer 1997 Springer C, Krappmann S, Künzler M, Zmasek C, Braus GH. Regulation of the yeast HIS7 gene by the global transcription factor Abf1p. <i>Mol Gen Genet</i> 1997;256(2):136–46.	doi:10.1007/s004380050555 PMID 9349705	–	1	1	1	1	–	1	EMSA
1419	Kitagaki 1997 Kitagaki H, Shimoi H, Itoh K. Identification and analysis of a static culture-specific cell wall protein, Tir1p/Srp1p in <i>Saccharomyces cerevisiae</i> . <i>Eur J Biochem</i> 1997;249(1):343–9.	doi:10.1111/j.1432-1033.1997.t01-1-00343.x PMID 9363789	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1420	Alepuz 1997 Alepuz PM, Cunningham KW, Estruch F. Glucose repression affects ion homeostasis in yeast through the regulation of the stress-activated ENA1 gene. <i>Mol Microbiol</i> 1997;26(1):91–8.	doi:10.1046/j.1365-2958.1997.5531917.x PMID 9383192	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1421	Byrd 1998 Byrd C, Turner GC, Varshavsky A. The N-end rule pathway controls the import of peptides through degradation of a transcriptional repressor. <i>EMBO J</i> 1998;17(1):269–77.	doi:10.1093/emboj/17.1.269 PMID 9427760 PMC1170377	–	1	1	1	–	1	1	Northern blot - wt vs TF overexpression
1422	Angermayr 1997 Angermayr M, Bandlow W. The general regulatory factor Reb1p controls basal, but not Gal4p-mediated, transcription of the GCY1 gene in yeast. <i>Mol Gen Genet</i> 1997;256(6):682–9.	doi:10.1007/s004380050616 PMID 9435793	–	1	1	1	1	–	1	EMSA
1423	Tennyson 1998 Tennyson CN, Lee J, Andrews BJ. A role for the Pcl9-Pho85 cyclin-cdk complex at the M/G1 boundary in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 1998;28(1):69–79.	doi:10.1046/j.1365-2958.1998.00773.x PMID 9593297	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1424	Wang 1998 Wang KL, Warner JR. Positive and negative autoregulation of REB1 transcription in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1998;18(7):4368–76.	doi:10.1128/MCB.18.7.4368 PMID 9632820 PMC109020	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1425	Menéndez 1998 Menéndez J, Gancedo C. Regulatory regions in the promoters of the <i>Saccharomyces cerevisiae</i> PYC1 and PYC2 genes encoding isoenzymes of pyruvate carboxylase. <i>FEMS Microbiol Lett</i> 1998;164(2):345–52.	doi:10.1111/j.1574-6968.1998.tb13108.x PMID 9682484	–	1	1	1	1	–	1	EMSA
1426	Schroeder 1998 Schroeder SC, Weil PA. Biochemical and genetic characterization of the dominant positive element driving transcription of the yeast TBP-encoding gene, SPT15. <i>Nucleic Acids Res</i> 1998;26(18):4186–95.	doi:10.1093/nar/26.18.4186 PMID 9722639 PMC147844	–	1	1	1	1	–	1	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1427	Song 1998 Song W, Carlson M. Srb/mediator proteins interact functionally and physically with transcriptional repressor Sfl1. <i>EMBO J</i> 1998;17(19):5757–65.	doi:10.1093/emboj/17.19.5757 PMID 9755175 PMC1170903	–	1	1	1	1	–	1	Oligonucleotide affinity chromatography
1428	Xu 1998 Xu Z, Norris D. The SFP1 gene product of <i>Saccharomyces cerevisiae</i> regulates G2/M transitions during the mitotic cell cycle and DNA-damage response. <i>Genetics</i> 1998;150(4):1419–28.	doi:10.1093/genetics/150.4.1419 PMID 9832520 PMC1460418	–	1	1	1	–	1	1	Northern blot - wt vs TF overexpression
1429	Iraqui 1999 Iraqui I, Vissers S, Bernard F, et al., André B. Amino acid signaling in <i>Saccharomyces cerevisiae</i> : a permease-like sensor of external amino acids and F-Box protein Grr1p are required for transcriptional induction of the AGP1 gene, which encodes a broad-specificity amino acid permease. <i>Mol Cell Biol</i> 1999;19(2):989–1001.	doi:10.1128/MCB.19.2.989 PMID 9891035 PMC116030	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1430	Zhu 1998 Zhu G, Davis TN. The fork head transcription factor Hcm1p participates in the regulation of SPC110, which encodes the calmodulin-binding protein in the yeast spindle pole body. <i>Biochim Biophys Acta</i> 1998;1448(2):236–44.	doi:10.1016/s0167-4889(98)00135-9 PMID 9920414	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1431	Park 1999 Park SH, Koh SS, Chun JH, Hwang HJ, Kang HS. Nrg1 is a transcriptional repressor for glucose repression of STA1 gene expression in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1999;19(3):2044–50.	doi:10.1128/MCB.19.3.2044 PMID 10022891 PMC83997	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1432	Avram 1999 Avram D, Leid M, Bakalinsky AT. Fzf1p of <i>Saccharomyces cerevisiae</i> is a positive regulator of SSU1 transcription and its first zinc finger region is required for DNA binding. <i>Yeast</i> 1999;15(6):473–80.	<a href="https://doi.org/10.1002/(SICI)1097-0061(199904)15:6<473::AID-YEA388>3.0.CO;2-Y">doi:10.1002/(SICI)1097-0061(199904)15:6<473::AID-YEA388>3.0.CO;2-Y PMID 10234785	–	1	1	1	1	–	1	DNA footprinting
1433	Zaman 1999 Zaman Z, Bowman SB, Kornfeld GD, Brown AJ, Dawes IW. Transcription factor GCN4 for control of amino acid biosynthesis also regulates the expression of the gene for lipoamide dehydrogenase. <i>Biochem J</i> 1999;340 (Pt 3)(Pt 3):855–62.	PMID 10359673 PMC1220320	–	1	1	1	1	–	1	EMSA
1434	Baumgartner 1999 Baumgartner U, Hamilton B, Piskacek M, Ruis H, Rottensteiner H. Functional analysis of the Zn(2)Cys(6) transcription factors Oaf1p and Pip2p. Different roles in fatty acid induction of beta-oxidation in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 1999;274(32):22208–16.	doi:10.1074/jbc.274.32.22208 PMID 10428786	–	1	1	1	1	–	1	DNA footprinting

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1435	Basrai 1999 Basrai MA, Velculescu VE, Kinzler KW, Hieter P. NORF5/HUG1 is a component of the MEC1-mediated checkpoint response to DNA damage and replication arrest in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 1999;19(10):7041–9.	doi:10.1128/MCB.19.10.7041 PMID 10490641 PMC84699	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1436	Graham 1999 Graham IR, Haw RA, Spink KG, Halden KA, Chambers A. In vivo analysis of functional regions within yeast Rap1p. <i>Mol Cell Biol</i> 1999;19(11):7481–90.	doi:10.1128/MCB.19.11.7481 PMID 10523636 PMC84746	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1437	Rahner 1999 Rahner A, Hiesinger M, Schüller HJ. Dereglulation of gluconeogenic structural genes by variants of the transcriptional activator Cat8p of the yeast <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 1999;34(1):146–56.	doi:10.1046/j.1365-2958.1999.01588.x PMID 10540293	–	1	1	1	–	1	1	Northern blot - wt vs TF overexpression
1438	Sato 1999 Sato T, Lopez MC, Sugioka S, Jigami Y, Baker HV, Uemura H. The E-box DNA binding protein Sgc1p suppresses the gcr2 mutation, which is involved in transcriptional activation of glycolytic genes in <i>Saccharomyces cerevisiae</i> . <i>FEBS Lett</i> 1999;463(3):307–11.	doi:10.1016/s0014-5793(99)01654-3 PMID 10606743	–	1	1	1	1	–	1	DNA footprinting
1439	Suh 2000 Suh JK, Robertus JD. Yeast flavin-containing monooxygenase is induced by the unfolded protein response. <i>Proc Natl Acad Sci U S A</i> 2000;97(1):121–6.	doi:10.1073/pnas.97.1.121 PMID 10618381 PMC26626	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1440	Lascaris 2000 Lascaris RF, Groot E, Hoen PB, Mager WH, Planta RJ. Different roles for abf1p and a T-rich promoter element in nucleosome organization of the yeast RPS28A gene. <i>Nucleic Acids Res</i> 2000;28(6):1390–6.	doi:10.1093/nar/28.6.1390 PMID 10684934 PMC111049	–	1	1	1	1	–	1	DNA footprinting
1441	Kunoh 2000 Kunoh T, Kaneko Y, Harashima S. YHP1 encodes a new homeoprotein that binds to the IME1 promoter in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 2000;16(5):439–49.	<a href="https://doi.org/10.1002/(SICI)1097-0061(20000330)16:5<439::AID-YEA536>3.0.CO;2-M">doi:10.1002/(SICI)1097-0061(20000330)16:5<439::AID-YEA536>3.0.CO;2-M PMID 10705372	–	1	1	1	–	1	1	Enzyme activity - wt vs TF mutant
1442	Eriksson 2000 Eriksson P, Alipour H, Adler L, Blomberg A. Rap1p-binding sites in the <i>saccharomyces cerevisiae</i> GPD1 promoter are involved in its response to NaCl. <i>J Biol Chem</i> 2000;275(38):29368–76.	doi:10.1074/jbc.M001663200 PMID 10842169	–	1	1	1	1	–	1	EMSA
1443	D'Alessio 2000 D'Alessio M, Brandriss MC. Cross-pathway regulation in <i>Saccharomyces cerevisiae</i> : activation of the proline utilization pathway by Gal1p in vivo. <i>J Bacteriol</i> 2000;182(13):3748–53.	doi:10.1128/JB.182.13.3748-3753.2000 PMID 10850990 PMC94546	–	1	1	1	–	1	1	lacZ - wt vs TF overexpression
1444	MacDiarmid 2000 MacDiarmid CW, Gaither LA, Eide D. Zinc transporters that regulate vacuolar zinc storage in <i>Saccharomyces cerevisiae</i> . <i>EMBO J</i> 2000;19(12):2845–55.	doi:10.1093/emboj/19.12.2845 PMID 10856230 PMC203372	–	1	1	1	–	1	1	lacZ - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1445	Bird 2000 Bird AJ, Zhao H, Luo H, et al., Eide DJ. A dual role for zinc fingers in both DNA binding and zinc sensing by the Zap1 transcriptional activator. <i>EMBO J</i> 2000;19(14):3704–13.	doi:10.1093/emboj/19.14.3704 PMID 10899124 PMC313982	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1446	Miyabe 2000 Miyabe S, Izawa S, Inoue Y. Expression of ZRC1 coding for suppressor of zinc toxicity is induced by zinc-starvation stress in Zap1-dependent fashion in <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2000;276(3):879–84.	doi:10.1006/bbrc.2000.3580 PMID 11027563	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1447	Nguyễn 2001 Nguyễn DT, Alarco AM, Raymond M. Multiple Yap1p-binding sites mediate induction of the yeast major facilitator FLR1 gene in response to drugs, oxidants, and alkylating agents. <i>J Biol Chem</i> 2001;276(2):1138–45.	doi:10.1074/jbc.M008377200 PMID 11056165	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1448	Chan 2001 Chan TF, Bertram PG, Ai W, Zheng XF. Regulation of APG14 expression by the GATA-type transcription factor Gln3p. <i>J Biol Chem</i> 2001;276(9):6463–7.	doi:10.1074/jbc.M008162200 PMID 11096087	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1449	Zhang 2001 Zhang X, Cui Z, Miyakawa T, Moye-Rowley WS. Cross-talk between transcriptional regulators of multidrug resistance in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2001;276(12):8812–9.	doi:10.1074/jbc.M010686200 PMID 11134057	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1450	Nginyi 2001 Nginyi JM, Duncan JL, Mellor DJ, et al., Gatongi PM. Epidemiology of parasitic gastrointestinal nematode infections of ruminants on smallholder farms in central Kenya. <i>Res Vet Sci</i> 2001;70(1):33–9.	doi:10.1053/rvsc.2000.0438 PMID 11170849	–	1	1	1	1	–	1	EMSA
1451	Maxon 2001 Maxon ME, Herskowitz I. Ash1p is a site-specific DNA-binding protein that actively represses transcription. <i>Proc Natl Acad Sci U S A</i> 2001;98(4):1495–500.	doi:10.1073/pnas.98.4.1495 PMID 11171979 PMC29285	–	1	1	1	1	–	1	EMSA
1452	Nishizawa 2001 Nishizawa M, Tanabe M, Yabuki N, Kitada K, Toh-E A. Pho85 kinase, a yeast cyclin-dependent kinase, regulates the expression of UGP1 encoding UDP-glucose pyrophosphorylase. <i>Yeast</i> 2001;18(3):239–49.	<a href="https://doi.org/10.1002/1097-0061(200102)18:3<239::AID-YEA664>3.0.CO;2-0">doi:10.1002/1097-0061(200102)18:3<239::AID-YEA664>3.0.CO;2-0 PMID 11180457	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1453	Li 2001 Li Q, Johnston SA. Are all DNA binding and transcription regulation by an activator physiologically relevant? <i>Mol Cell Biol</i> 2001;21(7):2467–74.	doi:10.1128/MCB.21.7.2467-2474.2001 PMID 11259595 PMC86879	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1454	Theodoris 2001 Theodoris G, Bisson LF. DDSE: downstream targets of the SNF3 signal transduction pathway. <i>FEMS Microbiol Lett</i> 2001;197(1):73–7.	doi:10.1111/j.1574-6968.2001.tb10585.x PMID 11287149	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1455	Miled 2001 Miled C, Mann C, Faye G. Xbp1-mediated repression of CLB gene expression contributes to the modifications of yeast cell morphology and cell cycle seen during nitrogen-limited growth. <i>Mol Cell Biol</i> 2001;21(11):3714–24.	doi:10.1128/MCB.21.11.3714-3724.2001 PMID 11340165 PMC87007	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1456	Huang 2001 Huang S, Jeffery DA, Anthony MD, O’Shea EK. Functional analysis of the cyclin-dependent kinase inhibitor Pho81 identifies a novel inhibitory domain. <i>Mol Cell Biol</i> 2001;21(19):6695–705.	doi:10.1128/MCB.21.19.6695-6705.2001 PMID 11533256 PMC99814	–	1	1	1	–	–	1	–
1457	Hollenhorst 2001 Hollenhorst PC, Pietz G, Fox CA. Mechanisms controlling differential promoter-occupancy by the yeast forkhead proteins Fkh1p and Fkh2p: implications for regulating the cell cycle and differentiation. <i>Genes Dev</i> 2001;15(18):2445–56.	doi:10.1101/gad.906201 PMID 11562353 PMC312786	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1458	Li 2002 Li S, Dean S, Li Z, Horecka J, Deschenes RJ, Fassler JS. The eukaryotic two-component histidine kinase Sln1p regulates OCH1 via the transcription factor, Skn7p. <i>Mol Biol Cell</i> 2002;13(2):412–24.	doi:10.1091/mbc.01-09-0434 PMID 11854400 PMC65637	–	1	1	1	1	–	1	EMSA
1459	Valerius 2002 Valerius O, Brendel C, Düvel K, Braus GH. Multiple factors prevent transcriptional interference at the yeast ARO4-HIS7 locus. <i>J Biol Chem</i> 2002;277(24):21440–5.	doi:10.1074/jbc.M201841200 PMID 11937506	–	1	1	1	1	–	1	Northern blot - wt vs mutant promoter sequence
1460	Tachibana 2002 Tachibana T, Astumi S, Shioda R, Ueno M, Uritani M, Ushimaru T. A novel non-conventional heat shock element regulates expression of MDJ1 encoding a DnaJ homolog in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2002;277(25):22140–6.	doi:10.1074/jbc.M201267200 PMID 11940587	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1461	Dormer 2002 Dormer UH, Westwater J, Stephen DW, Jamieson DJ. Oxidant regulation of the <i>Saccharomyces cerevisiae</i> GSH1 gene. <i>Biochim Biophys Acta</i> 2002;1576(1–2):23–9.	doi:10.1016/S00167-4781(02)00248-8 PMID 12031480	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1462	De Sanctis 2002 De Sanctis V, La Terra S, Bianchi A, et al., Negri R. In vivo topography of Rap1p-DNA complex at <i>Saccharomyces cerevisiae</i> TEF2 UAS(RPG) during transcriptional regulation. <i>J Mol Biol</i> 2002;318(2):333–49.	doi:10.1016/S0022-2836(02)00110-9 PMID 12051841	–	1	1	1	1	–	1	DNA footprinting
1463	Park 2002 Park YU, Hwang O, Kim J. Two-hybrid cloning and characterization of OSH3, a yeast oxysterol-binding protein homolog. <i>Biochem Biophys Res Commun</i> 2002;293(2):733–40.	doi:10.1016/S0006-291X(02)00288-7 PMID 12054531	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1464	Brons 2002 Brons JF, De Jong M, Valens M, Grivell LA, Bolotin-Fukuhara M, Blom J. Dissection of the promoter of the HAP4 gene in <i>S. cerevisiae</i> unveils a complex regulatory framework of transcriptional regulation. <i>Yeast</i> 2002;19(11):923–32.	doi:10.1002/yea.886 PMID 12125049	–	1	1	1	–	1	1	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1465	Tomás-Cobos 2002 Tomás-Cobos L, Sanz P. Active Snf1 protein kinase inhibits expression of the <i>Saccharomyces cerevisiae</i> HXT1 glucose transporter gene. <i>Biochem J</i> 2002;368(Pt 2):657–63.	doi:10.1042/BJ20020984 PMID 12220226 PMC1223017	–	1	1	1	–	–	1	–
1466	Zhu 2002 Zhu H, Pan S, Gu S, Bradbury EM, Chen X. Amino acid residue specific stable isotope labeling for quantitative proteomics. <i>Rapid Commun Mass Spectrom</i> 2002;16(22):2115–23.	doi:10.1002/rcm.831 PMID 12415544	–	1	1	1	–	1	1	Proteomics - wt vs TF mutant
1467	Yamamoto 2002 Yamamoto K, Jigami Y. Mutation of TRS130, which encodes a component of the TRAPP II complex, activates transcription of OCH1 in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 2002;42(2):85–93.	doi:10.1007/s00294-002-0336-5 PMID 12478387	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1468	Mosley 2003 Mosley AL, Lakshmanan J, Aryal BK, Ozcan S. Glucose-mediated phosphorylation converts the transcription factor Rgt1 from a repressor to an activator. <i>J Biol Chem</i> 2003;278(12):10322–7.	doi:10.1074/jbc.M212802200 PMID 12527758	–	1	1	1	1	–	1	ChIP
1469	MacDiarmid 2003 MacDiarmid CW, Milanick MA, Eide DJ. Induction of the ZRC1 metal tolerance gene in zinc-limited yeast confers resistance to zinc shock. <i>J Biol Chem</i> 2003;278(17):15065–72.	doi:10.1074/jbc.M300568200 PMID 12556516	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1470	Kren 2003 Kren A, Mamnun YM, Bauer BE, et al., Kuchler K. War1p, a novel transcription factor controlling weak acid stress response in yeast. <i>Mol Cell Biol</i> 2003;23(5):1775–85.	doi:10.1128/MCB.23.5.1775-1785.2003 PMID 12588995 PMC151711	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1471	Garay-Arroyo 2003 Garay-Arroyo A, Lledías F, Hansberg W, Covarrubias AA. Cu,Zn-superoxide dismutase of <i>Saccharomyces cerevisiae</i> is required for resistance to hyperosmosis. <i>FEBS Lett</i> 2003;539(1–3):68–72.	doi:10.1016/s0014-5793(03)00199-6 PMID 12650928	–	1	1	1	–	1	1	Slot blot - wt vs TF mutant
1472	Türkel 2003 Türkel S, Turgut T, López MC, Uemura H, Baker HV. Mutations in GCR1 affect SUC2 gene expression in <i>Saccharomyces cerevisiae</i> . <i>Mol Genet Genomics</i> 2003;268(6):825–31.	doi:10.1007/s00438-003-0808-4 PMID 12655409	–	1	1	1	1	–	1	DNA footprinting
1473	Kaadige 2003 Kaadige MR, Lopes JM. Opi1p, Ume6p and Sin3p control expression from the promoter of the INO2 regulatory gene via a novel regulatory cascade. <i>Mol Microbiol</i> 2003;48(3):823–32.	doi:10.1046/j.1365-2958.2003.03472.x PMID 12694624	–	1	1	1	–	1	1	Enzyme activity - wt vs TF mutant
1474	Alepuz 2003 Alepuz PM, de Nadal E, Zapater M, Ammerer G, Posas F. Osmostress-induced transcription by Hot1 depends on a Hog1-mediated recruitment of the RNA Pol II. <i>EMBO J</i> 2003;22(10):2433–42.	doi:10.1093/emboj/cdg243 PMID 12743037 PMC156001	–	1	1	1	1	–	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1475	Rutherford 2003 Rutherford JC, Jaron S, Winge DR. Aft1p and Aft2p mediate iron-responsive gene expression in yeast through related promoter elements. <i>J Biol Chem</i> 2003;278(30):27636–43.	doi:10.1074/jbc.M300076200 PMID 12756250	–	1	1	1	–	1	1	Microarray analysis - wt vs TF mutant
1476	Van Slyke 2003 Van Slyke C, Grayhack EJ. The essential transcription factor Reb1p interacts with the CLB2 UAS outside of the G2/M control region. <i>Nucleic Acids Res</i> 2003;31(15):4597–607.	doi:10.1093/nar/gkg638 PMID 12888520 PMC169905	–	1	1	1	1	–	1	EMSA
1477	Gardocki 2003 Gardocki ME, Lopes JM. Expression of the yeast PIS1 gene requires multiple regulatory elements including a Rox1p binding site. <i>J Biol Chem</i> 2003;278(40):38646–52.	doi:10.1074/jbc.M305251200 PMID 12890676	–	1	1	1	–	1	1	CAT - wt vs mutant promoter sequence
1478	Almaguer 2003 Almaguer C, Mantella D, Perez E, Patton-Vogt J. Inositol and phosphate regulate GIT1 transcription and glycerophosphoinositol incorporation in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2003;2(4):729–36.	doi:10.1128/EC.2.4.729-736.2003 PMID 12912892 PMC178388	–	1	1	1	–	1	1	Enzyme activity - wt vs TF mutant
1479	Wolfger 2004 Wolfger H, Mamnun YM, Kuchler K. The yeast Pdr15p ATP-binding cassette (ABC) protein is a general stress response factor implicated in cellular detoxification. <i>J Biol Chem</i> 2004;279(12):11593–9.	doi:10.1074/jbc.M311282200 PMID 14699125	–	1	1	1	1	–	1	DNA footprinting
1480	Prinz 2004 Prinz S, Avila-Campillo I, Aldridge C, et al., Galitski T. Control of yeast filamentous-form growth by modules in an integrated molecular network. <i>Genome Res</i> 2004;14(3):380–90.	doi:10.1101/gr.2020604 PMID 14993204 PMC353223	no GEO	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1481	Ogawa 2004 Ogawa N, Mori K. Autoregulation of the HAC1 gene is required for sustained activation of the yeast unfolded protein response. <i>Genes Cells</i> 2004;9(2):95–104.	doi:10.1111/j.1365-2443.2004.00704.x PMID 15009095	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1482	Alimardani 2004 Alimardani P, Régnacq M, Moreau-Vauzelle C, et al., Bergès T. SUT1-promoted sterol uptake involves the ABC transporter Aus1 and the mannoprotein Dan1 whose synergistic action is sufficient for this process. <i>Biochem J</i> 2004;381(Pt 1):195–202.	doi:10.1042/BJ20040297 PMID 15035656 PMC1133777	–	1	1	1	–	1	1	Microarray analysis - wt vs TF overexpression
1483	Lascaris 2004 Lascaris R, Piwowski J, van der Spek H, de Mattos JT, Grivell L, Blom J. Overexpression of HAP4 in glucose-derepressed yeast cells reveals respiratory control of glucose-regulated genes. <i>Microbiology (Reading)</i> 2004;150(Pt 4):929–934.	doi:10.1099/mic.0.26742-0 PMID 15073302	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1484	Schröder 2004 Schröder M, Clark R, Liu CY, Kaufman RJ. The unfolded protein response represses differentiation through the RPD3-SIN3 histone deacetylase. <i>EMBO J</i> 2004;23(11):2281–92.	doi:10.1038/sj.emboj.7600233 PMID 15141165 PMC419911	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1485	Güldener 2004 Güldener U, Koehler GJ, Haussmann C, et al., Hegemann JH. Characterization of the <i>Saccharomyces cerevisiae</i> Foll1 protein: starvation for C1 carrier induces pseudohyphal growth. <i>Mol Biol Cell</i> 2004;15(8):3811–28.	doi:10.1091/mbc.e03-09-0680 PMID 15169867 PMC491839	–	1	1	1	–	1	1	GFP - wt vs TF mutant
1486	Tsuzi 2004 Tsuzi D, Maeta K, Takatsume Y, Izawa S, Inoue Y. Distinct regulatory mechanism of yeast GPX2 encoding phospholipid hydroperoxide glutathione peroxidase by oxidative stress and a calcineurin/Crz1-mediated Ca ²⁺ signaling pathway. <i>FEBS Lett</i> 2004;569(1–3):301–6.	doi:10.1016/j.febslet.2004.05.077 PMID 15225652	–	1	1	1	1	–	1	EMSA
1487	Gao 2004 Gao C, Wang L, Milgrom E, Shen WC. On the mechanism of constitutive Pdr1 activator-mediated PDR5 transcription in <i>Saccharomyces cerevisiae</i> : evidence for enhanced recruitment of coactivators and altered nucleosome structures. <i>J Biol Chem</i> 2004;279(41):42677–86.	doi:10.1074/jbc.M406363200 PMID 15294907	–	1	1	1	–	1	1	Northern blot - wt vs TF overexpression
1488	Mollapour 2004 Mollapour M, Fong D, Balakrishnan K, et al., Piper PW. Screening the yeast deletant mutant collection for hypersensitivity and hyper-resistance to sorbate, a weak organic acid food preservative. <i>Yeast</i> 2004;21(11):927–46.	doi:10.1002/yea.1141 PMID 15334557	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1489	Ju 2004 Ju D, Wang L, Mao X, Xie Y. Homeostatic regulation of the proteasome via an Rpn4-dependent feedback circuit. <i>Biochem Biophys Res Commun</i> 2004;321(1):51–7.	doi:10.1016/j.bbrc.2004.06.105 PMID 15358214	–	1	1	1	–	1	1	Western blot - wt vs TF overexpression
1490	Coluccio 2004 Coluccio A, Bogengruber E, Conrad MN, Dresser ME, Briza P, Neiman AM. Morphogenetic pathway of spore wall assembly in <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2004;3(6):1464–75.	doi:10.1128/EC.3.6.1464-1475.2004 PMID 15590821 PMC539034	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1491	Velasco 2004 Velasco I, Tenreiro S, Calderon IL, André B. <i>Saccharomyces cerevisiae</i> Aqr1 is an internal-membrane transporter involved in excretion of amino acids. <i>Eukaryot Cell</i> 2004;3(6):1492–503.	doi:10.1128/EC.3.6.1492-1503.2004 PMID 15590823 PMC539038	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1492	Hon 2005 Hon T, Lee HC, Hu Z, Iyer VR, Zhang L. The heme activator protein Hap1 represses transcription by a heme-independent mechanism in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2005;169(3):1343–52.	doi:10.1534/genetics.104.037143 PMID 15654089 PMC1449556	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1493	Takatsume 2005 Takatsume Y, Maeta K, Izawa S, Inoue Y. Enrichment of yeast thioredoxin by green tea extract through activation of Yap1 transcription factor in <i>Saccharomyces cerevisiae</i> . <i>J Agric Food Chem</i> 2005;53(2):332–7.	doi:10.1021/jf048818h PMID 15656669	–	1	1	1	–	–	1	–

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1494	Moreno 2005 Moreno F, Ahuatzli D, Riera A, Palomino CA, Herrero P. Glucose sensing through the Hxk2-dependent signalling pathway. <i>Biochem Soc Trans</i> 2005;33(Pt 1):265–8.	doi:10.1042/BST0330265 PMID 15667322	–	1	1	1	1	–	1	EMSA
1495	Palomino 2005 Palomino A, Herrero P, Moreno F. Rgt1, a glucose sensing transcription factor, is required for transcriptional repression of the HXK2 gene in <i>Saccharomyces cerevisiae</i> . <i>Biochem J</i> 2005;388(Pt 2):697–703.	doi:10.1042/BJ20050160 PMID 15705057 PMC1138978	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1496	Han 2005 Han SH, Han GS, Iwanyshyn WM, Carman GM. Regulation of the PIS1-encoded phosphatidylinositol synthase in <i>Saccharomyces cerevisiae</i> by zinc. <i>J Biol Chem</i> 2005;280(32):29017–24.	doi:10.1074/jbc.M505881200 PMID 15980062 PMC1201514	–	1	1	1	1	–	1	EMSA
1497	Liang 2006 Liang S, Rossby SP, Liang P, et al., Sulser F. Detection of an mRNA polymorphism by differential display. <i>Methods Mol Biol</i> 2006;317:279–85.	doi:10.1385/1-59259-968-0:279 PMID 16264236	–	1	1	1	–	–	1	–
1498	Valenzuela 2006 Valenzuela L, Gangadharan S, Kamakaka RT. Analyses of SUM1-1-mediated long-range repression. <i>Genetics</i> 2006;172(1):99–112.	doi:10.1534/genetics.105.050427 PMID 16272409 PMC1456157	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1499	Taylor 2005 Taylor R, Kebaara BW, Nazarens T, et al., Atkin AL. Gene set coregulated by the <i>Saccharomyces cerevisiae</i> nonsense-mediated mRNA decay pathway. <i>Eukaryot Cell</i> 2005;4(12):2066–77.	doi:10.1128/EC.4.12.2066-2077.2005 PMID 16339724 PMC1317485	–	1	1	1	–	1	1	Northern blot - wt vs TF overexpression
1500	Todeschini 2006 Todeschini AL, Condon C, Bénard L. Sodium-induced GCN4 expression controls the accumulation of the 5' to 3' RNA degradation inhibitor, 3'-phosphoadenosine 5'-phosphate. <i>J Biol Chem</i> 2006;281(6):3276–82.	doi:10.1074/jbc.M511688200 PMID 16352596	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1501	Gur 2006 Gur Y, Breitbart H. Mammalian sperm translate nuclear-encoded proteins by mitochondrial-type ribosomes. <i>Genes Dev</i> 2006;20(4):411–6.	doi:10.1101/gad.367606 PMID 16449571 PMC1369042	–	1	1	1	–	–	1	–
1502	Palomino 2006 Palomino A, Herrero P, Moreno F. Tpk3 and Snf1 protein kinases regulate Rgt1 association with <i>Saccharomyces cerevisiae</i> HXK2 promoter. <i>Nucleic Acids Res</i> 2006;34(5):1427–38.	doi:10.1093/nar/gkl028 PMID 16528100 PMC1401511	–	1	1	1	1	–	1	ChIP
1503	Su 2006 Su X, Dowhan W. Regulation of cardiolipin synthase levels in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> 2006;23(4):279–91.	doi:10.1002/yea.1352 PMID 16544270 PMC4108995	–	1	1	1	1	–	1	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1504	Kersting 2006 Kersting MC, Carman GM. Regulation of the <i>Saccharomyces cerevisiae</i> EKI1-encoded ethanolamine kinase by zinc depletion. <i>J Biol Chem</i> 2006;281(19):13110–13116.	doi:10.1074/jbc.M601612200 PMID 16551612 PMC1779367	–	1	1	1	1	–	1	EMSA
1505	Azevedo 2007 Azevedo D, Nascimento L, Labarre J, Toledano MB, Rodrigues-Pousada C. The <i>S. cerevisiae</i> Yap1 and Yap2 transcription factors share a common cadmium-sensing domain. <i>FEBS Lett</i> 2007;581(2):187–95.	doi:10.1016/j.febslet.2006.11.083 PMID 17187783	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1506	Vargas 2007 Vargas RC, García-Salcedo R, Tenreiro S, et al., Sá-Correia I. <i>Saccharomyces cerevisiae</i> multidrug resistance transporter Qdr2 is implicated in potassium uptake, providing a physiological advantage to quinidine-stressed cells. <i>Eukaryot Cell</i> 2007;6(2):134–42.	doi:10.1128/EC.00290-06 PMID 17189489 PMC1797947	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1507	Di 2007 Di Y, Tamás MJ. Regulation of the arsenic-responsive transcription factor Yap8p involves the ubiquitin-proteasome pathway. <i>J Cell Sci</i> 2007;120(Pt 2):256–64.	doi:10.1242/jcs.03346 PMID 17200139	–	1	1	1	–	1	1	lacZ - wt vs TF overexpression
1508	Sarma 2007 Sarma NJ, Haley TM, Barbara KE, Buford TD, Willis KA, Santangelo GM. Glucose-responsive regulators of gene expression in <i>Saccharomyces cerevisiae</i> function at the nuclear periphery via a reverse recruitment mechanism. <i>Genetics</i> 2007;175(3):1127–35.	doi:10.1534/genetics.106.068932 PMID 17237508 PMC1840092	–	1	1	1	–	1	1	Enzyme activity - wt vs TF mutant
1509	Camarasa 2007 Camarasa C, Faucet V, Dequin S. Role in anaerobiosis of the isoenzymes for <i>Saccharomyces cerevisiae</i> fumarate reductase encoded by OSM1 and FRDS1. <i>Yeast</i> 2007;24(5):391–401.	doi:10.1002/yea.1467 PMID 17345583	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1510	Chen 2007 Chen M, Lopes JM. Multiple basic helix-loop-helix proteins regulate expression of the ENO1 gene of <i>Saccharomyces cerevisiae</i> . <i>Eukaryot Cell</i> 2007;6(5):786–96.	doi:10.1128/EC.00383-06 PMID 17351075 PMC1899243	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1511	Skoneczna 2007 Skoneczna A, Micialkiewicz A, Skoneczny M. <i>Saccharomyces cerevisiae</i> Hsp31p, a stress response protein conferring protection against reactive oxygen species. <i>Free Radic Biol Med</i> 2007;42(9):1409–20.	doi:10.1016/j.freeradbiomed.2007.01.042 PMID 17395014	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1512	Kundu 2007 Kundu S, Horn PJ, Peterson CL. SWI/SNF is required for transcriptional memory at the yeast GAL gene cluster. <i>Genes Dev</i> 2007;21(8):997–1004.	doi:10.1101/gad.1506607 PMID 17438002 PMC1847716	–	1	1	1	1	–	1	ChIP
1513	Freddie 2007 Freddie CT, Ji Z, Marais A, Sharrocks AD. Functional interactions between the Forkhead transcription factor FOXK1 and the MADS-box protein SRF. <i>Nucleic Acids Res</i> 2007;35(15):5203–12.	doi:10.1093/nar/gkm528 PMID 17670796 PMC1976435	–	1	1	1	1	–	1	EMSA

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1514	Shen 2007 Shen C, Lancaster CS, Shi B, Guo H, Thimmaiah P, Bjornsti MA. TOR signaling is a determinant of cell survival in response to DNA damage. <i>Mol Cell Biol</i> 2007;27(20):7007–17.	doi:10.1128/MCB.00290-07 PMID 17698581 PMC2168917	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1515	Xin 2007 Xin X, Lan C, Lee HC, Zhang L. Regulation of the HAP1 gene involves positive actions of histone deacetylases. <i>Biochem Biophys Res Commun</i> 2007;362(1):120–125.	doi:10.1016/j.bbrc.2007.07.156 PMID 17706600 PMC2030606	–	1	1	1	1	–	1	EMSA
1516	Okazaki 2007 Okazaki S, Tachibana T, Naganuma A, Mano N, Kuge S. Multistep disulfide bond formation in Yap1 is required for sensing and transduction of H2O2 stress signal. <i>Mol Cell</i> 2007;27(4):675–88.	doi:10.1016/j.molcel.2007.06.035 PMID 17707237	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1517	Fischer 2008 Fischer C, Valerius O, Rupprecht H, Dumkow M, Krappmann S, Braus GH. Posttranscriptional regulation of FLO11 upon amino acid starvation in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Res</i> 2008;8(2):225–36.	doi:10.1111/j.1567-1364.2007.00331.x PMID 17999676	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1518	Ghavidel 2007 Ghavidel A, Kislinger T, Pogoutse O, Sopko R, Jurisica I, Emili A. Impaired tRNA nuclear export links DNA damage and cell-cycle checkpoint. <i>Cell</i> 2007;131(5):915–26.	doi:10.1016/j.cell.2007.09.042 PMID 18045534	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1519	Fontanesi 2008 Fontanesi F, Jin C, Tzagoloff A, Barrientos A. Transcriptional activators HAP/NF-Y rescue a cytochrome c oxidase defect in yeast and human cells. <i>Hum Mol Genet</i> 2008;17(6):775–88.	doi:10.1093/hmg/ddm349 PMID 18045776	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1520	Li 2008 Li L, Bagley D, Ward DM, Kaplan J. Yap5 is an iron-responsive transcriptional activator that regulates vacuolar iron storage in yeast. <i>Mol Cell Biol</i> 2008;28(4):1326–37.	doi:10.1128/MCB.01219-07 PMID 18070921 PMC2258748	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1521	Dumlao 2008 Dumlao DS, Hertz N, Clarke S. Secreted 3-isopropylmalate methyl ester signals invasive growth during amino acid starvation in <i>Saccharomyces cerevisiae</i> . <i>Biochemistry</i> 2008;47(2):698–709.	doi:10.1021/bi7018157 PMID 18092814	–	1	1	1	–	1	1	Enzyme activity - wt vs TF mutant
1522	Kim 2008 Kim KY, Truman AW, Levin DE. Yeast Mpk1 mitogen-activated protein kinase activates transcription through Swi4/Swi6 by a noncatalytic mechanism that requires upstream signal. <i>Mol Cell Biol</i> 2008;28(8):2579–89.	doi:10.1128/MCB.01795-07 PMID 18268013 PMC2293104	–	1	1	1	1	–	1	ChIP
1523	Kumánovics 2008 Kumánovics A, Chen OS, Li L, et al., Kaplan J. Identification of FRA1 and FRA2 as genes involved in regulating the yeast iron regulon in response to decreased mitochondrial iron-sulfur cluster synthesis. <i>J Biol Chem</i> 2008;283(16):10276–86.	doi:10.1074/jbc.M801160200 PMID 18281282 PMC2447656	–	1	1	1	1	–	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1524	Mok 2008 Mok YG, Lee BD, Kim YJ, et al., Hwang S. The tobacco gene Ntscy07 confers arsenite tolerance in <i>Saccharomyces cerevisiae</i> by reducing the steady state levels of intracellular arsenic. <i>FEBS Lett</i> 2008;582(6):916–24.	doi:10.1016/j.febslet.2008.02.030 PMID 18294968	–	1	1	1	1	–	1	One-hybrid assay
1525	Ma 2008 Ma J, Dobry CJ, Krysan DJ, Kumar A. Unconventional genomic architecture in the budding yeast <i>saccharomyces cerevisiae</i> masks the nested antisense gene NAG1. <i>Eukaryot Cell</i> 2008;7(8):1289–98.	doi:10.1128/EC.00053-08 PMID 18310357 PMC2519765	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1526	Hiraishi 2008 Hiraishi H, Miyake T, Ono B. Transcriptional regulation of <i>Saccharomyces cerevisiae</i> CYS3 encoding cystathionine gamma-lyase. <i>Curr Genet</i> 2008;53(4):225–34.	doi:10.1007/s00294-008-0181-2 PMID 18317767 PMC2668581	–	1	1	1	1	–	1	EMSA
1527	Archer 2008 Archer CT, Delahodde A, Gonzalez F, Johnston SA, Kodadek T. Activation domain-dependent monoubiquitylation of Gal4 protein is essential for promoter binding in vivo. <i>J Biol Chem</i> 2008;283(18):12614–23.	doi:10.1074/jbc.M801050200 PMID 18326036 PMC2335349	–	1	1	1	1	–	1	EMSA
1528	Raithatha 2008 Raithatha SA, Stuart DT. The <i>Saccharomyces cerevisiae</i> CLB5 promoter contains two middle sporulation elements (MSEs) that are differentially regulated during sporulation. <i>Yeast</i> 2008;25(4):259–72.	doi:10.1002/yea.1585 PMID 18327887	–	1	1	1	1	–	1	Northern blot - wt vs mutant promoter sequence
1529	Jambhekar 2008 Jambhekar A, Amon A. Control of meiosis by respiration. <i>Curr Biol</i> 2008;18(13):969–75.	doi:10.1016/j.cub.2008.05.047 PMID 18595705 PMC2504020	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1530	Hollenbach 2008 Hollenbach M, Hintersdorf A, Huse K, et al., Birkenmeier G. Ethyl pyruvate and ethyl lactate down-regulate the production of pro-inflammatory cytokines and modulate expression of immune receptors. <i>Biochem Pharmacol</i> 2008;76(5):631–44.	doi:10.1016/j.bcp.2008.06.006 PMID 18625205	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1531	Tyler 2008 Tyler CW, Kontsevich LL, Ferree TC. Independent components in stimulus-related BOLD signals and estimation of the underlying neural responses. <i>Brain Res</i> 2008;1229:72–89.	doi:10.1016/j.brainres.2008.06.050 PMID 18625206 PMC6685289	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1532	Wilson 2008 Wilson HD, Uhelski ML, Fuchs PN. Examining the role of the medial thalamus in modulating the affective dimension of pain. <i>Brain Res</i> 2008;1229:90–9.	doi:10.1016/j.brainres.2008.06.009 PMID 18625207	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1533	Klimesch 2008 Klimesch W, Freunberger R, Sauseng P, Gruber W. A short review of slow phase synchronization and memory: evidence for control processes in different memory systems? <i>Brain Res</i> 2008;1235:31–44.	doi:10.1016/j.brainres.2008.06.049 PMID 18625208	–	1	1	1	–	1	1	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1534	Andermann 2008 Andermann ML, Moore CI. Mechanical resonance enhances the sensitivity of the vibrissa sensory system to near-threshold stimuli. <i>Brain Res</i> 2008;1235:74–81.	doi:10.1016/j.brainres.2008.06.054 PMID 18625209 PMC3655715	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1535	Dakubo 2008 Dakubo GD, Beug ST, Mazerolle CJ, Thurig S, Wang Y, Wallace VA. Control of glial precursor cell development in the mouse optic nerve by sonic hedgehog from retinal ganglion cells. <i>Brain Res</i> 2008;1228:27–42.	doi:10.1016/j.brainres.2008.06.058 PMID 18625210	–	1	1	1	–	1	1	lacZ - wt vs TF overexpression
1536	Price 2008 Price CJ, Samson WK, Ferguson AV. Nesfatin-1 inhibits NPY neurons in the arcuate nucleus. <i>Brain Res</i> 2008;1230:99–106.	doi:10.1016/j.brainres.2008.06.084 PMID 18625211 PMC2590930	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1537	Kerzerho 2008 Kerzerho S, Gentaz E, Streri A. Influence of visual contextual cues on haptic discrimination of orientations in 5-month-old infants. <i>Brain Res</i> 2008;1242:276–82.	doi:10.1016/j.brainres.2008.06.055 PMID 18625212	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1538	Xia 2008 Xia Z, Turner GC, Hwang CS, Byrd C, Varshavsky A. Amino acids induce peptide uptake via accelerated degradation of CUP9, the transcriptional repressor of the PTR2 peptide transporter. <i>J Biol Chem</i> 2008;283(43):28958–68.	doi:10.1074/jbc.M803980200 PMID 18708352 PMC2570885	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1539	Moldón 2008 Moldón A, Malapeira J, Gabrielli N, et al., Ayté J. Promoter-driven splicing regulation in fission yeast. <i>Nature</i> 2008;455(7215):997–1000.	doi:10.1038/nature07325 PMID 18815595	–	1	1	1	1	–	1	ChIP
1540	Choi 2008 Choi ID, Jeong MY, Ham MS, Sung HC, Yun CW. Novel Ree1 regulates the expression of ENO1 via the Snf1 complex pathway in <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2008;377(2):395–399.	doi:10.1016/j.bbrc.2008.09.146 PMID 18851946	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1541	Spitzner 2008 Spitzner A, Perzlmaier AF, Geillinger KE, Reihl P, Stolz J. The proline-dependent transcription factor Put3 regulates the expression of the riboflavin transporter MCH5 in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2008;180(4):2007–17.	doi:10.1534/genetics.108.094458 PMID 18940788 PMC2600938	–	1	1	1	1	–	1	EMSA
1542	Kim 2009 Kim JH. DNA-binding properties of the yeast Rgt1 repressor. <i>Biochimie</i> 2009;91(2):300–3.	doi:10.1016/j.biochi.2008.09.002 PMID 18950675 PMC2859070	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1543	Wang 2009 Wang Z, Wang Y, Hegg EL. Regulation of the heme A biosynthetic pathway: differential regulation of heme A synthase and heme O synthase in <i>Saccharomyces cerevisiae</i> . <i>J Biol Chem</i> 2009;284(2):839–47.	doi:10.1074/jbc.M804167200 PMID 18953022 PMC2613620	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1544	Shirra 2008 Shirra MK, McCartney RR, Zhang C, Shokat KM, Schmidt MC, Arndt KM. A chemical genomics study identifies Snf1 as a repressor of GCN4 translation. <i>J Biol Chem</i> 2008;283(51):35889–98.	doi:10.1074/jbc.M805325200 PMID 18955495 PMC2602918	GSE12061	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1545	Brown 2009 Brown V, Sabina J, Johnston M. Specialized sugar sensing in diverse fungi. <i>Curr Biol</i> 2009;19(5):436–41.	doi:10.1016/j.cub.2009.01.056 PMID 19249212 PMC2762733	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1546	Yang 2009 Yang HY, Tatebayashi K, Yamamoto K, Saito H. Glycosylation defects activate filamentous growth Kss1 MAPK and inhibit osmoregulatory Hog1 MAPK. <i>EMBO J</i> 2009;28(10):1380–91.	doi:10.1038/emboj.2009.104 PMID 19369942 PMC2688530	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1547	Wei 2009 Wei M, Fabrizio P, Madia F, et al., Longo VD. Tor1/Sch9-regulated carbon source substitution is as effective as calorie restriction in life span extension. <i>PLoS Genet</i> 2009;5(5):e1000467.	doi:10.1371/journal.pgen.1000467 PMID 19424415 PMC2669710	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1548	Madia 2009 Madia F, Wei M, Yuan V, et al., Longo VD. Oncogene homologue Sch9 promotes age-dependent mutations by a superoxide and Rev1/Polzeta-dependent mechanism. <i>J Cell Biol</i> 2009;186(4):509–23.	doi:10.1083/jcb.200906011 PMID 19687253 PMC2733759	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1549	Hector 2009 Hector RE, Bowman MJ, Skory CD, Cotta MA. The <i>Saccharomyces cerevisiae</i> YMR315W gene encodes an NADP(H)-specific oxidoreductase regulated by the transcription factor Stb5p in response to NADPH limitation. <i>N Biotechnol</i> 2009;26(3–4):171–80.	doi:10.1016/j.nbt.2009.08.008 PMID 19712762	–	1	1	1	1	–	1	lacZ - wt vs mutant promoter sequence
1550	Bumgarner 2009 Bumgarner SL, Dowell RD, Grisafi P, Gifford DK, Fink GR. Toggle involving cis-interfering noncoding RNAs controls variegated gene expression in yeast. <i>Proc Natl Acad Sci U S A</i> 2009;106(43):18321–6.	doi:10.1073/pnas.0909641106 PMID 19805129 PMC2775344	–	1	1	1	1	–	1	Chip-on-chip
1551	Wang 2009 Wang H, Carey LB, Cai Y, Wijnen H, Futcher B. Recruitment of Cln3 cyclin to promoters controls cell cycle entry via histone deacetylase and other targets. <i>PLoS Biol</i> 2009;7(9):e1000189.	doi:10.1371/journal.pbio.1000189 PMID 19823669 PMC2730028	–	1	1	1	1	–	1	ChIP
1552	Kato 2010 Kato M, Kito K, Ota K, Ito T. Remodeling of the SCF complex-mediated ubiquitination system by compositional alteration of incorporated F-box proteins. <i>Proteomics</i> 2010;10(1):115–23.	doi:10.1002/pmic.200900497 PMID 19882662	–	1	1	1	1	–	1	RT-PCR - wt vs mutant promoter sequence
1553	Lainé 2009 Lainé JP, Singh BN, Krishnamurthy S, Hampsey M. A physiological role for gene loops in yeast. <i>Genes Dev</i> 2009;23(22):2604–9.	doi:10.1101/gad.1823609 PMID 19933150 PMC2779762	–	1	1	1	1	–	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1554	Esposito 2010 Esposito M, Konarzewska P, Odeyale O, Shen CH. Gene-wide histone acetylation at the yeast INO1 requires the transcriptional activator Ino2p. <i>Biochem Biophys Res Commun</i> 2010;391(2):1285–90.	doi:10.1016/j.bbrc.2009.12.063 PMID 20018175	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1555	Piccirillo 2010 Piccirillo S, White MG, Murphy JC, Law DJ, Honigberg SM. The Rim101p/PacC pathway and alkaline pH regulate pattern formation in yeast colonies. <i>Genetics</i> 2010;184(3):707–16.	doi:10.1534/genetics.109.113480 PMID 20038633 PMC2845339	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1556	Iwai 2010 Iwai K, Naganuma A, Kuge S. Peroxiredoxin Ahp1 acts as a receptor for alkylhydroperoxides to induce disulfide bond formation in the Cad1 transcription factor. <i>J Biol Chem</i> 2010;285(14):10597–604.	doi:10.1074/jbc.M109.090142 PMID 20145245 PMC2856268	–	1	1	1	1	–	1	ChIP
1557	Xu 2010 Xu T, Shively CA, Jin R, et al., Kumar A. A profile of differentially abundant proteins at the yeast cell periphery during pseudohyphal growth. <i>J Biol Chem</i> 2010;285(20):15476–15488.	doi:10.1074/jbc.M110.114926 PMID 20228058 PMC2865295	GSE20351	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1558	Wolf 2010 Wolf JJ, Dowell RD, Mahony S, Rabani M, Gifford DK, Fink GR. Feed-forward regulation of a cell fate determinant by an RNA-binding protein generates asymmetry in yeast. <i>Genetics</i> 2010;185(2):513–22.	doi:10.1534/genetics.110.113944 PMID 20382833 PMC2881133	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1559	Casado 2010 Casado C, Yenush L, Melero C, et al., Ramos J. Regulation of Trk-dependent potassium transport by the calcineurin pathway involves the Hal5 kinase. <i>FEBS Lett</i> 2010;584(11):2415–20.	doi:10.1016/j.febslet.2010.04.042 PMID 20412803	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1560	Heo 2010 Heo DH, Baek IJ, Kang HJ, et al., Yun CW. Cadmium regulates copper homeostasis by inhibiting the activity of Mac1, a transcriptional activator of the copper regulon, in <i>Saccharomyces cerevisiae</i> . <i>Biochem J</i> 2010;431(2):257–65.	doi:10.1042/BJ20100638 PMID 20670216	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1561	Young 2010 Young BP, Shin JJ, Oriji R, et al., Loewen CJ. Phosphatidic acid is a pH biosensor that links membrane biogenesis to metabolism. <i>Science</i> 2010;329(5995):1085–8.	doi:10.1126/science.1191026 PMID 20798321	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1562	Strudwick 2010 Strudwick N, Brown M, Parmar VM, Schröder M. Ime1 and Ime2 are required for pseudohyphal growth of <i>Saccharomyces cerevisiae</i> on nonfermentable carbon sources. <i>Mol Cell Biol</i> 2010;30(23):5514–30.	doi:10.1128/MCB.00390-10 PMID 20876298 PMC2976428	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1563	Tyrrell 2010 Tyrrell A, Flick K, Kleiger G, Zhang H, Deshaies RJ, Kaiser P. Physiologically relevant and portable tandem ubiquitin-binding domain stabilizes polyubiquitylated proteins. <i>Proc Natl Acad Sci U S A</i> 2010;107(46):19796–801.	doi:10.1073/pnas.1010648107 PMID 21041680 PMC2993400	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1564	Rice 2010 Rice C, Cooke M, Treloar N, Vollbrecht P, Stuke J, McDonough V. A role for MGA2, but not SPT23, in activation of transcription of ERG1 in <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2010;403(3-4):293-7.	doi:10.1016/j.bbrc.2010.11.020 PMID 21075079	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1565	Herzog 2011 Herzog B, Streckfuss-Bömeke K, Braus GH. A feedback circuit between transcriptional activation and self-destruction of Gcn4 separates its metabolic and morphogenic response in diploid yeasts. <i>J Mol Biol</i> 2011;405(4):909-25.	doi:10.1016/j.jmb.2010.11.033 PMID 21111745	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1566	Liang 2011 Liang CY, Hsu PH, Chou DF, et al., Lo WS. The histone H3K36 demethylase Rph1/KDM4 regulates the expression of the photoreactivation gene PHR1. <i>Nucleic Acids Res</i> 2011;39(10):4151-65.	doi:10.1093/nar/gkr040 PMID 21296759 PMC3105397	–	1	1	1	1	–	1	ChIP
1567	Mira 2011 Mira NP, Henriques SF, Keller G, et al., Sá-Correia I. Identification of a DNA-binding site for the transcription factor Haa1, required for <i>Saccharomyces cerevisiae</i> response to acetic acid stress. <i>Nucleic Acids Res</i> 2011;39(16):6896-907.	doi:10.1093/nar/gkr228 PMID 21586585 PMC3167633	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1568	Phenix 2011 Phenix H, Morin K, Batenchuk C, et al., Kærn M. Quantitative epistasis analysis and pathway inference from genetic interaction data. <i>PLoS Comput Biol</i> 2011;7(5):e1002048.	doi:10.1371/journal.pcbi.1002048 PMID 21589890 PMC3093353	–	1	1	1	–	1	1	GFP - wt vs TF mutant
1569	Campbell 2011 Campbell RN, Westhorpe F, Reece RJ. Isolation of compensatory inhibitor domain mutants to novel activation domain variants using the split-ubiquitin screen. <i>Yeast</i> 2011;28(8):569-78.	doi:10.1002/yea.1861 PMID 21732556	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1570	Solé 2011 Solé C, Nadal-Ribelles M, Kraft C, Peter M, Posas F, de Nadal E. Control of Ubp3 ubiquitin protease activity by the Hog1 SAPK modulates transcription upon osmostress. <i>EMBO J</i> 2011;30(16):3274-84.	doi:10.1038/emboj.2011.227 PMID 21743437 PMC3160652	–	1	1	1	1	–	1	ChIP
1571	Promlek 2011 Promlek T, Ishiwata-Kimata Y, Shido M, Sakuramoto M, Kohno K, Kimata Y. Membrane aberrancy and unfolded proteins activate the endoplasmic reticulum stress sensor Ire1 in different ways. <i>Mol Biol Cell</i> 2011;22(18):3520-32.	doi:10.1091/mbc.E11-04-0295 PMID 21775630 PMC3172275	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1572	Sarode 2011 Sarode N, Miracle B, Peng X, Ryan O, Reynolds TB. Vacuolar protein sorting genes regulate mat formation in <i>Saccharomyces cerevisiae</i> by Flo11p-dependent and -independent mechanisms. <i>Eukaryot Cell</i> 2011;10(11):1516-26.	doi:10.1128/EC.05078-11 PMID 21908597 PMC3209053	–	1	1	1	–	1	1	Western blot - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1573	Xiao 2011 Xiao L, Kamau E, Donze D, Grove A. Expression of yeast high mobility group protein HMO1 is regulated by TOR signaling. <i>Gene</i> 2011;489(1):55–62.	doi:10.1016/j.gene.2011.08.017 PMID 21924331	–	1	1	1	1	–	1	ChIP
1574	Lissina 2011 Lissina E, Young B, Urbanus ML, et al., Nislow C. A systems biology approach reveals the role of a novel methyltransferase in response to chemical stress and lipid homeostasis. <i>PLoS Genet</i> 2011;7(10):e1002332.	doi:10.1371/journal.pgen.1002332 PMID 22028670 PMC3197675	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1575	Liang 2012 Liang H, Lim HH, Venkitaraman A, Surana U. Cdk1 promotes kinetochore bi-orientation and regulates Cdc20 expression during recovery from spindle checkpoint arrest. <i>EMBO J</i> 2012;31(2):403–16.	doi:10.1038/emboj.2011.385 PMID 22056777 PMC3261552	–	1	1	1	–	1	1	Northern blot - wt vs TF overexpression
1576	Uprety 2012 Uprety B, Lahudkar S, Malik S, Bhaumik SR. The 19S proteasome subcomplex promotes the targeting of NuA4 HAT to the promoters of ribosomal protein genes to facilitate the recruitment of TFIID for transcriptional initiation in vivo. <i>Nucleic Acids Res</i> 2012;40(5):1969–83.	doi:10.1093/nar/gkr977 PMID 22086954 PMC3300024	–	1	1	1	1	–	1	ChIP
1577	Raithatha 2012 Raithatha S, Su TC, Lourenco P, Goto S, Sadowski I. Cdk8 regulates stability of the transcription factor Phd1 to control pseudohyphal differentiation of <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2012;32(3):664–74.	doi:10.1128/MCB.05420-11 PMID 22124158 PMC3266601	–	1	1	1	1	–	1	DNA footprinting
1578	Soto-Cardalda 2012 Soto-Cardalda A, Fakas S, Pascual F, Choi HS, Carman GM. Phosphatidate phosphatase plays role in zinc-mediated regulation of phospholipid synthesis in yeast. <i>J Biol Chem</i> 2012;287(2):968–77.	doi:10.1074/jbc.M111.313130 PMID 22128164 PMC3256892	–	1	1	1	1	–	1	EMSA
1579	Radchenko 2011 Radchenko E, Rogoza T, Khokhrina M, Drozdova P, Mironova L. SUP35 expression is enhanced in yeast containing [ISP+], a prion form of the transcriptional regulator Sfp1. <i>Prion</i> 2011; 5(4):317–22.	doi:10.4161/pri.18426 PMID 22156729 PMC4012397	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1580	Ungar 2011 Ungar L, Harari Y, Toren A, Kupiec M. Tor complex 1 controls telomere length by affecting the level of Ku. <i>Curr Biol</i> 2011;21(24):2115–20.	doi:10.1016/j.cub.2011.11.024 PMID 22169538	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1581	Abramczyk 2012 Abramczyk D, Holden S, Page CJ, Reece RJ. Interplay of a ligand sensor and an enzyme in controlling expression of the <i>Saccharomyces cerevisiae</i> GAL genes. <i>Eukaryot Cell</i> 2012; 11(3):334–42.	doi:10.1128/EC.05294-11 PMID 22210830 PMC3294446	–	1	1	1	1	–	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1582	Shukla 2012 Shukla A, Lahudkar S, Durairaj G, Bhaumik SR. Sgf29p facilitates the recruitment of TATA box binding protein but does not alter SAGA's global structural integrity in vivo. <i>Biochemistry</i> 2012;51(2):706–14.	doi:10.1021/bi201708z PMID 22224423 PMC3264703	–	1	1	1	1	–	1	ChIP
1583	Albrecht 2012 Albrecht TR, Wagner EJ. snRNA 3' end formation requires heterodimeric association of integrator subunits. <i>Mol Cell Biol</i> 2012;32(6):1112–23.	doi:10.1128/MCB.06511-11 PMID 22252320 PMC3295014	–	1	1	1	1	–	1	ChIP
1584	Ott 2012 Ott C, Ross K, Straub S, et al., Kozjak-Pavlovic V. Sam50 functions in mitochondrial intermembrane space bridging and biogenesis of respiratory complexes. <i>Mol Cell Biol</i> 2012;32(6):1173–88.	doi:10.1128/MCB.06388-11 PMID 22252321 PMC3295012	–	1	1	1	1	–	1	ChIP
1585	Montefusco 2012 Montefusco DJ, Newcomb B, Gandy JL, et al., Hannun YA. Sphingoid bases and the serine catabolic enzyme CHA1 define a novel feedforward/feedback mechanism in the response to serine availability. <i>J Biol Chem</i> 2012;287(12):9280–9.	doi:10.1074/jbc.M111.313445 PMID 22277656 PMC3308826	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1586	Miceli 2011 Miceli MV, Jiang JC, Tiwari A, Rodriguez-Quinones JF, Jazwinski SM. Loss of mitochondrial membrane potential triggers the retrograde response extending yeast replicative lifespan. <i>Front Genet</i> 2011;2:102.	doi:10.3389/fgene.2011.00102 PMID 22303396 PMC3266616	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1587	Torres 2012 Torres AM, Maceiras ML, Belmonte ER, Naveira LN, Calvo MB, Cerdán ME. KIRox1p contributes to yeast resistance to metals and is necessary for KLYCF1 expression in the presence of cadmium. <i>Gene</i> 2012;497(1):27–37.	doi:10.1016/j.gene.2012.01.050 PMID 22310389	–	1	1	1	1	–	1	EMSA
1588	Mousley 2012 Mousley CJ, Yuan P, Gaur NA, et al., Bankaitis VA. A sterol-binding protein integrates endosomal lipid metabolism with TOR signaling and nitrogen sensing. <i>Cell</i> 2012;148(4):702–15.	doi:10.1016/j.cell.2011.12.026 PMID 22341443 PMC3285437	–	1	1	1	1	–	1	ChIP
1589	Ryo 2012 Ryo S, Ishii J, Iguchi Y, Fukuda N, Kondo A. Transplantation of the GAL regulon into G-protein signaling circuitry in yeast. <i>Anal Biochem</i> 2012;424(1):27–31.	doi:10.1016/j.ab.2012.02.005 PMID 22343189	–	1	1	1	–	1	1	GFP - wt vs TF overexpression
1590	Rosonina 2012 Rosonina E, Duncan SM, Manley JL. Sumoylation of transcription factor Gcn4 facilitates its Srb10-mediated clearance from promoters in yeast. <i>Genes Dev</i> 2012;26(4):350–5.	doi:10.1101/gad.184689.111 PMID 22345516 PMC3289883	–	1	1	1	1	–	1	ChIP
1591	Miles 2012 Miles WO, Tschöp K, Herr A, Ji JY, Dyson NJ. Pumilio facilitates miRNA regulation of the E2F3 oncogene. <i>Genes Dev</i> 2012;26(4):356–68.	doi:10.1101/gad.182568.111 PMID 22345517 PMC3289884	–	1	1	1	1	–	1	ChIP

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1592	Liu 2012 Liu L, Qi J, Yang Z, Peng L, Li C. Low-affinity copper transporter CTR2 is regulated by copper-sensing transcription factor Mac1p in <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2012;420(3):600–4.	doi:10.1016/j.bbrc.2012.03.040 PMID 22445756	–	1	1	1	1	–	1	ChIP
1593	Sun 2012 Sun K, Xiang L, Ishihara S, Matsuura A, Sakagami Y, Qi J. Anti-aging effects of hesperidin on <i>Saccharomyces cerevisiae</i> via inhibition of reactive oxygen species and UTH1 gene expression. <i>Biosci Biotechnol Biochem</i> 2012;76(4):640–5.	doi:10.1271/bbb.110535 PMID 22484922	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1594	Pfeiffer 2012 Pfeiffer V, Lingner J. TERRA promotes telomere shortening through exonuclease 1-mediated resection of chromosome ends. <i>PLoS Genet</i> 2012;8(6):e1002747.	doi:10.1371/journal.pgen.1002747 PMID 22719262 PMC3375253	–	1	1	1	1	–	1	ChIP
1595	Karunanithi 2012 Karunanithi S, Cullen PJ. The filamentous growth MAPK Pathway Responds to Glucose Starvation Through the Mig1/2 transcriptional repressors in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> 2012;192(3):869–87.	doi:10.1534/genetics.112.142661 PMID 22904036 PMC3522164	–	1	1	1	–	1	1	His3 - wt vs TF mutant
1596	Zelin 2012 Zelin E, Zhang Y, Toogun OA, Zhong S, Freeman BC. The p23 molecular chaperone and GCN5 acetylase jointly modulate protein-DNA dynamics and open chromatin status. <i>Mol Cell</i> 2012;48(3):459–70.	doi:10.1016/j.molcel.2012.08.026 PMID 23022381 PMC3496002	–	1	1	1	1	–	1	Fluorescence anisotropy
1597	Ogata 2012 Ogata T. Nitrogen starvation induces expression of Lg-FLO1 and flocculation in bottom-fermenting yeast. <i>Yeast</i> 2012; 29(11):487–94.	doi:10.1002/yea.2928 PMID 23065862	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1598	Zhao 2013 Zhao XF, Li M, Li YQ, Chen XD, Gao XD. The TEA/ATTS transcription factor YlTec1p represses the yeast-to-hypha transition in the dimorphic yeast <i>Yarrowia lipolytica</i> . <i>FEMS Yeast Res</i> 2013;13(1):50–61.	doi:10.1111/j.1567-1364.2012.12008.x PMID 23067114	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1599	Batista-Nascimento 2013 Batista-Nascimento L, Toledano MB, Thiele DJ, Rodrigues-Pousada C. Yeast protective response to arsenate involves the repression of the high affinity iron uptake system. <i>Biochim Biophys Acta</i> 2013;1833(5):997–1005.	doi:10.1016/j.bbamcr.2012.12.018 PMID 23295455	GSE33427	1	1	1	–	1	1	lacZ - wt vs TF mutant
1600	Bodvard 2013 Bodvard K, Jörhov A, Blomberg A, Molin M, Käll M. The yeast transcription factor Crz1 is activated by light in a Ca ²⁺ /calcineurin-dependent and PKA-independent manner. <i>PLoS One</i> 2013;8(1):e53404.	doi:10.1371/journal.pone.0053404 PMID 23335962 PMC3546054	–	1	1	1	–	1	1	GFP - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1601	Pincus 2013 Pincus D, Ryan CJ, Smith RD, Brent R, Resnekov O. Assigning quantitative function to post-translational modifications reveals multiple sites of phosphorylation that tune yeast pheromone signaling output. <i>PLoS One</i> 2013;8(3):e56544.	doi:10.1371/journal.pone.0056544 PMID 23554854 PMC3595240	–	1	1	1	–	1	1	mCherry fluorescence WT vs N-terminus fluorescence tag fusion
1602	Chang 2013 Chang M, Kang HJ, Baek IJ, Kang CM, Park YS, Yun CW. Kdx1 regulates RCK1 gene expression by interacting with Rlm1 in <i>Saccharomyces cerevisiae</i> . <i>Biochem Biophys Res Commun</i> 2013;435(3):350–5.	doi:10.1016/j.bbrc.2013.04.083 PMID 23660188	GSE45692	1	1	1	–	1	1	Northern blot - wt vs TF overexpression
1603	Bastajian 2013 Bastajian N, Friesen H, Andrews BJ. Bck2 acts through the MADS box protein Mcm1 to activate cell-cycle-regulated genes in budding yeast. <i>PLoS Genet</i> 2013;9(5):e1003507.	doi:10.1371/journal.pgen.1003507 PMID 23675312 PMC3649975	–	1	1	1	1	–	1	ChIP
1604	Montanari 2013 Montanari A, Zhou YF, D’Orsi MF, Bolotin-Fukuhara M, Frontali L, Francisci S. Analyzing the suppression of respiratory defects in the yeast model of human mitochondrial tRNA diseases. <i>Gene</i> 2013;527(1):1–9.	doi:10.1016/j.gene.2013.05.042 PMID 23727608	–	1	1	1	1	–	1	RT-PCR - wt vs mutant promoter sequence
1605	Paiva 2013 Paiva S, Strachotová D, Kučerová H, et al., Váchová L. The transport of carboxylic acids and important role of the Jen1p transporter during the development of yeast colonies. <i>Biochem J</i> 2013;454(3):551–8.	doi:10.1042/BJ20120312 PMID 23790185	–	1	1	1	–	1	1	GFP - wt vs TF mutant
1606	Carballo 2013 Carballo JA, Panizza S, Serrentino ME, et al., Cha RS. Budding yeast ATM/ATR control meiotic double-strand break (DSB) levels by down-regulating Rec114, an essential component of the DSB-machinery. <i>PLoS Genet</i> 2013;9(6):e1003545.	doi:10.1371/journal.pgen.1003545 PMID 23825959 PMC3694840	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1607	Malik 2013 Malik S, Durairaj G, Bhaumik SR. Mechanisms of antisense transcription initiation from the 3’ end of the GAL10 coding sequence in vivo. <i>Mol Cell Biol</i> 2013;33(18):3549–67.	doi:10.1128/MCB.01715-12 PMID 23836882 PMC3753873	–	1	1	1	1	–	1	RT-PCR - wt vs mutant promoter sequence
1608	Chakrabarti 2013 Chakrabarti P, Kim JY, Singh M, et al., Kandrор KV. Insulin inhibits lipolysis in adipocytes via the evolutionarily conserved mTORC1-Egr1-ATGL-mediated pathway. <i>Mol Cell Biol</i> 2013;33(18):3659–66.	doi:10.1128/MCB.01584-12 PMID 23858058 PMC3753874	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1609	Karpov 2013 Karpov DS, Spasskaya DS, Tutyaeva VV, Mironov AS, Karpov VL. Proteasome inhibition enhances resistance to DNA damage via upregulation of Rpn4-dependent DNA repair genes. <i>FEBS Lett</i> 2013;587(18):3108–14.	doi:10.1016/j.febslet.2013.08.007 PMID 23954292	–	1	1	1	1	–	1	DamID system Wild-type vs Mutated binding site

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1610	Dong 2013 Dong K, Addinall SG, Lydall D, Rutherford JC. The yeast copper response is regulated by DNA damage. <i>Mol Cell Biol</i> 2013;33(20):4041–50.	doi:10.1128/MCB.00116-13 PMID 23959798 PMC3811678	–	1	1	1	1	–	1	DNA footprinting
1611	Gonçalves 2014 Gonçalves IR, Conde e Silva N, Garay CL, Lesuisse E, Camadro JM, Blaiseau PL. The basis for evolution of DNA-binding specificity of the Aft1 transcription factor in yeasts. <i>Genetics</i> 2014;196(1):149–60.	doi:10.1534/genetics.113.157693 PMID 24172132 PMC3872181	–	1	1	1	1	–	1	EMSA
1612	Kang 2014 Kang HJ, Jeong SJ, Kim KN, et al., Yun CW. A novel protein, Pho92, has a conserved YTH domain and regulates phosphate metabolism by decreasing the mRNA stability of PHO4 in <i>Saccharomyces cerevisiae</i> . <i>Biochem J</i> 2014;457(3):391–400.	doi:10.1042/BJ20130862 PMID 24206186	GSE50086	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1613	Hanna 2014 Hanna J, Waterman D, Isasa M, et al., Finley D. Cuz1/Ynl155w, a zinc-dependent ubiquitin-binding protein, protects cells from metalloinduced proteotoxicity. <i>J Biol Chem</i> 2014;289(3):1876–85.	doi:10.1074/jbc.M113.534032 PMID 24297164 PMC3894362	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1614	Amaral 2013 Amaral C, Pimentel C, Matos RG, Arraiano CM, Matzapetakis M, Rodrigues-Pousada C. Two residues in the basic region of the yeast transcription factor Yap8 are crucial for its DNA-binding specificity. <i>PLoS One</i> 2013;8(12):e83328.	doi:10.1371/journal.pone.0083328 PMID 24358276 PMC3865217	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1615	Pascual 2014 Pascual F, Hsieh LS, Soto-Cardalda A, Carman GM. Yeast Pahlp phosphatidate phosphatase is regulated by proteasome-mediated degradation. <i>J Biol Chem</i> 2014;289(14):9811–22.	doi:10.1074/jbc.M114.550103 PMID 24563465 PMC3975026	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1616	Berra 2014 Berra S, Ayachi S, Ramotar D. Upregulation of the <i>Saccharomyces cerevisiae</i> efflux pump Tpo1 rescues an Imp2 transcription factor-deficient mutant from bleomycin toxicity. <i>Environ Mol Mutagen</i> 2014;55(6):518–24.	doi:10.1002/em.21865 PMID 24599794	–	1	1	1	–	1	1	Western blot - wt vs TF overexpression
1617	Ceschin 2014 Ceschin J, Saint-Marc C, Laporte J, et al., Pinson B. Identification of yeast and human 5-aminoimidazole-4-carboxamide-1- β -d-ribofuranoside (AICAr) transporters. <i>J Biol Chem</i> 2014;289(24):16844–54.	doi:10.1074/jbc.M114.551192 PMID 24778186 PMC4059127	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1618	Cho 2014 Cho BR, Lee P, Hahn JS. CK2-dependent inhibitory phosphorylation is relieved by Ppt1 phosphatase for the ethanol stress-specific activation of Hsf1 in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2014;93(2):306–16.	doi:10.1111/mmi.12660 PMID 24894977	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1619	Zhang 2014 Zhang L, Di J. The transcriptional activator Ino2p dissociates from the yeast INM1 promoter in induction. <i>DNA Cell Biol</i> 2014;33(12):863–8.	doi:10.1089/dna.2014.2397 PMID 25211324 PMC4248251	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1620	Surlov 2014 Surlov BA, Cooley BM, Needham PG, Brodsky JL, Patton-Vogt J. Loss of Ypk1, the yeast homolog to the human serum- and glucocorticoid-induced protein kinase, accelerates phospholipase B1-mediated phosphatidylcholine deacylation. <i>J Biol Chem</i> 2014;289(45):31591–604.	doi:10.1074/jbc.M114.581157 PMID 25258318 PMC4223355	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1621	Nussbaum 2014 Nussbaum I, Weindling E, Jubran R, Cohen A, Bar-Nun S. Deteriorated stress response in stationary-phase yeast: Sir2 and Yap1 are essential for Hsf1 activation by heat shock and oxidative stress, respectively. <i>PLoS One</i> 2014;9(10):e111505.	doi:10.1371/journal.pone.0111505 PMID 25356557 PMC4214751	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1622	Yarrington 2015 Yarrington RM, Rudd JS, Stillman DJ. Spatiotemporal cascade of transcription factor binding required for promoter activation. <i>Mol Cell Biol</i> 2015;35(4):688–98.	doi:10.1128/MCB.01285-14 PMID 25512608 PMC4301721	–	1	1	1	1	–	1	RT-PCR - wt vs mutant promoter sequence
1623	Kim 2015 Kim SR, Xu H, Lesmana A, et al., Jin YS. Deletion of PHO13, encoding haloacid dehalogenase type IIA phosphatase, results in upregulation of the pentose phosphate pathway in <i>Saccharomyces cerevisiae</i> . <i>Appl Environ Microbiol</i> 2015;81(5):1601–9.	doi:10.1128/AEM.03474-14 PMID 25527558 PMC4325158	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1624	Fu 2015 Fu J, Hou J, Chen L, et al., Bao X. The yeast BDF1 regulates endocytosis via LSP1 under salt stress. <i>Curr Microbiol</i> 2015;70(5):671–8.	doi:10.1007/s00284-014-0773-7 PMID 25572496	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1625	Weindling 2015 Weindling E, Bar-Nun S. Sir2 links the unfolded protein response and the heat shock response in a stress response network. <i>Biochem Biophys Res Commun</i> 2015;457(3):473–8.	doi:10.1016/j.bbrc.2015.01.021 PMID 25600811	–	1	1	1	–	1	1	Microarray analysis - wt vs TF mutant
1626	Liu 2015 Liu Y, Stuparevic I, Xie B, Becker E, Law MJ, Primig M. The conserved histone deacetylase Rpd3 and the DNA binding regulator Ume6 repress BOI1's meiotic transcript isoform during vegetative growth in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2015;96(4):861–74.	doi:10.1111/mmi.12976 PMID 25708805	–	1	1	1	1	–	1	ChIP
1627	Herrador 2015 Herrador A, Livas D, Soletto L, Becuwe M, Léon S, Vincent O. Casein kinase 1 controls the activation threshold of an α -arrestin by multisite phosphorylation of the interdomain hinge. <i>Mol Biol Cell</i> 2015;26(11):2128–38.	doi:10.1091/mbc.E14-11-1552 PMID 25851600 PMC4472021	–	1	1	1	–	1	1	Western blot - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1628	Saint-Marc 2015 Saint-Marc C, Hürlimann HC, Daignan-Fornier B, Pinson B. Serine hydroxymethyltransferase: a key player connecting purine, folate and methionine metabolism in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 2015;61(4):633–40.	doi:10.1007/s00294-015-0489-7 PMID 25893566	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1629	Li 2015 Li P, Shao Y, Jin H, Yu HG. Ndj1, a telomere-associated protein, regulates centrosome separation in budding yeast meiosis. <i>J Cell Biol</i> 2015;209(2):247–59.	doi:10.1083/jcb.201408118 PMID 25897084 PMC4411264	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1630	Wang 2015 Wang LC, Montalvo-Munoz F, Tsai YC, Liang CY, Chang CC, Lo WS. The Histone Acetyltransferase Gcn5 Regulates ncRNA-ICR1 and FLO11 Expression during Pseudohyphal Development in <i>Saccharomyces cerevisiae</i> . <i>Biomed Res Int</i> 2015;2015:284692.	doi:10.1155/2015/284692 PMID 25922832 PMC4398931	–	1	1	1	1	–	1	ChIP
1631	Mattiazzi Ušaj 2015 Mattiazzi Ušaj M, Prelec M, Brložnik M, et al., Petrovič U. Yeast <i>Saccharomyces cerevisiae</i> adiponectin receptor homolog Izh2 is involved in the regulation of zinc, phospholipid and pH homeostasis. <i>Metallomics</i> 2015;7(9):1338–51.	doi:10.1039/c5mt00095e PMID 26067383	–	1	1	1	–	1	1	Northern blot - wt vs TF mutant
1632	Yadav 2015 Yadav KK, Singh N, Rajasekharan R. PHO4 transcription factor regulates triacylglycerol metabolism under low-phosphate conditions in <i>Saccharomyces cerevisiae</i> . <i>Mol Microbiol</i> 2015;98(3):456–72.	doi:10.1111/mmi.13133 PMID 26179227	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1633	Ferreira 2015 Ferreira RT, Menezes RA, Rodrigues-Pousada C. E4-Ubiquitin ligase Ufd2 stabilizes Yap8 and modulates arsenic stress responses independent of the U-box motif. <i>Biol Open</i> 2015;4(9):1122–31.	doi:10.1242/bio.010405 PMID 26276098 PMC4582114	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1634	Taylor 2015 Taylor MB, Ehrenreich IM. Transcriptional Derepression Uncovers Cryptic Higher-Order Genetic Interactions. <i>PLoS Genet</i> 2015;11(10):e1005606.	doi:10.1371/journal.pgen.1005606 PMID 26484664 PMC4618523	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1635	Villasmil 2016 Villasmil ML, Francisco J, Gallo-Ebert C, et al., Nickels JT Jr. Ceramide signals for initiation of yeast mating-specific cell cycle arrest. <i>Cell Cycle</i> 2016;15(3):441–54.	doi:10.1080/15384101.2015.1127475 PMID 26726837 PMC4943704	–	1	1	1	–	1	1	lacZ - wt vs TF overexpression
1636	Zhao 2016 Zhao Y, Yan H, Happeck R, et al., Jiang L. The plasma membrane protein Rch1 is a negative regulator of cytosolic calcium homeostasis and positively regulated by the calcium/calcineurin signaling pathway in budding yeast. <i>Eur J Cell Biol</i> 2016;95(3–5):164–74.	doi:10.1016/j.ejcb.2016.01.001 PMID 26832117	–	1	1	1	–	1	1	lacZ - wt vs TF mutant

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1637	Berterame 2016 Berterame NM, Porro D, Ami D, Branduardi P. Protein aggregation and membrane lipid modifications under lactic acid stress in wild type and OPI1 deleted <i>Saccharomyces cerevisiae</i> strains. <i>Microb Cell Fact</i> 2016;15:39.	doi:10.1186/s12934-016-0438-2 PMID 26887851 PMC4756461	–	1	1	1	–	1	1	mRNA amplification Wild-type vs TF deletion mutant
1638	Salinas 2016 Salinas F, de Boer CG, Abarca V, et al., Cubillos FA. Natural variation in non-coding regions underlying phenotypic diversity in budding yeast. <i>Sci Rep</i> 2016;6:21849.	doi:10.1038/srep21849 PMID 26898953 PMC4761897	–	1	1	1	–	1	1	Luciferase - wt vs TF mutant
1639	Poos 2016 Poos AM, Maicher A, Dieckmann AK, et al., König R. Mixed Integer Linear Programming based machine learning approach identifies regulators of telomerase in yeast. <i>Nucleic Acids Res</i> 2016;44(10):e93.	doi:10.1093/nar/gkw111 PMID 26908654 PMC4889924	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1640	Yadav 2016 Yadav KK, Rajasekharan R. The transcription factor GCN4 regulates PHM8 and alters triacylglycerol metabolism in <i>Saccharomyces cerevisiae</i> . <i>Curr Genet</i> 2016;62(4):841–851.	doi:10.1007/s00294-016-0590-6 PMID 26979516	–	1	1	1	1	–	1	EMSA
1641	Guerra-Moreno 2016 Guerra-Moreno A, Hanna J. Tmc1 Is a Dynamically Regulated Effector of the Rpn4 Proteotoxic Stress Response. <i>J Biol Chem</i> 2016;291(28):14788–95.	doi:10.1074/jbc.M116.726398 PMID 27226598 PMC4938195	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1642	Kuznetsov 2016 Kuznetsov E, Váchová L, Palková Z. Cellular localization of Sun4p and its interaction with proteins in the yeast birth scar. <i>Cell Cycle</i> 2016;15(14):1898–907.	doi:10.1080/15384101.2016.1189043 PMID 27229769 PMC4968892	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1643	D’Urso 2016 D’Urso A, Takahashi YH, Xiong B, et al., Brickner JH. Set1/COMPASS and Mediator are repurposed to promote epigenetic transcriptional memory. <i>Elife</i> 2016;5.	doi:10.7554/eLife.16691 PMID 27336723 PMC4951200	–	1	1	1	1	–	1	ChIP
1644	Roy 2016 Roy K, Gabunilas J, Gillespie A, Ngo D, Chanfreau GF. Common genomic elements promote transcriptional and DNA replication roadblocks. <i>Genome Res</i> 2016;26(10):1363–1375.	doi:10.1101/gr.204776.116 PMID 27540088 PMC5052057	–	1	1	1	1	–	1	Northern blot - wt vs mutant promoter sequence
1645	Ramya 2016 Ramya V, Rajasekharan R. ATG15 encodes a phospholipase and is transcriptionally regulated by YAP1 in <i>Saccharomyces cerevisiae</i> . <i>FEBS Lett</i> 2016;590(18):3155–67.	doi:10.1002/1873-3468.12369 PMID 27543826	–	1	1	1	1	–	1	EMSA
1646	Ball 2016 Ball DA, Mehta GD, Salomon-Kent R, et al., Karpova TS. Single molecule tracking of Ace1p in <i>Saccharomyces cerevisiae</i> defines a characteristic residence time for non-specific interactions of transcription factors with chromatin. <i>Nucleic Acids Res</i> 2016;44(21):e160.	doi:10.1093/nar/gkw744 PMID 27566148 PMC5137432	–	1	1	1	1	–	1	SMT Single molecule tracking

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1647	Wang 2016 Wang Q, Donze D. Transcription factor Reb1 is required for proper transcriptional start site usage at the divergently transcribed TFC6-ESC2 locus in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> 2016;594(1):108–116.	doi:10.1016/j.gene.2016.09.004 PMID 27601258	–	1	1	1	1	–	1	EMSA
1648	Dugan 2016 Dugan A, Majmudar CY, Pricer R, et al., Mapp AK. Discovery of Enzymatic Targets of Transcriptional Activators via in Vivo Covalent Chemical Capture. <i>J Am Chem Soc</i> 2016; 138(38):12629–35.	doi:10.1021/jacs.6b07680 PMID 27611834 PMC5217703	–	1	1	1	1	–	1	ChIP
1649	Fujita 2017 Fujita KI, Ishikura T, Jono Y, et al., Tanaka T. Anethole potentiates dodecanol's fungicidal activity by reducing PDR5 expression in budding yeast. <i>Biochim Biophys Acta Gen Subj</i> 2017;1861(2):477–484.	doi:10.1016/j.bbagen.2016.09.010 PMID 27632201	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1650	Yadav 2016 Yadav KK, Singh N, Rajvanshi PK, Rajasekharan R. The RNA polymerase I subunit Rpa12p interacts with the stress-responsive transcription factor Msn4p to regulate lipid metabolism in budding yeast. <i>FEBS Lett</i> 2016;590(20): 3559–3573.	doi:10.1002/1873-3468.12422 PMID 27637775	–	1	1	1	1	–	1	EMSA
1651	Suzuki 2017 Suzuki H, Sakabe T, Hirose Y, Eki T. Development and evaluation of yeast-based GFP and luciferase reporter assays for chemical-induced genotoxicity and oxidative damage. <i>Appl Microbiol Biotechnol</i> 2017;101(2):659–671.	doi:10.1007/s00253-016-7911-z PMID 27766356	–	1	1	1	–	1	1	Luciferase - wt vs TF mutant
1652	Vizoso-Vázquez 2017 Vizoso-Vázquez A, Lamas-Maceiras M, Fernández-Leiro R, Rico-Díaz A, Becerra M, Cerdán ME. Dual function of Ixr1 in transcriptional regulation and recognition of cisplatin-DNA adducts is caused by differential binding through its two HMG-boxes. <i>Biochim Biophys Acta Gene Regul Mech</i> 2017; 1860(2):256–269.	doi:10.1016/j.bbagra.2016.11.005 PMID 27871851	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1653	Menezes 2017 Menezes RA, Pimentel C, Silva AR, et al., Rodrigues-Pousada C. Mediator, SWI/SNF and SAGA complexes regulate Yap8-dependent transcriptional activation of ACR2 in response to arsenate. <i>Biochim Biophys Acta Gene Regul Mech</i> 2017;1860(4):472–481.	doi:10.1016/j.bbagra.2017.02.001 PMID 28188921	–	1	1	1	1	–	1	ChIP
1654	Degreif 2017 Degreif D, de Rond T, Bertl A, Keasling JD, Budin I. Lipid engineering reveals regulatory roles for membrane fluidity in yeast flocculation and oxygen-limited growth. <i>Metab Eng</i> 2017;41:46–56.	doi:10.1016/j.ymben.2017.03.002 PMID 28323063	–	1	1	1	–	1	1	RT-PCR - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1655	Martins 2018 Martins TS, Pereira C, Canadell D, et al., Costa V. The Hog1p kinase regulates Aft1p transcription factor to control iron accumulation. <i>Biochim Biophys Acta Mol Cell Biol Lipids</i> 2018;1863(1):61–70.	doi:10.1016/j.bbali.2017.10.001 PMID 29032057	–	1	1	1	–	1	1	lacZ - wt vs TF mutant
1656	Delorme-Axford 2018 Delorme-Axford E, Abernathy E, Lennemann NJ, et al., Klionsky DJ. The exoribonuclease Xrn1 is a post-transcriptional negative regulator of autophagy. <i>Autophagy</i> 2018;14(5):898–912.	doi:10.1080/15548627.2018.1441648 PMID 29465287 PMC6070002	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1657	Moretto 2018 Moretto F, Wood NE, Kelly G, Doncic A, van Werven FJ. A regulatory circuit of two lncRNAs and a master regulator directs cell fate in yeast. <i>Nat Commun</i> 2018;9(1):780.	doi:10.1038/s41467-018-03213-z PMID 29472539 PMC5823921	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1658	Stuecker 2018 Stuecker TN, Scholes AN, Lewis JA. Linkage mapping of yeast cross protection connects gene expression variation to a higher-order organismal trait. <i>PLoS Genet</i> 2018;14(4):e1007335.	doi:10.1371/journal.pgen.1007335 PMID 29649251 PMC5978988	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1659	Sukegawa 2018 Sukegawa Y, Negishi T, Kikuchi Y, et al., Ohya Y. Genetic dissection of the signaling pathway required for the cell wall integrity checkpoint. <i>J Cell Sci</i> 2018;131(13).	doi:10.1242/jcs.219063 PMID 29853633	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1660	Dey 2018 Dey T, Krishna Rao K, Khatun J, Rangarajan PN. The nuclear transcription factor Rtg1p functions as a cytosolic, post-transcriptional regulator in the methylotrophic yeast <i>Pichia pastoris</i> . <i>J Biol Chem</i> 2018;293(43):16647–16660.	doi:10.1074/jbc.RA118.004486 PMID 30185617 PMC6204902	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1661	Mehta 2018 Mehta GD, Ball DA, Eriksson PR, et al., Karpova TS. Single-Molecule Analysis Reveals Linked Cycles of RSC Chromatin Remodeling and Ace1p Transcription Factor Binding in Yeast. <i>Mol Cell</i> 2018;72(5):875–887.e9.	doi:10.1016/j.molcel.2018.09.009 PMID 30318444 PMC6289719	–	1	1	1	1	–	1	HaloTag-TF co-localization
1662	Hamada 2019 Hamada K, Naito A, Hamaguchi Y, et al., Usui T. Ubr1p-Cup9p-Ptr2p pathway involves in the sensitivity to readthrough compounds negamycin derivatives in budding yeast. <i>Biosci Biotechnol Biochem</i> 2019;83(10):1889–1892.	doi:10.1080/09168451.2019.1625263 PMID 31159660	–	1	1	1	–	1	1	Western blot - wt vs TF mutant
1663	Mat Nanyan 2019 Mat Nanyan NSB, Watanabe D, Sugimoto Y, Takagi H. Involvement of the stress-responsive transcription factor gene MSN2 in the control of amino acid uptake in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Res</i> 2019;19(5).	doi:10.1093/femsyr/foz052 PMID 31328231	–	1	1	1	–	1	1	Western blot - wt vs TF overexpression

Table 4, continued

#	Publication	Links	SPELL	rows	TFs	pairs	direct	indirect	cond.	top assay
1664	Mat Nanyan 2020 Mat Nanyan NSB, Watanabe D, Sugimoto Y, Takagi H. Effect of the deubiquitination enzyme gene UBP6 on the stress-responsive transcription factor Msn2-mediated control of the amino acid permease Gnp1 in yeast. <i>J Biosci Bioeng</i> 2020;129(4):423–427.	doi:10.1016/j.jbiosc.2019.10.002 PMID 31640922	–	1	1	1	–	1	1	Western blot - wt vs TF overexpression
1665	Mukai 2019 Mukai Y, Kamei Y, Liu X, et al., Takagi H. Proline metabolism regulates replicative lifespan in the yeast <i>Saccharomyces cerevisiae</i> . <i>Microb Cell</i> 2019;6(10):482–490.	doi:10.15698/mic2019.10.694 PMID 31646149 PMC6780008	–	1	1	1	–	1	1	RT-PCR - wt vs TF overexpression
1666	Konduri 2020 Konduri PC, Wang T, Salamat N, Zhang L. Heme, A Metabolic Sensor, Directly Regulates the Activity of the KDM4 Histone Demethylase Family and Their Interactions with Partner Proteins. <i>Cells</i> 2020;9(3).	doi:10.3390/cells9030773 PMID 32235736 PMC7140707	–	1	1	1	–	1	1	LacZ - WT vs. TF double mutant
1667	Singh 2021 Singh S, Sahu RK, Tomar RS. The N-Terminal Tail of Histone H3 Regulates Copper Homeostasis in <i>Saccharomyces cerevisiae</i> . <i>Mol Cell Biol</i> 2021;41(2).	doi:10.1128/MCB.00210-20 PMID 33257505 PMC8093488	–	1	1	1	1	–	1	ChIP
1668	Mfarej 2020 Mfarej MG, Skibbens RV. DNA damage induces Yap5-dependent transcription of ECO1/CTF7 in <i>Saccharomyces cerevisiae</i> . <i>PLoS One</i> 2020;15(12):e0242968.	doi:10.1371/journal.pone.0242968 PMID 33373396 PMC7771704	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant
1669	Stewart 2021 Stewart T, Wolfe BE, Fuchs SM. Defining the role of the polyasparagine repeat domain of the <i>S. cerevisiae</i> transcription factor Azflp. <i>PLoS One</i> 2021;16(5):e0247285.	doi:10.1371/journal.pone.0247285 PMID 34019539 PMC8139511	–	1	1	1	–	1	1	RT-PCR - wt vs. TF mutant
1670	Vallabhaneni 2021 Vallabhaneni AR, Kabashi M, Haymowicz M, et al., Conrad-Webb H. HSF1 induces RNA polymerase II synthesis of ribosomal RNA in <i>S. cerevisiae</i> during nitrogen deprivation. <i>Curr Genet</i> 2021;67(6):937–951.	doi:10.1007/s00294-021-01197-w PMID 34363098 PMC8594204	–	1	1	1	1	–	1	ChIP
1671	Nishimura 2021 Nishimura A, Yoshikawa Y, Ichikawa K, Takemoto T, Tanahashi R, Takagi H. Longevity Regulation by Proline Oxidation in Yeast. <i>Microorganisms</i> 2021;9(8).	doi:10.3390/microorganisms9081650 PMID 34442729 PMC8400801	–	1	1	1	–	1	1	RT-PCR - wt vs TF mutant