


Figure 5: Supplement three: how profiles compose, and what the predicted ones carry. **a**, Emergent and masked gene rates of each public pair against how alike its two singles are, with the control-matched null beside them. The median emergent fraction is 0.355 against a null of 0.621, and the masked fraction 0.649; the only two near-replicate singles in the screen disagree on 0.536 to 0.603 of their own hits, which is the noise floor these rates sit inside. ✗● **b**, The one dose grid the public screen holds, methotrexate against fluorouracil: the weight the free linear fit puts on the first single moves from 0.128 to 0.757 across 9 dose pairs, while the mean wins at 9 of them. ✗● **c**, **d**, Profile similarity among the six inhibitors, measured where a measurement exists and then with every profile predicted. Measured, the strongest pair reads 0.924; predicted, 0.986, because the small acids have near-identical predicted profiles. One scale carries both panels. ✗● **e**, Similarity against the pair's departure from Bliss, both versions: 0.204 at a permutation p of 0.462 with measurements in, and -0.225 at 0.425 with everything predicted. ✗● **f**, The GO read: significant terms in the top hundred sensitive genes of the measured profile, of the ridge-predicted profile, and of the gene mean over the fitted compounds. The control carries 58 terms and the predicted profiles share 48 of them, which is what the predicted enrichment is. ✗●