


1. How many hydrogens does an alkane with 17 carbons have? How many carbons does an alkane with 74 hydrogens have?




36H and 36C

2. Draw the structure of octane and 2-methylheptane.



3. Draw the structure of C_5H_{12} that has a) one tertiary carbon b) no secondary carbons.



4. Draw the structures and name the four constitutional isomers for C_4H_9Br .



5. Draw the structure for each of the following:


a) 2,3-dimethylhexane



b) 4,4-diethyldecane



c) 2,4,5-trimethyl-4-(1-methylethyl) heptane



6. Convert the following structures to skeletal structures:

a) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$



b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$




c) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OCH}_3$



d) 3,4-diethyl-2-methylheptane




e) 2,2,5-trimethylhexane



7. Give the IUPAC name and indicate whether each is a primary, secondary or tertiary alcohol.

a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$

primary. 1-pentanol




© 2017 Pearson Education, Inc.

5-chloro-2-methyl-2-pentanol

Tertiary.

c. $\begin{array}{c} \text{CH}_3\text{CHCH}_2\text{CHCH}_2\text{CH}_3 \\ | \quad | \\ \text{CH}_3 \quad \text{OH} \end{array}$


Secondary. 5-methyl-3-hexanol





7-methyl-3,5-octandiol

Secondary

8. Are the following compounds primary, secondary or tertiary?



© 2017 Pearson Education, Inc.



(A) Tertiary alkyl halide. (B) Primary amine.

(C) Secondary alcohol. (D) Secondary amine.

9. Predict the approximate size of the following bond angles?

For Halogens and alcohols count the number of C attached to the next C IE, secondary C secondary alcohol for amines it only matters the amount of C the amine is directly attached too IE, nitrogen is attached to 3 C then it's a tertiary amine.

a) the C-O-C bond angle in an ether

b) The C-O-H bond angle in an alcohol

c) the C-N-C bond angle in a secondary amine

(A) The bond angle would be similar to the bond angle in water (104.5°)

(B) The bond angle would be similar to the bond angle in ammonia (107.3°)

(C) The bond angle would be similar to the bond angle in water (104.5°)

10. Which of the following compounds form hydrogen bonds between the molecules?

a) $\text{CH}_3\text{CH}_3\text{CH}_2\text{OH}$

b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{F}$



c) $\text{CH}_3\text{OCH}_2\text{CH}_3$





A only

The atom in question must have a direct contact with a hydrogen. Only A has this bond.

11. Convert the following Newman projections to skeletal structures and name them





© 2017 Pearson Education, Inc.





3-ethyl-2-pentanol

2-methyl-2-pentanamine

12. Draw the more stable conformer of cis-1-ethyl-2-methylcyclohexane and trans-1-ethyl-2-methylcyclohexane.



more stable



Less stable

13. Which is more stable: cis-1-ethyl-2-methylcyclohexane or trans-1-ethyl-2-methylcyclohexane?

Overall the trans conformer is more stable than the cis one since there is less steric hindrance (electrons and big groups are further apart).