The LUA-PHYSICAL library

Version 1.0

Thomas Jenni

May 7, 2019

Abstract

lua-physical is a pure Lua library, which provides functions and objects for the computation of physical quantities. A physical quantity is the product of a numerical value and a physical unit. The package has been written, to simplify the creation physics problem sets. The package provides units of the SI and the imperial system. Furthermore, an almost complete set of international currencies are supported, however without realtime exchange rates. In order to display the numbers with measurement uncertainties, the package is able to perform gaussian error propagation.

Contents

1	Intr	oduction	3
2	Loa 2.1	ding Dependencies	3
	2.2	License	4
3	Usa	${f ge}$	5
	3.1	Unit conversion	5
		3.1.1 Temperature Conversion	6
	3.2	Uncertainty Propagation	7
	3.3	Mathematical operations	10
4	Sup	ported Units	12
	4.1	Prefixes	12
	4.2	Base Units	13
	4.3	Constants	15
	4.4	Coherent derived units in the SI	16
	4.5	Non-SI units accepted for use with the SI	17
	4.6	Nominal Astronomical Units	17
	4.7	Other Non-SI units	18

	4.8	Imperial Units	20
	4.9	U.S. customary units	22
		International Currencies	
		4.10.1 Pegged International Currencies	31
5	Lua	Documentation	35
	5.1	physical.Quantity	35
	5.2	physical.Dimension	45
	5.3	physical.Unit	45
	5.4	physical.Number	46
Bi	bliog	raphy	49
In	dex o	of Units	50
In	dex o	of Currencies	53

1 Introduction

The author of this package is a physics teacher at the high school Kantonsschule Zug, Switzerland. The main use of this package is to write physics problem sets. LualateX does make it possible to integrate physical calculations directly. The package has been in use since 2016. Many bugs have been found and fixed. Nevertheless it still is possible, that some were not found yet. Therefore the author recommends not to use this package industry or science. If one does so, it's the responsability of the user to check results for plausability. If the user finds some bugs, they can be reported at github.com or directly to the author (thomas.jenni (at) ksz.ch).

2 Loading

This package is a pure Lua library. Therefore one has to require it explicitly by calling require("physical"). For printing physical quanties, the siunitx is supported. It's recommended to define a macro like \q to convert the lua quantity object to a siunitx expression.

The following Latex preamble loads the lua-physical package and creates a macro \q for printing physical quantities.

Listing 1: basic preamble

```
\usepackage{lua-physical}
                                                                    1
\usepackage{siunitx}
                                                                    2
                                                                    3
% configure siunitx
                                                                    4
\sisetup{
                                                                    5
 output-decimal-marker = {.},
                                                                    6
  per-mode = symbol,
                                                                    7
  separate-uncertainty = true,
                                                                    8
  add-decimal-zero = true,
  exponent-product = \cdot,
                                                                    10
  round-mode = off
                                                                    11
                                                                    12
                                                                    13
% load the lua-physical package
                                                                    14
\begin{luacode*}
                                                                    15
  physical = require("physical")
                                                                    16
  N = physical.Number
                                                                    17
\end{luacode*}
                                                                    18
                                                                    19
% print a physical quantity
                                                                    20
                                                                    21
\newcommand{\q}[1]{\%}
  \directlua{tex.print(physical.Quantity.tosiunitx(#1,"
                                                                    22
        scientific-notation=fixed,exponent-to-prefix=false"))}
}
                                                                    23
```

2.1 Dependencies

In principle this library is standalone, but it is compatible with the siunitx package. Calculation results can be written to LuaLATEX directly by calling the tosiunitx() method. If the preamble above is used, the printing is done be the \q{} macro.

2.2 License

This code is freely distributable under the terms of the MIT license.

Copyright (c) 2019 Thomas Jenni

Permission is hereby granted, free of charge, to any person obtaining a copy of this software and associated documentation files (the "Software"), to deal in the Software without restriction, including without limitation the rights to use, copy, modify, merge, publish, distribute, sublicense, and/or sell copies of the Software, and to permit persons to whom the Software is furnished to do so, subject to the following conditions:

The above copyright notice and this permission notice shall be included in all copies or substantial portions of the Software.

THE SOFTWARE IS PROVIDED "AS IS", WITHOUT WARRANTY OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO THE WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE AND NONINFRINGEMENT. IN NO EVENT SHALL THE AUTHORS OR COPYRIGHT HOLDERS BE LIABLE FOR ANY CLAIM, DAMAGES OR OTHER LIABILITY, WHETHER IN AN ACTION OF CONTRACT, TORT OR OTHERWISE, ARISING FROM, OUT OF OR IN CONNECTION WITH THE SOFTWARE OR THE USE OR OTHER DEALINGS IN THE SOFTWARE.

3 Usage

Given the basic preamble, units can be used in lua code directly. By convention, all units have an underscore in front of them, i.e. meter is _m, second is _s. All available units are listed in chapter 4. The following example illustrates the use of this library.

Listing 2: Velocity of a car.

```
\begin{luacode}
    s = 10 * _m
    t = 2 * _s
    v = s/t
    \end{luacode}

A car travels $\q{s}$ in $\q{t}$. calculate its velocity.

$$
    v = \frac{s}{t} = \frac{\q{s}}{\q{t}} = \ulline{\q{v}}

$$

1

2

4

4

4

4

4

4

6

A car travels $\q{s}$ in $\q{t}$. calculate its velocity.

7

$$

9

$$
```

A car travels 10 m in 2 s. Calculate its velocity.

$$v = \frac{s}{t} = \frac{10 \,\mathrm{m}}{2 \,\mathrm{s}} = \underline{\underline{5.0 \,\mathrm{m/s}}}$$

In the above Listing 2, the variable s stands for displacement and has the unit meter _m, the variable t stands for time and is given in seconds _s. If mathematical operations are done on them, new physical quantities are created. In the problem above, the velocity v is calculated by dividing s by t. The instance v has the derived unit m/s. By using the macro \q{} all quantities can be printed to the LuaLATEX code directly.

3.1 Unit conversion

Very often, the result of a calculation has to be converted to another unit. In the following Listing 3, the task is to calculate the volume of a cuboid. The length of the edges are given in different units. The volume is calculated by multipling all three lengths, the unit of the result is cm mm m. If the unit cm³ is preferred, it has to be converted explicitly. The conversion function is called to() and is available on all physical quantitiy instances. At first this looks a bit cumbersome. The reason of this behaviour is, that the software is not able to guess the unit of the result. In many cases, like in the example here, it's not clear what unit the result sould have. Therefore the user has always to give the target unit explicitly.

Listing 3: Volume of a cuboid.

```
\begin{luacode}
                                                                              1
  a = 12 * _cm
                                                                              2
  b = 150 * _mm
                                                                              3
  c = 1.5 * _m
                                                                              4
                                                                              5
  V = a*b*c
                                                                              6
\end{luacode}
                                                                              7
                                                                              8
Find the volume of a rectangular cuboid with lengths \q{a},
                                                                              9
                                                                              10
q{b}\ and \ q{c}\ .
  V= a \cdot b \cdot c
                                                                              12
  = \q{a} \cdot \q{b} \cdot \q{c}
                                                                              13
  = \q{V}
                                                                              14
    \displaystyle \left\{ \left( \frac{dm^3}{} \right) \right\}
                                                                              15
                                                                              16
```

Find the volume of a rectangular cuboid with lengths $12\,\mathrm{cm},\,150\,\mathrm{mm}$ and $1.5\,\mathrm{m}.$

```
V = a \cdot b \cdot c = 12 \,\mathrm{cm} \cdot 150 \,\mathrm{mm} \cdot 1.5 \,\mathrm{m} = 2700.0 \,\mathrm{cm} \,\mathrm{mm} \,\mathrm{m} = 27.0 \,\mathrm{dm}^3
```

3.1.1 Temperature Conversion

Most physical units transform linearly. Exceptions are the unit degree Celsius <code>_degC</code> and degree Fahrenheit <code>_degF</code>. These units are ambigous and can be interpreted as temperature differences or as an absolute temperatures. In the latter case, the conversion to base units is not a linear, but an affine transformation. This is because degree Celsius and degree Fahrenheit scales have their zero points at different temperatures compared to the unit Kelvin.

By default $_{\text{degC}}$ and $_{\text{degF}}$ units are temperature differences. If one wants to have it converted absolutely, it has to be done adding / subtracting $_{\text{degC}_0} = 273.15*_{\text{K}}$ or $_{\text{degF}_0} = (273.15 - 32*(5/9)) * _K$, the zero point temperatures of the scales.

In the following problem, Listing 4 , the task is to convert temperatures given in the unit degree Celsius and degree Fahrenheit to Kelvin.

Listing 4: Temperature conversion.

```
\begin{luacode}
                                                                    1
                                                                    2
 theta_1 = 110 * _degC
 T_1 = ( theta_1 + _degC_0 ):to(_K)
                                                                    3
                                                                    4
 T_2 = 100 * _K
                                                                    5
 theta_2 = (T_2 - degC_0):to(degC)
                                                                    6
                                                                    7
 theta_3 = 212 * _degF
                                                                    8
 T_3 = ( theta_3 + _degF_0 ):to(_K)
                                                                    9
```

```
10
 T_4 = 100 * _K
theta_4 = ( T_4 - _degF_0 ):to(_degF)
                                                                 11
                                                                 12
                                                                 13
  theta_5 = 100 * _degC
                                                                 14
  theta_6 = ( (theta_5 + _degC_0 ):to(_K) - _degF_0):to(_degF)
                                                                 15
\end{luacode}
                                                                 16
                                                                 17
\begin{align*}
                                                                 18
 19
                                                                 20
  \q\{theta_2\} \ \mathrel{\widehat{=}} \ \q\{T_2\} \ \
                                                                 21
                                                                 22
  \q\{theta_3\} \&\\mathrel\{\widehat\{=\}\} \q\{T_3\} \
                                                                 23
                                                                 24
  \q\{theta_4\} \&\mathrel\{\widehat\{=\}\} \q\{T_4\} \
                                                                 25
                                                                 26
  \q{theta_5}  &\mathbf{\d} \
                                                                 27
\end{align*}
                                                                 28
```

```
110 \,^{\circ}\text{C} \, \widehat{=} \, 383.15 \,\text{K}
-173.15 \,^{\circ}\text{C} \, \widehat{=} \, 100 \,\text{K}
212 \,^{\circ}\text{F} \, \widehat{=} \, 373.15 \,\text{K}
-279.67 \,^{\circ}\text{F} \, \widehat{=} \, 100 \,\text{K}
100 \,^{\circ}\text{C} \, \widehat{=} \, 212.0 \,^{\circ}\text{F}
```

3.2 Uncertainty Propagation

The package supports uncertainty propagation. To create a number with an uncertainty, an instance of physical.Number has to be created, see Listing 5. It has to be remembered, that N is a alias for physical.Number. The first argument of the constructor N(mean, uncertainty) is the mean value and the second one the uncertainty of the measurement. If the proposed preamble 1 is used, the uncertainty is by default separated from the mean value by a plus-minus sign. For the uncertainty propagation the gaussian formula

$$\Delta f = \sqrt{\left(\frac{\partial f}{x_1} \cdot \Delta x_1\right)^2 + \dots + \left(\frac{\partial f}{x_n} \cdot \Delta x_2\right)^2}$$

is used. This formula is a good estimation for the uncertainty Δf , if the quantities x_1, \ldots, x_n the function f depends on, have no correlation. Further, the function f has to change linear, if quantities x_i are changed in the range of their uncertainties.

Listing 5: Uncertainty in area calculation.

```
\begin{luacode}
                                                                                                                             1
   a = N(2,0.1) * _m
                                                                                                                             2
   b = N(3,0.1) * _m
                                                                                                                             3
                                                                                                                             4
   A = (a*b):to(_m^2)
                                                                                                                             5
\end{luacode}
                                                                                                                             6
                                                                                                                             7
Calculate the area of a rectangle with lengths \q{a}\ and \q{b}
                                                                                                                             8
                                                                                                                             9
$$
   A = a \setminus cdot b
                                                                                                                             10
   = \q{a} \cdot \q{b}
                                                                                                                             11
      \displaystyle \begin{array}{l} \displaystyle \begin{array}{l} \displaystyle \begin{array}{l} \displaystyle \begin{array}{l} \displaystyle \\ \displaystyle \end{array} \end{array} \end{array}
                                                                                                                             12
                                                                                                                             13
```

Calculate the area of a rectangle with lengths $(2.00 \pm 0.10)\,\mathrm{m}$ and $(3.00 \pm 0.10)\,\mathrm{m}$.

$$A = a \cdot b = (2.00 \pm 0.10) \,\mathrm{m} \cdot (3.00 \pm 0.10) \,\mathrm{m} = \underline{(6.0 \pm 0.4) \,\mathrm{m}^2}$$

Instead of printing the uncertainties, one can use the uncertainty calculation to provide significant digits and omit it.

In the following problem, listing 6, the task is to find the volume of an ideal gas. Given are pressure p in _bar, amount of substance n in _mol and temperature T in degree celsius _degC. In order to do the calculation, one has to convert T, which is given as an absolute temperature in degree celsius to the base unit Kelvin first. By setting N.omitUncertainty = true, all uncertainties are not printed.

Listing 6: Volume of an ideal gas.

```
\begin{luacode}
                                                                                                   1
   N.omitUncertainty = true
                                                                                                   2
   p = N(1.013, 0.0001) * _bar
                                                                                                   3
  n = N(1,0.01) * _mol
                                                                                                    4
  T = N(30, 0.1) * _degC
                                                                                                   5
                                                                                                   6
   V = (n * _R * (T + _degC_0):to(_K) / p):to(_L)
                                                                                                   7
\end{luacode}
                                                                                                   8
                                                                                                   9
An ideal gas (q{n}) has a pressure of q{p} and a
                                                                                                   10
         temperature of q{T}. Calculate the volume of the gas.
$$
                                                                                                   11
   V = \frac{ q\{n\} \cdot q\{n\} \cdot q\{R\} \cdot q\{(T + _degC_0):to(K)\} }{
                                                                                                   12
            \q{p} \
   = \q{V}
                                                                                                   13
     \displaystyle \begin{array}{l} \displaystyle \begin{array}{l} \displaystyle \begin{array}{l} \displaystyle \begin{array}{l} \displaystyle \\ \displaystyle \end{array} \end{array} \end{array} 
                                                                                                   14
$$
                                                                                                   15
```

An ideal gas (1.0 mol) has a pressure of 1.013 bar and a temperature of 30 $^{\circ}\mathrm{C}.$ Calculate the volume of the gas.

$$V = \frac{1.0\,{\rm mol} \cdot 8.31\,{\rm J/(mol\,K)} \cdot 303\,{\rm K}}{1.013\,{\rm bar}} = \underline{25\,{\rm L}}$$

This example shows, that the result has only two digits. If more digits are needed, the uncertainties of the given quantities should be smaller.

3.3 Mathematical operations

Two physical quantities with identical dimensions can be added or subtracted. The library checks the validity of those operations and throws an error if two addends haven't the same dimensions.

Listing 7: Lua console: Addition and subtraction

```
1_1 = 1 * _m
                                                                         1
1_{2} = 2 * _{cm}
                                                                         2
t = 2 * _s
                                                                         3
                                                                         4
1_1 + t
                                                                         5
Error: Cannot add '1* _{m}' to '2 * _{s}', because they have
                                                                         6
      different dimensions.
                                                                         7
1_1 + 1_2
                                                                         8
102.0 * _cm
                                                                         9
```

New physical quantities can be created by division and multiplication. As long as no division by zero is made, no errors should occur.

Listing 8: Lua console: Multiplication and Division

```
1_1 = 1 * _m

1_2 = 2 * _cm

(1_1 * 1_2):to(_m^2)

0.02 * _m^2

(1_1 / 1_2):to(_1)

50.0 * _1
```

Physical quantities can be exponentiated. The library doesn't check, if the result has units with non integer exponents.

Listing 9: Lua console: Exponentiation

```
1 = 5 * _m
A = 1^2
                                                                        2
                                                                        3
A:to(_m^2)
                                                                        4
25.0 * _m^2
                                                                        5
A:sqrt()
                                                                        7
5.0 * _m
                                                                        8
                                                                        9
A^0.5
                                                                        10
5.0 * _m
                                                                        11
```

Exponential functions an the logarithms should have dimensionless arguments. The library throws an error if that's not the case.

Listing 10: Lua console: Exponential function and logarithm

4 Supported Units

All supported units are listed in this chapter. Subchapter 4.2 lists the seven base units of the International System of Units (SI). In subchapter 4.3 mathematical and physical constants are defined. The subchapter 4.4 contains all coherent derived units from the SI system and 4.5 those which are accepted to use with the SI.

The subchapter 4.6 lists nominal astronomical units, which are proposed by [4]. Subchapter 4.7 lists units, which are common but outside of the SI system. The subchapters 4.8 and 4.9 are dedicated to imperial and U.S. customary units. The last subchapter 4.10 containts international currencies.

4.1 Prefixes

All SI units have prefixed versions, i.e. _us microsecond, _cm centimeter, _mN millinewton, see table 1. Some units of data processing, like _bit have prefixes which are powers of 2. They are called binary or IEC prefixes, see table 2. [2, 121]

Prefix	Symbol	Definition	Prefix	Symbol	Definition
yotta	Y	1e24	deci	d	1e-1
zetta	Z	1e21	centi	С	1e-2
exa	E	1e18	milli	m	1e-3
peta	P	1e15	micro	u	1e-6
tera	Т	1e12	nano	n	1e-9
giga	G	1e9	pico	p	1e-12
mega	M	1e6	femto	f	1e-15
kilo	k	1e3	atto	a	1e-18
hecto	h	1e2	zepto	z	1e-21
deca	da	1e1	yocto	У	1e-23

Table 1: SI prefixes [2, 121]

Prefix	Symbol	Definition
kibi	Ki	1024
mebi	Mi	1048576
gibi	Gi	1073741824
tebi	Ti	1099511627776
pebi	Pi	1125899906842624
exbi	Ei	1152921504606846976
zebi	Zi	1180591620717411303424
yobi	Yi	1208925819614629174706176

Table 2: IEC prefixes [2, 121]

4.2 Base Units

The lua-physical library has nine base quantities. These are the seven basis units or basis quantities of the SI system [3] and in addition the base quantity of information _bit and of currency _EUR. All other quantities are derived from these base units.

Quantity	Unit	Symbol	Dim.	Definition	
number 1	_	_1	1	The dimensionless number one.	
time	second	_s	Τ	The SI unit of time. It is defined by taking the fixed numerical value of the caesium frequency $\Delta\nu_{Cs}$, the unperturbed ground-state hyperfine transition frequency of the caesium 133 atom, to be 9 192 631 770 when expressed in the unit 1/s.	
length	meter	_m	L	The SI unit of length. It is defined by taking the fixed numercial value of the speed of light in vacuum c to be 299 792 458 when expressed in the unit of $1.0\mathrm{m/s}$.	

 $^{^1{\}rm The}$ number one is a unit with dimension zero. Stricly speaking it is not a base unit.

Quantity	Unit	Symbol	Dim.	Definition
mass	kilogram	_kg	M	The SI unit of mass. It is defined by taking the fixed numerical value of the Planck constant h to be $6.62607015\cdot10^{-34}$ when expressed in m ² kg/s.
electric current	ampere	_A	I	The SI unit of electric current. It is defined by taking the fixed numerical value of the elementary charge e to be $1.602176634\cdot10^{-19}$ when expressed in As.
thermody- namic temperature	kelvin	_K	K ¹	The SI unit of the thermodynamic temperature. It is defineed by taking the fixed numerical value of the Boltzmann constant k_B to be $1.380649\cdot10^{-23}$ when expressed in $1.0\mathrm{kg}\mathrm{m}^2/(\mathrm{s}^2\mathrm{K})$
amount of substance	mole	_mol	N	The SI unit of amount of substance. One mole contains exactly $6.02214076\cdot10^{23}$ elementary entities. This number is the fixed numerical value of the Avogadro constant N_A when expressed in 1/mol.
luminous intensity	candela	_cd	J	The SI unit of luminous intensity in a given direction. It is defined by taking the fixed numerical value of the luminous efficacy of monochromatic radiation of frequency $5.4 \cdot 10^{14} \mathrm{Hz}$, K_{cd} , to be 683 when expressed in the unit $\mathrm{cd} \mathrm{sr} \mathrm{s}^3 / (\mathrm{kg} \mathrm{m}^2)$.
informa- tion	bit	_bit	В	The smallest amount of information.
currency	euro	_EUR	С	The value of the currency Euro.

Table 3: Base units

 $^{^1}$ The SI symbol for the dimension of temperature is Θ , but all symbols of this library consist of roman letters, numbers and underscores only. Therefore the symbol for the dimension of the thermodynamic temperature is the letter K.

4.3 Constants

All physical constants are taken from the NIST webpage [1].

Name	Symbol	Definition
pi	_Pi	3.1415926535897932384626433832795028841971 * _1
eulersnumber	_E	2.7182818284590452353602874713526624977572 * _1
speedoflight	_c	299792458 * _m/_s
gravitationalconstant	_Gc	N(6.67408e-11,3.1e-15) * _m^3/(_kg*_s^2)
planckconstant	_h_P	6.62607015e-34 * _J*_s
${\it reduced planck constant}$	_h_Pbar	_h_P/(2*_Pi)
elementarycharge	_e	1.602176634e-19 * _C
vacuumpermeability	_u_0	4e-7*Pi * _N/_A^2
vacuumpermitivity	_e_0	1/(_u_0*_c^2)
atomicmassunit	_u	N(1.66053904e-27, 2e-35) * _kg
electronmass	_m_e	N(9.10938356e-31, 1.1e-38) * _kg
protonmass	_m_p	N(1.672621898e-27, 2.1e-35) * _kg
neutronmass	_m_n	N(1.674927471e-27, 2.1e-35) * _kg
bohrmagneton	_u_B	_e*_h_Pbar/(2*_m_e)
nuclearmagneton	_u_N	_e*_h_Pbar/(2*_m_p)
electronmagneticmoment	_u_e	N(-928.4764620e-26,5.7e-32) * _J/_T
protonmagneticmoment	_u_p	N(1.4106067873e-26,9.7e-35) * _J/_T
neutronmagneticmoment	_u_n	N(-0.96623650e-26,2.3e-26) * _J/_T
fine structure constant	_alpha	_u_0*_e^2*_c/(2*_h_P)
rydbergconstant	_Ry	_alpha^2*_m_e*_c/(2*_h_P)
avogadronumber	_N_A	6.02214076e23/_mol
boltzmannconstant	_k_B	1.380649e-23 * _J/_K
molargasconstant	_R	N(8.3144598, 4.8e-6) * _J/(_K*_mol)
${\it stefanboltz} mann constant$	_sigma	_Pi^2*_k_B^4/(60*_h_Pbar^3*_c^2)
standardgravity	_g_0	9.80665 * _m/_s^2

Table 4: Physical and mathematical constants

4.4 Coherent derived units in the SI

All units in this section are coherent derived units from the SI base units with special names, [2, 118].

Quantity	Unit	Symbol	Definition
Plane Angle ¹	radian	_rad	_1
Solid Angle ²	steradian	_sr	_rad^2
Frequency	hertz	_Hz	1/_s
Force	newton	_N	_kg*_m/_s^2
Pressure	pascal	_Pa	_N/_m^2
Energy	joule	_J	_N*_m
Power	watt	_W_	_J/_s
Electric Charge	$\operatorname{coulomb}$	_C	_A*_s
Electric Potential	volt	_V	_J/_C
Electric Capacitance	farad	_F	_C/_V
Electric Resistance	ohm	_Ohm	_V/_A
Electric Conductance 3	siemens	_S	_A/_V
Magnetic Flux	weber	_Wb	_V*_s
Magnetic Flux Density	tesla	_T	_Wb/_m^2
Inductance	henry	_H	_Wb/_A
$Temperature^4$	celsius	_degC	_K
Luminous Flux	lumen	_lm	_cd*_sr
Illuminance	lux	_lx	_lm/_m^2
Activity	becquerel	_Bq	1/_s
Absorbed Dose	gray	_Gy	_J/_kg
Dose Equivalent	sievert	_Sv	_J/_kg
Catalytic Activity	katal	_kat	_mol/_s

 $^{^{1}\}mathrm{In}$ the SI system, the quantity Plane Angle has the dimension of a number.

 $^{^2\}mathrm{In}$ the SI system, the quantity Solid Angle has the dimension of a number.

³The unit _PS stands for peta siemens and is in conflict with the metric version of the unit horsepower (german Pferdestärke). Since the latter is more common than peta siemens, _PS is defined to be the metric version of horsepower.

 $^{^4{\}rm The~unit~_degC}$ is by default interpreted as a temperature difference.

4.5 Non-SI units accepted for use with the SI

There are a few units with dimension 1. [2, 124].

Quantity	Unit	Symbol	Definition
Time	minute	_min	60 * _s
	hour	_h	60 * _min
	day	_d	24 * _h
Plane Angle	degree	_deg	(_Pi/180) * _rad
	arcminute	_arcmin	_deg/60
	arcsecond	_arcsec	_arcmin/60
Area	hectare	_hectare	1e4 * _m^2
Volume	liter	_L	1e-3 * _m^3
Mass	tonne	_t	1e3 * _kg

4.6 Nominal Astronomical Units

The nominal values of solar, terrestrial and jovial quantities are taken from IAU Resolution B3 [4].

Quantity	Unit	Symbol	Definition
Length	nomsolradius	_R_S_nom	6.957e8 * _m
Irradiance	${\bf nom solir radiance}$	_S_S_nom	1361 * _W/_m^2
Radiant Flux	nom solluminosity	$_{\tt L_S_nom}$	3.828e26 * _W
Temperature	nom so leff temperature	_T_S_nom	5772 * _K
Mass Parameter	nom sol mass parameter	_GM_S_nom	1.3271244e20 * _m^3*_s^-2
Length	nomterreqradius	_Re_E_nom	6.3781e6 * _m
Length	nomterrpolradius	_Rp_E_nom	6.3568e6 * _m
Mass Parameter	nom terr mass parameter	_GM_E_nom	3.986004e14 * _m^3*_s^-2
Length	nomjoveqradius	_Re_J_nom	7.1492e7 * _m
Length	nomjovpolradius	_Rp_J_nom	6.6854e7 * _m
Mass Parameter	${\bf nomjov mass parameter}$	_GM_J_nom	1.2668653e17 * _m^3*_s^-2

4.7 Other Non-SI units

The unit Bel is only available with prefix decibel, because _B is the unit byte.

Quantity	Unit	Symbol	Definition
Length	angstrom	_angstrom	1e-10 * _m
	fermi	_fermi	1e-15 * _m
Time	svedberg	_svedberg	1e-13 * _s
	week	_wk	7 * _d
	year	_a	365.25 * _d
	astronomicalunit	_au	149597870700 * _m
	lightsecond	_ls	_c*_s
	lightyear	_ly	_c*_a
	parsec	_pc	(648000/_Pi) * _au
Area	barn	_barn	1e-28 * _m^2
	are	_are	1e2 * _m^2
Volume	metricteaspoon	_tsp	5e-3 * _L
	metrictablespoon	_Tbsp	3 * _tsp
Plane Angle	gradian	_gon	(Pi/200) * _rad
	turn	_tr	2*Pi * _rad
Solid Angle	spat	_sp	4*Pi * _sr
Force	kilopond	_kp	_kg*_g_0
Pressure	bar	_bar	1e5 * _Pa
	standardatmosphere	_atm	101325 * _Pa
	technical atmosphere	_at	_kp/_cm^2
	millimeterofmercury	_mmHg	133.322387415 * _Pa
	torr	_Torr	(101325/760) * _Pa
Energy	thermochemical calorie	_cal	4.184 * _J
	internationalcalorie	_cal_IT	4.1868 * _J
	gramoftnt	_g_TNT	1e3 * _cal
	tonoftnt	_t_TNT	1e9 * _cal

Quantity	Unit	Symbol	Definition
	electronvolt	_eV	_e*_V
	wattsecond	_Ws	_W*_s
	watthour	_Wh	_W*_h
Power	voltampere	_VA	_V*_A
Electric Charge	amperesecond	_As	_A*_s
	amperehour	_Ah	_A*_h
Information	nibble	_nibble	4 * _bit
	byte	_B	8 * _bit
Information Transfer Rate	bitpersecond	_bps	_bit/_s
Number	percent	_percent	1e-2 * _1
	permille	_permille	1e-3 * _1
	partspermillion	_ppm	1e-6 * _1
	partsperbillion	_ppb	1e-9 * _1
	partspertrillion	_ppt	1e-12 * _1
	partsperquadrillion	_ppq	1e-15 * _1
	decibel	_dB	_1
Power	metrichorsepower	_PS	75 * _g_0*_kg*_m/_s
Activity	curie	_Ci	3.7e10 * _Bq
Absorbed Dose	rad	_Rad	1e-2 * _Gy
Dose Equivalent	rem	_rem	1e-2 * _Sv
Viscosity	poiseuille	_P1	_Pa*_s

4.8 Imperial Units

Quantity	Unit	Symbol	Definition
Length	inch	_in	2.54e-2 * _m
	thou	_th	1e-3 * _in
$DTP Point^1$	point	_pt	_in/72
	pica	_pica	12 * _pt
	hand	_hh	4 * _in
	foot	_ft	12 * _in
	yard	_yd	3 * _ft
	rod	_rd	5.5 * _yd
	chain	_ch	4 * _rd
	furlong	_fur	10 * _ch
	mile	_mi	8 * _fur
	league	_lea	3*_mi
	nautical mile	_nmi	1852 * _m
	${\it nautical league}$	_nlea	3 * _nmi
	cable	_cbl	0.1 * _nmi
	fathom	_ftm	6 * _ft
Velocity	knot	_kn	_nmi/_h
Area	acre	_ac	10 * _ch^2
Volume	gallon	_gal	4.54609*_L
	quart	_qt	_gal/4
	pint	_pint	_qt/2
	cup	_cup	_pint/2
	gill	_gi	_pint/4
	fluidounce	_fl_oz	_gi/5
	fluiddram	_fl_dr	_fl_oz/8

 $^{^{1}}$ The desktop publishing point or PostScript point is 1/72 of an international inch.

Quantity	Unit	Symbol	Definition
Mass	grain	_gr	64.79891*_mg
	pound	_1b	7000*_gr
	ounce	_oz	_lb/16
	dram	_dr	_1b/256
	stone	_st	14*_lb
	quarter	_qtr	2*_st
	hundredweight	_cwt	4*_qtr
	longton	_ton	20*_cwt
	troypound	_lb_t	5760*_gr
	troyounce	_oz_t	_lb_t/12
	pennyweight	_dwt	24*_gr
	firkin	_fir	56*_1b
Time	sennight	_sen	7*_d
	fortnight	_ftn	14*_d
${\bf Temperature^1}$	fahrenheit	_degF	(5/9)*_K
Force	poundforce	_lbf	_lb*_g_0
	poundal	_pdl	_lb*_ft/_s^2
Mass	slug	_slug	_lbf*_s^2/_ft
Pressure	poundforcepersquareinch	_psi	_lbf/_in^2
Energy, Torque	the hembritish thermal unit	_BTU	(1897.83047608/1.8)*_J
Energy, Torque	int british thermal unit	_BTU_it	1055.05585262 * _J
Power	horsepower	_hp	33000*_ft*_lbf/_min

 $^{^1\}mathrm{The~unit~_degF}$ is by default interpreted as a temperature difference.

4.9 U.S. customary units

In the U.S., the length units are bound to the meter differently than in the imperial system. The followin definitions are taken from https://en.wikipedia.org/wiki/United_States_customary_units.

Quantity	Unit	Symbol	Definition
Length	ussurveyinch	_in_US	_m/39.37
	ussurveyhand	_hh_US	4 * _in_US
	ussurveyfoot	_ft_US	3 * _hh_US
	ussurveylink	_li_US	0.66 * _ft_US
	ussurveyyard	_yd_US	3 * _ft_US
	ussurveyrod	_rd_US	5.5 * _yd_US
	ussurveychain	_ch_US	4 * _rd_US
	us survey furlong	_fur_US	10 * _ch_US
	ussurveymile	_mi_US	8 * _fur_US
	us survey league	_lea_US	3 * _mi_US
	us survey fathom	_ftm_US	72 * _in_US
	ussurveycable	_cbl_US	120 * _ftm_US
Area	ussurveyacre	_ac_US	_ch_US * _fur_US
Volume	usgallon	_gal_US	231 * _in^3
	usquart	_qt_US	_gal_US/4
	uspint	_pint_US	_qt_US/2
	uscup	_cup_US	_pint_US/2
	usgill	_gi_US	_pint_US/4
	usfluidounce	_fl_oz_US	_gi_US/4
	ustablespoon	_Tbsp_US	_fl_oz_US/2
	usteaspoon	_tsp_US	_Tbsp_US/3
	usfluiddram	_fl_dr_US	_fl_oz_US/8
Mass	usquarter	_qtr_US	25 * _lb
	ushundred weight	_cwt_US	4 * _qtr_US
	uston	_ton_US	20 * _cwt_US

4.10 International Currencies

International currency units based on exchange rates from 7.3.2019, 21:00 UTC.

Name	Symbol	Definition
AfghanAfghani	_AFN	0.012 * _EUR
AfghanPul	_cAFN	0.01 * _AFN
AlbanianLek	_ALL	0.008 * _EUR
ArmenianDram	_AMD	0.0018 * _EUR
ArmenianLuma	_cAMD	0.01 * _AMD
AngolanKwanza	_AOA	0.0028 * _EUR
AngolanCentimo	_cAOA	0.01 * _AOA
ArgentinePeso	_ARS	0.021 * _EUR
ArgentineCentavo	_cARS	0.01 * _ARS
AustralianDollar	_AUD	0.63 * _EUR
AustralianCent	_cAUD	0.01 * _AUD
AzerbaijaniManat	_AZN	0.63 * _EUR
AzerbaijaniQepik	_cAZN	0.01 * _AZN
BosnianMark	_BAM	0.51 * _EUR
BosnianFenings	_cBAM	0.01 * _BAM
BangladeshiTaka	_BDT	0.011 * _EUR
BangladeshiPoisha	_cBDT	0.01 * _BDT
BurundianFranc	_BIF	0.00049 * _EUR
BurundianCentime	_cBIF	0.01 * _BIF
BolivianBoliviano	_B0B	0.13 * _EUR
BolivianCentavo	_cBOB	0.01 * _BOB
BrazilianReal	_BRL	0.23 * _EUR
BrazilianCentavo	_cBRL	0.01 * _BRL
BotswanaPula	_BWP	0.083 * _EUR
BotswanaThebe	_cBWP	0.01 * _BWP
BelarusianRuble	_BYN	0.42 * _EUR
BelarusianKapiejka	_cBYN	0.01 * _BYN

Name	Symbol	Definition
CanadianDollar	_CAD	0.66 * _EUR
CanadianCent	_cCAD	0.01 * _CAD
CongoleseFranc	_CDF	0.00055 * _EUR
CongoleseCentime	_cCDF	0.01 * _CDF
SwissFranc	_CHF	0.88 * _EUR
SwissRappen	_cCHF	0.01 * _CHF
ChileanPeso	_CLP	0.0013 * _EUR
ChileanCentavo	_cCLP	0.01 * _CLP
ChineseRenminbiYuan	_CNY	0.13 * _EUR
ChineseRenminbiFen	_cCNY	0.01 * _CNY
ColombianPeso	_COP	0.00028 * _EUR
ColombianCentavo	_cCOP	0.01 * _COP
CostaRicanColon	_CRC	0.0015 * _EUR
CostaRicanCentimos	_cCRC	0.01 * _CRC
CzechKoruna	_CZK	0.039 * _EUR
CzechHaler	_cCZK	0.01 * _CZK
DanishKrone	_DKK	0.13 * _EUR
DanishOre	_cDKK	0.01 * _DKK
DominicanPeso	_DOP	0.018 * _EUR
DominicanCentavo	_cDOP	0.01 * _DOP
AlgerianDinar	_DZD	0.0074 * _EUR
AlgerianSanteem	_cDZD	0.01 * _DZD
EgyptianPound	_EGP	0.051 * _EUR
EgyptianPiastre	_cEGP	0.01 * _EGP
EthiopianBirr	_ETB	0.031 * _EUR
EthiopianSantim	_cETB	0.01 * _ETB
FijianDollar	_FJD	0.42 * _EUR
FijianCent	_cFJD	0.01 * _FJD

Name	Symbol	Definition
PoundSterling	_GBP	1.16 * _EUR
PennySterling	_cGBP	0.01 * _GBP
GeorgianLari	_GEL	0.33 * _EUR
GeorgianTetri	_cGEL	0.01 * _GEL
GhanaianCedi	_GHS	0.16 * _EUR
GhanaianPesewa	_cGHS	0.01 * _GHS
GambianDalasi	_GMD	0.018 * _EUR
GambianButut	_cGMD	0.01 * _GMD
GuineanFranc	_GNF	9.6e-05 * _EUR
GuineanCentime	_cGNF	0.01 * _GNF
${\bf Guatemal an Quetzal}$	_GTQ	0.12 * _EUR
GuatemalanCentavo	_cGTQ	0.01 * _GTQ
GuyaneseDollar	_GYD	0.0043 * _EUR
GuyaneseCent	_cGYD	0.01 * _GYD
${\bf Hong Kong Dollar}$	_HKD	0.11 * _EUR
HongKongCent	_cHKD	0.01 * _HKD
HonduranLempira	_HNL	0.036 * _EUR
HonduranCentavo	_cHNL	0.01 * _HNL
CroatianKuna	_HRK	0.13 * _EUR
CroatianLipa	_cHRK	0.01 * _HRK
HaitianGourde	_HTG	0.011 * _EUR
HaitianCentime	_cHTG	0.01 * _HTG
HungarianForint	_HUF	0.0032 * _EUR
HungarianFiller	_cHUF	0.01 * _HUF
IndonesianRupiah	_IDR	6.2e-05 * _EUR
IndonesianSen	_cIDR	0.01 * _IDR
IsraeliNewShekel	_ILS	0.25 * _EUR
${\bf Israeli New Agora}$	_cILS	0.01 * _ILS

Name	Symbol	Definition
IndianRupee	_INR	0.013 * _EUR
IndianPaisa	_cINR	0.01 * _INR
IraqiDinar	_IQD	0.00074 * _EUR
IraqiFils	_cIQD	0.001 * _IQD
IranianRial	_IRR	2.7e-05 * _EUR
IranianToman	_cIRR	10.0 * _IRR
IcelandicKrona	_ISK	0.0073 * _EUR
JamaicanDollar	_JMD	0.007 * _EUR
JamaicanCent	_cJMD	0.01 * _JMD
JapaneseYen	_JPY	0.008 * _EUR
KenyanShilling	_KES	0.0089 * _EUR
KenyanCent	_cKES	0.01 * _KES
KyrgyzstaniSom	_KGS	0.013 * _EUR
KyrgyzstaniTyiyn	_cKGS	0.01 * _KGS
CambodianRiel	_KHR	0.00022 * _EUR
NorthKoreanWon	_KPW	0.00099 * _EUR
NorthKoreanChon	_cKPW	0.01 * _KPW
SouthKoreanWon	_KRW	0.00078 * _EUR
South Korean Jeon	_cKRW	0.01 * _KRW
KuwaitiDinar	_KWD	2.93 * _EUR
KuwaitiFils	_cKWD	0.001 * _KWD
KazakhstaniTenge	_KZT	0.0023 * _EUR
KazakhstaniTiyn	_cKZT	0.01 * _KZT
LaoKip	_LAK	0.0001 * _EUR
LaoAtt	_cLAK	0.01 * _LAK
SriLankanRupee	_LKR	0.005 * _EUR
SriLankanCent	_cLKR	0.01 * _LKR
LiberianDollar	_LRD	0.0055 * _EUR
LiberianCent	_cLRD	0.01 * _LRD

Name	Symbol	Definition
LibyanDinar	_LYD	0.64 * _EUR
LibyanDirham	_cLYD	0.001 * _LYD
MoroccanDirham	_MAD	0.092 * _EUR
MoroccanSantim	_cMAD	0.01 * _MAD
MoldovanLeu	_MDL	0.052 * _EUR
MoldovanBan	_cMDL	0.01 * _MDL
MalagasyAriary	_MGA	0.00025 * _EUR
MalagasyIraimbilanja	_cMGA	0.2 * _MGA
MacedonianDenar	_MKD	0.016 * _EUR
MacedonianDeni	_cMKD	0.01 * _MKD
BurmeseKyat	_MMK	0.00059 * _EUR
BurmesePya	_cMMK	0.01 * _MMK
MongolianTogrog	_MNT	0.00034 * _EUR
MongolianMongo	_cMNT	0.01 * _MNT
MauritanianOuguiya	_MRU	0.025 * _EUR
MauritanianKhoums	_cMRU	0.2 * _MRU
MauritianRupee	_MUR	0.025 * _EUR
MauritianCent	_cMUR	0.01 * _MUR
MaldivianRufiyaa	_MVR	0.058 * _EUR
MaldivianLaari	_cMVR	0.01 * _MVR
MalawianKwacha	_MWK	0.0012 * _EUR
MalawianTambala	_cMWK	0.01 * _MWK
MexicanPeso	_MXN	0.046 * _EUR
MexicanCentavo	_cMXN	0.01 * _MXN
MalaysianRinggit	_MYR	0.22 * _EUR
MalaysianSen	_cMYR	0.01 * _MYR
MozambicanMetical	_MZN	0.014 * _EUR
MozambicanCentavo	_cMZN	0.01 * _MZN

Name	Symbol	Definition
NigerianNaira	_NGN	0.0025 * _EUR
NigerianKobo	_cNGN	0.01 * _NGN
NicaraguanCordoba	_NIO	0.027 * _EUR
NicaraguanCentavo	_cNIO	0.01 * _NIO
NorwegianKrone	_NOK	0.1 * _EUR
NorwegianOre	_cNOK	0.01 * _NOK
New Zeal and Dollar	_NZD	0.61 * _EUR
NewZealandCent	_cNZD	0.01 * _NZD
PeruvianSol	_PEN	0.27 * _EUR
PeruvianCentimo	_cPEN	0.01 * _PEN
PapuaNewGuineanKina	_PGK	0.26 * _EUR
PapuaNewGuineanToea	_cPGK	0.01 * _PGK
PhilippinePeso	_PHP	0.017 * _EUR
PhilippineSentimo	_cPHP	0.01 * _PHP
PakistaniRupee	_PKR	0.0064 * _EUR
PakistaniPaisa	_cPKR	0.01 * _PKR
PolishZloty	_PLN	0.23 * _EUR
PolishGrosz	_cPLN	0.01 * _PLN
ParaguayanGuarani	_PYG	0.00015 * _EUR
ParaguayanCentimo	_cPYG	0.01 * _PYG
QatariRiyal	_QAR	0.24 * _EUR
QatariDirham	_cQAR	0.01 * _QAR
RomanianLeu	_RON	0.21 * _EUR
RomanianBan	_cRON	0.01 * _RON
SerbianDinar	_RSD	0.0085 * _EUR
SerbianPara	_cRSD	0.01 * _RSD
RussianRuble	_RUB	0.013 * _EUR
RussianKopeyka	_cRUB	0.01 * _RUB

Name	Symbol	Definition
RwandanFranc	_RWF	0.00098 * _EUR
RwandanCentime	_cRWF	0.01 * _RWF
${\bf Solomon Islands Dollar}$	_SBD	0.11 * _EUR
SolomonIslandsCent	_cSBD	0.01 * _SBD
SeychelloisRupee	_SCR	0.065 * _EUR
SeychelloisCent	_cSCR	0.01 * _SCR
SudanesePound	_SDG	0.019 * _EUR
${\bf SudaneseQirsh}$	_cSDG	0.01 * _SDG
SwedishKrona	_SEK	0.094 * _EUR
SwedishOre	_cSEK	0.01 * _SEK
SingaporeDollar	_SGD	0.65 * _EUR
SingaporeCent	_cSGD	0.01 * _SGD
SierraLeoneanLeone	_SLL	0.0001 * _EUR
SierraLeoneanCent	_cSLL	0.01 * _SLL
SomalilandShilling	_SQS	0.00013 * _EUR
SomalilandCent	_cSQS	0.01 * _SQS
SomaliShilling	_SOS	0.0015 * _EUR
SomaliSenti	_cSOS	0.01 * _SOS
SurinameseDollar	_SRD	0.12 * _EUR
SurinameseCent	_cSRD	0.01 * _SRD
SyrianPound	_SYP	0.0017 * _EUR
SyrianPiastre	_cSYP	0.01 * _SYP
ThaiBaht	_THB	0.028 * _EUR
ThaiSatang	_cTHB	0.01 * _THB
TajikistaniSamani	_TJS	0.094 * _EUR
TajikistaniDiram	_cTJS	0.01 * _TJS
Tonganpaanga	_TOP	0.397 * _EUR
TonganSeniti	_cTOP	0.01 * _TOP

Name	Symbol	Definition
TurkishLira	_TRY	0.16 * _EUR
TurkishKurus	_cTRY	0.01 * _TRY
${\bf Trinidad And Tobago Dollar}$	_TTD	0.13 * _EUR
${\bf Trinidad And Tobago Cent}$	_cTTD	0.01 * _TTD
NewTaiwanDollar	_TWD	0.029 * _EUR
NewTaiwanCent	_cTWD	0.01 * _TWD
TanzanianShilling	_TZS	0.00038 * _EUR
TanzanianSenti	_cTZS	0.01 * _TZS
UkrainianHryvnia	_UAH	0.00038 * _EUR
UkrainianKopiyka	_cUAH	0.01 * _UAH
UgandanShilling	_UGX	0.00024 * _EUR
UgandanCent	_cUGX	0.01 * _UGX
USDollar	_USD	0.89 * _EUR
USCent	_cUSD	0.01 * _USD
UruguayanPeso	_UYU	0.027 * _EUR
UruguayanCentesimo	_cUYU	0.01 * _UYU
UzbekistaniSom	_UZS	0.00011 * _EUR
UzbekistaniTiyin	_cUZS	0.01 * _UZS
${\bf Venezuel an Bolivar Soberano}$	_VES	0.0003 * _EUR
Venezuelan Centimo Soberano	_cVES	0.01 * _VES
VietnameseDong	_VND	3.8e-05 * _EUR
VietnameseXu	_cVND	0.01 * _VND
SamoanTala	_WST	0.34 * _EUR
SamoanSene	_cWST	0.01 * _WST
YemeniRial	_YER	0.0036 * _EUR
YemeniDinar	_cYER	0.01 * _YER
SouthAfricanRand	_ZAR	0.062 * _EUR
SouthAfricanCent	_cZAR	0.01 * _ZAR
ZambianKwacha	_ZMW	0.074 * _EUR
ZambianNgwee	_cZMW	0.01 * _ZMW

4.10.1 Pegged International Currencies

International currency which are pegged to other currencies.

Name	Symbol	Definition
UnitedArabEmiratesDirham	_AED	(1/3.6725) * _USD
${\bf United Arab Emirates Fils}$	_cAED	0.01 * _AED
Netherlands Antillean Guilder	_ANG	(1/1.79) * _USD
${\bf Netherlands Antillean Cent}$	_cANG	0.01 * _ANG
ArubanFlorin	_AWG	(1/1.79) * _USD
ArubanCent	_cAWG	0.01 * _AWG
BarbadianDollar	_BBD	0.5 * _USD
BarbadianCent	_cBBD	0.01 * _BBD
BulgarianLev	_BGN	0.51129 * _EUR
BulgarianStotinka	_cBGN	0.01 * _BGN
BahrainiDinar	_BHD	(1/0.376) * _USD
BahrainiFils	_cBHD	0.001 * _BHD
BermudianDollar	_BMD	1 * _USD
BermudianCent	_cBMD	0.01 * _BMD
BruneiDollar	_BND	1 * _SGD
BruneiSen	_cBND	0.01 * _BND
BahamianDollar	_BSD	1 * _USD
BahamianCent	_cBSD	0.01 * _BSD
Bhutanese Ngultrum	_BTN	1 * _INR
Bhutanese Chhertum	_cBTN	0.01 * _BTN
BelizeDollar	_BZD	0.5 * _USD
BelizeCent	_cBZD	0.01 * _BZD
${\bf Cubano Convertible Peso}$	_CUC	1 * _USD
${\bf Cubano Convertible Centavo}$	_cCUC	0.01 * _CUC
CubanPeso	_CUP	(1/24) * _CUC
CubanCentavo	_cCUP	0.01 * _CUP

Name	Symbol	Definition
CapeVerdeanEscudo	_CVE	(1/110.265) * _EUR
CapeVerdeanCentavo	_cCVE	0.01 * _CVE
DjiboutianFranc	_DJF	(1/177.721) * _USD
DjiboutianCentime	_cDJF	0.01 * _DJF
EritreanNakfa	_ERN	(1/15) * _USD
EritreanCent	_cERN	0.01 * _ERN
FalklandIslandsPound	_FKP	1 * _GBP
FalklandIslandsPenny	_cFKP	0.01 * _FKP
GuernseyPound	_GGP	1 * _GBP
GuernseyPenny	_cGGP	0.01 * _GGP
GibraltarPound	_GIP	1 * _GBP
GibraltarPenny	_cGIP	0.01 * _GIP
ManxPound	_IMP	1 * _GBP
ManxPenny	_cIMP	0.01 * _IMP
JerseyPound	_JEP	1 * _GBP
JerseyPenny	_cJEP	0.01 * _JEP
JordanianDinar	_JOD	(1/0.708) * _USD
JordanianFils	_cJOD	0.001 * _JOD
KiribatiDollar	_KID	1 * _AUD
KiribatiCent	_cKID	0.01 * _KID
Comorianfranc	_KMF	(1/491.96775) * _EUR
ComorianCentime	_cKMF	0.01 * _KMF
CaymanIslandsDollar	_KYD	1.2 * _USD
CaymanIslandsCent	_cKYD	0.01 * _KYD
LebanesePound	_LBP	(1/1507.5) * _USD
LebaneseQeresh	_cLBP	0.01 * _LBP
MacanesePataca	_MOP	(1/1.03) * _HKD
MacaneseAvo	_cMOP	0.01 * _MOP

NT	C11	D-6-:::
Name		Definition
NamibianDollar	_NAD	1 * _ZAR
NamibianCent	_cNAD	O.O1 * _NAD
NepaleseRupee	_NPR	(1/1.6) * _INR
NepalesePaisa	_cNPR	0.01 * _NPR
OmaniRial	_OMR	(1/2.6008) * _USD
OmaniBaisa	_cOMR	0.001 * _OMR
PanamanianBalboa	_PAB	1 * _USD
PanamanianCentesimo	_cPAB	0.01 * _PAB
TransnistrianRuble	_PRB	(1/16.1) * _USD
TransnistrianKopeck	_cPRB	0.01 * _PRB
SaudiRiyal	_SAR	(1/3.75) * _USD
SaudiHalalah	_cSAR	0.01 * _SAR
SaintHelenaPound	_SHP	1 * _GBP
SaintHelenaPenny	_cSHP	0.01 * _SHP
SouthSudanesePound	_SSP	1 * _SDG
South Sudanese Piaster	_cSSP	0.01 * _SSP
${\bf Sao Tome And Principe Dobra}$	_STN	(1/24.5) * _EUR
${\bf Sao Tome And Principe Centimo}$	_cSTN	0.01 * _STN
SwaziLilangeni	_SZL	1 * _ZAR
SwaziCent	_cSZL	0.01 * _SZL
TurkmenistanManat	_TMT	(1/3.5) * _USD
TurkmenistanTenge	_cTMT	0.01 * _TMT
TuvaluanDollar	_TVD	1 * _AUD
TuvaluanCent	_cTVD	0.01 * _TVD
${\bf Central A frican CFA Franc}$	_XAF	(1/655.957) * _EUR
${\bf Central African CFA Centime}$	_cXAF	0.01 * _XAF
EasternCaribbeanDollar	_XCD	(1/2.7) * _USD
EasternCaribbeanCent	_cXCD	0.01 * _XCD

Name	Symbol	Definition
WestAfricanCFAFranc	_XOF	(1/655.957) * _USD
${\bf West A frican CFA Centime}$	_cXOF	0.01 * _XOF
CFPFranc	_XPF	(1000/8.38) * _EUR
CFPCentime	_cXPF	0.01 * _XPF
ZimbabweanBonds	_ZWL	1 * _USD
ZimbabweanCent	_cZWL	0.01 * _ZWL

5 Lua Documentation

In this chapter, the following shortcuts will be used.

```
local D = physical.Dimension

local U = physical.Unit

local N = physical.Number

local Q = physical.Quantity

1
```

The term number refers to a lua integer or a lua float number. By string a lua string is meant and by bool a lua boolean.

5.1 physical.Quantity

The quantity class is the main part of the library. Each physical Quantity and all units are represented by an instance of this class.

Q.new(q=nil)

Copy Constuctor

```
\mathbf{q}: \mathbf{Q}, \, \mathbf{number}, \, \mathbf{object}, \, \mathbf{nil} returns: \mathbf{Q}
```

As an argument it takes Q, number, object or nil. If an instance of Q is given, a copy is made and returned. If a number or an instance object of another class is given, the function creates a dimensionless quantity with the number or the instance as a value. In the case nil is given, a dimensionless quantity with value 1 is returned.

Example

```
print( Q() )
1
1
2
3
print( Q(42) )
42
5
print( Q(73*_m) )
73 * _m
8
```

Q.defineBase(symbol,name,dimension)

This function is used to declare base quantities from which all other quantities are derived from.

symbol: string

The symbol of the base quantity.

name: string

The name of the base quantity.

dimension: D

An instance of the D class, which represents the dimension of the quantity.

returns: Q

The created Q instance.

The function creates a global variable of the created base quantity. The name consist of an underscore concatenated with the symbol argument, i.e. the symbol m becomes the global variable _m.

The name is used for example in the siunitx conversion function, e.g meter will be converted to \meter.

Each quantity has a dimension associated with it. The argument dimension allows any dimension to be associated to base quantities.

Example

```
Q.defineBase("m", "meter", L)
Q.defineBase("kg", "kilogram", M)
```

Q.define(symbol, name, q)

Creates a new derived quantity from an expression of other quantities. Affine quantities like the absolute temperature in celsius are not supported.

symbol: string

Symbol of the base quantity

name: string, nil

The Name of the derived quantity.

q: physical.Quantity

The definition of the derived quantity.

return: Quantity

The created quantity.

The function creates a global variable of the created base quantity. The name consist of an underscore concatenated with the <code>symbol</code> argument, i.e. the symbol N becomes the global variable _N.

The name is used for example in the siunitx conversion function, e.g newton will be converted to \newton.

Example

```
Q.define("L", "liter", _dm^3) 1
Q.define("Pa", "pascal", _N/_m^2) 2
Q.define("C", "coulomb", _A*_s) 3
Q.define("degC", "celsius", _K) 5
```

Q.definePrefix(symbol,name,factor)

Defines a new prefix.

symbol: string

Symbol of the base quantity

name: string

Name of the base quantity

factor: number

The factor which corresponds to the prefix

```
Q.definePrefix("c", "centi", 1e-2)
Q.definePrefix("a", "atto", 1e-18)
```

Q.addPrefix(prefixes, units)

Create several units with prefixes from a given unit.

prefixes: string

A list of unit symbols.

units: Quantity

A list of quantities.

```
Q.addPrefix({"n","u","m","k","M","G"},{_m,_s,_A})
```

Q.to(self,q)

Converts the quantity self to the unit of the quantity q.

self: Quantity

A quantity.

q: Quantity The quantity to wich self will be converted to.

```
s = 1.9 * _km
                                                                 1
print( s:to(_m) )
                                                                 2
1900.0 * _m
                                                                 3
                                                                 4
T = 10 * _degC
                                                                 5
print( T:to(_K) )
                                                                 6
10.0 * _K
print( T:to(_K) )
                                                                 7
                                                                 8
                                                                 9
10.0 * _K
```

Q.tosiunitx(self,param,mode)

Converts the quantity into a siunity string.

self : Quantity
param : string

 $mode : Number, 0:\SI, 1:\num, 2:\si$

```
s = 1.9 * _km
                                                              1
                                                              2
print( s:tosiunitx() )
                                                              3
SI{1.9}{\kappailo\meter}
                                                              4
                                                              5
print( s:tosiunitx(nil,1) )
                                                              6
\num{1.9}
                                                              7
                                                              8
print( s:tosiunitx(nil,2) )
                                                              9
\si{\kilo\meter}
                                                              10
```

Q.isclose(self,q,r)

Checks if this quantity is close to another one. The argument ${\tt r}$ is the maximal relative deviation.

self : Quantity

 ${\tt q}: {\tt Quantity}, {\tt Number}$

r: Number

```
s_1 = 1.9 * _m
s_2 = 2.0 * _m
print( s_1:isclose(s_2,0.1) )
true
print( s_1:isclose(s_2,0.01) )
false

1
2
4
6
```

Q.min(q1, q2, ...)

Returns the smallest quantity of several given ones. The function returns q1 if the Quantities are equal.

q1: Quantity, Number, first argument

q2: Quantity, Number, second argument

```
s_1 = 15 * _m

s_2 = 5 * _m

print(s_1:min(s_2))

5 * _m
```

Q.max(q1, q2, ...)

Returns the biggest quantity of several given ones. The function returns q1 if the Quantities are equal.

q1: Quantity, Number, first argument

q2: Quantity, Number, second argument

```
s_1 = 15 * _m
s_2 = 5 * _m
print(s_1:max(s_2))
15 * _m
1
2
4
```

Q.abs(q)

Returns the absolute value of the given quantity q.

q: Q, N, number dimensionless argument

Returns: Q

the absolute value of ${\tt q}$

```
U = -5 * _V
print(U)
2
-5 * _V
3
print(U:abs())
5 * _V
```

Q.sqrt(q)

Returns the square root of the given quantity.

q: Q, N, number

dimensionless argument

Returns: Q

the square root of q

```
A = 25 * _m^2 1
s = A:sqrt() 2
print(s) 3
5.0 * _m 4
```

Q.log(q, base=nil)

Returns the logarithm of a given quantity to the given base. If no base is given, the natural logarithm is returned.

q: Q, N, number

dimensionless argument

 ${\tt base}:\, {\tt Q},\, {\tt N},\, {\tt number},\, {\tt nil}$

dimensionless argument

Returns: Q

logarithm of q to the base

```
I = 1 * _W/_m^2
I_0 = 1e-12 * _W/_m^2
print(10 * (I/I_0):log(10) * _dB)
120.0 * _dB
1
2
```

Q.exp(q)

Returns the value of the natural exponential function of the given quantitiy.

q: Q, N, number

dimensionless argument

Returns: Q

natural exponential of q

```
x = 2 * _1
print( x:exp() ) 2
7.3890560989307 3
```

Q.sin(q)

Returns the value of the sinus function of the given quantitiy.

 $\begin{array}{c} q:\, Q,\, N,\, number\\ & {\rm dimensionless}\,\, argument \end{array}$

Returns : Q sine of q

Q.cos(q)

Returns the value of the cosinus function of the given quantity. The quantity has to be dimensionless.

q: Q, N, number dimensionless argument

Returns : Q cosine of q

```
alpha = 60 * _deg
print( alpha:cos() )
0.5
1
2
3
```

Q.tan(q)

Returns the value of the tangent function of the given quantity. The quantity has to be dimensionless.

 $\begin{array}{c} q: \; Q, \; N, \; number \\ & \mathrm{dimensionless} \; \mathrm{argument} \end{array}$

 $\begin{array}{c} {\rm Returns}:\, Q \\ {\rm tangent} \,\, {\rm of} \,\, q \end{array}$

Q.asin(q)

Returns the value of the arcus sinus function of the given quantity. The quantity has to be dimensionless.

 ${\tt q}:{\tt Q},\,{\tt N},\,{\tt number}$

dimensionless argument

Returns: Q

inverse sine of q

```
x = 0.5 * _1
print( x:asin():to(_deg) )
2
30.0 * _deg
3
```

Q.acos(q)

Returns the value of the arcus cosinus function of the given quantity. The quantity has to be dimensionless.

q: Q, N, number

dimensionless argument

Returns: Q

inverse cosine of q

```
x = 0.5 * _1
print( x:acos():to(_deg) )
60.0 * _deg
1
2
```

Q.atan(q)

Returns the value of the arcus tangent function of the given quantity. The quantity has to be dimensionless.

q: Q, N, number

dimensionless argument

Returns: Q

inverse tangent of ${\tt q}$

```
x = 1 * _1
print( x:atan():to(_deg) )
2
45.0 * _deg
1
```

Q.sinh(q)

Returns the value of the hyperbolic sine function of the given quantity. The quantity has to be dimensionless. Since Lua doesn't implement the hyperbolic functions, the following formula is used

$$\sinh(x) = 0.5 \cdot e^x - 0.5/e^x \quad .$$

q: Q, N, number

dimensionless argument

Returns: Q

hyperbolic sine of q

```
x = 1 * _1
print( x:sinh() ) 2
1.1752011936438 3
```

Q.cosh(q)

Returns the value of the hyperbolic cosine function of the given quantity. The quantity has to be dimensionless. Since Lua doesn't implement the hyperbolic functions, the following formula is used

$$\cosh(x) = 0.5 \cdot e^x + 0.5/e^x \quad .$$

q: Q, N, number

dimensionless argument

Returns: Q

hyperbolic cosine of q

```
x = 1 * _1
print(x:cosh())
2
1.5430806348152
3
```

Q.tanh(q)

Returns the value of the hyperbolic tangent function of the given quantity. The quantity has to be dimensionless. Since Lua doesn't implement the hyperbolic functions, the following formula is used

$$\tanh(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}} \quad .$$

q: Q, N, number

dimensionless argument

Returns: Q

hyperbolic tangent of ${\tt q}$

```
x = 1 * _1
print(x:tanh()) 2
0.76159415595576 3
```

Q.asinh(q)

Returns the value of the inverse hyperbolic sine function of the given quantity. The quantity has to be dimensionless. Since Lua doesn't implement the hyperbolic functions, the following formula is used

$$asinh(x) = \ln\left(x + \sqrt{x^2 + 1}\right)$$
.

q: Q, N, number

dimensionless argument

Returns: Q

inverse hyperbolic sine of q

Q.acosh(q)

Returns the value of the inverse hyperbolic cosine function of the given quantity. The quantity has to be dimensionless. Since Lua doesn't implement the hyperbolic functions, the following formula is used

$$a\cosh(x) = \ln\left(x + \sqrt{x^2 - 1}\right) \quad , x > 1 \quad .$$

q: Q, N, number

dimensionless argument bigger or equal to one

Returns: Q

inverse hyperbolic cosine of q

Q.atanh(q)

Returns the value of the inverse hyperbolic tangent function of the given quantity. The quantity has to be dimensionless. Since Lua doesn't implement the hyperbolic functions, the following formula is used

$$\operatorname{atanh}(x) = \ln\left(\frac{1+x}{1-x}\right) \quad , -1 < x < 1 \quad .$$

q: Q, N, number

dimensionless argument with magnitude smaller than one

Returns: Q

inverse hyperbolic tangent of q

5.2 physical. Dimension

All physical quantities do have a physical dimension. For example the quantity Area has the dimension L^2 (length to the power of two). In the SI-System there are seven base dimensions, from which all other dimensions are derived. Each dimension is represented by an n-tuple, where n is the number of base dimensions. Each physical quantity has an associated dimension object. It is used to check equality and if addition or substraction is allowed.

Dimension.new(q=nil)

Constructor of the Dimension class.

q: Dimension or string, optional

The name or symbol of the dimension. If ${\sf q}$ is a dimension, a copy of it is made. If no argument ist given, a dimension ${\it zero}$ is created.

return: Dimension

The created Quantity object

Example

```
V_1 = D("Velocity")

L = D("L")

V_2 = D(L/T)

1
2
```

5.3 physical.Unit

The task of this class is keeping track of the unit term. The unit term is a fraction of units. The units in the enumerator and denominator can have an exponent.

Unit.new(u=nil)

Copy Constructor. It copies a given unit object. If nothing is given, an empty unit is created.

u: Unit

The unit object which will be copied.

return : Unit

The created Unit object

Unit.new(symbol, name, prefixsymbol=nil, prefixname=nil)

Constructor. A new Unit object with symbol is created. The prefixsymbol and prefixname are optional.

symbol: String

The symbol of the unit.

name: String

The name of the unit.

prefixsymbol : String

The optional symbol of the prefix.

prefixname : String

The optional name of the prefix.

return : Unit

The created Unit object

Unit.tosiunitx(self)

The unit term will be compiled into a string, which the LaTeX package siunitx can understand.

return: String

The siunitx representation of the unit term.

5.4 physical.Number

It does arithmetics with gaussian error propagation. A number instance has a mean value called x and an uncertainty called dx.

Number.new(n=nil)

This is the copy Constructor. It copies a given number object. If n is nil, an instance representing number zero with uncertainty zero is created.

n: Number

The number object to be copied.

return: Number

The created Number instance.

Number.new(x, dx)

This constructor, creates a new instance with mean value ${\tt x}$ and uncertainty ${\tt dx}.$

x: number mean value

dx: number

uncertainty value

return : Number

The created Number instance.

Example

```
n = N(12,0.1)
print(n)

1
2
```

Number.new(str)

This constructor creates a new instance from a string. It can parse strings of the form 3.4, 3.4e-3, 5.4e-3 +/- 2.4e-6, 5.45(7)e-23.

Parameters / Return

str:string

The number as a string.

return : Number

The created Number object

Example

```
n_1 = N("12.3e-6")
  print(n_1)

1
2

n_2 = N("12 +/- 0.1")
  print(n_2)

5

n_3 = N("12.0(1)")
  print(n_3)
8
```

Number.mean(n)

Returns the mean value

Parameters / Return

return: number
The mean value

Number.uncertainty(n)

Returns the uncertainty value

Parameters / Return

return : number

The uncertainty value

Number.abs(n)

Returns the absolute value of the number.

Parameters / Return

return : number

The absolute value

Number.sqrt(n)

Returns the square root of the number.

Parameters / Return

return : number

The square root

References

- [1] Webpage https://physics.nist.gov/cuu/index.html, August 2019.
- [2] Bureau International des Poids et Mesures. The international system of units (si), 2006.
- [3] Bureau International des Poids et Mesures. Resolutions of the 26th cgpm, November 2018.
- [4] Prša et al. Nominal values for selected solar and planetary quantities: Iau 2015 resolution b3. The Astronomical Journal, 152:41, August 2016.

Index of Units

acre _ac, 20 amperehour _Ah, 19 amperesecond _As, 19 angstrom _angstrom, 18 arcminute _arcmin, 17 arcsecond _arcsec, 17 are _are, 18 astronomicalunit _au, 18 atomicmassunit , 15 avogadronumber , 15 bar _bar, 18 barn _barn, 18	fluiddram _fl_dr, 20 fluidounce _fl_oz, 20 foot _ft, 20 fortnight _ftn, 21 furlong _fur, 20 gallon _gal, 20 gill _gi, 20 gradian _gon, 18 grain _gr, 21 gramoftnt _g_TNT, 18 gravitationalconstant, 15 gray _Gy, 16
becquerel _Bq, 16 bitpersecond _bps, 19 bohrmagneton , 15 boltzmannconstant , 15 byte _B, 19 cable _cbl, 20 celsius _degC, 16 chain _ch, 20 coulomb _C, 16 cup _cup, 20 curie _Ci, 19	hand _hh, 20 hectare _hectare, 17 henry _H, 16 hertz _Hz, 16 horsepower _hp, 21 hour _h, 17 hundredweight _cwt, 21 inch _in, 20 intbritishthermalunit _BTU_it, 21 internationalcalorie _cal_IT, 18 joule _J, 16
day _d, 17 decibel _dB, 19 degree _deg, 17 dram _dr, 21	katal _kat, 16 kilopond _kp, 18 knot _kn, 20
electronmagneticmoment , 15 electronmass , 15 electronvolt _eV, 19 elementarycharge , 15 eulersnumber , 15	league _lea, 20 lightsecond _ls, 18 lightyear _ly, 18 liter _L, 17 longton _ton, 21 lumen _lm, 16 lux lx, 16
fahrenheit _degF, 21 farad _F, 16 fathom _ftm, 20 fermi _fermi, 18 finestructureconstant , 15 firkin _fir, 21	metrichorsepower _PS, 19 metrictablespoon _Tbsp, 18 metricteaspoon _tsp, 18 mile _mi, 20 millimeterofmercury _mmHg, 18

minute _min, 17	poundforcepersquareinch _psi, 21
molargasconstant, 15	protonmagnetic moment, 15
,	protonmass, 15
nauticalleague _nlea, 20	
nauticalmile _nmi, 20	quart _qt, 20
neutronmagnetic moment , 15	quarter _qtr, 21
neutronmass, 15	1 7 1 10
newton _N, 16	rad _Rad, 19
nibble _nibble, 19	radian _rad, 16
nomjoveqradius _Re_J_nom, 17	reducedplanckconstant, 15
$nomjov mass parameter \verb _GM_J_nom ,$	rem _rem, 19
17	rod_rd, 20
nomjovpolradius _Rp_J_nom, 17	rydbergconstant, 15
nomsolefftemperature _T_S_nom, 17	sennight _sen, 21
nomsolirradiance _S_S_nom, 17	siemens _S, 16
nomsolluminosity _L_S_nom, 17	sievert _Sv, 16
nomsolmassparameter _GM_S_nom,	slug_slug, 21
17	spat _sp, 18
nomsolradius _R_S_nom, 17	speedoflight, 15
nomterreqradius _Re_E_nom, 17	standardatmosphere _atm, 18
nomterrmassparameter _GM_E_nom,	standardgravity, 15
17	stefanboltzmannconstant, 15
nomterrpolradius _Rp_E_nom, 17	steradian _sr, 16
nuclearmagneton, 15	stone _st, 21
	svedberg _svedberg, 18
ohm _Ohm, 16	svedberg _svedberg, 10
ounce _oz, 21	technicalatmosphere _at, 18
	tesla _T, 16
parsec _pc, 18	thchembritishthermalunit _BTU, 21
partsperbillion _ppb, 19	thermochemicalcalorie _cal, 18
partspermillion _ppm, 19	thou _th, 20
partsperquadrillion _ppq, 19	tonne_t, 17
partspertrillion _ppt, 19	tonoftnt _t_TNT, 18
pascal _Pa, 16	torr_Torr, 18
pennyweight _dwt, 21	troyounce _oz_t, 21
percent _percent, 19	troypound _lb_t, 21
permille _permille, 19	turn _tr, 18
pi , 15	
pica _pica, 20	uscup _cup_US, 22
pint _pint, 20	usfluiddram _fl_dr_US, 22
planckconstant, 15	usfluidounce _fl_oz_US, 22
point _pt, 20	usgallon _gal_US, 22
poiseuille P1, 19	usgill _gi_US, 22
pound _1b, 21	ushundredweight _cwt_US, 22
poundal _pdl, 21	uspint _pint_US, 22
poundforce _lbf, 21	usquart _qt_US, 22
·	

usquarter _qtr_US, 22 ussurveyacre _ac_US, 22 ussurveycable _cbl_US, 22 ussurveychain _ch_US, 22 ussurveyfathom _ftm_US, 22 ussurveyfurlong _fur_US, 22 ussurveyfurlong _fur_US, 22 ussurveyhand _hh_US, 22 ussurveyleague _lea_US, 22 ussurveylink _li_US, 22 ussurveymile _mi_US, 22 ussurveymile _mi_US, 22 ussurveyrod _rd_US, 22 ussurveyyard _yd_US, 22 ustablespoon _Tbsp_US, 22

 $\begin{array}{c} {\tt usteaspoon\ _tsp_US,\ 22} \\ {\tt uston\ _ton_US,\ 22} \end{array}$

 $\begin{array}{c} {\rm vacuumpermeability}\ ,\ 15\\ {\rm vacuumpermitivity}\ ,\ 15\\ {\rm volt}\ _{\tt V},\ 16\\ {\rm voltampere}\ _{\tt VA},\ 19 \\ \end{array}$

watt _W, 16 watthour _Wh, 19 wattsecond _Ws, 19 weber _Wb, 16 week _wk, 18

 $\begin{array}{c} \text{yard } _\text{yd},\, 20 \\ \text{year } _\text{a},\, 18 \end{array}$

Index of Currencies

Afghan Afghani _AFN, 23	BruneiDollar _BND, 31
AfghanPul_cAFN, 23	BruneiSen _cBND, 31
AlbanianLek _ALL, 23	BulgarianLev _BGN, 31
AlgerianDinar _DZD, 24	BulgarianStotinka _cBGN, 31
AlgerianSanteem _cDZD, 24	BurmeseKyat _MMK, 27
AngolanCentimo _cAOA, 23	BurmesePya _cMMK, 27
AngolanKwanza AOA, 23	BurundianCentime _cBIF, 23
ArgentineCentavo _cARS, 23	BurundianFranc_BIF, 23
ArgentinePeso _ARS, 23	- /
ArmenianDram _AMD, 23	CambodianRiel _KHR, 26
ArmenianLuma _cAMD, 23	CanadianCent _cCAD, 24
ArubanCent _cAWG, 31	CanadianDollar _CAD, 24
ArubanFlorin _AWG, 31	CapeVerdeanCentavo _cCVE, 32
AustralianCent _cAUD, 23	CapeVerdeanEscudo _CVE, 32
AustralianDollar _AUD, 23	CaymanIslandsCent _cKYD, 32
AzerbaijaniManat _AZN, 23	CaymanIslandsDollar _KYD, 32
AzerbaijaniQepik _cAZN, 23	CentralAfricanCFACentime _cXAF
BahamianCent _cBSD, 31	CentralAfricanCFAFranc _XAF, 33
BahamianDollar _BSD, 31	CFPCentime _cXPF, 34
BahrainiDinar _BHD, 31	CFPFranc _XPF, 34
BahrainiFils _cBHD, 31	ChileanCentavo _cCLP, 24
BangladeshiPoisha _cBDT, 23	ChileanPeso _CLP, 24
BangladeshiTaka _BDT, 23	ChineseRenminbiFen _cCNY, 24
BarbadianCent _cBBD, 31	ChineseRenminbiYuan _CNY, 24
BarbadianDollar _BBD, 31	ColombianCentavo _ccop, 24
BelarusianKapiejka _cBYN, 23	ColombianPeso _COP, 24
BelarusianRuble _BYN, 23	ComorianCentime _cKMF, 32
BelizeCent _cBZD, 31	Comorianfranc _KMF, 32
BelizeDollar _BZD, 31	CongoleseCentime _cCDF, 24
BermudianCent _cBMD, 31	CongoleseFranc _CDF, 24
BermudianDollar _BMD, 31	CostaRicanCentimos _cCRC, 24
BhutaneseChhertum _cBTN, 31	CostaRicanColon _CRC, 24
BhutaneseNgultrum _BTN, 31	CroatianKuna _HRK, 25
BolivianBoliviano _BOB, 23	CroatianLipa _cHRK, 25
BolivianCentavo _cBOB, 23	CubanCentavo _cCUP, 31
BosnianFenings _cBAM, 23	CubanoConvertibleCentavo _cCUC,
BosnianMark _BAM, 23	31
BotswanaPula _BWP, 23	CubanoConvertiblePeso _CUC, 31
BotswanaThebe _cBWP, 23	CubanPeso _CUP, 31
BrazilianCentavo _cBRL, 23	CzechHaler _cCZK, 24
BrazilianReal _BRL, 23	CzechKoruna _CZK, 24

DanishKrone _DKK, 24 DanishOre _cDKK, 24 DjiboutianCentime _cDJF, 32 DjiboutianFranc _DJF, 32 DominicanCentavo _cDOP, 24 DominicanPeso _DOP, 24

EasternCaribbeanCent _cXCD, 33 EasternCaribbeanDollar _XCD, 33 EgyptianPiastre _cEGP, 24 EgyptianPound _EGP, 24 EritreanCent _cERN, 32 EritreanNakfa _ERN, 32 EthiopianBirr _ETB, 24 EthiopianSantim _cETB, 24

FalklandIslandsPenny _cFKP, 32 FalklandIslandsPound _FKP, 32 FijianCent _cFJD, 24 FijianDollar _FJD, 24

GambianButut cGMD, 25 Gambian Dalasi GMD, 25 GeorgianLari_GEL, 25 GeorgianTetri_cGEL, 25 GhanaianCedi _GHS, 25 GhanaianPesewa _cGHS, 25 GibraltarPenny _cGIP, 32 GibraltarPound _GIP, 32 GuatemalanCentavo _cGTQ, 25 Guatemalan Quetzal _GTQ, 25 GuernseyPenny _cGGP, 32 GuernseyPound _GGP, 32 GuineanCentime _cGNF, 25 GuineanFranc _GNF, 25 GuyaneseCent _cGYD, 25 GuyaneseDollar GYD, 25

HaitianCentime _cHTG, 25 HaitianGourde _HTG, 25 HonduranCentavo _cHNL, 25 HonduranLempira _HNL, 25 HongKongCent _cHKD, 25 HongKongDollar _HKD, 25 HungarianFiller _cHUF, 25 HungarianForint _HUF, 25 IcelandicKrona _ISK, 26 IndianPaisa _cINR, 26 IndianRupee _INR, 26 IndonesianRupiah _IDR, 25 IndonesianSen _cIDR, 25 IranianRial _IRR, 26 IranianToman _cIRR, 26 IraqiDinar _IQD, 26 IraqiFils _cIQD, 26 IsraeliNewAgora _cILS, 25 IsraeliNewShekel _ILS, 25

JamaicanCent _cJMD, 26 JamaicanDollar _JMD, 26 JapaneseYen _JPY, 26 JerseyPenny _cJEP, 32 JerseyPound _JEP, 32 JordanianDinar _JOD, 32 JordanianFils _cJOD, 32

KazakhstaniTenge _KZT, 26 KazakhstaniTiyn _cKZT, 26 KenyanCent _cKES, 26 KenyanShilling _KES, 26 KiribatiCent _cKID, 32 KiribatiDollar _KID, 32 KuwaitiDinar _KWD, 26 KuwaitiFils _cKWD, 26 KyrgyzstaniSom _KGS, 26 KyrgyzstaniTyiyn _cKGS, 26

LaoAtt _cLAK, 26 LaoKip _LAK, 26 LebanesePound _LBP, 32 LebaneseQeresh _cLBP, 32 LiberianCent _cLRD, 26 LiberianDollar _LRD, 26 LibyanDinar _LYD, 27 LibyanDirham _cLYD, 27

MacaneseAvo _cMOP, 32 MacanesePataca _MOP, 32 MacedonianDenar _MKD, 27 MacedonianDeni _cMKD, 27 MalagasyAriary _MGA, 27 MalagasyIraimbilanja _cMGA, 27 MalawianKwacha MWK, 27 PakistaniRupee _PKR, 28 MalawianTambala _cMWK, 27 PanamanianBalboa _PAB, 33 MalaysianRinggit MYR, 27 Panamanian Centesimo cPAB, 33 MalaysianSen_cMYR, 27 PapuaNewGuineanKina PGK, 28 MaldivianLaari _cMVR, 27 PapuaNewGuineanToea cPGK, 28 MaldivianRufiyaa _MVR, 27 ParaguayanCentimo cPYG, 28 ManxPenny_cIMP, 32 ParaguayanGuarani PYG, 28 ManxPound _IMP, 32 PennySterling cGBP, 25 MauritanianKhoums cMRU, 27 PeruvianCentimo cPEN, 28 MauritanianOuguiya MRU, 27 PeruvianSol PEN, 28 MauritianCent cMUR, 27 PhilippinePeso_PHP, 28 MauritianRupee _MUR, 27 PhilippineSentimo cPHP, 28 MexicanCentavo cMXN, 27 PolishGrosz _cPLN, 28 MexicanPeso MXN, 27 PolishZloty PLN, 28 MoldovanBan cMDL, 27 PoundSterling GBP, 25 MoldovanLeu MDL, 27 QatariDirham _cQAR, 28 MongolianMongo _cMNT, 27 MongolianTogrog _MNT, 27 QatariRiyal QAR, 28 MoroccanDirham _MAD, 27 RomanianBan _cRON, 28 MoroccanSantim _cMAD, 27 RomanianLeu_RON, 28 MozambicanCentavo _cMZN, 27 RussianKopeyka cRUB, 28 MozambicanMetical MZN, 27 RussianRuble_RUB, 28 NamibianCent cNAD, 33 RwandanCentime cRWF, 29 NamibianDollar NAD, 33 RwandanFranc_RWF, 29 NepalesePaisa cNPR, 33 NepaleseRupee NPR, 33 SaintHelenaPenny _cSHP, 33 NetherlandsAntilleanCent cANG, 31 SaintHelenaPound _SHP, 33 NetherlandsAntilleanGuilder _ANG, SamoanSene _cWST, 30 31 SamoanTala_WST, 30 NewTaiwanCent _cTWD, 30 ${\bf Sao Tome And Principe Centimo}$ NewTaiwanDollar _TWD, 30 _cSTN, 33 NewZealandCent_cNZD, 28 SaoTomeAndPrincipeDobra _STN, NewZealandDollar _NZD, 28 Nicaraguan Centavo _cNIO, 28 SaudiHalalah _cSAR, 33 NicaraguanCordoba NIO, 28 SaudiRiyal _SAR, 33 NigerianKobo _cNGN, 28 SerbianDinar _RSD, 28 NigerianNaira _NGN, 28 SerbianPara cRSD, 28 NorthKoreanChon _cKPW, 26 SeychelloisCent cSCR, 29 NorthKoreanWon KPW, 26 SeychelloisRupee _SCR, 29 NorwegianKrone _NOK, 28 SierraLeoneanCent cSLL, 29 NorwegianOre cNOK, 28 SierraLeoneanLeone_SLL, 29 SingaporeCent cSGD, 29 OmaniBaisa _cOMR, 33 SingaporeDollar SGD, 29 OmaniRial _OMR, 33 SolomonIslandsCent cSBD, 29

SolomonIslandsDollar SBD, 29

PakistaniPaisa _cPKR, 28

SomalilandCent _cSQS, 29 TurkishKurus _cTRY, 30 SomalilandShilling _SQS, 29 TurkishLira_TRY, 30 SomaliSenti cSOS, 29 TurkmenistanManat TMT, 33 SomaliShilling SOS, 29 TurkmenistanTenge _cTMT, 33 SouthAfricanCent _cZAR, 30 TuvaluanCent _cTVD, 33 TuvaluanDollar _TVD, 33 SouthAfricanRand _ZAR, 30 SouthKoreanJeon_cKRW, 26 UgandanCent _cUGX, 30 SouthKoreanWon KRW, 26 UgandanShilling _UGX, 30 SouthSudanesePiaster _cSSP, 33 UkrainianHryvnia _UAH, 30 SouthSudanesePound SSP, 33 UkrainianKopiyka _cUAH, 30 SriLankanCent _cLKR, 26 UnitedArabEmiratesDirham _AED, SriLankanRupee _LKR, 26 SudanesePound_SDG, 29 UnitedArabEmiratesFils _cAED, 31 SudaneseQirsh cSDG, 29 UruguayanCentesimo _cUYU, 30 SurinameseCent _cSRD, 29 UruguayanPeso _UYU, 30 SurinameseDollar _SRD, 29 USCent_cUSD, 30 SwaziCent _cSZL, 33 USDollar _USD, 30 SwaziLilangeni _SZL, 33 UzbekistaniSom _UZS, 30 SwedishKrona _SEK, 29 UzbekistaniTiyin _cUZS, 30 SwedishOre _cSEK, 29 SwissFranc _CHF, 24 VenezuelanBolivarSoberano _VES, SwissRappen _cCHF, 24 SyrianPiastre _cSYP, 29 VenezuelanCentimoSoberano cVES, SyrianPound _SYP, 29 VietnameseDong_VND, 30 TajikistaniDiram _cTJS, 29 VietnameseXu _cVND, 30 TajikistaniSamani TJS, 29 TanzanianSenti _cTZS, 30 WestAfricanCFACentime _cXOF, 34 TanzanianShilling _TZS, 30 WestAfricanCFAFranc _XOF, 34 ThaiBaht THB, 29 ThaiSatang _cTHB, 29 YemeniDinar _cYER, 30 Tonganpaanga TOP, 29 YemeniRial _YER, 30 TonganSeniti _cTOP, 29 TransnistrianKopeck cPRB, 33 ZambianKwacha ZMW, 30 TransnistrianRuble _PRB, 33 ZambianNgwee _cZMW, 30

ZimbabweanBonds ZWL, 34

ZimbabweanCent _cZWL, 34

TrinidadAndTobagoCent cTTD, 30

TrinidadAndTobagoDollar TTD, 30