

geet 1a Mn
+ 1) S Norval: TNo S
1 frat | Te Spec: Heading [DATE])))) [NAME] |S
in i <NRP> le [NAME] || Subtle fm...) [NAME] E
wi nil Y f

Heading 1!

i! Styles
) 2 t [NAME]
i i 5 ACI ice

[HOSPITAL][DATE]) 41-32.00.

TECHNIQUE:

+ Asal dittusion study.

© Sagittal T2W's.

= Axial, [DATE], [DATE] and [HOSPITAL],
* Coronal TAM.

FINDINGS:

¢ Noareas of diffusion restriction (no MRI evidence of acute stroke).

[NAME] thin sheet of high T2 and [HOSPITAL] signal are seen

© Bilateral frontal and perietal small periventricular and subcortical foci
of high T2 and [HOSPITAL] signal are seen neither surrounded by edema
nor exerting mass effect. Some of the foci show CSF like signal.

Prominent cerebral ventricles 4nd mildly dilated cortical sulci and
extra axial [HOSPITAL] spaces.

Normal size (and) parenchymal signal, intensity pattern of the
cerebellum, and brain stem.

© No shift of the midline structures.

[NAME] of extra axial \$0.15.

e Noextra-axial collections,

Intact cranio-cervical junction:

aight manillary sinusitis and left mastoid.

e Small artery disease with bilateral cerebral ischemic foci and old
[LOCATION] infarcts.

© [HOSPITAL] involutional changes.