



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection?

The most distinctive comorbidities of 32 non-survivors from a group of 52 intensive care unit patients with novel coronavirus disease 2019 (COVID-19) in the study by Xiaobo Yang and colleagues¹ were cerebrovascular diseases (22%) and diabetes (22%). Another study² included 1099 patients with confirmed COVID-19, of whom 173 had severe disease with comorbidities of hypertension (23.7%), diabetes mellitus (16.2%), coronary heart diseases (5.8%), and cerebrovascular disease (2.3%). In a third study,³ of 140 patients who were admitted to hospital with COVID-19, 30% had hypertension and 12% had diabetes. Notably, the most frequent comorbidities reported in these three studies of patients with COVID-19 are often treated with angiotensin-converting enzyme (ACE) inhibitors; however, treatment was not assessed in either study.

Human pathogenic coronaviruses (severe acute respiratory syndrome coronavirus [SARS-CoV] and SARS-CoV-2) bind to their target cells through angiotensin-converting enzyme 2 (ACE2), which is expressed by epithelial cells of the lung, intestine, kidney, and blood vessels.⁴ The expression of ACE2 is substantially increased in patients with type 1 or type 2 diabetes, who are treated with ACE inhibitors and angiotensin II type-I receptor blockers (ARBs).⁴ Hypertension is also treated with ACE inhibitors and ARBs, which results in an upregulation of

ACE2.⁵ ACE2 can also be increased by thiazolidinediones and ibuprofen. These data suggest that ACE2 expression is increased in diabetes and treatment with ACE inhibitors and ARBs increases ACE2 expression. Consequently, the increased expression of ACE2 would facilitate infection with COVID-19. We therefore hypothesise that diabetes and hypertension treatment with ACE2-stimulating drugs increases the risk of developing severe and fatal COVID-19.

If this hypothesis were to be confirmed, it could lead to a conflict regarding treatment because ACE2 reduces inflammation and has been suggested as a potential new therapy for inflammatory lung diseases, cancer, diabetes, and hypertension. A further aspect that should be investigated is the genetic predisposition for an increased risk of SARS-CoV-2 infection, which might be due to ACE2 polymorphisms that have been linked to diabetes mellitus, cerebral stroke, and hypertension, specifically in Asian populations. Summarising this information, the sensitivity of an individual might result from a combination of both therapy and ACE2 polymorphism.

We suggest that patients with cardiac diseases, hypertension, or diabetes, who are treated with ACE2-increasing drugs, are at higher risk for severe COVID-19 infection and, therefore, should be monitored for ACE2-modulating medications, such as ACE inhibitors or ARBs. Based on a PubMed search on Feb 28, 2020, we did not find any evidence to suggest that antihypertensive calcium channel blockers increased ACE2 expression or activity, therefore these could be a suitable alternative treatment in these patients.

We declare no competing interests.


Lei Fang, George Karakioulakis,
*Michael Roth
michael.roth@usb.ch

Pulmonary Cell Research and Pneumology,
Department of Biomedicine and Internal
Medicine, University Hospital Basel,
CH-4031 Basel, Switzerland (LF, MR);
and Department of Pharmacology, School of
Medicine, Aristotle University of Thessaloniki,
Thessaloniki, Greece (GK)

- 1 Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 2020; published online Feb 24. [https://doi.org/10.1016/S2213-2600\(20\)30079-5](https://doi.org/10.1016/S2213-2600(20)30079-5).
- 2 Guan W, Ni Z, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020; published online Feb 28. DOI:10.1056/NEJMoa2002032.
- 3 Zhang JJ, Dong X, Cao YY, et al. Clinical characteristics of 140 patients infected by SARS-CoV-2 in Wuhan, China. *Allergy* 2020; published online Feb 19. DOI:10.1111/all.14238.
- 4 Wan Y, Shang J, Graham R, Baric RS, Li F. Receptor recognition by novel coronavirus from Wuhan: An analysis based on decade-long structural studies of SARS. *J Virology* 2020; published online Jan 29. DOI:10.1128/JVI.00127-20.
- 5 Li XC, Zhang J, Zhuo JL. The vasoprotective axes of the renin-angiotensin system: physiological relevance and therapeutic implications in cardiovascular, hypertensive and kidney diseases. *Pharmacol Res* 2017; **125**: 21–38.



Published Online
March 11, 2020
[https://doi.org/10.1016/S2213-2600\(20\)30116-8](https://doi.org/10.1016/S2213-2600(20)30116-8)



Juan Gantner/Science Photo Library