

CHAROTAR UNIVERSITY OF SCIENCE AND TECHNOLOGY

University Examination

Date: 06/06/18 Day: Wednesday

Time: 10:00 am to 10:30 am

Total Marks: 20

Student's ID No.									
Student's Name: _____									
Student's Sign : _____									
Supervisor's Name : _____									
Supervisor's Sign: _____									

Answer Sheet for Multiple Choice Questions (MCQs), True or false and Match the following	
Examination : MPT- June 2018	
Semester: 4	
Section : I	
Course Code : PT916	
Course Name: Paediatric Neurophysiotherapy	

Important Instructions:

- There are two sections in the paper. **SECTION –I** contains 20 Questions and **SECTION – II** contains 3 Questions.
- **SECTION –I** should be answered in the question paper itself & **SECTION – II** should be answered in separate answer book.
- Marks on the right side suggest the total marks of that question.
- Maximum time allotted for **SECTION – I** is 30 minutes Question paper for **SECTION – II** will be given once the **SECTION – I** is completed within the stipulated time and returned. However, Candidates who complete **SECTION – I** earlier than 30 minutes can collect main question paper for **SECTION –II** immediately after handing over the **SECTION-I** question paper with answers.
- **SECTION – I** should be attached with the main answer book at the end of the examination and should be submitted to the supervisor.

SECTION-I

Multiple Choice Questions: (MCQs)
(Encircle the correct answer)

(10 × 1 = 10)

1. Voluntary release comes at the age of:
 - a. 2 months
 - b. 3 months
 - c. 4 months
 - d. 5 months
2. APGAR score is used to assess all system Except:
 - a. Endocrine System
 - b. Nervous System
 - c. Cardiac system
 - d. Respiratory system

P.T.O.

3. Persistence of this reflex influence prognosis.
 - a. Righting reactions
 - b. Flexion withdrawal
 - c. Parachute reflex
 - d. Abdominal reflex
4. A patient reports to physical therapy with the diagnosis of a lesion in the lateral cord of the brachial plexus. Which of the following would you most likely detect upon the assessment of this patient?
 - a. Paralysis of the biceps, coracobrachialis, finger flexors.
 - b. Paralysis of the deltoid
 - c. Paralysis of wrist extension.
 - d. Paralysis of the hand.
5. _____ is the most common type of Spina Bifida.
 - a. Meningocele
 - b. Spina bifida occulta
 - c. Myelo meningocele
 - d. Lipo meningocele
6. What would you consider as the primary goal of child centered intervention when you are planning the program of a 5 year old child with spastic diplegic Level III CP with a sound cognitive function?
 - a. Prescribe orthosis and mobility aids to optimize function
 - b. Optimize his school participation and peer interaction
 - c. Improve his gait and posture
 - d. Increase his strength and endurance
7. Which of the following reflex may not be significant while examining an infant with Brachial Plexus Injury?
 - a. Moro reflex
 - b. Startle reflex
 - c. Grasp reflex
 - d. ATNR
8. Which of the following statements with regard to Physical therapy intervention of Duchene Muscular Dystrophy is incorrect?
 - a. Helps in maintaining the level of protein dystrophin and reduces CPK
 - b. Preserves or prolong patients' ambulatory status for as long as possible
 - c. Minimizes contractures and deformities
 - d. Improves the quality of life of the child and the parents
9. Which of the following statement is incorrect about neural tube formation
 - a. Neural groove appears as a dorsal focal thickening caused by the proliferation of ectodermal cells.
 - b. About 21st day cell adhesion causes formation of Neural tube
 - c. Superficial ectoderm separates and forms the neural arch and spinal vertebrae
 - d. The cephalic end closes first

10. Urinary symptoms in a lower lumbar spina bifida can be evidence by the following except.....
- Urinary retention and incontinence
 - Neurogenic bladder with decreased tone
 - Lack of awareness of bladder fullness
 - Reflex Bladder

Identify the sentence as True or False:
(Please tick mark the answer)

(5 × 1 = 5)

- | | True | False |
|---|--------------------------|--------------------------|
| 11. High-resistance strength training at submaximal and at near-maximal levels has short term benefits, in slowly progressive myopathic disorders | ☐ | ☐ |
| 12. Aeroplane splint is used to treat klumpke's paralysis | ☐ | ☐ |
| 13. Gower's sign is positive in all types of muscular dystrophy. | ☐ | ☐ |
| 14. Alberta Infant Motor scale is applicable until 18 months of a child | ☐ | ☐ |
| 15. Meningomyelocele can occur due to VP shunt | ☐ | ☐ |

Match the following

(5 × 1 = 5)

16.	Resiliency factor	A. Set of customary ways of reacting
17.	Motivation	B. Positive response from caregivers
18.	Mastery Motivation	C. Key determinant of developmental change
19.	Child behavior characteristics	D. Element of Physical danger
20.	Risk taking behavior	E. Lowers the level of motor skill development
		F. Originates without the need of external reward
		G. Contributes to individual adaptability

(16 -), (17-), (18-), (19-), (20-)

CHAROTAR UNIVERSITY OF SCIENCE & TECHNOLOGY

4th Semester of M.P.T. Examination

June 2018

PT916 Paediatric Neurophysiotherapy

Date: 06/06/18

Time: 10:30 am to 1:00 pm

Maximum Marks: 50

Instructions:

1. There are two Sections in the paper. **SECTION –I** contains 20 Questions and **SECTION – II** contains 3 Questions.
2. **SECTION –I** should be answered in the question paper itself & **SECTION – II** should be answered in separate answer book.
3. Marks on the right side suggest the total marks of that question.
4. Maximum time allotted for **SECTION – II** is 2 Hr 30 min.
5. Draw the figure where necessary.

SECTION-II

Q. 1. Short answers (5 out of 7) (Answer not exceeding 50 words) (5 × 2 = 10)

- a. Scarf sign.
- b. Athetoid Cerebral Palsy
- c. Myelomeningocele
- d. Erb's point
- e. Causes of spina bifida
- f. Classification of Muscular dystrophy
- g. Peri-natal causes for high risk infants.

Q. 2. Short Notes (4 out of 5) (Answer not exceeding 150 words) (4 × 5 = 20)

- a. Orthotic management for complete brachial plexus injury
- b. Approach to sharing an unbiased and complete information with parents about their child with myelomeningocele.
- c. Family is constant while the service systems and the personals in the system may fluctuate. Discuss this statement in relation to a child with CP.
- d. Pathophysiology of Duchene Muscular Dystrophy
- e. Barriers to family centered care.

Q. 3. Essay (2 out of 3) (2 × 10 = 20)

- a. Explain the phases of the multivariate model of the determinants of motor change in CP. Emphasize on the phase one of the model for a 3 year old child who is unable to stand due to spastic diplegia.

P.T.O.

- b. Describe the evidence based decision making and the collaborative model of service delivery for a 5 year old child with muscular dystrophy.
- c. A 10 year old boy has spastic tetraplegic CP, who can independently sit and perform basic bed mobility but unable to transfer independently. His school participation is limited, as he needs special care in transfer and seating arrangements. His cognitive function is sound. Using the ICF framework of clinical decision-making, plan a service intervention to optimize his participation in life and improve his quality of life.