

CHAROTAR UNIVERSITY OF SCIENCE & TECHNOLOGY
5th Semester of B. Pharm. Examination
University Theory Examination March 2018
PH315 Pharmacology-III

Date: 28.03.2018, Wednesday

Time: 10a.m. to 1 p.m.

Maximum Marks: 80

Instructions:

1. There are three sections in this question paper.
2. SECTION – I comprises of Question 1. Total marks for Section 1 are 20. There are 20 sub-questions (MCQ type). Answers to SECTION – I are to be given in Answer Sheet for MCQ type questions provided to you. Maximum time allotted for SECTION – I is 30 minutes. Answers to SECTION – I must be written during the first 30 minutes of the examination.
3. Answers to SECTION – II and SECTION – III are to be provided in separate Main Answer Books provided to you.
4. Figures to right indicate marks.
5. Draw neat labeled sketches wherever necessary.

SECTION - I

- | | | |
|------------|--|-----------|
| Q 1 | Attempt all questions. Each question is of one mark. | 20 |
| 1. | Hypnotic drugs are used to treat:
[A] Psychosis
[B] Sleep disorders
[C] Narcolepsy
[D] Parkinsonian disorders | |
| 2. | Select a hypnotic drug, which is a benzodiazepine derivative:
[A] Zolpidem
[B] Flurazepam
[C] Secobarbital
[D] Phenobarbitone | |
| 3. | Indicate an irreversible inhibitor of GABA aminotransferase (GABA-T):
[A] Diazepam
[B] Phenobarbital
[C] Vigabatrin
[D] Felbamate | |
| 4. | Valproate is very effective against:
[A] Absence seizures
[B] Myoclonic seizures
[C] Generalized tonic-clonic seizures
[D] All of the above | |
| 5. | Which of the following agents is related to tricyclic antidepressants?
[A] Nefazodon
[B] Amitriptyline
[C] Fluoxetine
[D] Isocarboxazid | |
| 6. | Which of the following drugs is used in the treatment of Parkinsonian disorders?
[A] Phenytoin
[B] Selegiline
[C] Haloperidol
[D] Fluoxetine | |
| 7. | Indicate the nootropic agent:
[A] Paracetamol | |

- [B] Digoxin
 - [C] Fluoxetine
 - [D] Piracetam
8. Which of the following benzodiazepines is preferred for elderly patients?
- [A] Clorazepate
 - [B] Clordiazepoxide
 - [C] Triazolam
 - [D] Prazepam
9. Indicate a peripheral dopa decarboxylase inhibitor:
- [A] Tolcapone
 - [B] Clozapine
 - [C] Carbidopa
 - [D] Selegiline
10. The combination of disulfiram and ethanol leads to accumulation of:
- [A] Formaldehyde
 - [B] Acetate
 - [C] Formic acid
 - [D] Acetaldehyde
11. Indicate the atypical antipsychotic drug:
- [A] Haloperidol
 - [B] Clozapine
 - [C] Thioridazine
 - [D] Thiothixene
12. Chemical mediators in the nociceptive pathway are all of the following EXCEPT:
- [A] Enkephalins
 - [B] Kinins
 - [C] Prostaglandins
 - [D] Substance P
13. Barbiturates are being replaced by hypnotic benzodiazepines because of:
- [A] Low therapeutic index
 - [B] Suppression in REM sleep
 - [C] High potential of physical dependence and abuse
 - [D] All of the above
14. Which of the following dopaminergic systems is most closely related to behavior?
- [A] The hypothalamic-pituitary system
 - [B] The extrapyramidal system
 - [C] The mesolimbic and mesofrontal systems
 - [D] The chemoreceptor trigger zone of the medulla
15. Which of the following agents may be used as an antidote for ethylene glycol and methanol poisoning?
- [A] Disulfiram
 - [B] Fomepizol
 - [C] Naltrexone
 - [D] Amphetamine
16. Most of non-narcotic analgetics have:
- [A] Anti-inflammatory effect
 - [B] Analgesic effect
 - [C] Antipyretic effect
 - [D] All of the above
17. Parkinsonian symptoms and tarditive dyskinesia are caused by blockade dopamine in:
- [A] The nigrostriatal system
 - [B] The mesolimbic and mesofrontal systems
 - [C] The chemoreceptor trigger zone of the medulla

- [D] The tuberoinfundibular system
18. Select the antiseizure drug with an analgesic component of effect:
 [A] Carbamazepine
 [B] Ethosuximide
 [C] Phenytoin
 [D] Clonazepam
19. Mu (μ) receptors are associated with:
 [A] Analgesia, euphoria, respiratory depression, physical dependence
 [B] Spinal analgesia, mydriasis, sedation, physical dependence
 [C] Dysphoria, hallucinations, respiratory and vasomotor stimulation
 [D] Analgesia, euphoria, respiratory stimulation, physical dependence
20. Which of the following phenothiazine derivatives may produce cardiac toxicity, including ventricular arrhythmias, cardiac conduction block, and sudden death?
 [A] Thioridazine
 [B] Chlorpromazine
 [C] Perphenazine
 [D] Fluphenazine

SECTION – II

- Q 2** Attempt any **TWO** of the following
- A** Describe mechanism of action and therapeutic uses of Nitrates. **05**
- B** Explain pathophysiology of hypertension. **05**
- C** Explain the role of corticosteroids as immunosuppressant. **05**
- Q 3** Attempt any **TWO** of the following
- A** Give a short note on Anticogaulants. **05**
- B** Define Arrhythmia and explain types of Arrhythmia. **05**
- C** Explain difference between loop and thiazide diuretics. **05**

SECTION – III

- Q 4** Attempt any **FOUR** of the following
- A** Explain in detail about role of dopamine as neurotransmitter in CNS. **05**
- B** Classify antiepileptic agents and give short mechanism of action of each class. **05**
- C** Write a note on CNS stimulant agents. **05**
- D** Explain pathogenesis of parkinsonism. Classify antiparkinsonian agents. **05**
- E** What are sedatives and hypnotics? Explain mechanism of action for benzodiazepines with their advantages over barbiturates. **05**
- F** Classify NSAIDs. Write their mechanism of actions and therapeutic uses. **05**
- Q 5** Attempt any **FOUR** of the following
- A** Write a note on ethanol. **05**
- B** Classify antipsychotic agents. Write advantages of atypical agents over typical agents. **05**
- C** Give stages of general anaesthesia with mechanism of action for each stage. **05**
- D** Explain pharmacology of morphine. **05**
- E** Describe pathogenesis of depression and classify antidepressant agents. **05**
- F** Give mechanism of action and therapeutic uses of lithium and ketamine. **05**