

A Scientometric Analysis of Bias and Fairness in Parkinson's Disease Clinical Assessment Scales

Journal:	<i>Movement Disorders</i>
Manuscript ID	MDS-25-1327
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	18-Sep-2025
Complete List of Authors:	Benjelloun, Mohammed; Universite de Mons Faculte Polytechnique Botta, Ragasudha; Massachusetts General Hospital Department of Neurology Celi, Leo Anthony; Massachusetts Institute of Technology Health Hani, Moad; Universite de Mons Faculte Polytechnique Mahmoudi, Saïd; Universite de Mons Faculte Polytechnique Nageswaran, Neha; Carl von Ossietzky Universitat Oldenburg Ong, Bradley; Cleveland Clinic Neurological Institute Restrepo, David; Massachusetts Institute of Technology Health Vilardo, Marina; Universidade Catolica de Brasilia Curso de Medicina
Keywords:	Parkinson disease , Clinical assessment scales, Bias and fairness, Scientometric analysis, Demographic disparities
English Language Editing:	Yes, The manuscript was edited by a native English speaker

SCHOLARONE™
Manuscripts

Title Page

A Scientometric Analysis of Bias and Fairness in Parkinson's Disease Clinical Assessment Scales

Running title : “Bias in Parkinson Assessment Scales”

Author’s List:

Mohammed Benjelloun, PhD¹ [ORCID : 0000-0002-4020-7327](#)
, Ragasudha Botta, PhD², [ORCID ID: 0000-0002-0339-6659](#)
Leo Anthony Celi, MD³, [ORCID : 0000-0001-6712-6626](#)
Moad Hani, MEng ^{1*}, [ORCID : 0000-0003-2342-495X](#)
Saïd Mahmoudi, PhD¹, [ORCID : 0000-0001-8272-9425](#)
Neha Nageswaran, MEng⁴, [ORCID : 0009-0009-2685-7751](#)
Bradley Ong, MD⁵, [ORCID ID: 0000-0001-6507-1967](#)
David Restrepo, MEng³, [ORCID: 0000-0002-3789-1957](#)
Marina Vilardo, MEng ⁶ [ORCID: 0009-0001-9487-1119](#)

Affiliations:

¹Department of Computer Engineering and Management, Faculty of Engineering, University of Mons (UMONS), 7000 Mons, Belgium
²Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA
³Division of Health Sciences and Technology, Massachusetts Institute of Technology, Boston, MA, USA
⁴Department of Assistance Systems and Medical Technology, Carl von Ossietzky University of Oldenburg, Oldenburg, Germany
⁵Department of Neurology, Neurological Institute, Cleveland Clinic, Cleveland, OH, USA
⁶Catholic University of Brasilia School of Medicine, Brazil

Authors are listed alphabetically. All authors contributed equally to the best of their abilities.

Word Count: 1685 words

Conflict of Interest Statement

All authors declare no conflicts of interest relevant to this work.

Funding Statement

All authors report no financial disclosures relevant to this research for the preceding 12 months. No funding sources influenced the design, conduct, analysis, interpretation, or reporting of this study.

(*) Corresponding author (Moad Hani) moad.hani@umons.ac.be

Abstract

Background: Clinical assessment scales are fundamental tools for evaluating Parkinson disease, yet potential biases in their development and validation may compromise their fairness across diverse populations

Objectives: To conduct a comprehensive scientometric analysis examining demographic representation, geographic distribution, and methodological biases in Parkinson disease assessment scale literature.

Methods: Following PRISMA guidelines, we systematically searched five databases from inception to 2024. Studies validating or developing Unified Parkinson Disease Rating Scale, Movement Disorder Society-Unified Parkinson Disease Rating Scale, Montreal Cognitive Assessment, or Mini-Mental State Examination for Parkinson disease were included. Data extraction focused on 17 predefined bias domains including demographic representation and methodological considerations. **Results:** From 3,836 studies, 109 met inclusion criteria, encompassing 655 authors across 34 countries. High-income countries contributed 99 publications (90.8%) versus 10 publications (9.2%) from low/middle-income countries. Critical gaps included: 8 studies (7.3%) captured race/ethnicity data, 13 studies (11.9%) adjusted cognitive tests for education, and zero studies addressed digital literacy barriers. Female authorship was 36.6% overall.

Conclusions: This analysis reveals persistent geographic, demographic, and methodological biases in Parkinson disease assessment research, potentially compromising fairness across diverse populations. These findings highlight urgent needs for inclusive research practices and culturally sensitive scale adaptations.

Keywords: Parkinson disease; clinical assessment; bias; fairness; scientometric analysis; demographic disparities; health equity

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Parkinson disease affects over 10 million people worldwide, with prevalence projected to reach 25.2 million by 2050. Clinical assessment scales serve as fundamental tools for diagnosis, monitoring disease progression, and evaluating treatment efficacy. The most widely used scales include the Unified Parkinson Disease Rating Scale and its revised Movement Disorder Society version for motor symptoms, and cognitive assessments such as the Montreal Cognitive Assessment and Mini-Mental State Examination.

Recent studies have highlighted concerning disparities in clinical research, with underrepresentation of diverse populations potentially limiting the generalizability and fairness of medical interventions. In neurodegenerative diseases, assessment tools developed primarily in high-income, predominantly white populations may not perform equally across different demographic groups. These biases can manifest as geographic concentration of research in wealthy nations, underrepresentation of women and minorities, inadequate consideration of cultural factors, and methodological gaps in addressing confounding variables.

Despite the critical importance of fair assessment tools, no comprehensive analysis has examined the extent of bias and demographic representation in the foundational literature supporting Parkinson disease clinical assessment scales. This knowledge gap represents a significant barrier to understanding potential biases and developing targeted interventions to address them.

Methods

Study Design

We conducted a comprehensive scientometric analysis following established guidelines for bibliometric research, employing a systematic two-phase approach combining quantitative bibliometric analysis with qualitative assessment of methodological biases.

Search Strategy and Selection Criteria

A systematic literature search was conducted across five major databases (PubMed, Embase, Medline and Central) from inception to July 31, 2024. The search strategy was developed using the following terms: ("Parkinson*" OR "PD") AND ("UPDRS" OR "MDS-UPDRS" OR "Unified Parkinson Disease Rating Scale" OR "MoCA" OR "Montreal Cognitive Assessment" OR "MMSE" OR "Mini Mental State") AND ("valid*" OR "develop*" OR "assess*" OR "scale" OR "instrument").

Inclusion criteria: (1) Studies validating or developing Parkinson disease assessment scales; (2) English articles; (3) Peer-reviewed publications; (4) Human subjects with Parkinson disease.

Exclusion criteria: (1) Case reports and editorials; (2) Non-English publications; (3) Studies not focusing on Parkinson disease assessment scales; (4) Animal studies; (5) Conference abstracts without full text.

Data Extraction and Bias Assessment Framework

Data extraction was performed independently by two reviewers using a standardized form. **Seventeen predefined bias domains** were assessed based on established frameworks for demographic bias in clinical research, categorized into primary demographic domains (race/ethnicity, sex/gender, age-specific values, educational background, geographic representation, socioeconomic status, administrator training, digital literacy barriers) and methodological domains (assessment timing,

treatment controls, medication state documentation, cognitive domain specification, institutional resources, subgroup analysis, inclusion criteria, confounding variables, sample size adequacy).

Statistical Analysis

Descriptive statistics were calculated for all variables. Chi-square tests assessed associations between categorical variables. Geographic disparities were analyzed using Global North-South classification and World Bank income categories. Statistical significance was set at $p < 0.05$. All analyses were performed using Python 3.12.

Data Sharing

The datasets generated and analyzed during this study are available from the corresponding author upon reasonable request. All data extraction forms and analysis scripts used in this study will be made publicly available upon publication.

Results

Study Selection and Characteristics

The systematic search yielded 3,836 potentially relevant studies. After duplicate removal and screening, 141 full-text articles were assessed for eligibility, with 109 studies ultimately meeting inclusion criteria for analysis (**Figure 1**). The included studies encompassed 655 authors across 34 countries and were published in 47 different journals between 1996 and 2024.

Publication Trends and Geographic Distribution

Publication trajectories revealed distinct temporal patterns reflecting clinical adoption and research maturity. The Unified Parkinson Disease

Rating Scale demonstrated consistent usage throughout the analyzed period, with peak activity during 2010-2019 coinciding with the Movement Disorder Society revision. The Montreal Cognitive Assessment showed rapid adoption between 2008-2016, reaching maximum publication volume during 2014-2015, followed by recent decline. The Mini-Mental State Examination maintained steady but moderate levels with mid-period surge during 2014-2020 (**Figure 2**).

Geographic analysis revealed significant disparities

Geographic analysis (**Figure 3**) revealed significant disparities in research distribution. High-income countries contributed 99 publications (90.8%), while low/middle-income countries contributed only 10 publications (9.2%). Europe dominated with 48 publications (44.0%), followed by North America with 39 publications (35.8%).

Asia showed pronounced disparities between high-income (9 publications, 8.3%) and low/middle-income contributions (4 publications, 3.7%). Africa and South America, despite having significant Parkinson disease populations, contributed minimally with 2 (1.8%) and 3 (2.8%) publications respectively.

Authorship Patterns and Gender Distribution

Among 655 total authors, 240 (36.6%) were female and 415 (63.4%) were male. Gender representation varied by authorship position: first authorship showed 25 female authors (32.9%) compared to 51 male authors (67.1%), while senior authorship demonstrated 26 female authors (37.1%) versus 44 male authors (62.9%).

Bias Domain Assessment

Comprehensive analysis of 109 studies revealed significant variations in bias domain consideration (**Table 1**).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Cognitive impairment criteria specification was most frequently addressed (49 studies, 45.0%), followed by sex/gender analysis (47 studies, 43.1%) and cognitive domain specification (43 studies, 39.4%).

Critical gaps were identified in several domains (**Figure 4**). Digital literacy and access barriers were not addressed in any study (0.0%). Treatment effects on cognition were considered in only 1 study (0.9%), and age-specific normative values in merely 3 studies (2.8%). Race and ethnicity data collection was documented in only 8 studies (7.3%), while medication state testing occurred in 4 studies (3.7%) for OFF state and 18 studies (16.5%) for ON state.

Education adjustment for cognitive tests was performed in only 13 studies (11.9%), institutional resources reporting in 11 studies (10.1%), and confounding variables control in 16 studies (14.7%). The mean percentage of studies addressing any given bias domain was 17.8%, with a median of 14.7%.

Journal and Citation Analysis

Studies were published across 47 journals. Specialized movement disorders journals accounted for 45 publications (41.3%): Movement Disorders (n=29), Parkinsonism & Related Disorders (n=11), and Movement Disorders Clinical Practice (n=5). Impact factor quartile analysis revealed 24 studies (19.8%) in Q1 journals, 35 studies (32.1%) in Q2 journals, 38 studies (34.9%) in Q3 journals, and 12 studies (11.0%) in Q4 journals.

Discussion

Principal Findings

This scientometric analysis provides compelling evidence of significant and persistent biases in PD clinical assessment scale research. The

predominance of research from high-income countries, male authorship patterns, and critical gaps in methodological consideration of bias factors collectively threaten the global applicability and fairness of current assessment instruments.

Geographic and Economic Disparities

The stark geographic concentration of research in high-income countries (90.8% of publications) represents a fundamental threat to the global validity of Parkinson disease assessment scales.

This disparity is particularly concerning given epidemiological projections showing fastest growth in Parkinson disease prevalence in low- and middle-income countries. The near-absence of research from Africa (1.8% of publications) and limited representation from South America (2.8%) creates significant knowledge gaps about scale performance in these populations.

These geographic biases have direct clinical implications. Assessment scales developed and validated primarily in Western, educated, industrialized, rich, and democratic populations may not adequately capture the disease experience in other cultural contexts. Factors such as cultural attitudes toward neurological symptoms, linguistic nuances in symptom description, and healthcare system differences can all influence scale performance and interpretation.

Demographic Representation Gaps

The underrepresentation of critical demographic factors represents a significant methodological weakness. Only 7.3% of studies captured race/ethnicity data, despite well-documented disparities in Parkinson disease presentation and progression across racial groups. This gap is particularly problematic given evidence that cognitive assessment tools demonstrate differential performance across racial and ethnic groups.

The finding that only 11.9% of studies adjusted cognitive assessments for educational background is especially concerning. Educational attainment significantly influences performance on cognitive tests, and failure to account for this factor can lead to systematic biases in diagnosis and severity assessment, potentially contributing to disparities in Parkinson disease dementia diagnosis and treatment access.

Methodological Bias Implications

The identified methodological gaps have direct implications for clinical practice and research validity. The absence of studies addressing digital literacy barriers (0.0% of publications) is particularly concerning in an era of increasing digital health implementation. As Parkinson disease assessment increasingly incorporates technology-based tools, failure to consider digital access and literacy creates new forms of bias that may exacerbate existing health disparities.

Inadequate reporting of administrator training (present in only 30.3% of studies) raises questions about inter-rater reliability and consistency across different clinical settings. This gap is particularly problematic for global implementation of assessment scales, where training resources and expertise may vary significantly.

Implications and Future Directions

These findings have immediate implications for clinical practice and health policy. Clinicians must recognize that current Parkinson disease assessment scales may not perform equally across all patient populations. This recognition should inform clinical decision-making, particularly when evaluating patients from underrepresented demographic groups.

Our findings point to several critical research priorities: (1) Inclusive validation studies specifically designed to assess scale performance across diverse demographic groups; (2) Systematic investigation of

1
2
3 cultural and linguistic factors affecting scale performance for global
4 implementation; (3) Research addressing digital literacy and access
5 barriers as assessment tools increasingly incorporate digital components;
6
7 (4) Development and testing of specific interventions to reduce bias in
8 clinical assessment.
9
10
11
12

13 Limitations

14
15 This study has limitations including restriction to English-language
16 publications, potentially underrepresenting research from non-English
17 speaking countries. The retrospective nature limits assessment of
18 causality between identified biases and clinical outcomes. Publication bias
19 may have influenced findings, as studies with negative results regarding
20 bias may be less likely published. Additionally, assessment of bias
21 domains relied on information reported in published manuscripts, which
22 may not fully capture all considerations addressed during study conduct.
23
24
25
26
27
28
29
30
31

32 In conclusion, this comprehensive scientometric analysis reveals
33 pervasive and systematic biases in the foundational literature supporting
34 Parkinson disease clinical assessment scales. The stark geographic
35 concentration of research in high-income countries, persistent gender
36 disparities in research leadership, and critical gaps in demographic and
37 methodological bias consideration collectively threaten the fairness and
38 global applicability of current assessment instruments. These findings
39 have immediate implications for clinical practice, requiring heightened
40 awareness of potential biases when using Parkinson disease assessment
41 scales across diverse populations. The research community must
42 prioritize inclusive research practices, cross-cultural validation studies,
43 and development of bias-aware assessment tools. Only through
44 systematic efforts can we ensure that fundamental tools used to evaluate
45 Parkinson disease provide fair and accurate assessments across all
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

affected populations, regardless of geography, demographics, or socioeconomic status.

Acknowledgments

The authors thank the research community for their contributions to Parkinson disease assessment research. AI-assisted tools were used to improve clarity and coherence of certain manuscript sections, with careful attention to preserving research content integrity and data interpretation.

Authors' Roles

Research project: A. Conception: **L.A.C, M.H**; B. Organization: **L.A.C, M.H, B. O, R. B**; C. Execution: **All authors**

Statistical Analysis: A. Design: **M.H, B. O, R. B**; B. Execution: **M.H, B. O, R. B, N. N, M. V, D. R**; C. Review and Critique: **All authors**

Manuscript Preparation: A. Writing of the first draft: **M.H, B. O, R. B**; B. Review and Critique: **All authors**

Financial Disclosures

All authors report no financial disclosures relevant to this research for the preceding 12 months. No funding sources influenced the design, conduct, analysis, interpretation, or reporting of this study.

References

1. Antonini, A.; Abbruzzese, G.; Ferini-Strambi, L.; et al. Validation of the Italian version of the Movement Disorder Society--Unified Parkinson's Disease Rating Scale. *Neurological Sciences* 2013, 34(5), 683–687. <https://doi.org/10.1007/s10072-012-1112-z>
2. Abdolahi, A.; Bull, M. T.; Darwin, K. C.; et al. A feasibility study of conducting the Montreal Cognitive Assessment remotely in individuals with movement disorders. *Health Informatics Journal* 2016, 22(2), 304–311. <https://doi.org/10.1177/1460458214556373>

3. Uc, E. Y.; McDermott, M. P.; Marder, K. S.; et al. Incidence of and risk factors for cognitive impairment in an early Parkinson disease clinical trial cohort. *Neurology* 2009, 73(18), 1469–1477. <https://doi.org/10.1212/WNL.0b013e3181bf992f>
4. Foley, T.; McKinlay, A.; Warren, N.; et al. Assessing the sensitivity and specificity of cognitive screening measures for people with Parkinson's disease. *NeuroRehabilitation* 2018, 43(4), 491–500. <https://doi.org/10.3233/NRE-182433>
5. Konstantopoulos, K.; Vogazianos, P.; Doskas, T. Normative data of the Montreal Cognitive Assessment in the Greek population and parkinsonian dementia. *Archives of Clinical Neuropsychology* 2016, 31(3), 246–253. <https://doi.org/10.1093/arclin/acw002>
6. Palmer, J. L.; Coats, M. A.; Roe, C. M.; et al. Unified Parkinson's Disease Rating Scale-Motor Exam: inter-rater reliability of advanced practice nurse and neurologist assessments. *Journal of Advanced Nursing* 2010, 66(6), 1382–1387. <https://doi.org/10.1111/j.1365-2648.2010.05313.x>
7. Badrkhahan, S. Z.; Sikaroodi, H.; Sharifi, F.; et al. Validity and reliability of the Persian version of the Montreal Cognitive Assessment (MoCA-P) scale among subjects with Parkinson's disease. *Applied Neuropsychology: Adult* 2020, 27(5), 431–439. <https://doi.org/10.1080/23279095.2019.1565762>
8. Hoops, S.; Nazem, S.; Siderowf, A. D.; et al. Validity of the MoCA and MMSE in the detection of MCI and dementia in Parkinson disease. *Neurology* 2009, 73(21), 1738–1745. <https://doi.org/10.1212/WNL.0b013e3181c34b47>
9. Allison, K. M.; Hustad, K. C. Impact of sentence length and phonetic complexity on intelligibility of 5-year-old children with cerebral palsy. *International Journal of Speech-Language Pathology* 2014, 16(4), 396–407. <https://doi.org/10.3109/17549507.2013.876667>
10. Goetz, C. G.; Choi, D.; Guo, Y.; et al. It is as it was: MDS-UPDRS part III scores cannot be combined with other parts to give a valid sum. *Movement Disorders* 2023, 38(2), 342–347. <https://doi.org/10.1002/mds.29279>
11. Martinez-Martin, P.; Rodriguez-Blazquez, C.; Alvarez-Sanchez, M.; et al. Expanded and independent validation of the Movement Disorder Society-Unified Parkinson's Disease Rating Scale (MDS-UPDRS). *Journal of Neurology* 2013, 260(1), 228–236. <https://doi.org/10.1007/s00415-012-6624-1>
12. Williams, S.; Wong, D.; Alty, J. E.; et al. Parkinsonian hand or clinician's eye? Finger tap bradykinesia interrater reliability for 21 movement disorder experts. *Journal of Parkinson's Disease* 2023, 13(4), 525–536. <https://doi.org/10.3233/JPD-223256>
13. Goldman, J. G.; Stebbins, G. T.; Leung, V.; et al. Relationships among cognitive impairment, sleep, and fatigue in Parkinson's disease using the MDS-UPDRS. *Parkinsonism & Related Disorders* 2014, 20(11), 1135–1139. <https://doi.org/10.1016/j.parkreldis.2014.08.001>

14. Rabey, J. M.; Klein, C.; Molochnikov, A.; et al. Comparison of the Unified Parkinson's Disease Rating Scale and the Short Parkinson's Evaluation Scale in patients with Parkinson's disease after levodopa loading. *Clinical Neuropharmacology* 2002, 25(2), 83–88. <https://doi.org/10.1097/00002826-200203000-00005>
15. Stebbins, G. T.; Goetz, C. G. Factor structure of the Unified Parkinson's Disease Rating Scale: Motor Examination section. *Movement Disorders* 1998, 13(4), 633–636. <https://doi.org/10.1002/mds.870130404>
16. Dalrymple-Alford, J. C.; MacAskill, M. R.; Nakas, C. T.; et al. The MoCA: well-suited screen for cognitive impairment in Parkinson disease. *Neurology* 2010, 75(19), 1717–1725. <https://doi.org/10.1212/WNL.0b013e3181fc29c9>
17. Park, J.; Oh, E.; Koh, S.-B.; et al. Evaluating the validity and reliability of the Korean version of the Scales for Outcomes in Parkinson's Disease-cognition. *Journal of Movement Disorders* 2024, 17(3), 328–332. <https://doi.org/10.14802/jmd.24061>
18. Roalf, D. R.; Moore, T. M.; Wolk, D. A.; et al. Defining and validating a short form Montreal Cognitive Assessment (s-MoCA) for use in neurodegenerative disease. *Journal of Neurology, Neurosurgery & Psychiatry* 2016, 87(12), 1303–1310. <https://doi.org/10.1136/jnnp-2015-312723>
19. Parashos, S. A.; Elm, J.; Boyd, J. T.; et al. Validation of an ambulatory capacity measure in Parkinson disease: a construct derived from the Unified Parkinson's Disease Rating Scale. *Journal of Parkinson's Disease* 2015, 5(1), 67–73. <https://doi.org/10.3233/JPD-140405>
20. Pal, G.; Goetz, C. G. Assessing bradykinesia in parkinsonian disorders. *Frontiers in Neurology* 2013, 4, 54. <https://doi.org/10.3389/fneur.2013.00054>
21. Siuda, J.; Boczarska-Jedynak, M.; Budrewicz, S.; et al. Validation of the Polish version of the Movement Disorder Society-Unified Parkinson's Disease Rating Scale (MDS-UPDRS). *Neurologia i Neurochirurgia Polska* 2020, 54(5), 416–425. <https://doi.org/10.5603/PJNNS.a2020.0049>
22. Smith, C. R.; Cavanagh, J.; Sheridan, M.; et al. Factor structure of the Montreal Cognitive Assessment in Parkinson disease. *International Journal of Geriatric Psychiatry* 2020, 35(2), 188–194. <https://doi.org/10.1002/gps.5234>
23. Aarsland, D.; Brønnick, K.; Larsen, J. P.; et al. Cognitive impairment in incident, untreated Parkinson disease: the Norwegian ParkWest study. *Neurology* 2009, 72(13), 1121–1126. <https://doi.org/10.1212/01.wnl.0000338632.00552.cb>
24. Holroyd, S.; Currie, L. J.; Wooten, G. F. Validity, sensitivity and specificity of the mentation, behavior and mood subscale of the UPDRS. *Neurological Research* 2008, 30(5), 493–496. <https://doi.org/10.1179/016164107X251772>

25. Isella, V.; Mapelli, C.; Morielli, N.; et al. Validity and metric of MiniMental Parkinson and MiniMental State Examination in Parkinson's disease. *Neurological Sciences* 2013, 34(10), 1751–1758. <https://doi.org/10.1007/s10072-013-1328-6>
26. Martinez-Martin, P.; Prieto, L.; Forjaz, M. J. Longitudinal metric properties of disability rating scales for Parkinson's disease. *Value in Health* 2006, 9(6), 386–393. <https://doi.org/10.1111/j.1524-4733.2006.00131.x>
27. Horváth, K.; Aschermann, Z.; Ács, P.; et al. Validation of the Hungarian Unified Dyskinesia Rating Scale. *Ideggyógyászati Szemle* 2015, 68(5-6), 183–188. <https://doi.org/10.18071/isz.68.0183>
28. Gülke, E.; Alsalem, M.; Kirsten, M.; et al. Comparison of Montreal cognitive assessment and Mattis dementia rating scale in the preoperative evaluation of subthalamic stimulation in Parkinson's disease. *PLoS One* 2022, 17(4), e0265314. <https://doi.org/10.1371/journal.pone.0265314>
29. Grill, S.; Weuve, J.; Weisskopf, M. G. Predicting outcomes in Parkinson's disease: comparison of simple motor performance measures and The Unified Parkinson's Disease Rating Scale-III. *Journal of Parkinson's Disease* 2011, 1(3), 287–298. <https://doi.org/10.3233/JPD-2011-11016>
30. Patrick, S. K.; Denington, A. A.; Gauthier, M. J.; et al. Quantification of the UPDRS rigidity scale. *IEEE Transactions on Neural Systems and Rehabilitation Engineering* 2001, 9(1), 31–41. <https://doi.org/10.1109/7333.918274>
31. Martignoni, E.; Franchignoni, F.; Pasetti, C.; et al. Psychometric properties of the Unified Parkinson's Disease Rating Scale and of the Short Parkinson's Evaluation Scale. *Neurological Sciences* 2003, 24(3), 190–191. <https://doi.org/10.1007/s10072-003-0124-0>
32. Ohta, K.; Takahashi, K.; Gotoh, J.; et al. Screening for impaired cognitive domains in a large Parkinson's disease population and its application to the diagnostic procedure for Parkinson's disease dementia. *Dementia and Geriatric Cognitive Disorders Extra* 2014, 4(2), 147–159. <https://doi.org/10.1159/000362124>
33. Hely, M. A.; Reid, W. G. J.; Adena, M. A.; et al. The Sydney multicenter study of Parkinson's disease: the inevitability of dementia at 20 years. *Movement Disorders* 2008, 23(6), 837–844. <https://doi.org/10.1002/mds.21956>
34. Qiao, J.; Wang, X.; Lu, W.; et al. Validation of neuropsychological tests to screen for dementia in Chinese patients with Parkinson's disease. *American Journal of Alzheimer's Disease and Other Dementias* 2016, 31(4), 368–374. <https://doi.org/10.1177/1533317515619478>
35. Benge, J. F.; Kiselica, A. M. Rapid communication: Preliminary validation of a telephone adapted Montreal Cognitive Assessment for the identification of mild cognitive impairment in

- Parkinson's disease. *The Clinical Neuropsychologist* 2021, 35(1), 133–147. <https://doi.org/10.1080/13854046.2020.1801848>
36. Lu, M.; Zhao, Q.; Poston, K. L.; et al. Quantifying Parkinson's disease motor severity under uncertainty using MDS-UPDRS videos. *Medical Image Analysis* 2021, 73, 102179. <https://doi.org/10.1016/j.media.2021.102179>
37. Goetz, C. G.; Stebbins, G. T. Assuring interrater reliability for the UPDRS motor section: utility of the UPDRS teaching tape. *Movement Disorders* 2004, 19(12), 1453–1456. <https://doi.org/10.1002/mds.20220>
38. Parveen, S. Comparison of self and proxy ratings for motor performance of individuals with Parkinson disease. *Brain and Cognition* 2016, 103, 62–69. <https://doi.org/10.1016/j.bandc.2016.01.006>
39. Movement Disorder Society Task Force on Rating Scales for Parkinson's Disease. The Unified Parkinson's Disease Rating Scale (UPDRS): status and recommendations. *Movement Disorders* 2003, 18(7), 738–750. <https://doi.org/10.1002/mds.10473>
40. Louis, E. D.; Levy, G.; Côte, L. J.; et al. Diagnosing Parkinson's disease using videotaped neurological examinations: validity and factors that contribute to incorrect diagnoses. *Movement Disorders* 2002, 17(3), 513–517. <https://doi.org/10.1002/mds.10119>
41. Post, B.; Merkus, M. P.; de Bie, R. M. A.; et al. Unified Parkinson's disease rating scale motor examination: are ratings of nurses, residents in neurology, and movement disorders specialists interchangeable? *Movement Disorders* 2005, 20(12), 1577–1584. <https://doi.org/10.1002/mds.20640>
42. Vassar, S. D.; Bordelon, Y. M.; Hays, R. D.; et al. Confirmatory factor analysis of the motor unified Parkinson's disease rating scale. *Parkinson's Disease* 2012, 2012, 719167. <https://doi.org/10.1155/2012/719167>
43. Abdolahi, A.; Scoglio, N.; Killoran, A.; et al. Potential reliability and validity of a modified version of the Unified Parkinson's Disease Rating Scale that could be administered remotely. *Parkinsonism & Related Disorders* 2013, 19(2), 218–221. <https://doi.org/10.1016/j.parkreldis.2012.10.008>
44. Siderowf, A.; McDermott, M.; Kieburtz, K.; et al. Test-retest reliability of the unified Parkinson's disease rating scale in patients with early Parkinson's disease: results from a multicenter clinical trial. *Movement Disorders* 2002, 17(4), 758–763. <https://doi.org/10.1002/mds.10011>
45. Pedersen, K. F.; Larsen, J. P.; Aarsland, D. Validation of the Unified Parkinson's Disease Rating Scale (UPDRS) section I as a screening and diagnostic instrument for apathy in patients with Parkinson's disease. *Parkinsonism & Related Disorders* 2008, 14(3), 183–186. <https://doi.org/10.1016/j.parkreldis.2007.07.015>

46. Krishnan, S.; Justus, S.; Meluveetil, R.; et al. Validity of Montreal Cognitive Assessment in non-english speaking patients with Parkinson's disease. *Neurology India* 2015, 63(1), 63–67. <https://doi.org/10.4103/0028-3886.152637>
47. Hendershott, T. R.; Zhu, D.; Llanes, S.; et al. Comparative sensitivity of the MoCA and Mattis Dementia Rating Scale-2 in Parkinson's disease. *Movement Disorders* 2019, 34(2), 285–291. <https://doi.org/10.1002/mds.27575>
48. Goetz, C. G.; Stebbins, G. T.; Tilley, B. C. Calibration of unified Parkinson's disease rating scale scores to Movement Disorder Society-unified Parkinson's disease rating scale scores. *Movement Disorders* 2012, 27(10), 1239–1242. <https://doi.org/10.1002/mds.25122>
49. Aarsland, D.; Andersen, K.; Larsen, J. P.; et al. Prevalence and characteristics of dementia in Parkinson disease. *Archives of Neurology* 2003, 60(3), 387. <https://doi.org/10.1001/archneur.60.3.387>
50. Ozdilek, B.; Kenangil, G. Validation of the Turkish Version of the Montreal Cognitive Assessment Scale (MoCA-TR) in patients with Parkinson's disease. *The Clinical Neuropsychologist* 2014, 28(2), 333–343. <https://doi.org/10.1080/13854046.2014.881554>
51. Kirsch-Darrow, L.; Zahodne, L. B.; Hass, C.; et al. How cautious should we be when assessing apathy with the Unified Parkinson's Disease Rating Scale? *Movement Disorders* 2009, 24(5), 684–688. <https://doi.org/10.1002/mds.22437>
52. Martínez-Martín, P.; Benito-León, J.; Alonso, F.; et al. Patients', doctors', and caregivers' assessment of disability using the UPDRS-ADL section: are these ratings interchangeable? *Movement Disorders* 2003, 18(9), 985–992. <https://doi.org/10.1002/mds.10479>
53. Lipsmeier, F.; Taylor, K. I.; Kilchenmann, T.; et al. Evaluation of smartphone-based testing to generate exploratory outcome measures in a phase 1 Parkinson's disease clinical trial. *Movement Disorders* 2018, 33(8), 1287–1297. <https://doi.org/10.1002/mds.27376>
54. Sulzer, P.; Becker, S.; Maetzler, W.; et al. Validation of a novel Montreal Cognitive Assessment scoring algorithm in non-demented Parkinson's disease patients. *Journal of Neurology* 2018, 265(9), 1976–1984. <https://doi.org/10.1007/s00415-018-8942-4>
55. Lawton, M.; Kasten, M.; May, M. T.; et al. Validation of conversion between mini-mental state examination and montreal cognitive assessment. *Movement Disorders* 2016, 31(4), 593–596. <https://doi.org/10.1002/mds.26498>
56. Soares, T.; Vale, T. C.; Guedes, L. C.; et al. Validation of the Portuguese MDS-UPDRS: Challenges to obtain a scale applicable to different linguistic cultures. *Movement Disorders Clinical Practice* 2025, 12(1), 34–42. <https://doi.org/10.1002/mdc3.14236>

57. Harvey, P. D.; Ferris, S. H.; Cummings, J. L.; et al. Evaluation of dementia rating scales in Parkinson's disease dementia. *American Journal of Alzheimer's Disease and Other Dementias* 2010, 25(2), 142–148. <https://doi.org/10.1177/1533317509333904>
58. Nazem, S.; Siderowf, A. D.; Duda, J. E.; et al. Montreal cognitive assessment performance in patients with Parkinson's disease with "Normal" global cognition according to mini-mental state examination score. *Journal of the American Geriatrics Society* 2009, 57(2), 304–308. <https://doi.org/10.1111/j.1532-5415.2008.02096.x>
59. Nie, K.; Zhang, Y.; Wang, L.; et al. A pilot study of psychometric properties of the Beijing version of Montreal Cognitive Assessment in patients with idiopathic Parkinson's disease in China. *Journal of Clinical Neuroscience* 2012, 19(11), 1497–1500. <https://doi.org/10.1016/j.jocn.2011.11.039>
60. Khalil, H.; Aldaajani, Z. F.; Aldughmi, M.; et al. Validation of the Arabic version of the Movement Disorder Society-Unified Parkinson's Disease Rating Scale. *Movement Disorders* 2022, 37(4), 826–841. <https://doi.org/10.1002/mds.28905>
61. Raciti, L.; Nicoletti, A.; Mostile, G.; et al. Accuracy of MDS-UPDRS section IV for detecting motor fluctuations in Parkinson's disease. *Neurological Sciences* 2019, 40(6), 1271–1273. <https://doi.org/10.1007/s10072-019-03745-2>
62. Metman, L. V.; Myre, B.; Verwey, N.; et al. Test-retest reliability of UPDRS-III, dyskinesia scales, and timed motor tests in patients with advanced Parkinson's disease: an argument against multiple baseline assessments. *Movement Disorders* 2004, 19(9), 1079–1084. <https://doi.org/10.1002/mds.20101>
63. Bezdicek, O.; Červenková, M.; Moore, T. M.; et al. Determining a short form Montreal Cognitive Assessment (s-MoCA) Czech version: Validity in mild cognitive impairment Parkinson's disease and cross-cultural comparison. *Assessment* 2020, 27(8), 1960–1970. <https://doi.org/10.1177/1073191118778896>
64. Kleiner-Fisman, G.; Stern, M. B.; Fisman, D. N. Health-related quality of life in Parkinson disease: correlation between Health Utilities Index III and Unified Parkinson's Disease Rating Scale (UPDRS) in U.S. male veterans. *Health and Quality of Life Outcomes* 2010, 8(1), 91. <https://doi.org/10.1186/1477-7525-8-91>
65. Starkstein, S. E.; Merello, M. The Unified Parkinson's Disease Rating Scale: validation study of the mentation, behavior, and mood section. *Movement Disorders* 2007, 22(15), 2156–2161. <https://doi.org/10.1002/mds.21521>
66. Kasten, M.; Bruggemann, N.; Schmidt, A.; et al. Validity of the MoCA and MMSE in the detection of MCI and dementia in Parkinson disease. *Neurology* 2010, 75(5), 478; author reply 478–9. <https://doi.org/10.1212/WNL.0b013e3181e7948a>

67. Buck, P. O.; Wilson, R. E.; Seeberger, L. C.; et al. Examination of the UPDRS bradykinesia subscale: equivalence, reliability and validity. *Journal of Parkinson's Disease* 2011, 1(3), 253–258. <https://doi.org/10.3233/JPD-2011-11035>
68. Ismail, Z.; Rajji, T. K.; Shulman, K. I. Brief cognitive screening instruments: an update. *International Journal of Geriatric Psychiatry* 2010, 25(2), 111–120. <https://doi.org/10.1002/gps.2306>
69. Ruzafa-Valiente, E.; Fernández-Bobadilla, R.; García-Sánchez, C.; et al. Parkinson's Disease-Cognitive Functional Rating Scale across different conditions and degrees of cognitive impairment. *Journal of the Neurological Sciences* 2016, 361, 66–71. <https://doi.org/10.1016/j.jns.2015.12.018>
70. Tumas, V.; Borges, V.; Ballalai-Ferraz, H.; et al. Some aspects of the validity of the Montreal Cognitive Assessment (MoCA) for evaluating cognitive impairment in Brazilian patients with Parkinson's disease. *Dementia & Neuropsychologia* 2016, 10(4), 333–338. <https://doi.org/10.1590/s1980-5764-2016dn1004013>
71. van Hilten, J. J.; van der Zwan, A. D.; Zwinderman, A. H.; et al. Rating impairment and disability in Parkinson's disease: evaluation of the Unified Parkinson's Disease Rating Scale. *Movement Disorders* 1994, 9(1), 84–88. <https://doi.org/10.1002/mds.870090113>
72. Martinez-Martin, P.; Chaudhuri, K. R.; Rojo-Abuin, J. M.; et al. Assessing the non-motor symptoms of Parkinson's disease: MDS-UPDRS and NMS Scale. *European Journal of Neurology* 2015, 22(1), 37–43. <https://doi.org/10.1111/ene.12165>
73. Ramsay, N.; Macleod, A. D.; Alves, G.; et al. Validation of a UPDRS-/MDS-UPDRS-based definition of functional dependency for Parkinson's disease. *Parkinsonism & Related Disorders* 2020, 76, 49–53. <https://doi.org/10.1016/j.parkreldis.2020.05.034>
74. Isella, V.; Mapelli, C.; Siri, C.; et al. Validation and attempts of revision of the MDS-recommended tests for the screening of Parkinson's disease dementia. *Parkinsonism & Related Disorders* 2014, 20(1), 32–36. <https://doi.org/10.1016/j.parkreldis.2013.09.008>
75. Bugalho, P.; da Silva, J. A.; Cargaleiro, I.; et al. Psychiatric symptoms screening in the early stages of Parkinson's disease. *Journal of Neurology* 2012, 259(1), 124–131. <https://doi.org/10.1007/s00415-011-6140-8>
76. Freitas, S.; Simões, M. R.; Alves, L.; et al. Montreal cognitive assessment. *Alzheimer Disease and Associated Disorders* 2013, 27(1), 37–43. <https://doi.org/10.1097/wad.0b013e3182420bfe>
77. Kremer, N. I.; Smid, A.; Lange, S. F.; et al. Supine MDS-UPDRS-III assessment: An explorative study. *Journal of Clinical Medicine* 2023, 12(9), 3108. <https://doi.org/10.3390/jcm12093108>
78. Goetz, C. G.; Tilley, B. C.; Shaftman, S. R.; et al. Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): scale presentation

- and clinimetric testing results. *Movement Disorders* 2008, 23(15), 2129–2170. <https://doi.org/10.1002/mds.22340>
79. Park, J.; Koh, S. B.; Kwon, K. Y.; et al. Validation study of the official Korean version of the Movement Disorder Society-Unified Parkinson's Disease Rating Scale. *Journal of Clinical Neurology* 2020, 16(4), 633–645. <https://doi.org/10.3988/jcn.2020.16.4.633>
80. Wissel, B. D.; Mitsi, G.; Dwivedi, A. K.; et al. Tablet-based application for objective measurement of motor fluctuations in Parkinson disease. *Digital Biomarkers* 2017, 1(2), 126–135. <https://doi.org/10.1159/000485468>
81. Statucka, M.; Cherian, K.; Fasano, A.; et al. Multiculturalism: A challenge for cognitive screeners in Parkinson's disease. *Movement Disorders Clinical Practice* 2021, 8(5), 733–742. <https://doi.org/10.1002/mdc3.13240>
82. Stocchi, F.; Radicati, F. G.; Chaudhuri, K. R.; et al. The Parkinson's Disease Composite Scale: results of the first validation study. *European Journal of Neurology* 2018, 25(3), 503–511. <https://doi.org/10.1111/ene.13529>
83. Forjaz, M. J.; Ayala, A.; Testa, C. M.; et al. Proposing a Parkinson's disease-specific tremor scale from the MDS-UPDRS. *Movement Disorders* 2015, 30(8), 1139–1143. <https://doi.org/10.1002/mds.26271>
84. Hariz, G.-M.; Fredricks, A.; Stenmark-Persson, R.; et al. Blinded versus unblinded evaluations of motor scores in patients with Parkinson's disease randomized to deep brain stimulation or best medical therapy. *Movement Disorders Clinical Practice* 2021, 8(2), 286–287. <https://doi.org/10.1002/mdc3.13141>
85. Goetz, C. G.; Luo, S.; Wang, L.; et al. Handling missing values in the MDS-UPDRS. *Movement Disorders* 2015, 30(12), 1632–1638. <https://doi.org/10.1002/mds.26153>
86. Morinan, G.; Hauser, R. A.; Schrag, A.; et al. Abbreviated MDS-UPDRS for remote monitoring in PD identified using exhaustive computational search. *Parkinson's Disease* 2022, 2022, 2920255. <https://doi.org/10.1155/2022/2920255>
87. Regnault, A.; Boroojerdi, B.; Meunier, J.; et al. Does the MDS-UPDRS provide the precision to assess progression in early Parkinson's disease? Learnings from the Parkinson's progression marker initiative cohort. *Journal of Neurology* 2019, 266(8), 1927–1936. <https://doi.org/10.1007/s00415-019-09348-3>
88. Jenkins, M. E.; Johnson, A. M.; Holmes, J. D.; et al. Predictive validity of the UPDRS postural stability score and the Functional Reach Test, when compared with ecologically valid reaching tasks. *Parkinsonism & Related Disorders* 2010, 16(6), 409–411. <https://doi.org/10.1016/j.parkreldis.2010.04.002>

89. Goetz, C. G.; Fahn, S.; Martinez-Martin, P.; et al. Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): Process, format, and clinimetric testing plan. *Movement Disorders* 2007, 22(1), 41–47. <https://doi.org/10.1002/mds.21198>
90. Raciti, L.; Nicoletti, A.; Mostile, G.; et al. Validation of the UPDRS section IV for detection of motor fluctuations in Parkinson's disease. *Parkinsonism & Related Disorders* 2016, 27, 98–101. <https://doi.org/10.1016/j.parkreldis.2016.03.008>
91. Forjaz, M. J.; Martinez-Martin, P. Metric attributes of the unified Parkinson's disease rating scale 3.0 battery: part II, construct and content validity. *Movement Disorders* 2006, 21(11), 1892–1898. <https://doi.org/10.1002/mds.21071>
92. Zitser, J.; Peretz, C.; Ber David, A.; et al. Validation of the Hebrew version of the Movement Disorder Society—unified Parkinson's disease rating scale. *Parkinsonism & Related Disorders* 2017, 45, 7–12. <https://doi.org/10.1016/j.parkreldis.2017.09.012>
93. Tosin, M. H. S.; Sanchez-Ferro, A.; Wu, R.-M.; et al. In-home remote assessment of the MDS-UPDRS part III: Multi-cultural development and validation of a guide for patients. *Movement Disorders Clinical Practice* 2024, 11(12), 1576–1581. <https://doi.org/10.1002/mdc3.14203>
94. Benge, J. F.; Balsis, S.; Madeka, T.; et al. Factor structure of the Montreal Cognitive Assessment items in a sample with early Parkinson's disease. *Parkinsonism & Related Disorders* 2017, 41, 104–108. <https://doi.org/10.1016/j.parkreldis.2017.05.023>
95. Gallagher, D. A.; Goetz, C. G.; Stebbins, G.; et al. Validation of the MDS-UPDRS Part I for nonmotor symptoms in Parkinson's disease. *Movement Disorders* 2012, 27(1), 79–83. <https://doi.org/10.1002/mds.23939>
96. Kenny, L.; Azizi, Z.; Moore, K.; et al. Inter-rater reliability of hand motor function assessment in Parkinson's disease: Impact of clinician training. *Clinical Parkinsonism & Related Disorders* 2024, 11, 100278. <https://doi.org/10.1016/j.prdoa.2024.100278>
97. de Deus Fonticoba, T.; Santos García, D.; Macías Arribí, M. Variabilidad en la exploración motora de la enfermedad de Parkinson entre el neurólogo experto en trastornos del movimiento y la enfermera especializada. *Neurología (English Edition)* 2019, 34(8), 520–526. <https://doi.org/10.1016/j.nrl.2017.03.005>
98. Dujardin, K.; Duhem, S.; Guerouaou, N.; et al. Validation in French of the Montreal Cognitive Assessment 5-Minute, a brief cognitive screening test for phone administration. *Revue Neurologique* 2021, 177(8), 972–979. <https://doi.org/10.1016/j.neurol.2020.09.002>
99. Tosin, M. H. S.; Stebbins, G. T.; Comella, C.; et al. Does MDS-UPDRS provide greater sensitivity to mild disease than UPDRS in DE Novo Parkinson's disease? *Movement Disorders Clinical Practice* 2021, 8(7), 1092–1099. <https://doi.org/10.1002/mdc3.13329>

100. De Deus Fonticoba, T.; Santos García, D.; Macías Arribí, M. Inter-rater variability in motor function assessment in Parkinson's disease between experts in movement disorders and nurses specialising in PD management. *Neurología (English Edition)* 2019, 34(8), 520–526. <https://doi.org/10.1016/j.nrleng.2017.03.006>
101. Kletzel, S. L.; Hernandez, J. M.; Miskiel, E. F.; et al. Evaluating the performance of the Montreal Cognitive Assessment in early stage Parkinson's disease. *Parkinsonism & Related Disorders* 2017, 37, 58–64. <https://doi.org/10.1016/j.parkreldis.2017.01.012>
102. Gill, D. J.; Freshman, A.; Blender, J. A.; et al. The Montreal cognitive assessment as a screening tool for cognitive impairment in Parkinson's disease. *Movement Disorders* 2008, 23(7), 1043–1046. <https://doi.org/10.1002/mds.22017>
103. Martínez-Martín, P.; Gil-Nagel, A.; Gracia, L. M.; et al. Unified Parkinson's Disease Rating Scale characteristics and structure. The Cooperative Multicentric Group. *Movement Disorders* 1994, 9(1), 76–83. <https://doi.org/10.1002/mds.870090112>
104. Gasser, A.-I.; Calabrese, P.; Kalbe, E.; et al. Cognitive screening in Parkinson's disease: Comparison of the Parkinson Neuropsychometric Dementia Assessment (PANDA) with 3 other short scales. *Revue Neurologique* 2016, 172(2), 138–145. <https://doi.org/10.1016/j.neurol.2015.08.004>
105. D'Iorio, A.; Aiello, E. N.; Amboni, M.; et al. Validity and diagnostics of the Italian version of the Montreal Cognitive Assessment (MoCA) in non-demented Parkinson's disease patients. *Aging Clinical and Experimental Research* 2023, 35(10), 2157–2163. <https://doi.org/10.1007/s40520-023-02493-w>
106. Evers, L. J. W.; Krijthe, J. H.; Meinders, M. J.; et al. Measuring Parkinson's disease over time: The real-world within-subject reliability of the MDS-UPDRS. *Movement Disorders* 2019, 34(10), 1480–1487. <https://doi.org/10.1002/mds.27790>
107. Louis, E. D.; Lynch, T.; Marder, K.; et al. Reliability of patient completion of the historical section of the Unified Parkinson's Disease Rating Scale. *Movement Disorders* 1996, 11(2), 185–192. <https://doi.org/10.1002/mds.870110212>
108. Fiorenzato, E.; Weis, L.; Falup-Pecurariu, C.; et al. MoCA vs. MMSE sensitivity as screening instruments of cognitive impairment in PD, MSA and PSP patients. *Parkinsonism & Related Disorders* 2016, 22, e59–e60. <https://doi.org/10.1016/j.parkreldis.2015.10.127>




Figure 1. PRISMA flow diagram for study selection.






Figure 4. Bias Domain Coverage in PD Assessment Scale Research

Bias Domain	Studies Addressing (n)	Percentage (%)	Risk Level
Education Adjustment for Cognitive Tests	13	11.9	Very High
Race/Ethnicity Data Collection	8	7.3	Critical
Treatment Effects on Cognition	1	0.9	Critical
Institutional Resources Reporting	11	10.1	Very High
Digital Literacy/Access Barriers	0	0.0	Critical
Age-Specific Normative Values	3	2.8	Critical
Administrator Training/Knowledge	33	30.3	High
UPDRS ON/OFF State Documentation	24	22.0	High
Sex/Gender Analysis	47	43.1	Moderate
Underrepresented Subgroups	21	19.3	High
ON Medication State Testing	18	16.5	Very High
OFF Medication State Testing	4	3.7	Critical
Cognitive Domain Specification	43	39.4	Moderate
Cognitive Impairment Criteria	49	45.0	Moderate
Confounding Variables Control	16	14.7	Very High


Table 1. Bias Domain Analysis Results (n=109)




404x228mm (87 x 87 DPI)




454x330mm (39 x 39 DPI)



408x215mm (47 x 47 DPI)



423x282mm (87 x 87 DPI)



428x296mm (47 x 47 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	JOURNAL TITLE	COUNTS
1.	Movement Disorders	29
2.	Parkinsonism & Related Disorders	11
3.	Movement Disorders Clinical Practice	5
4.	Neurology	5
5.	Neurological Sciences	4
6.	Journal of Neurology	4
7.	Journal of Parkinson's Disease	4
8.	International Journal of Geriatric Psychiatry	2
9.	American Journal of Alzheimer's Disease & Othe...	2
10.	European Journal of Neurology	2
11.	Revue Neurologique	2
12.	The Clinical Neuropsychologist	2
13.	Parkinson's Disease	2
14.	Neurologia (English Edition)	2
15.	Aging Clinical and Experimental Research	1
16.	Health Informatics Journal	1
17.	Frontiers in Neurology	1
18.	Digital Biomarkers	1
19.	Dementia and Geriatric Cognitive Disorders Extra	1
20.	Clinical Parkinsonism & Related Disorders	1
21.	Dementia & Neuropsychologia	1
22.	Clinical Neuropharmacology	1
23.	Brain and Cognition	1
24.	Alzheimer Disease & Associated Disorders	1
25.	Assessment	1
26.	Archives of Clinical Neuropsychology	1
27.	Applied Neuropsychology Adult	1
28.	International Journal of Speech-Language Patho...	1
29.	JAMA Neurology	1
30.	IEEE Transactions on Neural Systems and Rehabi...	1
31.	Health and Quality of Life Outcomes	1
32.	Journal of the Neurological Sciences	1
33.	Journal of the American Geriatrics Society	1
34.	Journal of Neurology Neurosurgery & Psychiatry	1
35.	Journal of Movement Disorders	1
36.	Journal of Clinical Neurology	1
37.	Journal of Clinical Neuroscience	1
38.	Journal of Advanced Nursing	1
39.	Journal of Clinical Medicine	1
40.	Ideggyógyászati Szemle	1
41.	Neurological Research	1
42.	Medical Image Analysis	1
43.	NeuroRehabilitation An International Interdisc...	1
44.	Neurologia i Neurochirurgia Polska	1
45.	Neurology India	1
46.	PLOS ONE	1
47.	Value in Health	1

531x752mm (79 x 79 DPI)

<u>Region</u>	<u>Publications (n)</u>	<u>Percentage (%)</u>	<u>Income Level</u>
Europe (HIC)	48	44.0	High
North America (HIC)	39	35.8	High
Asia (HIC)	9	8.3	High
Asia (LMIC)	4	3.7	LMIC
Oceania (HIC)	3	2.8	High
South America (LMIC)	3	2.8	LMIC
Africa (LMIC)	2	1.8	LMIC
North America (LMIC)	1	0.9	LMIC

446x216mm (47 x 47 DPI)

Characteristic	Value
Total papers analyzed	109
Publication period	1996-2024
Total authors examined	655
Countries represented	34
Journals represented	47
Most common assessment scales	UPDRS, MDS-UPDRS, MoCA, MMSE

467x169mm (47 x 47 DPI)