

# Homework 2

Dr. Purna Gamage

9/19/2022

## Problem 1 (30 Points)

Consider the random variable defined by counting the number of failures until the first success, for independent trials with success probability  $p$ . Given  $p$  between 0 and 1, the **R** commands `myattempts(p)` and `rgeom(1,p)` both simulate this random variable. Find a way of demonstrating that the two commands indeed give the same results, for five different values of  $p$ . *You may find the `table()` function useful.*

Choose five  $p$ 's of your choice, e.g.  $p \in \{.1, .3, .5, .7, .9\}$ . Run `myattempts` and `rgeom` each 10000 times.

- For the first  $p$ , store the fraction of outcomes in the simulation in the columns of a suitable data frame and compare (Hint: use 3 columns to compare `rgeom()`, `myattempts()` and `dgeom()`) (10 points)
- For the second  $p$ , Compare distribution by computing statistics such as mean and standard deviation. (5 points)
- For the third  $p$ , plot both distributions as histograms in the same plot. (5 points)
- For the fourth  $p$ , make side-by-side box plots. (5 points)
- For the fifth  $p$ , plot the two empirical distribution functions in the same plot. (5 points)

## Problem 2 (10 Points)

Consider the following random experiment: draw a uniformly distributed random number  $X_1$  from the interval  $(0, 1)$ . Next, draw a uniformly distributed random number  $X_2$  from the interval  $(0, 1 + X_1)$ , a uniformly distributed random number  $X_3$  from the interval  $(0, 1 + X_2)$  and so on until  $X_{20}$ .

Use a monte carlo simulation to give an approximate answer to What is the mean value of  $X_{20}$ ? and use a histogram to identify the distribution of  $X_{20}$ .

$$X_1 \sim \text{unif}(0, 1) \quad X_2 \sim \text{unif}(0, 1 + X_1) \quad X_3 \sim \text{unif}(0, 1 + X_2) \quad \dots \quad X_{20} \sim \text{unif}(0, 1 + X_{19})$$

The **R** command for drawing a uniformly distributed random number from the interval  $(0, b)$  is `runif(1, min = 0, max = b)`.

## Problem 3 (15 Points)

Suppose that the daily power consumption of a major city,  $X$ , has a Gamma distribution with shape parameter  $r = 4$  and scale parameter  $\rho = 2$ . Use **R** to compute the following quantities: \

- $\text{Prob}(X \leq 12)$  (3 Points)
- $\text{Prob}(X > 5)$  (3 Points)
- $\text{Prob}(|X - 8| < 1)$  (4.5 Points)

d.  $z$  such that  $\text{Prob}(X < z) = .95$  (4.5 Points)

(Hint:  $|X - 8| < 1$  is equivalent to  $7 < X < 9$ )

### Problem 4 (15 Points)

Probability theory says that a binomial distribution,  $B(n, p)$  is close to that of a normal distribution with mean  $np$  and standard deviation  $\sqrt{np(1-p)}$ , if  $np$  and  $n(1-p)$  are both sufficiently large, e.g. at least 10.

Check this by plotting both cumulative distribution functions in the same figure, using a staircase plot for the binomial distribution and a line plot for the normal distribution, for three different cases: a case where both  $np$  and  $n(1-p)$  are large, a case where  $np$  is large and  $n(1-p) < 10$ , and a case where  $np < 10$  and  $n(1-p) < 10$ .

Describe what happens in all three cases. In what sense are the cdf's not close in cases 2 and 3? (Hint: Compare with a cdf of the normal distribution)

(Hint: You can specify a staircase plot using the `lines()` function and passing the optional parameter `type=s`)

### Problem 5 (15 Points)

A graphical technique for checking whether a sample has an approximate normal distribution is a “quantile-quantile” plot. The **R** command is `qqnorm(x)`, where  $x$  is the vector of sample values. If the plot is approximately a straight line, then this suggests that the sample comes from a normal distribution. Use the dataset from the `openintro` package to find out which of the the four distributions (three exams and course grade) is the closest to a normal distribution? You can load the dataset with the following commands `library(openintro)` and `data(exam_grades)`. Explore this by making `qqnorm()` and `qqline()` plots of the four distributions. How close to straight lines are the plots in each case? How do the plots differ from straight lines? Hint: Make sure to remove the NA values.

### Problem 6 (15 Points)

If  $X$  has a continuous distribution with cumulative distribution function  $F$ , then the new random variable  $U = F(X)$  has a uniform  $U(0, 1)$  distribution. Verify this with simulations for three different continuous distributions of your choice, by making a random sample of sufficient size, sorting it, plugging it into the cdf  $F$ , and plotting the result.

### Bonus Problem (20 Points)

Suppose for  $X = X_1 + X_2$  is the sum of two exponentially distributed random variables with the same parameter  $\lambda$ . Then  $X^\alpha$  is very nearly normally distributed for a suitable choice of  $\alpha$ . Determine an approximate value for  $\alpha$  (within 0.05), using a simulation and `qqnorm()` plots for each of your choices of  $\alpha$ .