

SBML Model Report

Model name: “Heldt2018 - Proliferation-quiescence decision in response to DNA damage”



May 17, 2018

1 General Overview

This is a document in SBML Level 2 Version 4 format. This model was created by the following two authors: Frank Stefan Heldt¹ and Matthieu MAIRE² at March first 2017 at 11:13 a. m. and last time modified at May 17th 2018 at 4:39 p. m. Table 1 provides an overview of the quantities of all components of this model.

Table 1: Number of components in this model, which are described in the following sections.

Element	Quantity	Element	Quantity
compartment types	0	compartments	1
species types	0	species	30
events	0	constraints	0
reactions	49	function definitions	24
global parameters	56	unit definitions	3
rules	7	initial assignments	0

Model Notes

Heldt2018 - Proliferation-quiescence decision in response to DNA damage

¹University of Oxford, stefan.heldt@bioch.ox.ac.uk

²EMBL-EBI, matt.maire@free.fr

This model is described in the article: [A comprehensive model for the proliferation-quiescence decision in response to endogenous DNA damage in human cells](#). Heldt FS, Barr AR, Cooper S, Bakal C, Novk B. Proc. Natl. Acad. Sci. U.S.A. 2018 Feb; :

Abstract:

Human cells that suffer mild DNA damage can enter a reversible state of growth arrest known as quiescence. This decision to temporarily exit the cell cycle is essential to prevent the propagation of mutations, and most cancer cells harbor defects in the underlying control system. Here we present a mechanistic mathematical model to study the proliferation-quiescence decision in nontransformed human cells. We show that two bistable switches, the restriction point (RP) and the G1/S transition, mediate this decision by integrating DNA damage and mitogen signals. In particular, our data suggest that the cyclin-dependent kinase inhibitor p21 (Cip1/Waf1), which is expressed in response to DNA damage, promotes quiescence by blocking positive feedback loops that facilitate G1 progression downstream of serum stimulation. Intriguingly, cells exploit bistability in the RP to convert graded p21 and mitogen signals into an all-or-nothing cell-cycle response. The same mechanism creates a window of opportunity where G1 cells that have passed the RP can revert to quiescence if exposed to DNA damage. We present experimental evidence that cells gradually lose this ability to revert to quiescence as they progress through G1 and that the onset of rapid p21 degradation at the G1/S transition prevents this response altogether, insulating S phase from mild, endogenous DNA damage. Thus, two bistable switches conspire in the early cell cycle to provide both sensitivity and robustness to external stimuli.

This model is hosted on [BioModels Database](#) and identified by: [MODEL1703030000](#).

To cite BioModels Database, please use: [Chelliah V et al. BioModels: ten-year anniversary. Nucl. Acids Res. 2015, 43\(Database issue\):D542-8.](#)

To the extent possible under law, all copyright and related or neighbouring rights to this encoded model have been dedicated to the public domain worldwide. Please refer to [CC0 Public Domain Dedication](#) for more information.

2 Unit Definitions

This is an overview of five unit definitions of which two are predefined by SBML and not mentioned in the model.

2.1 Unit volume

Name volume

Definition ml

2.2 Unit time

Name time

Definition 60 s

2.3 Unit substance

Name substance

Definition mmol

2.4 Unit area

Notes Square metre is the predefined SBML unit for area since SBML Level 2 Version 1.

Definition m²

2.5 Unit length

Notes Metre is the predefined SBML unit for length since SBML Level 2 Version 1.

Definition m

3 Compartment

This model contains one compartment.

Table 2: Properties of all compartments.

Id	Name	SBO	Spatial Dimensions	Size	Unit	Constant	Outside
Cell	cell		3	1	litre	<input checked="" type="checkbox"/>	

3.1 Compartment Cell

This is a three dimensional compartment with a constant size of one ml.

Name cell

4 Species

This model contains 30 species. The boundary condition of seven of these species is set to true so that these species' amount cannot be changed by any reaction. Section 9 provides further details and the derived rates of change of each species.

Table 3: Properties of each species.

Id	Name	Compartment	Derived Unit	Constant	Boundary Condition
Rb	Retinoblastoma_protein	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
pRb	Retinoblastoma_protein- _hyperphosphorylated	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
E2f	E2f_active	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
RbE2f	Retinoblastoma_protein_E2f_complex- _inactive	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
E1	Emi1	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
C1	Cdh1_C_APC_active	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
pC1	C_Cdh1_APC_phosphorylated_inactive	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
E1C1	Emi1_C_Cdh1_complex_inactive	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
P21	P21	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
Ce	CyclinE_Cdk2_active	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
Ca	CyclinA_Cdk2	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
CeP21	CyclinE_Cdk2_P21_complex_inactive	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
CaP21	CyclinA_Cdk2_P21_complex_inactive	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
aPcna	Pcna_nuclear_active	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
iPcna	PCNA_Nuclear_inactive	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
Rc	Pre_Replication_complex	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
pRc	Pre_Replication_complex_primed	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
aRc	Pre_Replication_complex_active	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>

Id	Name	Compartment	Derived Unit	Constant	Boundary Condition
iRc	Pre_Replication_complex_inactive	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
Dna	Dna	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
P53	P53	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
Dam	Dna_damage	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
Pr	Activity_probe_of_APC_C_Cdh1	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input type="checkbox"/>
tRb	Retinoblastoma_protein_total	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input checked="" type="checkbox"/>
tE2f	E2f_total	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input checked="" type="checkbox"/>
tE1	Emi1_total	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input checked="" type="checkbox"/>
tC1	C_Cdh1_APC_total	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input checked="" type="checkbox"/>
tCe	CyclinE_Cdk2_total	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input checked="" type="checkbox"/>
tCa	CyclinA_Cdk2_total	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input checked="" type="checkbox"/>
tP21	P21_total	Cell	$\text{mmol} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input checked="" type="checkbox"/>

5 Parameters

This model contains 56 global parameters.

Table 4: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
Cd	Cd		0.650		✓
Skp2	Skp2		1.000		✓
Cdt2	Cdt2		1.000		✓
kSyE2f	kSyE2f		0.030		✓
kSyE2fE2f	kSyE2fE2f		0.040		✓
jSyE2f	jSyE2f		0.200		✓
kAsRbE2f	kAsRbE2f		5.000		✓
kDsRbE2f	kDsRbE2f		0.005		✓
kDeE2f	kDeE2f		0.050		✓
kPhRbCd	kPhRbCd		0.200		✓
kPhRbCe	kPhRbCe		0.300		✓
kPhRbCa	kPhRbCa		0.300		✓
kDpRb	kDpRb		0.050		✓
kSyE1	kSyE1		0.005		✓
kDeE1C1	kDeE1C1		0.005		✓
kDeE1	kDeE1		$5 \cdot 10^{-4}$		✓
kPhC1	kPhC1		0.000		✓
kPhC1Ce	kPhC1Ce		0.010		✓
kPhC1Ca	kPhC1Ca		1.000		✓
kDpC1	kDpC1		0.050		✓
kAsE1C1	kAsE1C1		10.000		✓
kDsE1C1	kDsE1C1		0.010		✓
kSyP21	kSyP21		0.002		✓
kSyP21P53	kSyP21P53		0.008		✓
kDeP21	kDeP21		0.003		✓
kDeP21Cy	kDeP21Cy		0.007		✓
kDeP21aRc	kDeP21aRc		1.000		✓
kSyCe	kSyCe		0.010		✓
kSyCa	kSyCa		0.020		✓
kAsCyP21	kAsCyP21		1.000		✓
kDsCyP21	kDsCyP21		0.050		✓
kDeCe	kDeCe		0.004		✓
kDeCa	kDeCa		0.010		✓
kDeCeCa	kDeCeCa		0.015		✓
kDeCaC1	kDeCaC1		2.000		✓
kImPc	kImPc		0.003		✓

Id	Name	SBO	Value	Unit	Constant
kExPc	kExPc		0.006		✓
kPhRc	kPhRc		0.100		✓
kDpRc	kDpRc		0.050		✓
jCy	jCy		1.800		✓
n	n		6.000		✓
kAsRcPc	kAsRcPc		0.010		✓
kDsRcPc	kDsRcPc		0.001		✓
kAsPcP21	kAsPcP21		100.000		✓
kDsPcP21	kDsPcP21		0.010		✓
kSyDna	kSyDna		0.009		✓
kSyP53	kSyP53		0.050		✓
kDeP53	kDeP53		0.050		✓
jP53	jP53		0.010		✓
kGeDam	kGeDam		0.001		✓
kGeDamArc	kGeDamArc		0.012		✓
kReDam	kReDam		0.001		✓
kReDamP53	kReDamP53		0.005		✓
jDam	jDam		0.500		✓
kSyPr	kSyPr		0.010		✓
kDePr	kDePr		10^{-4}		✓

6 Function definitions

This is an overview of 24 function definitions.

6.1 Function definition `Constant_flux__irreversible`

Name Constant flux (irreversible)

Argument v

Mathematical Expression

$$v \quad (1)$$

6.2 Function definition `rPhRb_2`

Name `rPhRb_2`

Arguments $[Ca]$, Cd , $[Ce]$, $[RbE2f]$, $kPhRbCa$, $kPhRbCd$, $kPhRbCe$

Mathematical Expression

$$(kPhRbCd \cdot Cd + kPhRbCe \cdot [Ce] + kPhRbCa \cdot [Ca]) \cdot [RbE2f] \quad (2)$$

6.3 Function definition $r_{\text{SyE2f_1}}$

Name $r_{\text{SyE2f_1}}$

Arguments $[\text{E2f}]$, j_{SyE2f} , k_{SyE2f} , k_{SyE2fE2f}

Mathematical Expression

$$k_{\text{SyE2f}} + \frac{k_{\text{SyE2fE2f}} \cdot [\text{E2f}]}{j_{\text{SyE2f}} + [\text{E2f}]} \quad (3)$$

6.4 Function definition $r_{\text{DeP21_1}}$

Name $r_{\text{DeP21_1}}$

Arguments $[\text{Ca}]$, Cdt2 , $[\text{Ce}]$, $[\text{CeP21}]$, Skp2 , $[\text{aRc}]$, k_{DeP21} , k_{DeP21Cy} , k_{DeP21aRc}

Mathematical Expression

$$(k_{\text{DeP21}} + k_{\text{DeP21Cy}} \cdot \text{Skp2} \cdot ([\text{Ce}] + [\text{Ca}]) + k_{\text{DeP21aRc}} \cdot \text{Cdt2} \cdot [\text{aRc}]) \cdot [\text{CeP21}] \quad (4)$$

6.5 Function definition $r_{\text{DeP53_1}}$

Name $r_{\text{DeP53_1}}$

Arguments $[\text{Dam}]$, $[\text{P53}]$, j_{P53} , k_{DeP53}

Mathematical Expression

$$\frac{k_{\text{DeP53}}}{j_{\text{P53}} + [\text{Dam}]} \cdot [\text{P53}] \quad (5)$$

6.6 Function definition $r_{\text{DeCa_2}}$

Name $r_{\text{DeCa_2}}$

Arguments $[\text{C1}]$, $[\text{CaP21}]$, k_{DeCa} , k_{DeCaC1}

Mathematical Expression

$$(k_{\text{DeCa}} + k_{\text{DeCaC1}} \cdot [\text{C1}]) \cdot [\text{CaP21}] \quad (6)$$

6.7 Function definition $r_{\text{DeP21_5}}$

Name $r_{\text{DeP21_5}}$

Arguments $[\text{Ca}]$, Cdt2 , $[\text{Ce}]$, Skp2 , $[\text{aRc}]$, $[\text{iRc}]$, k_{DeP21} , k_{DeP21Cy} , k_{DeP21aRc}

Mathematical Expression

$$(k_{\text{DeP21}} + k_{\text{DeP21Cy}} \cdot \text{Skp2} \cdot ([\text{Ce}] + [\text{Ca}]) + k_{\text{DeP21aRc}} \cdot \text{Cdt2} \cdot [\text{aRc}]) \cdot [\text{iRc}] \quad (7)$$

6.8 Function definition `rPhRb_1`

Name `rPhRb_1`

Arguments `[Ca]`, `Cd`, `[Ce]`, `[Rb]`, `kPhRbCa`, `kPhRbCd`, `kPhRbCe`

Mathematical Expression

$$(kPhRbCd \cdot Cd + kPhRbCe \cdot [Ce] + kPhRbCa \cdot [Ca]) \cdot [Rb] \quad (8)$$

6.9 Function definition `rSyP21_1`

Name `rSyP21_1`

Arguments `[P53]`, `kSyP21`, `kSyP21P53`

Mathematical Expression

$$kSyP21 + kSyP21P53 \cdot [P53] \quad (9)$$

6.10 Function definition `rPhC1_1`

Name `rPhC1_1`

Arguments `[C1]`, `[Ca]`, `[Ce]`, `kPhC1`, `kPhC1Ca`, `kPhC1Ce`

Mathematical Expression

$$(kPhC1 + kPhC1Ce \cdot [Ce] + kPhC1Ca \cdot [Ca]) \cdot [C1] \quad (10)$$

6.11 Function definition `rDeP21_3`

Name `rDeP21_3`

Arguments `[Ca]`, `Cdt2`, `[Ce]`, `[P21]`, `Skp2`, `[aRc]`, `kDeP21`, `kDeP21Cy`, `kDeP21aRc`

Mathematical Expression

$$(kDeP21 + kDeP21Cy \cdot Skp2 \cdot ([Ce] + [Ca]) + kDeP21aRc \cdot Cdt2 \cdot [aRc]) \cdot [P21] \quad (11)$$

6.12 Function definition `rDsRc_3`

Name `rDsRc_3`

Arguments `[Dna]`, `[aRc]`

Mathematical Expression

$$\begin{cases} 0 & \text{if } [Dna] < 1 \\ \begin{cases} 1 \cdot [aRc] & \text{if } [Dna] > 1 \\ 0.5 \cdot [aRc] & \text{otherwise} \end{cases} & \text{otherwise} \end{cases} \quad (12)$$

6.13 Function definition `rDeCe_2`

Name `rDeCe_2`

Arguments `[Ca]`, `[CeP21]`, `kDeCe`, `kDeCeCa`

Mathematical Expression

$$(kDeCe + kDeCeCa \cdot [Ca]) \cdot [CeP21] \quad (13)$$

6.14 Function definition `rDeCe_1`

Name `rDeCe_1`

Arguments `[Ca]`, `[Ce]`, `kDeCe`, `kDeCeCa`

Mathematical Expression

$$(kDeCe + kDeCeCa \cdot [Ca]) \cdot [Ce] \quad (14)$$

6.15 Function definition `rReDam_1`

Name `rReDam_1`

Arguments `[Dam]`, `[P53]`, `jDam`, `kReDam`, `kReDamP53`

Mathematical Expression

$$\left(kReDam + \frac{kReDamP53 \cdot [P53]}{jDam + [Dam]} \right) \cdot [Dam] \quad (15)$$

6.16 Function definition `rPhC1_2`

Name `rPhC1_2`

Arguments `[Ca]`, `[Ce]`, `[E1C1]`, `kPhC1`, `kPhC1Ca`, `kPhC1Ce`

Mathematical Expression

$$(kPhC1 + kPhC1Ce \cdot [Ce] + kPhC1Ca \cdot [Ca]) \cdot [E1C1] \quad (16)$$

6.17 Function definition `rDeP21_2`

Name `rDeP21_2`

Arguments `[Ca]`, `[CaP21]`, `Cdt2`, `[Ce]`, `Skp2`, `[aRc]`, `kDeP21`, `kDeP21Cy`, `kDeP21aRc`

Mathematical Expression

$$(kDeP21 + kDeP21Cy \cdot Skp2 \cdot ([Ce] + [Ca]) + kDeP21aRc \cdot Cdt2 \cdot [aRc]) \cdot [CaP21] \quad (17)$$

6.18 Function definition `rPhRc_1`

Name `rPhRc_1`

Arguments `[Ca]`, `[Ce]`, `[Rc]`, `jCy`, `kPhRc`, `n`

Mathematical Expression

$$\frac{kPhRc \cdot ([Ce] + [Ca])^n}{jCy^n + ([Ce] + [Ca])^n} \cdot [Rc] \quad (18)$$

6.19 Function definition `rDeCa_1`

Name `rDeCa_1`

Arguments `[C1]`, `[Ca]`, `kDeCa`, `kDeCaC1`

Mathematical Expression

$$(kDeCa + kDeCaC1 \cdot [C1]) \cdot [Ca] \quad (19)$$

6.20 Function definition `rDsRc_4`

Name `rDsRc_4`

Arguments `[Dna]`, `[iRc]`

Mathematical Expression

$$\begin{cases} 0 & \text{if } [Dna] < 1 \\ \begin{cases} 1 \cdot [iRc] & \text{if } [Dna] > 1 \\ 0.5 \cdot [iRc] & \text{otherwise} \end{cases} & \text{otherwise} \end{cases} \quad (20)$$

6.21 Function definition `rDeP21_4`

Name `rDeP21_4`

Arguments `[Ca]`, `Cdt2`, `[Ce]`, `Skp2`, `[aRc]`, `[iPcna]`, `kDeP21`, `kDeP21Cy`, `kDeP21aRc`

Mathematical Expression

$$(kDeP21 + kDeP21Cy \cdot Skp2 \cdot ([Ce] + [Ca]) + kDeP21aRc \cdot Cdt2 \cdot [aRc]) \cdot [iPcna] \quad (21)$$

6.22 Function definition r_{DsRc_1}

Name r_{DsRc_1}

Arguments $[Dna]$, $[Rc]$

Mathematical Expression

$$\begin{cases} 0 & \text{if } [Dna] < 1 \\ \begin{cases} 1 \cdot [Rc] & \text{if } [Dna] > 1 \\ 0.5 \cdot [Rc] & \text{otherwise} \end{cases} & \text{otherwise} \end{cases} \quad (22)$$

6.23 Function definition r_{DePr_1}

Name r_{DePr_1}

Arguments $[C1]$, $[Pr]$, k_{DeCaC1} , k_{DePr}

Mathematical Expression

$$(k_{DePr} + k_{DeCaC1} \cdot [C1]) \cdot [Pr] \quad (23)$$

6.24 Function definition r_{DsRc_2}

Name r_{DsRc_2}

Arguments $[Dna]$, $[pRc]$

Mathematical Expression

$$\begin{cases} 0 & \text{if } [Dna] < 1 \\ \begin{cases} 1 \cdot [pRc] & \text{if } [Dna] > 1 \\ 0.5 \cdot [pRc] & \text{otherwise} \end{cases} & \text{otherwise} \end{cases} \quad (24)$$

7 Rules

This is an overview of seven rules.

7.1 Rule t_{Rb}

Rule t_{Rb} is an assignment rule for species t_{Rb} :

$$t_{Rb} = [Rb] + [pRb] + [RbE2f] \quad (25)$$

Derived unit $\text{mmol} \cdot \text{ml}^{-1}$

7.2 Rule $tE2f$

Rule $tE2f$ is an assignment rule for species $tE2f$:

$$tE2f = [E2f] + [RbE2f] \quad (26)$$

Derived unit $\text{mmol} \cdot \text{ml}^{-1}$

7.3 Rule $tE1$

Rule $tE1$ is an assignment rule for species $tE1$:

$$tE1 = [E1] + [E1C1] \quad (27)$$

Derived unit $\text{mmol} \cdot \text{ml}^{-1}$

7.4 Rule $tC1$

Rule $tC1$ is an assignment rule for species $tC1$:

$$tC1 = [C1] + [pC1] + [E1C1] \quad (28)$$

Derived unit $\text{mmol} \cdot \text{ml}^{-1}$

7.5 Rule tCe

Rule tCe is an assignment rule for species tCe :

$$tCe = [Ce] + [CeP21] \quad (29)$$

Derived unit $\text{mmol} \cdot \text{ml}^{-1}$

7.6 Rule tCa

Rule tCa is an assignment rule for species tCa :

$$tCa = [Ca] + [CaP21] \quad (30)$$

Derived unit $\text{mmol} \cdot \text{ml}^{-1}$

7.7 Rule $tP21$

Rule $tP21$ is an assignment rule for species $tP21$:

$$tP21 = [P21] + [CeP21] + [CaP21] + [iPcna] + [iRc] \quad (31)$$

Derived unit $\text{mmol} \cdot \text{ml}^{-1}$

8 Reactions

This model contains 49 reactions. All reactions are listed in the following table and are subsequently described in detail. If a reaction is affected by a modifier, the identifier of this species is written above the reaction arrow.

Table 5: Overview of all reactions

Nº	Id	Name	Reaction Equation	SBO
1	Phosphorylation- _of_Rb	Phosphorylation of Rb	$\text{Rb} \xrightarrow{\text{Ce, Ca}} \text{pRb}$	
2	Phosphorylation- _Rb_in_Rb_E2F- _complexes	Phosphorylation Rb in Rb:E2F complexes	$\text{RbE2f} \xrightarrow{\text{Ce, Ca}} \text{pRb} + \text{E2f}$	
3	Dephosphorylation _of_Rb	Dephosphorylation of Rb	$\text{pRb} \longrightarrow \text{Rb}$	
4	Synthesis_of- _E2F	Synthesis of E2F	$\emptyset \xrightarrow{\text{E2f}} \text{E2f}$	
5	Degradation_of- _E2F	Degradation of E2F	$\text{E2f} \longrightarrow \emptyset$	
6	Degradation_of- _E2F_in_Rb_E2F- _complexes	Degradation of E2F in Rb:E2F complexes	$\text{RbE2f} \longrightarrow \text{Rb}$	
7	Association- _dissociation- _of_Rb_and_E2F	Association/dissociation of Rb and E2F	$\text{Rb} + \text{E2f} \rightleftharpoons \text{RbE2f}$	
8	Synthesis_of- _p21	Synthesis of p21	$\emptyset \xrightarrow{\text{P53}} \text{P21}$	
9	Synthesis_of- _CycE	Synthesis of CycE	$\text{E2f} \longrightarrow \text{E2f} + \text{Ce}$	

Nº	Id	Name	Reaction Equation	SBO
10	Synthesis_of- _CycA	Synthesis of CycA	$E2f \longrightarrow E2f + Ca$	
11	Association- _dissociation- _of_CycE_Cdk2- _and_p21	Association/dissociation of CycE:Cdk2 and p21	$Ce + P21 \rightleftharpoons CeP21$	
12	Association- _dissociation- _of_CycA_Cdk2- _and_p21	Association/dissociation of CycA:Cdk2 and p21	$Ca + P21 \rightleftharpoons CaP21$	
13	Degradation- _of_p21_in- _CycE_Cdk2_p21- _complexes	Degradation of p21 in CycE:Cdk2:p21 complexes	$CeP21 \xrightarrow{Ce, Ca, aRc} Ce$	
14	Degradation- _of_p21_in- _CycA_Cdk2_p21- _complexes	Degradation of p21 in CycA:Cdk2:p21 complexes	$CaP21 \xrightarrow{Ce, Ca, aRc} Ca$	
15	Degradation_of- _CycE_in_CycE- _Cdk2_complexes	Degradation of CycE in CycE:Cdk2 complexes	$Ce \xrightarrow{Ca} \emptyset$	
16	Degradation- _of_CycE_in- _CycE_Cdk2_p21- _complexes	Degradation of CycE in CycE:Cdk2:p21 complexes	$CeP21 \xrightarrow{Ca} P21$	

Nº	Id	Name	Reaction Equation	SBO
17	Degradation_of- _CycA_in_CycA- _Cdk2_complexes	Degradation of CycA in CycA:Cdk2 complexes	$\text{Ca} \xrightarrow{\text{C1}} \emptyset$	
18	Degradation- _of_CycA_in- _CycA_Cdk2_p21- _complexes	Degradation of CycA in CycA:Cdk2:p21 complexes	$\text{CaP21} \xrightarrow{\text{C1}} \text{P21}$	
19	Degradation_of- _free_p21	Degradation of free p21	$\text{P21} \xrightarrow{\text{Ce, Ca, aRc}} \emptyset$	
20	Synthesis_of- _Emi1	Synthesis of Emi1	$\text{E2f} \longrightarrow \text{E2f} + \text{E1}$	
21	Degradation_of- _Emi1	Degradation of Emi1	$\text{E1} \longrightarrow \emptyset$	
22	Association- _dissociation- _of_Emi1_APC_C- _Cdh1_complexes	Association/dissociation of Emi1:APC/C- Cdh1 complexes	$\text{E1} + \text{C1} \rightleftharpoons \text{E1C1}$	
23	Degradation_of- _Emi1_in_Emi1- _APC_C_Cdh1- _complexes	Degradation of Emi1 in Emi1:APC/C Cdh1 complexes	$\text{E1C1} \longrightarrow \text{C1}$	
24	Phosphorylation- _of_free_APC_C- _Cdh1	Phosphorylation of free APC/C Cdh1	$\text{C1} \xrightarrow{\text{Ce, Ca}} \text{pC1}$	

Nº	Id	Name	Reaction Equation	SBO
25	Phosphorylation- _of_APC_C_Cdh1- _in_Emi1_APC_C- _Cdh1_complexes	Phosphorylation of APC/C ^{Emi1} complexes	$E1C1 \xrightarrow{Ce, Ca} E1 + pC1$	
26	Dephosphorylation _of_APC_C_Cdh1	Dephosphorylation of APC/C ^{Emi1}	$pC1 \longrightarrow C1$	
27	Nuclear_import- _of_active_PCNA	Nuclear import of active PCNA	$\emptyset \longrightarrow aPcna$	
28	Nuclear_export- _of_active_PCNA	Nuclear export of active PCNA	$aPcna \longrightarrow \emptyset$	
29	Nuclear_export- _of_inactive- _PCNA	Nuclear export of inactive PCNA	$iPcna \longrightarrow P21$	
30	Association- _dissociation- _of_PCNA_and_p21	Association/dissociation of PCNA and p21	$aPcna + P21 \rightleftharpoons iPcna$	
31	Degradation_of- _p21_in_PCNA- _p21_complexes	Degradation of p21 in PCNA:p21 complexes	$iPcna \xrightarrow{Ce, Ca, aRc} aPcna$	
32	Association- _dissociation- _of_active- _PCNA_and- _replication- _complexes	Association/dissociation of active PCNA and replication complexes	$aPcna + pRc \rightleftharpoons aRc$	

Nº	Id	Name	Reaction Equation	SBO
33	Association- _dissociation- _of_inactive- _PCNA_and- _replication- _complexes	Association/dissociation of inactive PCNA and replication complexes	$iPcna + pRc \rightleftharpoons iRc$	
34	Phosphorylation- _priming_of- _replication- _complexes	Phosphorylation/priming of replication complexes	$Rc \xrightarrow{Ce, Ca} pRc$	
35	Dephosphorylation- _of- _replication- _complexes	Dephosphorylation of replication complexes	$pRc \longrightarrow Rc$	
36	Association- _dissociation- _of_p21_and- _replication- _complexes	Association/dissociation of p21 and replication complexes	$aRc + P21 \rightleftharpoons iRc$	
37	Degradation- _of_p21_in- _inactive- _replication- _complexes	Degradation of p21 in inactive replication complexes	$iRc \xrightarrow{Ce, Ca, aRc} aRc$	
38	Dissassembly- _of_RC	Dissassembly of RC	$Rc \xrightarrow{Dna} \emptyset$	

Nº	Id	Name	Reaction Equation	SBO
39	Dissasembly- _of_pRC	Dissasembly of pRC	$\text{pRc} \xrightarrow{\text{Dna}} \emptyset$	
40	Dissasembly- _of_aRC	Dissasembly of aRC	$\text{aRc} \xrightarrow{\text{Dna}} \text{aPcna}$	
41	Dissasembly- _of_iRC	Dissasembly of iRC	$\text{iRc} \xrightarrow{\text{Dna}} \text{iPcna}$	
42	Synthesis_of- _DNA	Synthesis of DNA	$\text{aRc} \longrightarrow \text{aRc} + \text{Dna}$	
43	Synthesis_of- _p53	Synthesis of p53	$\emptyset \longrightarrow \text{P53}$	
44	Degradation_of- _p53	Degradation of p53	$\text{P53} \xrightarrow{\text{Dam}} \emptyset$	
45	Induction_of- _DNA_damage	Induction of DNA damage	$\emptyset \longrightarrow \text{Dam}$	
46	Induction_of- _DNA_damage_by- _replication	Induction of DNA damage by replication	$\text{aRc} \longrightarrow \text{aRc} + \text{Dam}$	
47	Synthesis_of- _APC_C_Cdh1- _activity_probe	Synthesis of APC/C ^{dh1} activity probe	$\emptyset \longrightarrow \text{Pr}$	
48	Degradation- _of_APC_C_Cdh1- _activity_probe	Degradation of APC/C ^{dh1} activity probe	$\text{Pr} \xrightarrow{\text{C1}} \emptyset$	
49	Repair_of_DNA- _damage	Repair of DNA damage	$\text{Dam} \xrightarrow{\text{P53}} \emptyset$	

8.1 Reaction Phosphorylation_of_Rb

This is an irreversible reaction of one reactant forming one product influenced by two modifiers.

Name Phosphorylation of Rb

Reaction equation



Reactant

Table 6: Properties of each reactant.

Id	Name	SBO
Rb	Retinoblastoma_protein	

Modifiers

Table 7: Properties of each modifier.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	
Ca	CyclinA_Cdk2	

Product

Table 8: Properties of each product.

Id	Name	SBO
pRb	Retinoblastoma_protein_hyperphosphorylated	

Kinetic Law

Derived unit contains undeclared units

$$v_1 = \text{vol}(\text{Cell}) \cdot \text{rPhRb_1}([\text{Ca}], \text{Cd}, [\text{Ce}], [\text{Rb}], \text{kPhRbCa}, \text{kPhRbCd}, \text{kPhRbCe}) \quad (33)$$

$$\begin{aligned} &\text{rPhRb_1}([\text{Ca}], \text{Cd}, [\text{Ce}], [\text{Rb}], \text{kPhRbCa}, \text{kPhRbCd}, \text{kPhRbCe}) \\ &= (\text{kPhRbCd} \cdot \text{Cd} + \text{kPhRbCe} \cdot [\text{Ce}] + \text{kPhRbCa} \cdot [\text{Ca}]) \cdot [\text{Rb}] \end{aligned} \quad (34)$$

$$\begin{aligned} & rPhRb_1 ([Ca], Cd, [Ce], [Rb], kPhRbCa, kPhRbCd, kPhRbCe) \\ & = (kPhRbCd \cdot Cd + kPhRbCe \cdot [Ce] + kPhRbCa \cdot [Ca]) \cdot [Rb] \end{aligned} \quad (35)$$

8.2 Reaction Phosphorylation_Rb_in_Rb_E2F_complexes

This is an irreversible reaction of one reactant forming two products influenced by two modifiers.

Name Phosphorylation Rb in Rb:E2F complexes

Reaction equation



Reactant

Table 9: Properties of each reactant.

Id	Name	SBO
RbE2f	Retinoblastoma_protein_E2f_complex_inactive	

Modifiers

Table 10: Properties of each modifier.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	
Ca	CyclinA_Cdk2	

Products

Table 11: Properties of each product.

Id	Name	SBO
pRb	Retinoblastoma_protein_hyperphosphorylated	
E2f	E2f_active	

Kinetic Law

Derived unit contains undeclared units

$$v_2 = \text{vol}(\text{Cell}) \cdot \text{rPhRb_2}([\text{Ca}], \text{Cd}, [\text{Ce}], [\text{RbE2f}], \text{kPhRbCa}, \text{kPhRbCd}, \text{kPhRbCe}) \quad (37)$$

$$\begin{aligned} & \text{rPhRb_2}([\text{Ca}], \text{Cd}, [\text{Ce}], [\text{RbE2f}], \text{kPhRbCa}, \text{kPhRbCd}, \text{kPhRbCe}) \\ &= (\text{kPhRbCd} \cdot \text{Cd} + \text{kPhRbCe} \cdot [\text{Ce}] + \text{kPhRbCa} \cdot [\text{Ca}]) \cdot [\text{RbE2f}] \end{aligned} \quad (38)$$

$$\begin{aligned} & \text{rPhRb_2}([\text{Ca}], \text{Cd}, [\text{Ce}], [\text{RbE2f}], \text{kPhRbCa}, \text{kPhRbCd}, \text{kPhRbCe}) \\ &= (\text{kPhRbCd} \cdot \text{Cd} + \text{kPhRbCe} \cdot [\text{Ce}] + \text{kPhRbCa} \cdot [\text{Ca}]) \cdot [\text{RbE2f}] \end{aligned} \quad (39)$$

8.3 Reaction Dephosphorylation_of_Rb

This is an irreversible reaction of one reactant forming one product.

Name Dephosphorylation of Rb

Reaction equation



Reactant

Table 12: Properties of each reactant.

Id	Name	SBO
pRb	Retinoblastoma_protein_hyperphosphorylated	

Product

Table 13: Properties of each product.

Id	Name	SBO
Rb	Retinoblastoma_protein	

Kinetic Law

Derived unit contains undeclared units

$$v_3 = \text{vol}(\text{Cell}) \cdot k_{\text{DpRb}} \cdot [\text{pRb}] \quad (41)$$

8.4 Reaction *Synthesis_of_E2F*

This is an irreversible reaction of no reactant forming one product influenced by one modifier.

Name Synthesis of E2F

Reaction equation



Modifier

Table 14: Properties of each modifier.

Id	Name	SBO
E2f	E2f_active	

Product

Table 15: Properties of each product.

Id	Name	SBO
E2f	E2f_active	

Kinetic Law

Derived unit contains undeclared units

$$v_4 = \text{vol}(\text{Cell}) \cdot r_{\text{SyE2f}_1}([\text{E2f}], j_{\text{SyE2f}}, k_{\text{SyE2f}}, k_{\text{SyE2fE2f}}) \quad (43)$$

$$r_{\text{SyE2f}_1}([\text{E2f}], j_{\text{SyE2f}}, k_{\text{SyE2f}}, k_{\text{SyE2fE2f}}) = k_{\text{SyE2f}} + \frac{k_{\text{SyE2fE2f}} \cdot [\text{E2f}]}{j_{\text{SyE2f}} + [\text{E2f}]} \quad (44)$$

$$r_{\text{SyE2f}_1}([\text{E2f}], j_{\text{SyE2f}}, k_{\text{SyE2f}}, k_{\text{SyE2fE2f}}) = k_{\text{SyE2f}} + \frac{k_{\text{SyE2fE2f}} \cdot [\text{E2f}]}{j_{\text{SyE2f}} + [\text{E2f}]} \quad (45)$$

8.5 Reaction *Degradation_of_E2F*

This is an irreversible reaction of one reactant forming no product.

Name Degradation of E2F

Reaction equation



Reactant

Table 16: Properties of each reactant.

Id	Name	SBO
E2f	E2f_active	

Kinetic Law

Derived unit contains undeclared units

$$v_5 = \text{vol}(\text{Cell}) \cdot k_{\text{DeE2f}} \cdot [\text{E2f}] \quad (47)$$

8.6 Reaction Degradation_of_E2F_in_Rb_E2F_complexes

This is an irreversible reaction of one reactant forming one product.

Name Degradation of E2F in Rb:E2F complexes

Reaction equation



Reactant

Table 17: Properties of each reactant.

Id	Name	SBO
RbE2f	Retinoblastoma_protein_E2f_complex_inactive	

Product

Table 18: Properties of each product.

Id	Name	SBO
Rb	Retinoblastoma_protein	

Kinetic Law

Derived unit contains undeclared units

$$v_6 = \text{vol}(\text{Cell}) \cdot k_{\text{DeE2f}} \cdot [\text{RbE2f}] \quad (49)$$

8.7 Reaction Association_dissociation_of_Rb_and_E2F

This is a reversible reaction of two reactants forming one product.

Name Association/dissociation of Rb and E2F

Reaction equation



Reactants

Table 19: Properties of each reactant.

Id	Name	SBO
Rb	Retinoblastoma_protein	
E2f	E2f_active	

Product

Table 20: Properties of each product.

Id	Name	SBO
RbE2f	Retinoblastoma_protein_E2f_complex_inactive	

Kinetic Law

Derived unit contains undeclared units

$$v_7 = \text{vol}(\text{Cell}) \cdot (k_{\text{AsRbE2f}} \cdot [\text{Rb}] \cdot [\text{E2f}] - k_{\text{DsRbE2f}} \cdot [\text{RbE2f}]) \quad (51)$$

8.8 Reaction Synthesis_of_p21

This is an irreversible reaction of no reactant forming one product influenced by one modifier.

Name Synthesis of p21

Reaction equation



Modifier

Table 21: Properties of each modifier.

Id	Name	SBO
P53	P53	

Product

Table 22: Properties of each product.

Id	Name	SBO
P21	P21	

Kinetic Law

Derived unit contains undeclared units

$$v_8 = \text{vol}(\text{Cell}) \cdot \text{rSyP21_1}([P53], k\text{SyP21}, k\text{SyP21P53}) \quad (53)$$

$$\text{rSyP21_1}([P53], k\text{SyP21}, k\text{SyP21P53}) = k\text{SyP21} + k\text{SyP21P53} \cdot [P53] \quad (54)$$

$$\text{rSyP21_1}([P53], k\text{SyP21}, k\text{SyP21P53}) = k\text{SyP21} + k\text{SyP21P53} \cdot [P53] \quad (55)$$

8.9 Reaction *Synthesis_of_CycE*

This is an irreversible reaction of one reactant forming two products.

Name Synthesis of CycE

Reaction equation



Reactant

Table 23: Properties of each reactant.

Id	Name	SBO
E2f	E2f_active	

Products

Table 24: Properties of each product.

Id	Name	SBO
E2f	E2f_active	
Ce	CyclinE_Cdk2_active	

Kinetic Law

Derived unit contains undeclared units

$$v_9 = \text{vol}(\text{Cell}) \cdot k_{\text{SyCe}} \cdot [\text{E2f}] \quad (57)$$

8.10 Reaction *Synthesis_of_CycA*

This is an irreversible reaction of one reactant forming two products.

Name Synthesis of CycA

Reaction equation



Reactant

Table 25: Properties of each reactant.

Id	Name	SBO
E2f	E2f_active	

Products

Table 26: Properties of each product.

Id	Name	SBO
E2f	E2f_active	
Ca	CyclinA_Cdk2	

Kinetic Law

Derived unit contains undeclared units

$$v_{10} = \text{vol}(\text{Cell}) \cdot k_{\text{SyCa}} \cdot [\text{E2f}] \quad (59)$$

8.11 Reaction Association_dissociation_of_CycE_Cdk2_and_p21

This is a reversible reaction of two reactants forming one product.

Name Association/dissociation of CycE:Cdk2 and p21

Reaction equation



Reactants

Table 27: Properties of each reactant.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	
P21	P21	

Product

Table 28: Properties of each product.

Id	Name	SBO
CeP21	CyclinE_Cdk2_P21_complex_inactive	

Kinetic Law

Derived unit contains undeclared units


$$v_{11} = \text{vol}(\text{Cell}) \cdot (k_{\text{AsCyP21}} \cdot [\text{Ce}] \cdot [\text{P21}] - k_{\text{DsCyP21}} \cdot [\text{CeP21}]) \quad (61)$$

8.12 Reaction Association_dissociation_of_CycA_Cdk2_and_p21

This is a reversible reaction of two reactants forming one product.

Name Association/dissociation of CycA:Cdk2 and p21

Reaction equation



Reactants

Table 29: Properties of each reactant.

Id	Name	SBO
Ca	CyclinA_Cdk2	
P21	P21	

Product

Table 30: Properties of each product.

Id	Name	SBO
CaP21	CyclinA_Cdk2_P21_complex_inactive	

Kinetic Law

Derived unit contains undeclared units

$$v_{12} = \text{vol}(\text{Cell}) \cdot (k_{\text{AsCyP21}} \cdot [\text{Ca}] \cdot [\text{P21}] - k_{\text{DsCyP21}} \cdot [\text{CaP21}]) \quad (63)$$

8.13 Reaction Degradation_of_p21_in_CycE_Cdk2_p21_complexes

This is an irreversible reaction of one reactant forming one product influenced by three modifiers.

Name Degradation of p21 in CycE:Cdk2:p21 complexes

Reaction equation



Reactant

Table 31: Properties of each reactant.

Id	Name	SBO
CeP21	CyclinE_Cdk2_P21_complex_inactive	

Modifiers

Table 32: Properties of each modifier.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	
Ca	CyclinA_Cdk2	
aRc	Pre_Replication_complex_active	

Product

Table 33: Properties of each product.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	

Kinetic Law

Derived unit contains undeclared units

$$v_{13} = \text{vol}(\text{Cell}) \cdot r_{\text{DeP21_1}}([Ca], Cdt2, [Ce], [CeP21], Skp2, [aRc], k_{\text{DeP21}}, k_{\text{DeP21Cy}}, k_{\text{DeP21aRc}}) \quad (65)$$

$$\begin{aligned} & r_{\text{DeP21_1}}([Ca], Cdt2, [Ce], [CeP21], Skp2, [aRc], k_{\text{DeP21}}, k_{\text{DeP21Cy}}, k_{\text{DeP21aRc}}) \\ &= (k_{\text{DeP21}} + k_{\text{DeP21Cy}} \cdot Skp2 \cdot ([Ce] + [Ca]) + k_{\text{DeP21aRc}} \cdot Cdt2 \cdot [aRc]) \cdot [CeP21] \end{aligned} \quad (66)$$

$$\begin{aligned} & r_{\text{DeP21_1}}([Ca], Cdt2, [Ce], [CeP21], Skp2, [aRc], k_{\text{DeP21}}, k_{\text{DeP21Cy}}, k_{\text{DeP21aRc}}) \\ &= (k_{\text{DeP21}} + k_{\text{DeP21Cy}} \cdot Skp2 \cdot ([Ce] + [Ca]) + k_{\text{DeP21aRc}} \cdot Cdt2 \cdot [aRc]) \cdot [CeP21] \end{aligned} \quad (67)$$

8.14 Reaction Degradation_of_p21_in_CycA_Cdk2_p21_complexes

This is an irreversible reaction of one reactant forming one product influenced by three modifiers.

Name Degradation of p21 in CycA:Cdk2:p21 complexes

Reaction equation



Reactant

Table 34: Properties of each reactant.

Id	Name	SBO
CaP21	CyclinA_Cdk2_P21_complex_inactive	

Modifiers

Table 35: Properties of each modifier.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	
Ca	CyclinA_Cdk2	
aRc	Pre_Replication_complex_active	

Product

Table 36: Properties of each product.

Id	Name	SBO
Ca	CyclinA_Cdk2	

Kinetic Law

Derived unit contains undeclared units

$$v_{14} = \text{vol}(\text{Cell}) \cdot \text{rDeP21_2}([\text{Ca}], [\text{CaP21}], \text{Cdt2}, [\text{Ce}], \text{Skp2}, [\text{aRc}], \text{kDeP21}, \text{kDeP21Cy}, \text{kDeP21aRc}) \quad (69)$$

$$\begin{aligned} &\text{rDeP21_2}([\text{Ca}], [\text{CaP21}], \text{Cdt2}, [\text{Ce}], \text{Skp2}, [\text{aRc}], \text{kDeP21}, \text{kDeP21Cy}, \text{kDeP21aRc}) \\ &= (\text{kDeP21} + \text{kDeP21Cy} \cdot \text{Skp2} \cdot ([\text{Ce}] + [\text{Ca}] + \text{kDeP21aRc} \cdot \text{Cdt2} \cdot [\text{aRc}]) \cdot [\text{CaP21}] \end{aligned} \quad (70)$$

$$\begin{aligned} &\text{rDeP21_2}([\text{Ca}], [\text{CaP21}], \text{Cdt2}, [\text{Ce}], \text{Skp2}, [\text{aRc}], \text{kDeP21}, \text{kDeP21Cy}, \text{kDeP21aRc}) \\ &= (\text{kDeP21} + \text{kDeP21Cy} \cdot \text{Skp2} \cdot ([\text{Ce}] + [\text{Ca}] + \text{kDeP21aRc} \cdot \text{Cdt2} \cdot [\text{aRc}]) \cdot [\text{CaP21}] \end{aligned} \quad (71)$$

8.15 Reaction Degradation_of_CycE_in_CycE_Cdk2_complexes

This is an irreversible reaction of one reactant forming no product influenced by one modifier.

Name Degradation of CycE in CycE:Cdk2 complexes

Reaction equation



Reactant

Table 37: Properties of each reactant.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	

Modifier

Table 38: Properties of each modifier.

Id	Name	SBO
Ca	CyclinA_Cdk2	

Kinetic Law

Derived unit contains undeclared units

$$v_{15} = \text{vol}(\text{Cell}) \cdot \text{rDeCe_1}([\text{Ca}], [\text{Ce}], \text{kDeCe}, \text{kDeCeCa}) \quad (73)$$

$$\text{rDeCe_1}([\text{Ca}], [\text{Ce}], \text{kDeCe}, \text{kDeCeCa}) = (\text{kDeCe} + \text{kDeCeCa} \cdot [\text{Ca}]) \cdot [\text{Ce}] \quad (74)$$

$$\text{rDeCe_1}([\text{Ca}], [\text{Ce}], \text{kDeCe}, \text{kDeCeCa}) = (\text{kDeCe} + \text{kDeCeCa} \cdot [\text{Ca}]) \cdot [\text{Ce}] \quad (75)$$

8.16 Reaction Degradation_of_CycE_in_CycE_Cdk2_p21_complexes

This is an irreversible reaction of one reactant forming one product influenced by one modifier.

Name Degradation of CycE in CycE:Cdk2:p21 complexes

Reaction equation



Reactant

Table 39: Properties of each reactant.

Id	Name	SBO
CeP21	CyclinE_Cdk2_P21_complex_inactive	

Modifier

Table 40: Properties of each modifier.

Id	Name	SBO
Ca	CyclinA_Cdk2	

Product

Table 41: Properties of each product.

Id	Name	SBO
P21	P21	

Kinetic Law

Derived unit contains undeclared units

$$v_{16} = \text{vol}(\text{Cell}) \cdot \text{rDeCe_2}([\text{Ca}], [\text{CeP21}], k\text{DeCe}, k\text{DeCeCa}) \quad (77)$$

$$\text{rDeCe_2}([\text{Ca}], [\text{CeP21}], k\text{DeCe}, k\text{DeCeCa}) = (k\text{DeCe} + k\text{DeCeCa} \cdot [\text{Ca}]) \cdot [\text{CeP21}] \quad (78)$$

$$\text{rDeCe_2}([\text{Ca}], [\text{CeP21}], k\text{DeCe}, k\text{DeCeCa}) = (k\text{DeCe} + k\text{DeCeCa} \cdot [\text{Ca}]) \cdot [\text{CeP21}] \quad (79)$$

8.17 Reaction Degradation_of_CycA_in_CycA_Cdk2_complexes

This is an irreversible reaction of one reactant forming no product influenced by one modifier.

Name Degradation of CycA in CycA:Cdk2 complexes

Reaction equation



Reactant

Table 42: Properties of each reactant.

Id	Name	SBO
Ca	CyclinA_Cdk2	

Modifier

Table 43: Properties of each modifier.

Id	Name	SBO
C1	Cdh1_C-APC_active	

Kinetic Law

Derived unit contains undeclared units

$$v_{17} = \text{vol}(\text{Cell}) \cdot \text{rDeCa_1}([C1], [Ca], kDeCa, kDeCaC1) \quad (81)$$

$$\text{rDeCa_1}([C1], [Ca], kDeCa, kDeCaC1) = (kDeCa + kDeCaC1 \cdot [C1]) \cdot [Ca] \quad (82)$$

$$\text{rDeCa_1}([C1], [Ca], kDeCa, kDeCaC1) = (kDeCa + kDeCaC1 \cdot [C1]) \cdot [Ca] \quad (83)$$

8.18 Reaction Degradation of CycA in CycA-Cdk2-p21 complexes

This is an irreversible reaction of one reactant forming one product influenced by one modifier.

Name Degradation of CycA in CycA:Cdk2:p21 complexes

Reaction equation



Reactant

Table 44: Properties of each reactant.

Id	Name	SBO
CaP21	CyclinA_Cdk2_P21_complex_inactive	

Modifier

Table 45: Properties of each modifier.

Id	Name	SBO
C1	Cdh1_C_APC_active	

Product

Table 46: Properties of each product.

Id	Name	SBO
P21	P21	

Kinetic Law

Derived unit contains undeclared units

$$v_{18} = \text{vol}(\text{Cell}) \cdot \text{rDeCa_2}([C1], [CaP21], kDeCa, kDeCaC1) \quad (85)$$

$$\text{rDeCa_2}([C1], [CaP21], kDeCa, kDeCaC1) = (kDeCa + kDeCaC1 \cdot [C1]) \cdot [CaP21] \quad (86)$$

$$\text{rDeCa_2}([C1], [CaP21], kDeCa, kDeCaC1) = (kDeCa + kDeCaC1 \cdot [C1]) \cdot [CaP21] \quad (87)$$

8.19 Reaction Degradation_of_free_p21

This is an irreversible reaction of one reactant forming no product influenced by three modifiers.

Name Degradation of free p21

Reaction equation



Reactant

Table 47: Properties of each reactant.

Id	Name	SBO
P21	P21	

Modifiers

Table 48: Properties of each modifier.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	
Ca	CyclinA_Cdk2	
aRc	Pre_Replication_complex_active	

Kinetic Law

Derived unit contains undeclared units

$$v_{19} = \text{vol}(\text{Cell}) \cdot r_{\text{DeP21_3}}([\text{Ca}], \text{Cdt2}, [\text{Ce}], [\text{P21}], \text{Sklp2}, [\text{aRc}], k_{\text{DeP21}}, k_{\text{DeP21Cy}}, k_{\text{DeP21aRc}}) \quad (89)$$

$$\begin{aligned} & r_{\text{DeP21_3}}([\text{Ca}], \text{Cdt2}, [\text{Ce}], [\text{P21}], \text{Sklp2}, [\text{aRc}], k_{\text{DeP21}}, k_{\text{DeP21Cy}}, k_{\text{DeP21aRc}}) \\ &= (k_{\text{DeP21}} + k_{\text{DeP21Cy}} \cdot \text{Sklp2} \cdot ([\text{Ce}] + [\text{Ca}]) + k_{\text{DeP21aRc}} \cdot \text{Cdt2} \cdot [\text{aRc}]) \cdot [\text{P21}] \end{aligned} \quad (90)$$

$$\begin{aligned} & r_{\text{DeP21_3}}([\text{Ca}], \text{Cdt2}, [\text{Ce}], [\text{P21}], \text{Sklp2}, [\text{aRc}], k_{\text{DeP21}}, k_{\text{DeP21Cy}}, k_{\text{DeP21aRc}}) \\ &= (k_{\text{DeP21}} + k_{\text{DeP21Cy}} \cdot \text{Sklp2} \cdot ([\text{Ce}] + [\text{Ca}]) + k_{\text{DeP21aRc}} \cdot \text{Cdt2} \cdot [\text{aRc}]) \cdot [\text{P21}] \end{aligned} \quad (91)$$

8.20 Reaction Synthesis_of_Emi1

This is an irreversible reaction of one reactant forming two products.

Name Synthesis of Emi1

Reaction equation



Reactant

Table 49: Properties of each reactant.

Id	Name	SBO
E2f	E2f_active	

Products

Table 50: Properties of each product.

Id	Name	SBO
E2f	E2f_active	
E1	Emi1	

Kinetic Law

Derived unit contains undeclared units

$$v_{20} = \text{vol}(\text{Cell}) \cdot k_{\text{SyE1}} \cdot [\text{E2f}] \quad (93)$$

8.21 Reaction Degradation_of_Emi1

This is an irreversible reaction of one reactant forming no product.

Name Degradation of Emi1

Reaction equation



Reactant

Table 51: Properties of each reactant.

Id	Name	SBO
E1	Emi1	

Kinetic Law

Derived unit contains undeclared units

$$v_{21} = \text{vol}(\text{Cell}) \cdot k_{\text{DeE1}} \cdot [\text{E1}] \quad (95)$$

8.22 Reaction Association/dissociation of Emi1-APC-Cdh1 complexes

This is a reversible reaction of two reactants forming one product.

Name Association/dissociation of Emi1:APC/Cdh1 complexes

Reaction equation



Reactants

Table 52: Properties of each reactant.

Id	Name	SBO
E1	Emi1	
C1	Cdh1-APC-active	

Product

Table 53: Properties of each product.

Id	Name	SBO
E1C1	Emi1-APC-Cdh1-complex-inactive	

Kinetic Law

Derived unit contains undeclared units

$$v_{22} = \text{vol}(\text{Cell}) \cdot (k_{AsE1C1} \cdot [E1] \cdot [C1] - k_{DsE1C1} \cdot [E1C1]) \quad (97)$$

8.23 Reaction Degradation of Emi1 in Emi1-APC-Cdh1 complexes

This is an irreversible reaction of one reactant forming one product.

Name Degradation of Emi1 in Emi1:APC/Cdh1 complexes

Reaction equation



Reactant

Table 54: Properties of each reactant.

Id	Name	SBO
E1C1	Emi1_C_Cdh1_complex_inactive	

Product

Table 55: Properties of each product.

Id	Name	SBO
C1	Cdh1_C_APC_active	

Kinetic Law

Derived unit contains undeclared units

$$v_{23} = \text{vol}(\text{Cell}) \cdot k_{\text{DeE1C1}} \cdot [\text{E1C1}] \quad (99)$$

8.24 Reaction Phosphorylation_of_free_APC_C_Cdh1

This is an irreversible reaction of one reactant forming one product influenced by two modifiers.

Name Phosphorylation of free APC/C $\hat{\text{Cdh1}}$

Reaction equation



Reactant

Table 56: Properties of each reactant.

Id	Name	SBO
C1	Cdh1_C_APC_active	

Modifiers

Table 57: Properties of each modifier.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	
Ca	CyclinA_Cdk2	

Product

Table 58: Properties of each product.

Id	Name	SBO
pC1	C_Cdh1_APC_phosphorylated_inactive	

Kinetic Law

Derived unit contains undeclared units

$$v_{24} = \text{vol}(\text{Cell}) \cdot \text{rPhC1_1}([C1], [Ca], [Ce], kPhC1, kPhC1Ca, kPhC1Ce) \quad (101)$$

$$\begin{aligned} &\text{rPhC1_1}([C1], [Ca], [Ce], kPhC1, kPhC1Ca, kPhC1Ce) \\ &= (kPhC1 + kPhC1Ce \cdot [Ce] + kPhC1Ca \cdot [Ca]) \cdot [C1] \end{aligned} \quad (102)$$

$$\begin{aligned} &\text{rPhC1_1}([C1], [Ca], [Ce], kPhC1, kPhC1Ca, kPhC1Ce) \\ &= (kPhC1 + kPhC1Ce \cdot [Ce] + kPhC1Ca \cdot [Ca]) \cdot [C1] \end{aligned} \quad (103)$$


8.25 Reaction

Phosphorylation of APC/C_Cdh1 in Emi1-APC/C_Cdh1 complexes

This is an irreversible reaction of one reactant forming two products influenced by two modifiers.

Name Phosphorylation of APC/C_Cdh1 in Emi1-APC/C_Cdh1 complexes

Reaction equation



Reactant

Table 59: Properties of each reactant.

Id	Name	SBO
E1C1	Emi1_C_Cdh1_complex_inactive	

Modifiers

Table 60: Properties of each modifier.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	
Ca	CyclinA_Cdk2	

Products

Table 61: Properties of each product.

Id	Name	SBO
E1	Emi1	
pC1	C_Cdh1_APC_phosphorylated_inactive	

Kinetic Law

Derived unit contains undeclared units

$$v_{25} = \text{vol}(\text{Cell}) \cdot \text{rPhC1_2}([\text{Ca}], [\text{Ce}], [\text{E1C1}], \text{kPhC1}, \text{kPhC1Ca}, \text{kPhC1Ce}) \quad (105)$$

$$\begin{aligned} &\text{rPhC1_2}([\text{Ca}], [\text{Ce}], [\text{E1C1}], \text{kPhC1}, \text{kPhC1Ca}, \text{kPhC1Ce}) \\ &= (\text{kPhC1} + \text{kPhC1Ce} \cdot [\text{Ce}] + \text{kPhC1Ca} \cdot [\text{Ca}]) \cdot [\text{E1C1}] \end{aligned} \quad (106)$$

$$\begin{aligned} &\text{rPhC1_2}([\text{Ca}], [\text{Ce}], [\text{E1C1}], \text{kPhC1}, \text{kPhC1Ca}, \text{kPhC1Ce}) \\ &= (\text{kPhC1} + \text{kPhC1Ce} \cdot [\text{Ce}] + \text{kPhC1Ca} \cdot [\text{Ca}]) \cdot [\text{E1C1}] \end{aligned} \quad (107)$$

8.26 Reaction Dephosphorylation_of_APC_C_Cdh1

This is an irreversible reaction of one reactant forming one product.

Name Dephosphorylation of APC/Cdh1

Reaction equation



Reactant

Table 62: Properties of each reactant.

Id	Name	SBO
pC1	C_Cdh1_APC_phosphorylated_inactive	

Product

Table 63: Properties of each product.

Id	Name	SBO
C1	Cdh1_C_APC_active	

Kinetic Law

Derived unit contains undeclared units

$$v_{26} = \text{vol}(\text{Cell}) \cdot k_{\text{DpC1}} \cdot [\text{pC1}] \quad (109)$$

8.27 Reaction `Nuclear_import_of_active_PCNA`

This is an irreversible reaction of no reactant forming one product.

Name Nuclear import of active PCNA

Reaction equation



Product

Table 64: Properties of each product.

Id	Name	SBO
aPcna	Pcna_nuclear_active	

Kinetic Law

Derived unit contains undeclared units

$$v_{27} = \text{vol}(\text{Cell}) \cdot \text{Constant_flux_irreversible}(\text{kImPc}) \quad (111)$$

$$\text{Constant_flux_irreversible}(v) = v \quad (112)$$

$$\text{Constant_flux_irreversible}(v) = v \quad (113)$$

8.28 Reaction `Nuclear_export_of_active_PCNA`

This is an irreversible reaction of one reactant forming no product.

Name Nuclear export of active PCNA

Reaction equation



Reactant

Table 65: Properties of each reactant.

Id	Name	SBO
aPcna	Pcna_nuclear_active	

Kinetic Law

Derived unit contains undeclared units

$$v_{28} = \text{vol}(\text{Cell}) \cdot \text{kExPc} \cdot [\text{aPcna}] \quad (115)$$

8.29 Reaction `Nuclear_export_of_inactive_PCNA`

This is an irreversible reaction of one reactant forming one product.

Name Nuclear export of inactive PCNA

Reaction equation



Reactant

Table 66: Properties of each reactant.

Id	Name	SBO
iPcna	PCNA_Nuclear_inactive	

Product

Table 67: Properties of each product.

Id	Name	SBO
P21	P21	

Kinetic Law

Derived unit contains undeclared units


$$v_{29} = \text{vol}(\text{Cell}) \cdot k_{\text{ExpC}} \cdot [\text{iPcna}] \quad (117)$$

8.30 Reaction Association_dissociation_of_PCNA_and_p21

This is a reversible reaction of two reactants forming one product.

Name Association/dissociation of PCNA and p21

Reaction equation



Reactants

Table 68: Properties of each reactant.

Id	Name	SBO
aPcna	Pcna_nuclear_active	
P21	P21	

Product

Table 69: Properties of each product.

Id	Name	SBO
iPcna	PCNA_Nuclear_inactive	

Kinetic Law

Derived unit contains undeclared units

$$v_{30} = \text{vol}(\text{Cell}) \cdot (k_{\text{AsPcP21}} \cdot [\text{aPcna}] \cdot [\text{P21}] - k_{\text{DsPcP21}} \cdot [\text{iPcna}]) \quad (119)$$

8.31 Reaction Degradation_of_p21_in_PCNA_p21_complexes

This is an irreversible reaction of one reactant forming one product influenced by three modifiers.

Name Degradation of p21 in PCNA:p21 complexes

Reaction equation



Reactant

Table 70: Properties of each reactant.

Id	Name	SBO
iPcna	PCNA_Nuclear_inactive	

Modifiers

Table 71: Properties of each modifier.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	
Ca	CyclinA_Cdk2	
aRc	Pre_Replication_complex_active	

Product

Table 72: Properties of each product.

Id	Name	SBO
aPcna	Pcna_nuclear_active	

Kinetic Law

Derived unit contains undeclared units

$$v_{31} = \text{vol}(\text{Cell}) \cdot \text{rDeP21_4}([\text{Ca}], \text{Cdt2}, [\text{Ce}], \text{Skp2}, [\text{aRc}], [\text{iPcna}], \text{kDeP21}, \text{kDeP21Cy}, \text{kDeP21aRc})$$

$$\begin{aligned} & \text{rDeP21_4}([\text{Ca}], \text{Cdt2}, [\text{Ce}], \text{Skp2}, [\text{aRc}], [\text{iPcna}], \text{kDeP21}, \text{kDeP21Cy}, \text{kDeP21aRc}) \\ &= (\text{kDeP21} + \text{kDeP21Cy} \cdot \text{Skp2} \cdot ([\text{Ce}] + [\text{Ca}]) + \text{kDeP21aRc} \cdot \text{Cdt2} \cdot [\text{aRc}]) \cdot [\text{iPcna}] \end{aligned} \quad (122)$$

$$\begin{aligned} & \text{rDeP21_4}([\text{Ca}], \text{Cdt2}, [\text{Ce}], \text{Skp2}, [\text{aRc}], [\text{iPcna}], \text{kDeP21}, \text{kDeP21Cy}, \text{kDeP21aRc}) \\ &= (\text{kDeP21} + \text{kDeP21Cy} \cdot \text{Skp2} \cdot ([\text{Ce}] + [\text{Ca}]) + \text{kDeP21aRc} \cdot \text{Cdt2} \cdot [\text{aRc}]) \cdot [\text{iPcna}] \end{aligned} \quad (123)$$

8.32 Reaction

[Association/dissociation of active PCNA and replication complexes](#)

This is a reversible reaction of two reactants forming one product.

Name Association/dissociation of active PCNA and replication complexes

Reaction equation



Reactants

Table 73: Properties of each reactant.

Id	Name	SBO
aPcna	Pcna_nuclear_active	
pRc	Pre_Replication_complex_primed	

Product

Table 74: Properties of each product.

Id	Name	SBO
aRc	Pre_Replication_complex_active	

Kinetic Law

Derived unit contains undeclared units

$$v_{32} = \text{vol}(\text{Cell}) \cdot (k_{\text{AsRcPc}} \cdot [\text{aPcna}] \cdot [\text{pRc}] - k_{\text{DsRcPc}} \cdot [\text{aRc}]) \quad (125)$$

8.33 Reaction Association_dissociation_of_inactive_PCNA_and-replication_complexes

This is a reversible reaction of two reactants forming one product.

Name Association/dissociation of inactive PCNA and replication complexes

Reaction equation



Reactants

Table 75: Properties of each reactant.

Id	Name	SBO
iPcna	PCNA_Nuclear_inactive	
pRc	Pre_Replication_complex_primed	

Product

Table 76: Properties of each product.

Id	Name	SBO
iRc	Pre_Replication_complex_inactive	

Kinetic Law

Derived unit contains undeclared units


$$v_{33} = \text{vol}(\text{Cell}) \cdot (k_{\text{AsRcPc}} \cdot [\text{iPcna}] \cdot [\text{pRc}] - k_{\text{DsRcPc}} \cdot [\text{iRc}]) \quad (127)$$

8.34 Reaction Phosphorylation priming of replication complexes

This is an irreversible reaction of one reactant forming one product influenced by two modifiers.

Name Phosphorylation/priming of replication complexes

Reaction equation



Reactant

Table 77: Properties of each reactant.

Id	Name	SBO
Rc	Pre_Replication_complex	

Modifiers

Table 78: Properties of each modifier.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	
Ca	CyclinA_Cdk2	

Product

Table 79: Properties of each product.

Id	Name	SBO
pRc	Pre_Replication_complex_primed	

Kinetic Law

Derived unit contains undeclared units

$$v_{34} = \text{vol}(\text{Cell}) \cdot r_{\text{PhRc}_1}([\text{Ca}], [\text{Ce}], [\text{Rc}], j_{\text{Cy}}, k_{\text{PhRc}}, n) \quad (129)$$

$$r_{\text{PhRc}_1}([\text{Ca}], [\text{Ce}], [\text{Rc}], j_{\text{Cy}}, k_{\text{PhRc}}, n) = \frac{k_{\text{PhRc}} \cdot ([\text{Ce}] + [\text{Ca}])^n}{j_{\text{Cy}}^n + ([\text{Ce}] + [\text{Ca}])^n} \cdot [\text{Rc}] \quad (130)$$

$$r_{\text{PhRc}_1}([\text{Ca}], [\text{Ce}], [\text{Rc}], j_{\text{Cy}}, k_{\text{PhRc}}, n) = \frac{k_{\text{PhRc}} \cdot ([\text{Ce}] + [\text{Ca}])^n}{j_{\text{Cy}}^n + ([\text{Ce}] + [\text{Ca}])^n} \cdot [\text{Rc}] \quad (131)$$

8.35 Reaction [Dephosphorylation_of_replication_complexes](#)

This is an irreversible reaction of one reactant forming one product.

Name Dephosphorylation of replication complexes

Reaction equation



Reactant

Table 80: Properties of each reactant.

Id	Name	SBO
pRc	Pre_Replication_complex_primed	

Product

Table 81: Properties of each product.

Id	Name	SBO
Rc	Pre_Replication_complex	

Kinetic Law

Derived unit contains undeclared units

$$v_{35} = \text{vol}(\text{Cell}) \cdot k_{\text{DpRc}} \cdot [\text{pRc}] \quad (133)$$

8.36 Reaction

[Association_dissociation_of_p21_and_replication_complexes](#)

This is a reversible reaction of two reactants forming one product.

Name Association/dissociation of p21 and replication complexes

Reaction equation



Reactants

Table 82: Properties of each reactant.

Id	Name	SBO
aRc	Pre_Replication_complex_active	
P21	P21	

Product

Table 83: Properties of each product.

Id	Name	SBO
iRc	Pre_Replication_complex_inactive	

Kinetic Law

Derived unit contains undeclared units

$$v_{36} = \text{vol}(\text{Cell}) \cdot (k_{\text{AsPcP21}} \cdot [\text{aRc}] \cdot [\text{P21}] - k_{\text{DsPcP21}} \cdot [\text{iRc}]) \quad (135)$$

8.37 Reaction [Degradation_of_p21_in_inactive_replication_complexes](#)

This is an irreversible reaction of one reactant forming one product influenced by three modifiers.

Name Degradation of p21 in inactive replication complexes

Reaction equation



Reactant

Table 84: Properties of each reactant.

Id	Name	SBO
iRc	Pre_Replication_complex_inactive	

Modifiers

Table 85: Properties of each modifier.

Id	Name	SBO
Ce	CyclinE_Cdk2_active	
Ca	CyclinA_Cdk2	
aRc	Pre_Replication_complex_active	

Product

Table 86: Properties of each product.

Id	Name	SBO
aRc	Pre_Replication_complex_active	

Kinetic Law

Derived unit contains undeclared units

$$v_{37} = \text{vol}(\text{Cell}) \cdot \text{rDeP21_5}([\text{Ca}], \text{Cdt2}, [\text{Ce}], \text{Skp2}, [\text{aRc}], [\text{iRc}], \text{kDeP21}, \text{kDeP21Cy}, \text{kDeP21aRc}) \quad (137)$$

$$\begin{aligned} & \text{rDeP21_5}([\text{Ca}], \text{Cdt2}, [\text{Ce}], \text{Skp2}, [\text{aRc}], [\text{iRc}], \text{kDeP21}, \text{kDeP21Cy}, \text{kDeP21aRc}) \\ &= (\text{kDeP21} + \text{kDeP21Cy} \cdot \text{Skp2} \cdot ([\text{Ce}] + [\text{Ca}]) + \text{kDeP21aRc} \cdot \text{Cdt2} \cdot [\text{aRc}]) \cdot [\text{iRc}] \end{aligned} \quad (138)$$

$$\begin{aligned} & \text{rDeP21_5}([\text{Ca}], \text{Cdt2}, [\text{Ce}], \text{Skp2}, [\text{aRc}], [\text{iRc}], \text{kDeP21}, \text{kDeP21Cy}, \text{kDeP21aRc}) \\ &= (\text{kDeP21} + \text{kDeP21Cy} \cdot \text{Skp2} \cdot ([\text{Ce}] + [\text{Ca}]) + \text{kDeP21aRc} \cdot \text{Cdt2} \cdot [\text{aRc}]) \cdot [\text{iRc}] \end{aligned} \quad (139)$$

8.38 Reaction Dissassembly_of_RC

This is an irreversible reaction of one reactant forming no product influenced by one modifier.

Name Dissassembly of RC

Reaction equation



Reactant

Table 87: Properties of each reactant.

Id	Name	SBO
Rc	Pre_Replication_complex	

Modifier

Table 88: Properties of each modifier.

Id	Name	SBO
Dna	Dna	

Kinetic Law

Derived unit contains undeclared units

$$v_{38} = \text{vol}(\text{Cell}) \cdot \text{rDsRc}_1([\text{Dna}], [\text{Rc}]) \quad (141)$$

$$\text{rDsRc}_1([\text{Dna}], [\text{Rc}]) = \begin{cases} 0 & \text{if } [\text{Dna}] < 1 \\ \begin{cases} 1 \cdot [\text{Rc}] & \text{if } [\text{Dna}] > 1 \\ 0.5 \cdot [\text{Rc}] & \text{otherwise} \end{cases} & \text{otherwise} \end{cases} \quad (142)$$

$$\text{rDsRc}_1([\text{Dna}], [\text{Rc}]) = \begin{cases} 0 & \text{if } [\text{Dna}] < 1 \\ \begin{cases} 1 \cdot [\text{Rc}] & \text{if } [\text{Dna}] > 1 \\ 0.5 \cdot [\text{Rc}] & \text{otherwise} \end{cases} & \text{otherwise} \end{cases} \quad (143)$$

8.39 Reaction Dissassembly_of_pRC

This is an irreversible reaction of one reactant forming no product influenced by one modifier.

Name Dissassembly of pRC

Reaction equation



Reactant

Table 89: Properties of each reactant.

Id	Name	SBO
pRc	Pre_Replication_complex_primed	

Modifier

Table 90: Properties of each modifier.

Id	Name	SBO
Dna	Dna	

Kinetic Law

Derived unit contains undeclared units

$$v_{39} = \text{vol}(\text{Cell}) \cdot \text{rDsRc}_2([\text{Dna}], [\text{pRc}]) \quad (145)$$

$$\text{rDsRc}_2([\text{Dna}], [\text{pRc}]) = \begin{cases} 0 & \text{if } [\text{Dna}] < 1 \\ \begin{cases} 1 \cdot [\text{pRc}] & \text{if } [\text{Dna}] > 1 \\ 0.5 \cdot [\text{pRc}] & \text{otherwise} \end{cases} & \text{otherwise} \end{cases} \quad (146)$$

$$\text{rDsRc}_2([\text{Dna}], [\text{pRc}]) = \begin{cases} 0 & \text{if } [\text{Dna}] < 1 \\ \begin{cases} 1 \cdot [\text{pRc}] & \text{if } [\text{Dna}] > 1 \\ 0.5 \cdot [\text{pRc}] & \text{otherwise} \end{cases} & \text{otherwise} \end{cases} \quad (147)$$

8.40 Reaction Dissassembly_of_aRC

This is an irreversible reaction of one reactant forming one product influenced by one modifier.

Name Dissasembly of aRC

Reaction equation



Reactant

Table 91: Properties of each reactant.

Id	Name	SBO
aRc	Pre_Replication_complex_active	

Modifier

Table 92: Properties of each modifier.

Id	Name	SBO
Dna	Dna	

Product

Table 93: Properties of each product.

Id	Name	SBO
aPcna	Pcna_nuclear_active	

Kinetic Law

Derived unit contains undeclared units

$$v_{40} = \text{vol}(\text{Cell}) \cdot \text{rDsRc}_3([\text{Dna}], [\text{aRc}]) \quad (149)$$

$$\text{rDsRc}_3([\text{Dna}], [\text{aRc}]) = \begin{cases} 0 & \text{if } [\text{Dna}] < 1 \\ \begin{cases} 1 \cdot [\text{aRc}] & \text{if } [\text{Dna}] > 1 \\ 0.5 \cdot [\text{aRc}] & \text{otherwise} \end{cases} & \text{otherwise} \end{cases} \quad (150)$$

$$\text{rDsRc}_3([\text{Dna}], [\text{aRc}]) = \begin{cases} 0 & \text{if } [\text{Dna}] < 1 \\ \begin{cases} 1 \cdot [\text{aRc}] & \text{if } [\text{Dna}] > 1 \\ 0.5 \cdot [\text{aRc}] & \text{otherwise} \end{cases} & \text{otherwise} \end{cases} \quad (151)$$

8.41 Reaction *Dissassembly_of_iRC*

This is an irreversible reaction of one reactant forming one product influenced by one modifier.

Name Dissasembly of iRC

Reaction equation



Reactant

Table 94: Properties of each reactant.

Id	Name	SBO
iRc	Pre_Replication_complex_inactive	

Modifier

Table 95: Properties of each modifier.

Id	Name	SBO
Dna	Dna	

Product

Table 96: Properties of each product.

Id	Name	SBO
iPcna	PCNA_Nuclear_inactive	

Kinetic Law

Derived unit contains undeclared units

$$v_{41} = \text{vol}(\text{Cell}) \cdot \text{rDsRc_4}([\text{Dna}], [\text{iRc}]) \quad (153)$$

$$\text{rDsRc_4}([\text{Dna}], [\text{iRc}]) = \begin{cases} 0 & \text{if } [\text{Dna}] < 1 \\ 1 \cdot [\text{iRc}] & \text{if } [\text{Dna}] > 1 \\ 0.5 \cdot [\text{iRc}] & \text{otherwise} \end{cases} \quad (154)$$

$$rDsRc_4([Dna], [iRc]) = \begin{cases} 0 & \text{if } [Dna] < 1 \\ \begin{cases} 1 \cdot [iRc] & \text{if } [Dna] > 1 \\ 0.5 \cdot [iRc] & \text{otherwise} \end{cases} & \text{otherwise} \end{cases} \quad (155)$$

8.42 Reaction Synthesis_of_DNA

This is an irreversible reaction of one reactant forming two products.

Name Synthesis of DNA

Reaction equation



Reactant

Table 97: Properties of each reactant.

Id	Name	SBO
aRc	Pre_Replication_complex_active	

Products

Table 98: Properties of each product.

Id	Name	SBO
aRc	Pre_Replication_complex_active	
Dna	Dna	

Kinetic Law

Derived unit contains undeclared units

$$v_{42} = \text{vol}(\text{Cell}) \cdot k\text{SyDna} \cdot [aRc] \quad (157)$$

8.43 Reaction Synthesis_of_p53

This is an irreversible reaction of no reactant forming one product.

Name Synthesis of p53

Reaction equation



Product

Table 99: Properties of each product.

Id	Name	SBO
P53	P53	

Kinetic Law

Derived unit contains undeclared units

$$v_{43} = \text{vol}(\text{Cell}) \cdot \text{Constant_flux_irreversible}(\text{kSyP53}) \quad (159)$$

$$\text{Constant_flux_irreversible}(v) = v \quad (160)$$


$$\text{Constant_flux_irreversible}(v) = v \quad (161)$$

8.44 Reaction *Degradation_of_p53*

This is an irreversible reaction of one reactant forming no product influenced by one modifier.

Name Degradation of p53

Reaction equation



Reactant

Table 100: Properties of each reactant.

Id	Name	SBO
P53	P53	

Modifier

Table 101: Properties of each modifier.

Id	Name	SBO
Dam	Dna_damage	

Kinetic Law

Derived unit contains undeclared units

$$v_{44} = \text{vol}(\text{Cell}) \cdot \text{rDeP53_1}([\text{Dam}], [\text{P53}], \text{jP53}, \text{kDeP53}) \quad (163)$$

$$\text{rDeP53_1}([\text{Dam}], [\text{P53}], \text{jP53}, \text{kDeP53}) = \frac{\text{kDeP53}}{\text{jP53} + [\text{Dam}]} \cdot [\text{P53}] \quad (164)$$

$$\text{rDeP53_1}([\text{Dam}], [\text{P53}], \text{jP53}, \text{kDeP53}) = \frac{\text{kDeP53}}{\text{jP53} + [\text{Dam}]} \cdot [\text{P53}] \quad (165)$$

8.45 Reaction *Induction_of_DNA_damage*

This is an irreversible reaction of no reactant forming one product.

Name Induction of DNA damage

Reaction equation



Product

Table 102: Properties of each product.

Id	Name	SBO
Dam	Dna_damage	

Kinetic Law

Derived unit contains undeclared units

$$v_{45} = \text{vol}(\text{Cell}) \cdot \text{Constant_flux_irreversible}(\text{kGeDam}) \quad (167)$$

$$\text{Constant_flux_irreversible}(v) = v \quad (168)$$

$$\text{Constant_flux_irreversible}(v) = v \quad (169)$$

8.46 Reaction [Induction_of_DNA_damage_by_replication](#)

This is an irreversible reaction of one reactant forming two products.

Name Induction of DNA damage by replication

Reaction equation



Reactant

Table 103: Properties of each reactant.

Id	Name	SBO
aRc	Pre_Replication_complex_active	

Products

Table 104: Properties of each product.

Id	Name	SBO
aRc	Pre_Replication_complex_active	
Dam	Dna_damage	

Kinetic Law

Derived unit contains undeclared units

$$v_{46} = \text{vol}(\text{Cell}) \cdot k_{\text{GeDamArc}} \cdot [\text{aRc}] \quad (171)$$

8.47 Reaction [Synthesis_of_APC_C_Cdh1_activity_probe](#)

This is an irreversible reaction of no reactant forming one product.

Name Synthesis of APC/C[^]Cdh1 activity probe

Reaction equation



Product

Table 105: Properties of each product.

Id	Name	SBO
Pr	Activity_probe_of_APC_C_Cdh1	

Kinetic Law

Derived unit contains undeclared units

$$v_{47} = \text{vol}(\text{Cell}) \cdot \text{Constant_flux_irreversible}(\text{kSyPr}) \quad (173)$$

$$\text{Constant_flux_irreversible}(v) = v \quad (174)$$

$$\text{Constant_flux_irreversible}(v) = v \quad (175)$$

8.48 Reaction Degradation_of_APC_C_Cdh1_activity_probe

This is an irreversible reaction of one reactant forming no product influenced by one modifier.

Name Degradation of APC/C[^]Cdh1 activity probe

Reaction equation



Reactant

Table 106: Properties of each reactant.

Id	Name	SBO
Pr	Activity_probe_of_APC_C_Cdh1	

Modifier

Table 107: Properties of each modifier.

Id	Name	SBO
C1	Cdh1_C_APC_active	

Kinetic Law

Derived unit contains undeclared units

$$v_{48} = \text{vol}(\text{Cell}) \cdot \text{rDePr_1}([C1], [Pr], kDeCaC1, kDePr) \quad (177)$$

$$\text{rDePr_1}([C1], [Pr], kDeCaC1, kDePr) = (kDePr + kDeCaC1 \cdot [C1]) \cdot [Pr] \quad (178)$$

$$\text{rDePr_1}([C1], [Pr], kDeCaC1, kDePr) = (kDePr + kDeCaC1 \cdot [C1]) \cdot [Pr] \quad (179)$$

8.49 Reaction Repair_of_DNA_damage

This is an irreversible reaction of one reactant forming no product influenced by one modifier.

Name Repair of DNA damage

Reaction equation



Reactant

Table 108: Properties of each reactant.

Id	Name	SBO
Dam	Dna_damage	

Modifier

Table 109: Properties of each modifier.

Id	Name	SBO
P53	P53	

Kinetic Law

Derived unit contains undeclared units

$$v_{49} = \text{vol}(\text{Cell}) \cdot \text{rReDam}_1([\text{Dam}], [\text{P53}], j_{\text{Dam}}, k_{\text{ReDam}}, k_{\text{ReDamP53}}) \quad (181)$$

$$\begin{aligned} & \text{rReDam}_1([\text{Dam}], [\text{P53}], j_{\text{Dam}}, k_{\text{ReDam}}, k_{\text{ReDamP53}}) \\ &= \left(k_{\text{ReDam}} + \frac{k_{\text{ReDamP53}} \cdot [\text{P53}]}{j_{\text{Dam}} + [\text{Dam}]} \right) \cdot [\text{Dam}] \end{aligned} \quad (182)$$

$$\begin{aligned} & \text{rReDam}_1([\text{Dam}], [\text{P53}], j_{\text{Dam}}, k_{\text{ReDam}}, k_{\text{ReDamP53}}) \\ &= \left(k_{\text{ReDam}} + \frac{k_{\text{ReDamP53}} \cdot [\text{P53}]}{j_{\text{Dam}} + [\text{Dam}]} \right) \cdot [\text{Dam}] \end{aligned} \quad (183)$$

9 Derived Rate Equations

When interpreted as an ordinary differential equation framework, this model implies the following set of equations for the rates of change of each species.

Identifiers for kinetic laws highlighted in gray cannot be verified to evaluate to units of SBML substance per time. As a result, some SBML interpreters may not be able to verify the consistency of the units on quantities in the model. Please check if

- parameters without an unit definition are involved or
- volume correction is necessary because the `hasOnlySubstanceUnits` flag may be set to `false` and `spacialDimensions` > 0 for certain species.

9.1 Species Rb

Name Retinoblastoma_protein

Notes (AU) hypo-phosphorylated Rb

Initial concentration 0 mmol · ml⁻¹

This species takes part in four reactions (as a reactant in [Phosphorylation_of_Rb](#), [Association-dissociation_of_Rb_and_E2F](#) and as a product in [Dephosphorylation_of_Rb](#), [Degradation-of_E2F_in_Rb_E2F_complexes](#)).

$$\frac{d}{dt}\text{Rb} = v_3 + v_6 - v_1 - v_7 \quad (184)$$

9.2 Species pRb

Name Retinoblastoma_protein_hyperphosphorylated

Notes (AU) hyper-phosphorylated Rb

Initial concentration 5 mmol · ml⁻¹

This species takes part in three reactions (as a reactant in [Dephosphorylation_of_Rb](#) and as a product in [Phosphorylation_of_Rb](#), [Phosphorylation_Rb_in_Rb_E2F_complexes](#)).

$$\frac{d}{dt}pRb = v_1 + v_2 - v_3 \quad (185)$$

9.3 Species E2f

Name E2f_active

Notes (AU) free, active E2F

Initial concentration 0 mmol · ml⁻¹

This species takes part in eleven reactions (as a reactant in [Degradation_of_E2F](#), [Association-dissociation_of_Rb_and_E2F](#), [Synthesis_of_CycE](#), [Synthesis_of_CycA](#), [Synthesis_of_Emi1](#) and as a product in [Phosphorylation_Rb_in_Rb_E2F_complexes](#), [Synthesis_of_E2F](#), [Synthesis_of_CycE](#), [Synthesis_of_CycA](#), [Synthesis_of_Emi1](#) and as a modifier in [Synthesis_of_E2F](#)).

$$\frac{d}{dt}E2f = v_2 + v_4 + v_9 + v_{10} + v_{20} - v_5 - v_7 - v_9 - v_{10} - v_{20} \quad (186)$$

9.4 Species RbE2f

Name Retinoblastoma_protein_E2f_complex_inactive

Notes (AU) Rb:E2F complexes (inactive)

Initial concentration 0 mmol · ml⁻¹

This species takes part in three reactions (as a reactant in [Phosphorylation_Rb_in_Rb_E2F_complexes](#), [Degradation_of_E2F_in_Rb_E2F_complexes](#) and as a product in [Association-dissociation_of_Rb_and_E2F](#)).

$$\frac{d}{dt}RbE2f = v_7 - v_2 - v_6 \quad (187)$$

9.5 Species E1

Name Emi1

Notes (AU) free Emi1

Initial concentration 0 mmol · ml⁻¹

This species takes part in four reactions (as a reactant in [Degradation_of_Emi1](#), [Association-dissociation_of_Emi1-APC-C-Cdh1_complexes](#) and as a product in [Synthesis_of_Emi1](#), [Phosphorylation_of-APC-C-Cdh1_in_Emi1-APC-C-Cdh1_complexes](#)).

$$\frac{d}{dt}E1 = v_{20} + v_{25} - v_{21} - v_{22} \quad (188)$$

9.6 Species C1

Name Cdh1_C-APC_active

Notes (AU) free, active APC/C~Cdh1

Initial concentration 0 mmol · ml⁻¹

This species takes part in seven reactions (as a reactant in [Association-dissociation_of_Emi1-APC-C-Cdh1_complexes](#), [Phosphorylation_of-free-APC-C-Cdh1](#) and as a product in [Degradation_of_Emi1_in_Emi1-APC-C-Cdh1_complexes](#), [Dephosphorylation_of-APC-C-Cdh1](#) and as a modifier in [Degradation_of-CycA_in-CycA-Cdk2_complexes](#), [Degradation_of-CycA_in-CycA-Cdk2_p21_complexes](#), [Degradation_of-APC-C-Cdh1_activity_probe](#)).

$$\frac{d}{dt}C1 = v_{23} + v_{26} - v_{22} - v_{24} \quad (189)$$

9.7 Species pC1

Name C-Cdh1-APC_phosphorylated_inactive

Notes (AU) phosphorylated APC/C~Cdh1 (inactive)

Initial concentration 1 mmol · ml⁻¹

This species takes part in three reactions (as a reactant in [Dephosphorylation_of-APC-C-Cdh1](#) and as a product in [Phosphorylation_of-free-APC-C-Cdh1](#), [Phosphorylation_of-APC-C-Cdh1_in_Emi1-APC-C-Cdh1_complexes](#)).

$$\frac{d}{dt}pC1 = v_{24} + v_{25} - v_{26} \quad (190)$$

9.8 Species E1C1

Name Emi1_C_Cdh1_complex_inactive

Notes (AU) Emi1:APC/C⁺Cdh1 complexes (inactive) ; Emi1 is an APC_C_Cdh1 inhibitor

Initial concentration 0 mmol · ml⁻¹

This species takes part in three reactions (as a reactant in [Degradation_of_Emi1_in_Emi1-APC_C_Cdh1_complexes](#), [Phosphorylation_of_APC_C_Cdh1_in_Emi1-APC_C_Cdh1_complexes](#) and as a product in [Association_dissociation_of_Emi1-APC_C_Cdh1_complexes](#)).

$$\frac{d}{dt}E1C1 = v_{22} - v_{23} - v_{25} \quad (191)$$

9.9 Species P21

Name P21

Notes (AU) free p21

Initial concentration 0.6 mmol · ml⁻¹

This species takes part in nine reactions (as a reactant in [Association_dissociation_of_CycE_Cdk2_and_p21](#), [Association_dissociation_of_CycA_Cdk2_and_p21](#), [Degradation_of_free_p21](#), [Association_dissociation_of_PCNA_and_p21](#), [Association_dissociation_of_p21_and_replication_complexes](#) and as a product in [Synthesis_of_p21](#), [Degradation_of_CycE_in_CycE_Cdk2_p21_complexes](#), [Degradation_of_CycA_in_CycA_Cdk2_p21_complexes](#), [Nuclear_export_of_inactive_PCNA](#)).

$$\frac{d}{dt}P21 = v_8 + v_{16} + v_{18} + v_{29} - v_{11} - v_{12} - v_{19} - v_{30} - v_{36} \quad (192)$$

9.10 Species Ce

Name CyclinE_Cdk2_active

Notes (AU) free, active CycE:Cdk2

Initial concentration 0.5 mmol · ml⁻¹

This species takes part in 14 reactions (as a reactant in [Association_dissociation_of_CycE_Cdk2_and_p21](#), [Degradation_of_CycE_in_CycE_Cdk2_complexes](#) and as a product in [Synthesis_of_CycE](#), [Degradation_of_p21_in_CycE_Cdk2_p21_complexes](#) and as a modifier in [Phosphorylation_of_Rb](#), [Phosphorylation_Rb_in_Rb_E2F_complexes](#), [Degradation_of_p21_in_CycE_Cdk2_p21_complexes](#), [Degradation_of_p21_in_CycA_Cdk2_p21_complexes](#), [Degradation_of_free_p21](#), [Phosphorylation_of_free_APC_C_Cdh1](#), [Phosphorylation_of_APC_C_Cdh1_in_Emi1-APC_C_Cdh1_complexes](#), [Degradation_of_p21_in_PCNA_p21_complexes](#),

Phosphorylation priming of replication complexes, Degradation of p21 in inactive replication complexes).

$$\frac{d}{dt}Ce = v_9 + v_{13} - v_{11} - v_{15} \quad (193)$$

9.11 Species Ca

Name CyclinA_Cdk2

Notes (AU) free, active CycA:Cdk2

Initial concentration 1.2 mmol · ml⁻¹

This species takes part in 16 reactions (as a reactant in [Association dissociation of CycA_Cdk2 and p21](#), [Degradation of CycA in CycA_Cdk2 complexes](#) and as a product in [Synthesis of CycA](#), [Degradation of p21 in CycA_Cdk2_p21 complexes](#) and as a modifier in [Phosphorylation of Rb](#), [Phosphorylation Rb in Rb_E2F complexes](#), [Degradation of p21 in CycE_Cdk2_p21 complexes](#), [Degradation of p21 in CycA_Cdk2_p21 complexes](#), [Degradation of CycE in CycE_Cdk2 complexes](#), [Degradation of CycE in CycE_Cdk2_p21 complexes](#), [Degradation of free p21](#), [Phosphorylation of free APC_C_Cdh1](#), [Phosphorylation of APC_C_Cdh1 in Emi1_APC_C_Cdh1 complexes](#), [Degradation of p21 in PCNA_p21 complexes](#), [Phosphorylation priming of replication complexes](#), [Degradation of p21 in inactive replication complexes](#)).

$$\frac{d}{dt}Ca = v_{10} + v_{14} - v_{12} - v_{17} \quad (194)$$

9.12 Species CeP21

Name CyclinE_Cdk2_P21_complex_inactive

Notes (AU) CycE:Cdk2:p21 complexes (inactive)

Initial concentration 0 mmol · ml⁻¹

This species takes part in three reactions (as a reactant in [Degradation of p21 in CycE_Cdk2_p21 complexes](#), [Degradation of CycE in CycE_Cdk2_p21 complexes](#) and as a product in [Association dissociation of CycE_Cdk2 and p21](#)).

$$\frac{d}{dt}CeP21 = v_{11} - v_{13} - v_{16} \quad (195)$$

9.13 Species CaP21

Name CyclinA_Cdk2_P21_complex_inactive

Notes (AU) CycA:Cdk2:p21 complexes (inactive)

Initial concentration 0 mmol · ml⁻¹

This species takes part in three reactions (as a reactant in [Degradation_of_p21_in_CycA_Cdk2_p21_complexes](#), [Degradation_of_CycA_in_CycA_Cdk2_p21_complexes](#) and as a product in [Association_dissociation_of_CycA_Cdk2_and_p21](#)).

$$\frac{d}{dt}\text{CaP21} = v_{12} - v_{14} - v_{18} \quad (196)$$

9.14 Species aPcna

Name Pcna_nuclear_active

Notes (AU) active, nuclear PCNA

Initial concentration 0.5 mmol · ml⁻¹

This species takes part in six reactions (as a reactant in [Nuclear_export_of_active_PCNA](#), [Association_dissociation_of_PCNA_and_p21](#), [Association_dissociation_of_active_PCNA_and_replication_complexes](#) and as a product in [Nuclear_import_of_active_PCNA](#), [Degradation_of_p21_in_PCNA_p21_complexes](#), [Dissassembly_of_aRC](#)).

$$\frac{d}{dt}\text{aPcna} = v_{27} + v_{31} + v_{40} - v_{28} - v_{30} - v_{32} \quad (197)$$

9.15 Species iPcna

Name PCNA_Nuclear_inactive

Notes (AU) inactive, nuclear PCNA

Initial concentration 0 mmol · ml⁻¹

This species takes part in five reactions (as a reactant in [Nuclear_export_of_inactive_PCNA](#), [Degradation_of_p21_in_PCNA_p21_complexes](#), [Association_dissociation_of_inactive_PCNA_and_replication_complexes](#) and as a product in [Association_dissociation_of_PCNA_and_p21](#), [Dissassembly_of_iRC](#)).

$$\frac{d}{dt}\text{iPcna} = v_{30} + v_{41} - v_{29} - v_{31} - v_{33} \quad (198)$$

9.16 Species Rc

Name Pre_Replication_complex

Notes (AU) pre-replication complexes

Initial concentration 1 mmol · ml⁻¹

This species takes part in three reactions (as a reactant in [Phosphorylation priming of replication complexes](#), [Dissassembly of RC](#) and as a product in [Dephosphorylation of replication complexes](#)).

$$\frac{d}{dt}Rc = v_{35} - v_{34} - v_{38} \quad (199)$$

9.17 Species pRc

Name Pre_Replication_complex_primed

Notes (AU) primed replication complexes

Initial concentration 0 mmol · ml⁻¹

This species takes part in five reactions (as a reactant in [Association dissociation of active PCNA and replication complexes](#), [Association dissociation of inactive PCNA and replication complexes](#), [Dephosphorylation of replication complexes](#), [Dissassembly of pRC](#) and as a product in [Phosphorylation priming of replication complexes](#)).

$$\frac{d}{dt}pRc = v_{34} - v_{32} - v_{33} - v_{35} - v_{39} \quad (200)$$

9.18 Species aRc

Name Pre_Replication_complex_active

Notes (AU) active replication complexes

Initial concentration 0 mmol · ml⁻¹

This species takes part in 13 reactions (as a reactant in [Association dissociation of p21 and replication complexes](#), [Dissassembly of aRC](#), [Synthesis of DNA](#), [Induction of DNA damage by replication](#) and as a product in [Association dissociation of active PCNA and replication complexes](#), [Degradation of p21 in inactive replication complexes](#), [Synthesis of DNA](#), [Induction of DNA damage by replication](#) and as a modifier in [Degradation of p21 in CycE-Cdk2 p21 complexes](#), [Degradation of p21 in CycA-Cdk2 p21 complexes](#), [Degradation of free p21](#), [Degradation of p21 in PCNA p21 complexes](#), [Degradation of p21 in inactive replication complexes](#)).

$$\frac{d}{dt}aRc = v_{32} + v_{37} + v_{42} + v_{46} - v_{36} - v_{40} - v_{42} - v_{46} \quad (201)$$

9.19 Species iRc

Name Pre_Replication_complex_inactive

Notes (AU) inactive replication complexes

Initial concentration 0 mmol · ml⁻¹

This species takes part in four reactions (as a reactant in [Degradation_of_p21_in_inactive-replication_complexes](#), [Dissassembly_of_iRC](#) and as a product in [Association_dissociation_of_inactive_PCNA_and_replication_complexes](#), [Association_dissociation_of_p21_and_replication_complexes](#)).

$$\frac{d}{dt}iRc = v_{33} + v_{36} - v_{37} - v_{41} \quad (202)$$

9.20 Species Dna

Name Dna

Notes (AU) DNA

Initial concentration 0 mmol · ml⁻¹

This species takes part in five reactions (as a product in [Synthesis_of_DNA](#) and as a modifier in [Dissassembly_of_RC](#), [Dissassembly_of_pRC](#), [Dissassembly_of_aRC](#), [Dissassembly_of_iRC](#)).

$$\frac{d}{dt}Dna = v_{42} \quad (203)$$

9.21 Species P53

Name P53

Notes (AU) p53 transcription factor

Initial concentration 0 mmol · ml⁻¹

This species takes part in four reactions (as a reactant in [Degradation_of_p53](#) and as a product in [Synthesis_of_p53](#) and as a modifier in [Synthesis_of_p21](#), [Repair_of_DNA_damage](#)).

$$\frac{d}{dt}P53 = v_{43} - v_{44} \quad (204)$$

9.22 Species [Dam](#)

Name Dna_damage

Notes (AU) DNA damage

Initial concentration 0 mmol · ml⁻¹

This species takes part in four reactions (as a reactant in [Repair_of_DNA_damage](#) and as a product in [Induction_of_DNA_damage](#), [Induction_of_DNA_damage_by_replication](#) and as a modifier in [Degradation_of_p53](#)).

$$\frac{d}{dt}\text{Dam} = v_{45} + v_{46} - v_{49} \quad (205)$$

9.23 Species [Pr](#)

Name Activity_probe_of_APC_C_Cdh1

Notes (AU) APC/C[^]Cdh1 activity probe

Initial concentration 0.5 mmol · ml⁻¹

This species takes part in two reactions (as a reactant in [Degradation_of_APC_C_Cdh1_activity_probe](#) and as a product in [Synthesis_of_APC_C_Cdh1_activity_probe](#)).

$$\frac{d}{dt}\text{Pr} = v_{47} - v_{48} \quad (206)$$

9.24 Species [tRb](#)

Name Retinoblastoma_protein_total

Notes (AU) total Rb protein

Initial concentration 5 mmol · ml⁻¹

Involved in rule [tRb](#)

One rule determines the species' quantity.

9.25 Species [tE2f](#)

Name E2f_total

Notes (AU) total E2F protein

Initial concentration 0 mmol · ml⁻¹

Involved in rule [tE2f](#)

One rule determines the species' quantity.

9.26 Species [tE1](#)

Name Emi1_total

Notes (AU) total Emi1 protein

Initial concentration 0 mmol · ml⁻¹

Involved in rule [tE1](#)

One rule determines the species' quantity.

9.27 Species [tC1](#)

Name C_Cdh1_APC_total

Notes (AU) total APC/C[^]Cdh1 protein ; APC/C[^]Cdh1 protein ubiquitin E3 ligase that is act.

Initial concentration 1 mmol · ml⁻¹

Involved in rule [tC1](#)

One rule determines the species' quantity.

9.28 Species [tCe](#)

Name CyclinE_Cdk2_total

Notes (AU) total CycE:Cdk2

Initial concentration 0.5 mmol · ml⁻¹

Involved in rule [tCe](#)

One rule determines the species' quantity.

9.29 Species [tCa](#)

Name CyclinA_Cdk2_total

Notes (AU) total CycA:Cdk2

Initial concentration 1.2 mmol · ml⁻¹

Involved in rule [tCa](#)

One rule determines the species' quantity.

9.30 Species `tP21`

Name `P21_total`

Notes (AU) total p21 protein

Initial concentration $0.6 \text{ mmol} \cdot \text{ml}^{-1}$

Involved in rule `tP21`

One rule determines the species' quantity.

SBML²TeX was developed by Andreas Dräger^a, Hannes Planatscher^a, Dieudonné M Wouamba^a, Adrian Schröder^a, Michael Hucka^b, Lukas Endler^c, Martin Golebiewski^d and Andreas Zell^a. Please see <http://www.ra.cs.uni-tuebingen.de/software/SBML2LaTeX> for more information.

^aCenter for Bioinformatics Tübingen (ZBIT), Germany

^bCalifornia Institute of Technology, Beckman Institute BNMC, Pasadena, United States

^cEuropean Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, United Kingdom

^dEML Research gGmbH, Heidelberg, Germany