

SBML Model Report

Model name: “Kallenberger2014 - CD95L induced apoptosis initiated by caspase-8, wild-type HeLa cells (cis/trans-cis/trans variant)”



May 5, 2016

1 General Overview

This is a document in SBML Level 2 Version 4 format. This model was created by the following two authors: Vijayalakshmi Chelliah¹ and Stefan Kallenberger² at March 14th 2014 at 12:53 a. m. and last time modified at February 25th 2015 at 12:39 a. m. Table 1 gives an overview of the quantities of all components of this model.

Table 1: Number of components in this model, which are described in the following sections.

Element	Quantity	Element	Quantity
compartment types	0	compartments	1
species types	0	species	18
events	0	constraints	0
reactions	19	function definitions	0
global parameters	15	unit definitions	0
rules	1	initial assignments	0

¹EMBL-EBI, viji@ebi.ac.uk

²German Cancer Research Center, s.kallenberger@dkfz.de

Model Notes

Kallenberger2014 - CD95L induced apoptosis initiated by caspase-8, wild-type HeLa cells (cis/trans-cis/trans variant)

The paper describes a new approach that combines single cell and population data in the same model. The model consists of a large number of single cell models, which are fitted to single cell data. Simultaneously, ensemble averages are fitted to population data. It is assumed that the kinetics in each cell can be described with the same kinetic parameters. Therefore, cell-to-cell variability is explained by variable initial protein concentrations.

There are four variants of the model (with [CD95L]=500ng/ml = 16.6nM), i) cistrans (in CD95-HeLa cells) [[MODEL1403050000](#)], ii) cistrans (in wild-type HeLa cells) [[MODEL1403050001](#)], iii) cistrans-cistrans (in CD95-HeLa cells) [[MODEL1403050002](#)], and iv) cistrans-cistrans (in wild-type HeLa cells) [[MODEL1403050003](#)].

These model contain the equations for one „average cell,, with median initial concentrations for CD95, FADD, p55, BID, PrNES_mCherry and PrER_mGFP. By integrating the model, it should be possible to obtain trajectories for PrER_mGFP, PrNES_mCherry, p43 and p18 similar as in Figure 4A (CD95-HeLa cells) and Figure 4B (wild-type HeLa cells).

This model is described in the article:[Intra- and Interdimeric Caspase-8 Self-Cleavage Controls Strength and Timing of CD95-Induced Apoptosis](#) Stefan M. Kallenberger, Jol Beaudouin, Juliane Claus, Carmen Fischer, Peter K. Sorger, Stefan Legewie, and Roland Eils 11 March 2014: Vol. 7, Issue 316, p. ra23

Abstract:

Apoptosis in response to the ligand CD95L (also known as Fas ligand) is initiated by caspase-8, which is activated by dimerization and self-cleavage at death-inducing signaling complexes (DISCs). Previous work indicated that the degree of substrate cleavage by caspase-8 determines whether a cell dies or survives in response to a death stimulus. To determine how a death ligand stimulus is effectively translated into caspase-8 activity, we assessed this activity over time in single cells with compartmentalized probes that are cleaved by caspase-8 and used multiscale modeling to simultaneously describe single-cell and population data with an ensemble of single-cell models. We derived and experimentally validated a minimal model in which cleavage of caspase-8 in the enzymatic domain occurs in an interdimeric manner through interaction between DISCs, whereas prodomain cleavage sites are cleaved in an intradimeric manner within DISCs. Modeling indicated that sustained membrane-bound caspase-8 activity is followed by transient cytosolic activity, which can be interpreted as a molecular timer mechanism reflected by a limited lifetime of active caspase-8. The activation of caspase-8 by combined intra- and interdimeric cleavage ensures weak signaling at low concentrations of CD95L and strongly accelerated activation at higher ligand concentrations, thereby contributing to precise control of apoptosis.

This model is hosted on [BioModels Database](#) and identified by: [BIOMD0000000526](#).

To cite BioModels Database, please use: [BioModels Database: An enhanced, curated and annotated resource for published quantitative kinetic models](#).

To the extent possible under law, all copyright and related or neighbouring rights to this encoded model have been dedicated to the public domain worldwide. Please refer to [CC0 Public Domain Dedication](#) for more information.

2 Unit Definitions

This is an overview of five unit definitions which are all predefined by SBML and not mentioned in the model.

2.1 Unit substance

Notes Mole is the predefined SBML unit for substance.

Definition mol

2.2 Unit volume

Notes Litre is the predefined SBML unit for volume.

Definition l

2.3 Unit area

Notes Square metre is the predefined SBML unit for area since SBML Level 2 Version 1.

Definition m²

2.4 Unit length

Notes Metre is the predefined SBML unit for length since SBML Level 2 Version 1.

Definition m

2.5 Unit time

Notes Second is the predefined SBML unit for time.

Definition s

3 Compartment

This model contains one compartment.

Table 2: Properties of all compartments.

Id	Name	SBO	Spatial Dimensions	Size	Unit	Constant	Outside
cell	cell		3	1	litre	<input checked="" type="checkbox"/>	

3.1 Compartment `cell`

This is a three dimensional compartment with a constant size of one litre.

Name `cell`

4 Species

This model contains 18 species. Section 8 provides further details and the derived rates of change of each species.

Table 3: Properties of each species.

Id	Name	Compartment	Derived Unit	Constant	Boundary Condition
CD95	CD95	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
FADD	FADD	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
p55free	p55free	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
Bid	Bid	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
PrNES_mCherry	PrNES_mCherry	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
PrER_mGFP	PrER_mGFP	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
DISC	DISC	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
DISCp55	DISCp55	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
p30	p30	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
p43	p43	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
p18	p18	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
p18inactive	p18inactive	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
tBid	tBid	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
PrNES	PrNES	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
mCherry	mCherry	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
PrER	PrER	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
mGFP	mGFP	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
CD95L	CD95L	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square

5 Parameters

This model contains 15 global parameters.

Table 4: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
kon_FADD			0.001		<input checked="" type="checkbox"/>
koff_FADD			0.001		<input checked="" type="checkbox"/>
kDISC			$3.64965874405544 \cdot 10^{-4}$		<input checked="" type="checkbox"/>
kD216			0.006		<input checked="" type="checkbox"/>
kD216trans- _p55			$2.23246421372882 \cdot 10^{-4}$		<input checked="" type="checkbox"/>
kD216trans- _p43			$5.29906975294056 \cdot 10^{-5}$		<input checked="" type="checkbox"/>
kD374			$6.44612643975149 \cdot 10^{-4}$		<input checked="" type="checkbox"/>
kD374trans- _p55			$5.43518631342483 \cdot 10^{-4}$		<input checked="" type="checkbox"/>
kD374trans- _p43			0.004		<input checked="" type="checkbox"/>
kdiss_p18			0.065		<input checked="" type="checkbox"/>
kBid			$5.2134055139547 \cdot 10^{-4}$		<input checked="" type="checkbox"/>
kD374probe			0.002		<input checked="" type="checkbox"/>
KDR			57.205		<input checked="" type="checkbox"/>
KDL			30.006		<input checked="" type="checkbox"/>
CD95act			0.000		<input type="checkbox"/>

6 Rule

This is an overview of one rule.

6.1 Rule CD95act

Rule CD95act is an assignment rule for parameter CD95act:

$$\begin{aligned}
 &\text{CD95act} && (1) \\
 = & \frac{[\text{CD95}]^3 \cdot \text{KDL}^2 \cdot [\text{CD95L}]}{([\text{CD95L}] + \text{KDL}) \cdot ([\text{CD95}]^2 \cdot \text{KDL}^2 + \text{KDR} \cdot [\text{CD95L}]^2 + 2 \cdot \text{KDR} \cdot \text{KDL} \cdot [\text{CD95L}] + \text{KDR} \cdot \text{KDL}^2)}
 \end{aligned}$$

7 Reactions

This model contains 19 reactions. All reactions are listed in the following table and are subsequently described in detail. If a reaction is affected by a modifier, the identifier of this species is written above the reaction arrow.

Table 5: Overview of all reactions

Nº	Id	Name	Reaction Equation	SBO
1	reaction_1		$\text{FADD} \xrightarrow{\text{FADD}} \text{DISC}$	
2	reaction_2		$\text{DISC} \xrightarrow{\text{DISC}} \text{FADD}$	
3	reaction_3		$\text{p55free} + \text{DISC} \xrightarrow{\text{p55free, DISC}} \text{DISCp55}$	
4	reaction_4		$\text{DISCp55} \xrightarrow{\text{DISCp55}} \text{p30}$	
5	reaction_5		$\text{DISCp55} \xrightarrow{\text{DISCp55, p30, DISCp55, p30}} \text{p30}$	
6	reaction_6		$\text{DISCp55} \xrightarrow{\text{p43, DISCp55, p43}} \text{p30}$	
7	reaction_7		$\text{p43} \xrightarrow{\text{p43}} \text{p18} + \text{DISC}$	
8	reaction_8		$\text{p43} \xrightarrow{\text{DISCp55, p30, p43, DISCp55, p30}} \text{p18} + \text{DISC}$	
9	reaction_9		$\text{p43} \xrightarrow{\text{p43, p43}} \text{p18} + \text{DISC}$	
10	reaction_10		$\text{DISCp55} \xrightarrow{\text{DISCp55}} \text{p43}$	
11	reaction_11		$\text{DISCp55} \xrightarrow{\text{DISCp55, p30, DISCp55, p30}} \text{p43}$	
12	reaction_12		$\text{DISCp55} \xrightarrow{\text{p43, DISCp55, p43}} \text{p43}$	
13	reaction_13		$\text{p30} \xrightarrow{\text{p30}} \text{p18} + \text{DISC}$	
14	reaction_14		$\text{p30} \xrightarrow{\text{DISCp55, p30, p30, DISCp55}} \text{p18} + \text{DISC}$	
15	reaction_15		$\text{p30} \xrightarrow{\text{p43, p30, p43}} \text{p18} + \text{DISC}$	

Nº	Id	Name	Reaction Equation	SBO
16	reaction_16		$p18 \xrightarrow{p18} p18_{inactive}$	
17	reaction_17		$Bid \xrightarrow{p43, p18, Bid, p43, p18} tBid$	
18	reaction_18		$PrNES_mCherry \xrightarrow{p43, p18, PrNES_mCherry, p43, p18} PrNES + mCherry$	
19	reaction_19		$PrER_mGFP \xrightarrow{p18, PrER_mGFP, p18} PrER + mGFP$	

7.1 Reaction [reaction_1](#)

This is an irreversible reaction of one reactant forming one product influenced by one modifier.

Reaction equation



Reactant

Table 6: Properties of each reactant.

Id	Name	SBO
FADD	FADD	

Modifier

Table 7: Properties of each modifier.

Id	Name	SBO
FADD	FADD	

Product

Table 8: Properties of each product.

Id	Name	SBO
DISC	DISC	

Kinetic Law

Derived unit contains undeclared units

$$v_1 = \text{kon_FADD} \cdot \text{CD95act} \cdot [\text{FADD}] \cdot \text{vol}(\text{cell}) \quad (3)$$

7.2 Reaction [reaction_2](#)

This is an irreversible reaction of one reactant forming one product influenced by one modifier.

Reaction equation



Reactant

Table 9: Properties of each reactant.

Id	Name	SBO
DISC	DISC	

Modifier

Table 10: Properties of each modifier.

Id	Name	SBO
DISC	DISC	

Product

Table 11: Properties of each product.

Id	Name	SBO
FADD	FADD	

Kinetic Law

Derived unit contains undeclared units

$$v_2 = \text{koff_FADD} \cdot [\text{DISC}] \cdot \text{vol}(\text{cell}) \quad (5)$$

7.3 Reaction `reaction_3`

This is an irreversible reaction of two reactants forming one product influenced by two modifiers.

Reaction equation



Reactants

Table 12: Properties of each reactant.

Id	Name	SBO
p55free	p55free	

Id	Name	SBO
DISC	DISC	

Modifiers

Table 13: Properties of each modifier.

Id	Name	SBO
p55free	p55free	
DISC	DISC	

Product

Table 14: Properties of each product.

Id	Name	SBO
DISCp55	DISCp55	

Kinetic Law


Derived unit contains undeclared units

$$v_3 = k_{\text{DISC}} \cdot [\text{p55free}] \cdot [\text{DISC}] \cdot \text{vol}(\text{cell}) \quad (7)$$

7.4 Reaction `reaction_4`

This is an irreversible reaction of one reactant forming one product influenced by one modifier.

Reaction equation



Reactant

Table 15: Properties of each reactant.

Id	Name	SBO
DISCp55	DISCp55	

Modifier

Table 16: Properties of each modifier.

Id	Name	SBO
DISCp55	DISCp55	

Product

Table 17: Properties of each product.

Id	Name	SBO
p30	p30	

Kinetic Law


Derived unit contains undeclared units

$$v_4 = kD216 \cdot [\text{DISCp55}] \cdot \text{vol}(\text{cell}) \quad (9)$$

7.5 Reaction `reaction_5`

This is an irreversible reaction of one reactant forming one product influenced by four modifiers.

Reaction equation



Reactant

Table 18: Properties of each reactant.

Id	Name	SBO
DISCp55	DISCp55	

Modifiers

Table 19: Properties of each modifier.

Id	Name	SBO
DISCp55	DISCp55	
p30	p30	
DISCp55	DISCp55	

Id	Name	SBO
p30	p30	

Product

Table 20: Properties of each product.

Id	Name	SBO
p30	p30	

Kinetic Law


Derived unit contains undeclared units

$$v_5 = \text{kD216trans.p55} \cdot [\text{DISCp55}] \cdot ([\text{DISCp55}] + [\text{p30}]) \cdot \text{vol}(\text{cell}) \quad (11)$$

7.6 Reaction `reaction_6`

This is an irreversible reaction of one reactant forming one product influenced by three modifiers.

Reaction equation



Reactant

Table 21: Properties of each reactant.

Id	Name	SBO
DISCp55	DISCp55	

Modifiers

Table 22: Properties of each modifier.

Id	Name	SBO
p43	p43	
DISCp55	DISCp55	
p43	p43	

Product

Table 23: Properties of each product.

Id	Name	SBO
p30	p30	

Kinetic Law

Derived unit contains undeclared units

$$v_6 = kD216trans_p43 \cdot [DISCp55] \cdot [p43] \cdot vol(cell) \quad (13)$$

7.7 Reaction `reaction_7`

This is an irreversible reaction of one reactant forming two products influenced by one modifier.

Reaction equation



Reactant

Table 24: Properties of each reactant.

Id	Name	SBO
p43	p43	

Modifier

Table 25: Properties of each modifier.

Id	Name	SBO
p43	p43	

Products

Table 26: Properties of each product.

Id	Name	SBO
p18	p18	

Id	Name	SBO
DISC	DISC	

Kinetic Law


Derived unit contains undeclared units

$$v_7 = kD216 \cdot [p43] \cdot \text{vol}(\text{cell}) \quad (15)$$

7.8 Reaction `reaction_8`

This is an irreversible reaction of one reactant forming two products influenced by five modifiers.

Reaction equation



Reactant

Table 27: Properties of each reactant.

Id	Name	SBO
p43	p43	

Modifiers

Table 28: Properties of each modifier.

Id	Name	SBO
DISCp55	DISCp55	
p30	p30	
p43	p43	
DISCp55	DISCp55	
p30	p30	

Products

Table 29: Properties of each product.

Id	Name	SBO
p18	p18	

Id	Name	SBO
DISC	DISC	

Kinetic Law

Derived unit contains undeclared units

$$v_8 = kD216trans.p55 \cdot [p43] \cdot ([DISCp55] + [p30]) \cdot vol(cell) \quad (17)$$

7.9 Reaction `reaction_9`

This is an irreversible reaction of one reactant forming two products influenced by two modifiers.

Reaction equation



Reactant

Table 30: Properties of each reactant.

Id	Name	SBO
p43	p43	

Modifiers

Table 31: Properties of each modifier.

Id	Name	SBO
p43	p43	
p43	p43	

Products

Table 32: Properties of each product.

Id	Name	SBO
p18	p18	
DISC	DISC	

Kinetic Law

Derived unit contains undeclared units

$$v_9 = kD216trans_p43 \cdot [p43] \cdot [p43] \cdot vol (cell) \quad (19)$$

7.10 Reaction `reaction_10`

This is an irreversible reaction of one reactant forming one product influenced by one modifier.

Reaction equation



Reactant

Table 33: Properties of each reactant.

Id	Name	SBO
DISCp55	DISCp55	

Modifier

Table 34: Properties of each modifier.

Id	Name	SBO
DISCp55	DISCp55	

Product

Table 35: Properties of each product.

Id	Name	SBO
p43	p43	

Kinetic Law


Derived unit contains undeclared units

$$v_{10} = kD374 \cdot [DISCp55] \cdot vol (cell) \quad (21)$$

7.11 Reaction `reaction_11`

This is an irreversible reaction of one reactant forming one product influenced by four modifiers.

Reaction equation



Reactant

Table 36: Properties of each reactant.

Id	Name	SBO
DISCp55	DISCp55	

Modifiers

Table 37: Properties of each modifier.

Id	Name	SBO
DISCp55	DISCp55	
p30	p30	
DISCp55	DISCp55	
p30	p30	

Product

Table 38: Properties of each product.

Id	Name	SBO
p43	p43	

Kinetic Law


Derived unit contains undeclared units

$$v_{11} = k_{D374trans_p55} \cdot [\text{DISCp55}] \cdot ([\text{DISCp55}] + [\text{p30}]) \cdot \text{vol}(\text{cell}) \quad (23)$$

7.12 Reaction `reaction_12`

This is an irreversible reaction of one reactant forming one product influenced by three modifiers.

Reaction equation



Reactant

Table 39: Properties of each reactant.

Id	Name	SBO
DISCp55	DISCp55	

Modifiers

Table 40: Properties of each modifier.

Id	Name	SBO
p43	p43	
DISCp55	DISCp55	
p43	p43	

Product

Table 41: Properties of each product.

Id	Name	SBO
p43	p43	

Kinetic Law

Derived unit contains undeclared units

$$v_{12} = kD374\text{trans_p43} \cdot [\text{DISCp55}] \cdot [\text{p43}] \cdot \text{vol}(\text{cell}) \quad (25)$$

7.13 Reaction `reaction_13`

This is an irreversible reaction of one reactant forming two products influenced by one modifier.

Reaction equation



Reactant

Table 42: Properties of each reactant.

Id	Name	SBO
p30	p30	

Modifier

Table 43: Properties of each modifier.

Id	Name	SBO
p30	p30	

Products

Table 44: Properties of each product.

Id	Name	SBO
p18	p18	
DISC	DISC	

Kinetic Law


Derived unit contains undeclared units

$$v_{13} = kD374 \cdot [p30] \cdot \text{vol}(\text{cell}) \quad (27)$$

7.14 Reaction `reaction_14`

This is an irreversible reaction of one reactant forming two products influenced by four modifiers.

Reaction equation



Reactant

Table 45: Properties of each reactant.

Id	Name	SBO
p30	p30	

Modifiers

Table 46: Properties of each modifier.

Id	Name	SBO
DISCp55	DISCp55	
p30	p30	
p30	p30	
DISCp55	DISCp55	

Products

Table 47: Properties of each product.

Id	Name	SBO
p18	p18	
DISC	DISC	

Kinetic Law

Derived unit contains undeclared units

$$v_{14} = kD374trans_p55 \cdot [p30] \cdot ([DISCp55] + [p30]) \cdot vol(cell) \quad (29)$$

7.15 Reaction `reaction_15`

This is an irreversible reaction of one reactant forming two products influenced by three modifiers.

Reaction equation



Reactant

Table 48: Properties of each reactant.

Id	Name	SBO
p30	p30	

Modifiers

Table 49: Properties of each modifier.

Id	Name	SBO
p43	p43	
p30	p30	
p43	p43	

Products

Table 50: Properties of each product.

Id	Name	SBO
p18	p18	
DISC	DISC	

Kinetic Law

Derived unit contains undeclared units

$$v_{15} = kD374trans_p43 \cdot [p30] \cdot [p43] \cdot vol(cell) \quad (31)$$

7.16 Reaction [reaction_16](#)

This is an irreversible reaction of one reactant forming one product influenced by one modifier.

Reaction equation



Reactant

Table 51: Properties of each reactant.

Id	Name	SBO
p18	p18	

Modifier

Table 52: Properties of each modifier.

Id	Name	SBO
p18	p18	

Product

Table 53: Properties of each product.

Id	Name	SBO
p18inactive	p18inactive	

Kinetic Law

Derived unit contains undeclared units

$$v_{16} = k_{\text{diss_p18}} \cdot [\text{p18}] \cdot \text{vol}(\text{cell}) \quad (33)$$

7.17 Reaction `reaction_17`

This is an irreversible reaction of one reactant forming one product influenced by five modifiers.

Reaction equation



Reactant

Table 54: Properties of each reactant.

Id	Name	SBO
Bid	Bid	

Modifiers

Table 55: Properties of each modifier.

Id	Name	SBO
p43	p43	
p18	p18	
Bid	Bid	
p43	p43	
p18	p18	

Product

Table 56: Properties of each product.

Id	Name	SBO
tBid	tBid	

Kinetic Law


Derived unit contains undeclared units

$$v_{17} = k_{\text{Bid}} \cdot [\text{Bid}] \cdot ([\text{p43}] + [\text{p18}]) \cdot \text{vol}(\text{cell}) \quad (35)$$

7.18 Reaction [reaction_18](#)

This is an irreversible reaction of one reactant forming two products influenced by five modifiers.

Reaction equation



Reactant

Table 57: Properties of each reactant.

Id	Name	SBO
PrNES_mCherry	PrNES_mCherry	

Modifiers

Table 58: Properties of each modifier.

Id	Name	SBO
p43	p43	
p18	p18	
PrNES_mCherry	PrNES_mCherry	
p43	p43	
p18	p18	

Products

Table 59: Properties of each product.

Id	Name	SBO
PrNES	PrNES	
mCherry	mCherry	

Kinetic Law


Derived unit contains undeclared units

$$v_{18} = \text{kD374probe} \cdot [\text{PrNES_mCherry}] \cdot ([\text{p43}] + [\text{p18}]) \cdot \text{vol}(\text{cell}) \quad (37)$$

7.19 Reaction [reaction_19](#)

This is an irreversible reaction of one reactant forming two products influenced by three modifiers.

Reaction equation



Reactant

Table 60: Properties of each reactant.

Id	Name	SBO
PrER_mGFP	PrER_mGFP	

Modifiers

Table 61: Properties of each modifier.

Id	Name	SBO
p18	p18	
PrER_mGFP	PrER_mGFP	
p18	p18	

Products

Table 62: Properties of each product.

Id	Name	SBO
PrER	PrER	
mGFP	mGFP	

Kinetic Law

Derived unit contains undeclared units

$$v_{19} = kD374probe \cdot [PrER_mGFP] \cdot [p18] \cdot vol(cell) \quad (39)$$

8 Derived Rate Equations

When interpreted as an ordinary differential equation framework, this model implies the following set of equations for the rates of change of each species.

Identifiers for kinetic laws highlighted in gray cannot be verified to evaluate to units of SBML substance per time. As a result, some SBML interpreters may not be able to verify the consistency of the units on quantities in the model. Please check if

- parameters without an unit definition are involved or
- volume correction is necessary because the `hasOnlySubstanceUnits` flag may be set to `false` and `spacialDimensions` > 0 for certain species.

8.1 Species CD95

Name CD95

Initial concentration $12 \text{ mol} \cdot \text{l}^{-1}$

This species does not take part in any reactions. Its quantity does hence not change over time:

$$\frac{d}{dt}CD95 = 0 \quad (40)$$

8.2 Species FADD

Name FADD

Initial concentration 90 mol · l⁻¹

This species takes part in three reactions (as a reactant in [reaction_1](#) and as a product in [reaction_2](#) and as a modifier in [reaction_1](#)).

$$\frac{d}{dt}\text{FADD} = v_2 - v_1 \quad (41)$$

8.3 Species p55free

Name p55free

Initial concentration 127 mol · l⁻¹

This species takes part in two reactions (as a reactant in [reaction_3](#) and as a modifier in [reaction_3](#)).

$$\frac{d}{dt}\text{p55free} = -v_3 \quad (42)$$

8.4 Species Bid

Name Bid

Initial concentration 224 mol · l⁻¹

This species takes part in two reactions (as a reactant in [reaction_17](#) and as a modifier in [reaction_17](#)).

$$\frac{d}{dt}\text{Bid} = -v_{17} \quad (43)$$

8.5 Species PrNES_mCherry

Name PrNES_mCherry

Initial concentration 1909 mol · l⁻¹

This species takes part in two reactions (as a reactant in [reaction_18](#) and as a modifier in [reaction_18](#)).

$$\frac{d}{dt}\text{PrNES_mCherry} = -v_{18} \quad (44)$$

8.6 Species PrER_mGFP

Name PrER_mGFP

Initial concentration 3316 mol · l⁻¹

This species takes part in two reactions (as a reactant in [reaction_19](#) and as a modifier in [reaction_19](#)).

$$\frac{d}{dt}\text{PrER_mGFP} = -v_{19} \quad (45)$$

8.7 Species DISC

Name DISC

Initial concentration 0 mol · l⁻¹

This species takes part in eleven reactions (as a reactant in [reaction_2](#), [reaction_3](#) and as a product in [reaction_1](#), [reaction_7](#), [reaction_8](#), [reaction_9](#), [reaction_13](#), [reaction_14](#), [reaction_15](#) and as a modifier in [reaction_2](#), [reaction_3](#)).

$$\frac{d}{dt}\text{DISC} = v_1 + v_7 + v_8 + v_9 + v_{13} + v_{14} + v_{15} - v_2 - v_3 \quad (46)$$

8.8 Species DISCp55

Name DISCp55

Initial concentration 0 mol · l⁻¹

This species takes part in 19 reactions (as a reactant in [reaction_4](#), [reaction_5](#), [reaction_6](#), [reaction_10](#), [reaction_11](#), [reaction_12](#) and as a product in [reaction_3](#) and as a modifier in [reaction_4](#), [reaction_5](#), [reaction_5](#), [reaction_6](#), [reaction_8](#), [reaction_8](#), [reaction_10](#), [reaction_11](#), [reaction_11](#), [reaction_12](#), [reaction_14](#), [reaction_14](#)).

$$\frac{d}{dt}\text{DISCp55} = v_3 - v_4 - v_5 - v_6 - v_{10} - v_{11} - v_{12} \quad (47)$$

8.9 Species p30

Name p30

Initial concentration 0 mol · l⁻¹

This species takes part in 16 reactions (as a reactant in [reaction_13](#), [reaction_14](#), [reaction_15](#) and as a product in [reaction_4](#), [reaction_5](#), [reaction_6](#) and as a modifier in [reaction_5](#), [reaction_5](#), [reaction_8](#), [reaction_8](#), [reaction_11](#), [reaction_11](#), [reaction_13](#), [reaction_14](#), [reaction_14](#), [reaction_15](#)).

$$\frac{d}{dt}\text{p30} = v_4 + v_5 + v_6 - v_{13} - v_{14} - v_{15} \quad (48)$$

8.10 Species p43

Name p43

Initial concentration 0 mol · l⁻¹

This species takes part in 20 reactions (as a reactant in [reaction_7](#), [reaction_8](#), [reaction_9](#) and as a product in [reaction_10](#), [reaction_11](#), [reaction_12](#) and as a modifier in [reaction_6](#), [reaction_6](#), [reaction_7](#), [reaction_8](#), [reaction_9](#), [reaction_9](#), [reaction_12](#), [reaction_12](#), [reaction_15](#), [reaction_15](#), [reaction_17](#), [reaction_17](#), [reaction_18](#), [reaction_18](#)).

$$\frac{d}{dt}p43 = v_{10} + v_{11} + v_{12} - v_7 - v_8 - v_9 \quad (49)$$

8.11 Species p18

Name p18

Initial concentration 0 mol · l⁻¹

This species takes part in 14 reactions (as a reactant in [reaction_16](#) and as a product in [reaction_7](#), [reaction_8](#), [reaction_9](#), [reaction_13](#), [reaction_14](#), [reaction_15](#) and as a modifier in [reaction_16](#), [reaction_17](#), [reaction_17](#), [reaction_18](#), [reaction_18](#), [reaction_19](#), [reaction_19](#)).

$$\frac{d}{dt}p18 = v_7 + v_8 + v_9 + v_{13} + v_{14} + v_{15} - v_{16} \quad (50)$$

8.12 Species p18inactive

Name p18inactive

Initial concentration 0 mol · l⁻¹

This species takes part in one reaction (as a product in [reaction_16](#)).

$$\frac{d}{dt}p18inactive = v_{16} \quad (51)$$

8.13 Species tBid

Name tBid

Initial concentration 0 mol · l⁻¹

This species takes part in one reaction (as a product in [reaction_17](#)).

$$\frac{d}{dt}tBid = v_{17} \quad (52)$$

8.14 Species PrNES

Name PrNES

Initial concentration $0 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in one reaction (as a product in [reaction_18](#)).

$$\frac{d}{dt}\text{PrNES} = v_{18} \quad (53)$$

8.15 Species mCherry

Name mCherry

Initial concentration $0 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in one reaction (as a product in [reaction_18](#)).

$$\frac{d}{dt}\text{mCherry} = v_{18} \quad (54)$$

8.16 Species PrER

Name PrER

Initial concentration $0 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in one reaction (as a product in [reaction_19](#)).

$$\frac{d}{dt}\text{PrER} = v_{19} \quad (55)$$

8.17 Species mGFP

Name mGFP

Initial concentration $0 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in one reaction (as a product in [reaction_19](#)).

$$\frac{d}{dt}\text{mGFP} = v_{19} \quad (56)$$

8.18 Species CD95L

Name CD95L

Initial concentration $16.6 \text{ mol} \cdot \text{l}^{-1}$

This species does not take part in any reactions. Its quantity does hence not change over time:

$$\frac{d}{dt}\text{CD95L} = 0 \quad (57)$$

SBML²TeX was developed by Andreas Dräger^a, Hannes Planatscher^a, Dieudonné M Wouamba^a, Adrian Schröder^a, Michael Hucka^b, Lukas Endler^c, Martin Golebiewski^d and Andreas Zell^a. Please see <http://www.ra.cs.uni-tuebingen.de/software/SBML2LaTeX> for more information.

^aCenter for Bioinformatics Tübingen (ZBIT), Germany

^bCalifornia Institute of Technology, Beckman Institute BNMC, Pasadena, United States

^cEuropean Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, United Kingdom

^dEML Research gGmbH, Heidelberg, Germany