

SBML Model Report

Model name:
“Fujita2010_Akt_Signalling_EGF”



May 5, 2016

1 General Overview

This is a document in SBML Level 2 Version 4 format. This model was created by the following two authors: Lukas Endler¹ and Kazuhiro Fujita² at August 24th 2010 at 11:49 a. m. and last time modified at February 21st 2014 at 11:14 a. m. Table 1 provides an overview of the quantities of all components of this model.

Table 1: Number of components in this model, which are described in the following sections.

Element	Quantity	Element	Quantity
compartment types	0	compartments	1
species types	0	species	11
events	0	constraints	0
reactions	11	function definitions	0
global parameters	12	unit definitions	8
rules	4	initial assignments	1

Model Notes

EGF dependent Akt pathway model
made by Kazuhiro A. Fujita.

This is the EGF dependent Akt pathway model described in:

Decoupling of receptor and downstream signals in the Akt pathway by its low-pass filter

¹EMBL-EBI, lukas@ebi.ac.uk

²University of Tokyo, kazuhiro.fujita@gmail.com

characteristics.

Fujita KA, Toyoshima Y, Uda S, Ozaki Y, Kubota H, and Kuroda S. *Sci Signal*. 2010 Jul 27;3(132):ra56. PMID: [20664065](#) ; DOI: [10.1126/scisignal.2000810](#)

Abstract:

In cellular signal transduction, the information in an external stimulus is encoded in temporal patterns in the activities of signaling molecules; for example, pulses of a stimulus may produce an increasing response or may produce pulsatile responses in the signaling molecules. Here, we show how the Akt pathway, which is involved in cell growth, specifically transmits temporal information contained in upstream signals to downstream effectors. We modeled the epidermal growth factor (EGF)dependent Akt pathway in PC12 cells on the basis of experimental results. We obtained counterintuitive results indicating that the sizes of the peak amplitudes of receptor and downstream effector phosphorylation were decoupled; weak, sustained EGF receptor (EGFR) phosphorylation, rather than strong, transient phosphorylation, strongly induced phosphorylation of the ribosomal protein S6, a molecule downstream of Akt. Using frequency response analysis, we found that a three-component Akt pathway exhibited the property of a low-pass filter and that this property could explain decoupling of the peak amplitudes of receptor phosphorylation and that of downstream effectors. Furthermore, we found that lapatinib, an EGFR inhibitor used as an anticancer drug, converted strong, transient Akt phosphorylation into weak, sustained Akt phosphorylation, and, because of the low-pass filter characteristics of the Akt pathway, this led to stronger S6 phosphorylation than occurred in the absence of the inhibitor. Thus, an EGFR inhibitor can potentially act as a downstream activator of some effectors.

The different versions of input, step, pulse and ramp, can be simulated using the parameters EGF_conc_pulse , EGF_conc_step and EGF_conc_ramp . Depending on which one is set unequal to 0, either a continuous pulse with value EGF_conc_pulse , a 60 second step with EGF_conc_step or a signal increasing from 0 to EGF_conc_pulse over a time period of 3600 seconds are used as input. In case more than one parameter are set to values greater than 0 these input profiles are added to each other. The pulse time and the time over which the ramp input increases can be set by pulse_time and ramp_time .

This model originates from BioModels Database: A Database of Annotated Published Models. It is copyright (c) 2005-2010 The BioModels Team.

For more information see the [terms of use](#) .

To cite BioModels Database, please use [Le Novre N., Bornstein B., Broicher A., Courtot M., Donizelli M., Dharuri H., Li L., Sauro H., Schilstra M., Shapiro B., Snoep J.L., Hucka M. \(2006\) BioModels Database: A Free, Centralized Database of Curated, Published, Quantitative Kinetic Models of Biochemical and Cellular Systems Nucleic Acids Res., 34: D689-D691.](#)

2 Unit Definitions

This is an overview of ten unit definitions of which two are predefined by SBML and not mentioned in the model.

2.1 Unit substance

Name arbitrary_amount

Definition dimensionless

2.2 Unit conc

Name arbitrary_conc

Definition dimensionless · ml⁻¹

2.3 Unit time

Name seconds

Definition s

2.4 Unit volume

Name ml

Definition ml

2.5 Unit per_sec

Name per second

Definition s⁻¹

2.6 Unit ng

Name ng

Definition ng

2.7 Unit ng_per_ml

Name ng_per_ml

Definition ng · ml⁻¹

2.8 Unit per_conc_per_sec

Name per conc per second

Definition ml · dimensionless · s⁻¹

2.9 Unit area

Notes Square metre is the predefined SBML unit for area since SBML Level 2 Version 1.

Definition m^2

2.10 Unit length

Notes Metre is the predefined SBML unit for length since SBML Level 2 Version 1.

Definition m

3 Compartment

This model contains one compartment.

Table 2: Properties of all compartments.

Id	Name	SBO	Spatial Dimensions	Size	Unit	Constant	Outside
Cell	Cell	0000290	3	1	litre	<input checked="" type="checkbox"/>	

3.1 Compartment Cell

This is a three dimensional compartment with a constant size of one ml.

Name Cell

SBO:0000290 physical compartment

4 Species

This model contains eleven species. The boundary condition of two of these species is set to `true` so that these species' amount cannot be changed by any reaction. Section 9 provides further details and the derived rates of change of each species.

Table 3: Properties of each species.

Id	Name	Compartment	Derived Unit	Constant	Boundary Condition
EGF	EGF	Cell	$\text{ng} \cdot \text{ml}^{-1}$	<input type="checkbox"/>	<input checked="" type="checkbox"/>
EGFR	EGFR	Cell	dimensionless ml^{-1}	<input type="checkbox"/>	<input type="checkbox"/>
pEGFR	pEGFR	Cell	dimensionless ml^{-1}	<input type="checkbox"/>	<input type="checkbox"/>
pEGFR_Akt	pEGFR_Akt	Cell	dimensionless ml^{-1}	<input type="checkbox"/>	<input type="checkbox"/>
Akt	Akt	Cell	dimensionless ml^{-1}	<input type="checkbox"/>	<input type="checkbox"/>
pAkt	pAkt	Cell	dimensionless ml^{-1}	<input type="checkbox"/>	<input type="checkbox"/>
S6	S6	Cell	dimensionless ml^{-1}	<input type="checkbox"/>	<input type="checkbox"/>
pAkt_S6	pAkt_S6	Cell	dimensionless ml^{-1}	<input type="checkbox"/>	<input type="checkbox"/>
pS6	pS6	Cell	dimensionless ml^{-1}	<input type="checkbox"/>	<input type="checkbox"/>
pro_EGFR	pro_EGFR	Cell	dimensionless ml^{-1}	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
EGF_EGFR	EGF_EGFR	Cell	dimensionless ml^{-1}	<input type="checkbox"/>	<input type="checkbox"/>

Id	Name	Compartment	Derived Unit	Constant	Boundary Condi- tion

5 Parameters

This model contains twelve global parameters.

Table 4: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
pEGFR_total	pEGFR_total		0.000	ng · ml ⁻¹	<input type="checkbox"/>
pAkt_total	pAkt_total		0.000	ng · ml ⁻¹	<input type="checkbox"/>
pEGFR- _scaleFactor	pEGFR- _scaleFactor	1.81734813832032 · 10 ⁻⁴		ng	<input checked="" type="checkbox"/>
pAkt- _scaleFactor	pAkt_scaleFactor		60.059	ng	<input checked="" type="checkbox"/>
pS6- _scaleFactor	pS6_scaleFactor		49886.231	ng	<input checked="" type="checkbox"/>
pS6_total	pS6_total		0.000	ng · ml ⁻¹	<input type="checkbox"/>
EGF_conc- _step	EGF_conc_step		0.000	ng · ml ⁻¹	<input checked="" type="checkbox"/>
EGF_conc- _impulse	EGF_conc_impulse		0.000	ng · ml ⁻¹	<input checked="" type="checkbox"/>
EGF_conc- _ramp	EGF_conc_ramp		30.000	ng · ml ⁻¹	<input checked="" type="checkbox"/>
EGFR- _turnover	EGFR_turnover	1.06386129269658 · 10 ⁻⁴		s ⁻¹	<input checked="" type="checkbox"/>
pulse_time	pulse_time		60.000	s	<input checked="" type="checkbox"/>
ramp_time	ramp_time		3600.000	s	<input checked="" type="checkbox"/>

6 Initialassignment

This is an overview of one initialassignment.

6.1 Initialassignment EGFR

Derived unit ml⁻¹

Math [pro_EGFR]

7 Rules

This is an overview of four rules.

7.1 Rule EGF

Rule EGF is an assignment rule for species EGF:

$$\text{EGF} = \text{EGF_conc_step} + \begin{cases} \text{EGF_conc_impulse} & \text{if } \text{time} \leq \text{pulse_time} \\ 0 & \text{otherwise} \end{cases} + \frac{\text{EGF_conc_ramp} \cdot \text{time}}{\text{ramp_time}} \quad (1)$$

7.2 Rule pEGFR_total

Rule pEGFR_total is an assignment rule for parameter pEGFR_total:

$$\text{pEGFR_total} = ([\text{pEGFR}] + [\text{pEGFR_Akt}]) \cdot \text{pEGFR_scaleFactor} \quad (2)$$

Derived unit $\text{ml}^{-1} \cdot \text{ng}$

7.3 Rule pAkt_total

Rule pAkt_total is an assignment rule for parameter pAkt_total:

$$\text{pAkt_total} = ([\text{pAkt}] + [\text{pAkt_S6}]) \cdot \text{pAkt_scaleFactor} \quad (3)$$

Derived unit $\text{ml}^{-1} \cdot \text{ng}$

7.4 Rule pS6_total

Rule pS6_total is an assignment rule for parameter pS6_total:

$$\text{pS6_total} = [\text{pS6}] \cdot \text{pS6_scaleFactor} \quad (4)$$

Derived unit $\text{ml}^{-1} \cdot \text{ng}$

8 Reactions

This model contains eleven reactions. All reactions are listed in the following table and are subsequently described in detail. If a reaction is affected by a modifier, the identifier of this species is written above the reaction arrow.

Table 5: Overview of all reactions

Nº	Id	Name	Reaction Equation	SBO
1	reaction_1	EGF+EGFR	$\text{EGF} + \text{EGFR} \rightleftharpoons \text{EGF_EGFR}$	0000177
2	reaction_2	pEGFR+Akt	$\text{pEGFR} + \text{Akt} \rightleftharpoons \text{pEGFR_Akt}$	0000177
3	reaction_3	Akt_phosphorylation	$\text{pEGFR_Akt} \longrightarrow \text{pEGFR} + \text{pAkt}$	0000216
4	reaction_4	pEGFR_degradation	$\text{pEGFR} \longrightarrow \emptyset$	0000179
5	reaction_5	pAkt+S6	$\text{pAkt} + \text{S6} \rightleftharpoons \text{pAkt_S6}$	0000179
6	reaction_6	S6_phosphorylation	$\text{pAkt_S6} \longrightarrow \text{pAkt} + \text{pS6}$	0000179
7	reaction_7	pAkt_dephospho	$\text{pAkt} \longrightarrow \text{Akt}$	0000330
8	reaction_8	pS6_dephospho	$\text{pS6} \longrightarrow \text{S6}$	0000330
9	reaction_9	EGFR_synthesis	$\text{pro_EGFR} \longrightarrow \text{EGFR}$	0000184
10	reaction_10	EGFR_phosphorylation	$\text{EGF_EGFR} \longrightarrow \text{pEGFR}$	0000179
11	reaction_11	EGFR_degradation	$\text{EGFR} \longrightarrow \emptyset$	0000179

8.1 Reaction [reaction_1](#)

This is a reversible reaction of two reactants forming one product.

Name EGF+EGFR

SBO:0000177 non-covalent binding

Reaction equation



Reactants

Table 6: Properties of each reactant.

Id	Name	SBO
EGF	EGF	
EGFR	EGFR	

Product

Table 7: Properties of each product.

Id	Name	SBO
EGF_EGFR	EGF_EGFR	

Kinetic Law

Derived unit contains undeclared units

$$v_1 = \text{vol}(\text{Cell}) \cdot (k_1 \cdot [\text{EGF}] \cdot [\text{EGFR}] - k_2 \cdot [\text{EGF_EGFR}]) \quad (6)$$

Table 8: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1	k1		0.007		✓
k2	k2		0.041	s ⁻¹	✓


8.2 Reaction [reaction_2](#)

This is a reversible reaction of two reactants forming one product.

Name pEGFR+Akt

SBO:0000177 non-covalent binding

Reaction equation



Reactants

Table 9: Properties of each reactant.

Id	Name	SBO
pEGFR	pEGFR	
Akt	Akt	

Product

Table 10: Properties of each product.

Id	Name	SBO
pEGFR_Akt	pEGFR_Akt	

Kinetic Law

Derived unit s^{-1}

$$v_2 = \text{vol}(\text{Cell}) \cdot (k_1 \cdot [\text{pEGFR}] \cdot [\text{Akt}] - k_2 \cdot [\text{pEGFR_Akt}]) \quad (8)$$

Table 11: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1	k1		$1.5543 \cdot 10^{-5}$	$\text{ml} \cdot \text{dimensionless} \cdot \text{s}^{-1}$	<input checked="" type="checkbox"/>
k2	k2		0.005	s^{-1}	<input checked="" type="checkbox"/>


8.3 Reaction reaction_3

This is an irreversible reaction of one reactant forming two products.

Name Akt_phosphorylation

SBO:0000216 phosphorylation

Reaction equation



Reactant

Table 12: Properties of each reactant.

Id	Name	SBO
pEGFR_Akt	pEGFR_Akt	

Products

Table 13: Properties of each product.

Id	Name	SBO
pEGFR	pEGFR	
pAkt	pAkt	

Kinetic Law

Derived unit s^{-1}

$$v_3 = \text{vol}(\text{Cell}) \cdot k1 \cdot [\text{pEGFR_Akt}] \quad (10)$$

Table 14: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1	k1		0.031	s^{-1}	<input checked="" type="checkbox"/>

8.4 Reaction [reaction_4](#)

This is an irreversible reaction of one reactant forming no product.

Name pEGFR_degradation

SBO:0000179 degradation

Reaction equation



Reactant

Table 15: Properties of each reactant.

Id	Name	SBO
pEGFR	pEGFR	

Kinetic Law

Derived unit s^{-1}

$$v_4 = \text{vol}(\text{Cell}) \cdot k_1 \cdot [\text{pEGFR}] \quad (12)$$

Table 16: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1	k1		0.100	s^{-1}	<input checked="" type="checkbox"/>

8.5 Reaction `reaction_5`

This is a reversible reaction of two reactants forming one product.

Name pAkt+S6

SBO:0000179 degradation

Reaction equation



Reactants

Table 17: Properties of each reactant.

Id	Name	SBO
pAkt	pAkt	
S6	S6	

Product

Table 18: Properties of each product.

Id	Name	SBO
pAkt_S6	pAkt_S6	

Kinetic Law

Derived unit s^{-1}

$$v_5 = \text{vol}(\text{Cell}) \cdot (k_1 \cdot [\text{pAkt}] \cdot [\text{S6}] - k_2 \cdot [\text{pAkt_S6}]) \quad (14)$$

Table 19: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1	k1		$2.10189 \cdot 10^{-6}$	$\text{ml} \cdot \text{dimensionless} \cdot \text{s}^{-1}$	<input checked="" type="checkbox"/>
k2	k2		$5.1794 \cdot 10^{-15}$	s^{-1}	<input checked="" type="checkbox"/>

8.6 Reaction `reaction_6`

This is an irreversible reaction of one reactant forming two products.

Name S6_phosphorylation

SBO:0000179 degradation

Reaction equation



Reactant

Table 20: Properties of each reactant.

Id	Name	SBO
pAkt_S6	pAkt_S6	

Products

Table 21: Properties of each product.

Id	Name	SBO
pAkt	pAkt	
pS6	pS6	

Kinetic Law

Derived unit s^{-1}

$$v_6 = \text{vol}(\text{Cell}) \cdot k1 \cdot [\text{pAkt_S6}] \quad (16)$$

Table 22: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1	k1		0.001	s^{-1}	<input checked="" type="checkbox"/>

8.7 Reaction [reaction_7](#)

This is an irreversible reaction of one reactant forming one product.

Name pAkt_dephospho

SBO:0000330 dephosphorylation

Reaction equation



Reactant

Table 23: Properties of each reactant.

Id	Name	SBO
pAkt	pAkt	

Product

Table 24: Properties of each product.

Id	Name	SBO
Akt	Akt	

Id	Name	SBO
----	------	-----

Kinetic Law

Derived unit s^{-1}

$$v_7 = \text{vol}(\text{Cell}) \cdot k_1 \cdot [\text{pAkt}] \quad (18)$$

Table 25: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1	k1		0.033	s^{-1}	<input checked="" type="checkbox"/>

8.8 Reaction [reaction_8](#)

This is an irreversible reaction of one reactant forming one product.

Name pS6.dephospho

SBO:0000330 dephosphorylation

Reaction equation



Reactant

Table 26: Properties of each reactant.

Id	Name	SBO
pS6	pS6	

Product

Table 27: Properties of each product.

Id	Name	SBO
S6	S6	

Kinetic Law

Derived unit s^{-1}

$$v_8 = \text{vol}(\text{Cell}) \cdot k1 \cdot [\text{pS6}] \quad (20)$$

Table 28: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1	k1		0.001	s^{-1}	<input checked="" type="checkbox"/>

8.9 Reaction `reaction_9`

This is an irreversible reaction of one reactant forming one product.

Name EGFR_synthesis

SBO:0000184 translation

Reaction equation



Reactant

Table 29: Properties of each reactant.

Id	Name	SBO
pro_EGFR	pro_EGFR	

Product

Table 30: Properties of each product.

Id	Name	SBO
EGFR	EGFR	

Kinetic Law

Derived unit s^{-1}

$$v_9 = \text{vol}(\text{Cell}) \cdot \text{EGFR_turnover} \cdot [\text{pro_EGFR}] \quad (22)$$

8.10 Reaction [reaction_10](#)

This is an irreversible reaction of one reactant forming one product.

Name EGFR_phosphorylation

SBO:0000179 degradation

Reaction equation



Reactant

Table 31: Properties of each reactant.

Id	Name	SBO
EGF_EGFR	EGF_EGFR	

Product

Table 32: Properties of each product.

Id	Name	SBO
pEGFR	pEGFR	

Kinetic Law

Derived unit s^{-1}

$$v_{10} = \text{vol}(\text{Cell}) \cdot k_1 \cdot [\text{EGF_EGFR}] \quad (24)$$

Table 33: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
k1	k1		0.019	s^{-1}	<input checked="" type="checkbox"/>


8.11 Reaction [reaction_11](#)

This is an irreversible reaction of one reactant forming no product.

Name EGFR_degradation

SBO:0000179 degradation

Reaction equation



Reactant

Table 34: Properties of each reactant.

Id	Name	SBO
EGFR	EGFR	

Kinetic Law

Derived unit s^{-1}

$$v_{11} = \text{vol}(\text{Cell}) \cdot \text{EGFR_turnover} \cdot [\text{EGFR}] \quad (26)$$

9 Derived Rate Equations

When interpreted as an ordinary differential equation framework, this model implies the following set of equations for the rates of change of each species.

Identifiers for kinetic laws highlighted in gray cannot be verified to evaluate to units of SBML substance per time. As a result, some SBML interpreters may not be able to verify the consistency of the units on quantities in the model. Please check if

- parameters without an unit definition are involved or
- volume correction is necessary because the `hasOnlySubstanceUnits` flag may be set to `false` and `spacialDimensions` > 0 for certain species.

9.1 Species EGF

Name EGF

SBO:0000252 polypeptide chain

Initial amount 0 *Unknownunitng*

Involved in rule EGF

This species takes part in one reaction (as a reactant in [reaction_1](#)). Not this but one rule determines the species' quantity because this species is on the boundary of the reaction system.

9.2 Species EGFR

Name EGFR

SBO:0000297 protein complex

Initial amount 68190.1837333797 dimensionless

Initial assignment EGFR

This species takes part in three reactions (as a reactant in [reaction_1](#), [reaction_11](#) and as a product in [reaction_9](#)).

$$\frac{d}{dt}\text{EGFR} = v_9 - v_1 - v_{11} \quad (27)$$

9.3 Species pEGFR

Name pEGFR

SBO:0000297 protein complex

Initial amount 0 dimensionless

This species takes part in four reactions (as a reactant in [reaction_2](#), [reaction_4](#) and as a product in [reaction_3](#), [reaction_10](#)).

$$\frac{d}{dt}\text{pEGFR} = v_3 + v_{10} - v_2 - v_4 \quad (28)$$

9.4 Species pEGFR_Akt

Name pEGFR_Akt

SBO:0000297 protein complex

Initial amount 0 dimensionless

This species takes part in two reactions (as a reactant in [reaction_3](#) and as a product in [reaction_2](#)).

$$\frac{d}{dt}\text{pEGFR_Akt} = v_2 - v_3 \quad (29)$$

9.5 Species Akt

Name Akt

SBO:0000252 polypeptide chain

Initial amount 0.0433090165709309 dimensionless

This species takes part in two reactions (as a reactant in [reaction_2](#) and as a product in [reaction_7](#)).

$$\frac{d}{dt}Akt = v_7 - v_2 \quad (30)$$

9.6 Species pAkt

Name pAkt

SBO:0000252 polypeptide chain

Initial amount 0 dimensionless

This species takes part in four reactions (as a reactant in [reaction_5](#), [reaction_7](#) and as a product in [reaction_3](#), [reaction_6](#)).

$$\frac{d}{dt}pAkt = v_3 + v_6 - v_5 - v_7 \quad (31)$$

9.7 Species S6

Name S6

SBO:0000252 polypeptide chain

Initial amount 3.54316740542218 dimensionless

This species takes part in two reactions (as a reactant in [reaction_5](#) and as a product in [reaction_8](#)).

$$\frac{d}{dt}S6 = v_8 - v_5 \quad (32)$$

9.8 Species pAkt_S6

Name pAkt_S6

SBO:0000297 protein complex

Initial amount 0 dimensionless

This species takes part in two reactions (as a reactant in [reaction_6](#) and as a product in [reaction_5](#)).

$$\frac{d}{dt}pAkt.S6 = v_5 - v_6 \quad (33)$$

9.9 Species pS6

Name pS6

SBO:0000252 polypeptide chain

Initial amount 0 dimensionless

This species takes part in two reactions (as a reactant in [reaction_8](#) and as a product in [reaction_6](#)).

$$\frac{d}{dt} \text{pS6} = v_6 - v_8 \quad (34)$$

9.10 Species pro_EGFR

Name pro_EGFR

SBO:0000297 protein complex

Initial amount 68190.1837333797 dimensionless

This species takes part in one reaction (as a reactant in [reaction_9](#)), which does not influence its rate of change because this constant species is on the boundary of the reaction system:

$$\frac{d}{dt} \text{pro_EGFR} = 0 \quad (35)$$

9.11 Species EGF_EGFR

Name EGF_EGFR

SBO:0000297 protein complex

Initial amount 0 dimensionless

This species takes part in two reactions (as a reactant in [reaction_10](#) and as a product in [reaction_1](#)).

$$\frac{d}{dt} \text{EGF_EGFR} = v_1 - v_{10} \quad (36)$$

A Glossary of Systems Biology Ontology Terms

SBO:0000177 non-covalent binding: Interaction between several biochemical entities that results in the formation of a non-covalent complex

SBO:0000179 degradation: Complete disappearance of a physical entity

- SBO:0000184 translation:** Process in which a polypeptide chain is produced from a messenger RNA
- SBO:0000216 phosphorylation:** Addition of a phosphate group ($\text{-H}_2\text{PO}_4$) to a chemical entity
- SBO:0000252 polypeptide chain:** Naturally occurring macromolecule formed by the repetition of amino-acid residues linked by peptidic bonds. A polypeptide chain is synthesized by the ribosome. CHEBI:1654
- SBO:0000290 physical compartment:** Specific location of space, that can be bounded or not. A physical compartment can have 1, 2 or 3 dimensions
- SBO:0000297 protein complex:** Macromolecular complex containing one or more polypeptide chains possibly associated with simple chemicals. CHEBI:3608
- SBO:0000330 dephosphorylation:** Removal of a phosphate group ($\text{-H}_2\text{PO}_4$) from a chemical entity.

SBML²TeX was developed by Andreas Dräger^a, Hannes Planatscher^a, Dieudonné M Wouamba^a, Adrian Schröder^a, Michael Hucka^b, Lukas Endler^c, Martin Golebiewski^d and Andreas Zell^a. Please see <http://www.ra.cs.uni-tuebingen.de/software/SBML2LaTeX> for more information.

^aCenter for Bioinformatics Tübingen (ZBIT), Germany

^bCalifornia Institute of Technology, Beckman Institute BNMC, Pasadena, United States

^cEuropean Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, United Kingdom

^dEML Research gGmbH, Heidelberg, Germany