

SBML Model Report

Model name:
“Clancy2002_CardiacSodiumChannel_WT”



May 6, 2016

1 General Overview

This is a document in SBML Level 2 Version 1 format. This model was created by Enuo He¹ at July 15th 2007 at 1:35 p. m. and last time modified at October tenth 2014 at 10:23 a. m. Table 1 gives an overview of the quantities of all components of this model.

Table 1: Number of components in this model, which are described in the following sections.

Element	Quantity	Element	Quantity
compartment types	0	compartments	1
species types	0	species	9
events	2	constraints	0
reactions	11	function definitions	0
global parameters	24	unit definitions	2
rules	16	initial assignments	0

Model Notes

The model is according to the paper *Na⁺ Channel Mutation That Causes Both Brugada and Long-QT Syndrome Phenotypes: A Simulation Study of Mechanism* Original model comes from ModelDB with accession number: 62661. This is the wide type model. All the values and reactions obtained from Data Supplement6: Appendix of the paper. Figure3 has been reproduced by MathSBML. The stimulus $v = -30\text{mV}$ during the time from 5ms to 20 ms displayed in the event.

¹BNMC, enuo@caltech.edu

The meaning for the keyword, C: Close states; O: Open states; IF: Fast inactivation states; IC: Closed-Inactivation states; IM: Intermediat Inactivation states.

To the extent possible under law, all copyright and related or neighbouring rights to this encoded model have been dedicated to the public domain worldwide. Please refer to [CC0 Public Domain Dedication](#) for more information.

In summary, you are entitled to use this encoded model in absolutely any manner you deem suitable, verbatim, or with modification, alone or embedded it in a larger context, redistribute it, commercially or not, in a restricted way or not.

To cite BioModels Database, please use: [Li C, Donizelli M, Rodriguez N, Dharuri H, Endler L, Chelliah V, Li L, He E, Henry A, Stefan MI, Snoep JL, Hucka M, Le Novre N, Laibe C \(2010\) BioModels Database: An enhanced, curated and annotated resource for published quantitative kinetic models. BMC Syst Biol., 4:92.](#)

2 Unit Definitions

This is an overview of seven unit definitions of which five are predefined by SBML and not mentioned in the model.

2.1 Unit `unitDefinition_0000001`

Name time(ms)

Definition ms

2.2 Unit `unitDefinition_0000002`

Name voltage(mV)

Definition mV

2.3 Unit `substance`

Notes Mole is the predefined SBML unit for substance.

Definition mol

2.4 Unit `volume`

Notes Litre is the predefined SBML unit for volume.

Definition l

2.5 Unit `area`

Notes Square metre is the predefined SBML unit for `area` since SBML Level 2 Version 1.

Definition m^2

2.6 Unit `length`

Notes Metre is the predefined SBML unit for `length` since SBML Level 2 Version 1.

Definition m

2.7 Unit `time`

Notes Second is the predefined SBML unit for `time`.

Definition s

3 Compartment

This model contains one compartment.

Table 2: Properties of all compartments.

Id	Name	SBO	Spatial Dimensions	Size	Unit	Constant	Outside
cell			3	1	litre	<input checked="" type="checkbox"/>	

3.1 Compartment `cell`

This is a three dimensional compartment with a constant size of one litre.

4 Species

This model contains nine species. Section 9 provides further details and the derived rates of change of each species.

Table 3: Properties of each species.

Id	Name	Compartment	Derived Unit	Constant	Boundary Condi- tion
C1	open states	cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
C2		cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
C3		cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
IC3		cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
IC2		cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
IM1		cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
IM2		cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
O		cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square
IF		cell	$\text{mol} \cdot \text{l}^{-1}$	\square	\square

5 Parameters

This model contains 24 global parameters.

Table 4: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
nao	[Na]out		140.0		<input checked="" type="checkbox"/>
nai	[Na]in		15.0		<input checked="" type="checkbox"/>
scale			1000.0		<input checked="" type="checkbox"/>
v			−80.0		<input type="checkbox"/>
Ena			0.0		<input type="checkbox"/>
a11			0.0		<input type="checkbox"/>
a12			0.0		<input type="checkbox"/>
a13			0.0		<input type="checkbox"/>
b11			0.0		<input type="checkbox"/>
b12			0.0		<input type="checkbox"/>
b13			0.0		<input type="checkbox"/>
a2			0.0		<input type="checkbox"/>
b2			0.0		<input type="checkbox"/>
a3			0.0		<input type="checkbox"/>
b3			0.0		<input type="checkbox"/>
a4			0.0		<input type="checkbox"/>
b4			0.0		<input type="checkbox"/>
a5			0.0		<input type="checkbox"/>
b5			0.0		<input type="checkbox"/>
Gna			23.5		<input checked="" type="checkbox"/>
Rk			8314.0		<input checked="" type="checkbox"/>
Fara			96485.0		<input checked="" type="checkbox"/>
Temp			310.0		<input checked="" type="checkbox"/>
Ina	I_Na		0.0		<input type="checkbox"/>

6 Rules

This is an overview of 16 rules.

6.1 Rule a11

Rule a11 is an assignment rule for parameter a11:

$$a11 = \frac{3.802}{0.1027 \cdot \exp\left(\frac{v}{17}\right) + 0.2 \cdot \exp\left(\frac{v}{150}\right)} \quad (1)$$

6.2 Rule a12

Rule a12 is an assignment rule for parameter a12:

$$a12 = \frac{3.802}{0.1027 \cdot \exp\left(\frac{v}{15}\right) + 0.23 \cdot \exp\left(\frac{v}{150}\right)} \quad (2)$$

6.3 Rule a13

Rule a13 is an assignment rule for parameter a13:

$$a13 = \frac{3.802}{0.1027 \cdot \exp\left(\frac{v}{12}\right) + 0.25 \cdot \exp\left(\frac{v}{150}\right)} \quad (3)$$

6.4 Rule a2

Rule a2 is an assignment rule for parameter a2:

$$a2 = 9.178 \cdot \exp\left(\frac{v}{29.68}\right) \quad (4)$$

6.5 Rule a3

Rule a3 is an assignment rule for parameter a3:

$$a3 = 3.7933 \cdot 1.0E - 7 \cdot \exp\left(\frac{v}{7.7}\right) \quad (5)$$

6.6 Rule b3

Rule b3 is an assignment rule for parameter b3:

$$b3 = 0.0084 + 2.0E - 5 \cdot v \quad (6)$$

6.7 Rule a4

Rule a4 is an assignment rule for parameter a4:

$$a4 = \frac{a2}{100} \quad (7)$$

6.8 Rule b4

Rule b4 is an assignment rule for parameter b4:

$$b4 = a3 \quad (8)$$

6.9 Rule a5

Rule a5 is an assignment rule for parameter a5:

$$a5 = \frac{a2}{9.5 \cdot 10000} \quad (9)$$

6.10 Rule b5

Rule b5 is an assignment rule for parameter b5:

$$b5 = \frac{a3}{50} \quad (10)$$

6.11 Rule Ena

Rule Ena is an assignment rule for parameter Ena:

$$Ena = \frac{Rk \cdot Temp}{Fara} \cdot \left(\frac{nao}{nai} \right) \quad (11)$$

6.12 Rule Ina

Rule Ina is an assignment rule for parameter Ina:

$$Ina = \frac{Gna \cdot \frac{[O]}{[IC3]+[IC2]+[IF]+[IM1]+[IM2]+[C3]+[C2]+[C1]+[O]} \cdot (v - Ena)}{scale} \quad (12)$$

6.13 Rule b11

Rule b11 is an assignment rule for parameter b11:

$$b11 = 0.1917 \cdot \exp\left(\frac{v}{20.3}\right) \quad (13)$$

6.14 Rule b12

Rule b12 is an assignment rule for parameter b12:

$$b12 = 0.2 \cdot \exp\left(\frac{(v - 5)}{20.3}\right) \quad (14)$$

6.15 Rule b13

Rule b13 is an assignment rule for parameter b13:

$$b13 = 0.22 \cdot \exp\left(\frac{(v - 10)}{20.3}\right) \quad (15)$$

6.16 Rule b2

Rule b2 is an assignment rule for parameter b2:

$$b2 = \frac{a13 \cdot a2 \cdot a3}{b13 \cdot b3} \quad (16)$$

7 Events

This is an overview of two events. Each event is initiated whenever its trigger condition switches from `false` to `true`. A delay function postpones the effects of an event to a later time point. At the time of execution, an event can assign values to species, parameters or compartments if these are not set to constant.

7.1 Event `event_0000001`

Trigger condition

$$(\text{time} \geq 5) \wedge (\text{time} \leq 20) \quad (17)$$

Assignment

$$v = -30 \quad (18)$$

7.2 Event `event_0000002`

Trigger condition

$$\text{time} > 20 \quad (19)$$

Assignment

$$v = -80 \quad (20)$$

8 Reactions

This model contains eleven reactions. All reactions are listed in the following table and are subsequently described in detail. If a reaction is affected by a modifier, the identifier of this species is written above the reaction arrow.

Table 5: Overview of all reactions

Nº	Id	Name	Reaction Equation	SBO
1	reaction- _0000001	IC3=IC2	$IC3 \rightleftharpoons IC2$	
2	reaction- _0000002	IC2=IF	$IC2 \rightleftharpoons IF$	
3	reaction- _0000003	IF=IM1	$IF \rightleftharpoons IM1$	
4	reaction- _0000004	IM1=IM2	$IM1 \rightleftharpoons IM2$	
5	reaction- _0000005	C3=IC3	$C3 \rightleftharpoons IC3$	
6	reaction- _0000006	C2=C3	$C2 \rightleftharpoons C3$	
7	reaction- _0000007	C2=IC2	$C2 \rightleftharpoons IC2$	
8	reaction- _0000008	C1=C2	$C1 \rightleftharpoons C2$	
9	reaction- _0000009	C1=IF	$C1 \rightleftharpoons IF$	
10	reaction- _0000010	IF=O	$IF \rightleftharpoons O$	
11	reaction- _0000011	O=C1	$O \rightleftharpoons C1$	

8.1 Reaction [reaction_0000001](#)

This is a reversible reaction of one reactant forming one product.

Name IC3=IC2

Reaction equation



Reactant

Table 6: Properties of each reactant.

Id	Name	SBO
IC3		

Product

Table 7: Properties of each product.

Id	Name	SBO
IC2		

Kinetic Law

Derived unit contains undeclared units

$$v_1 = \text{vol}(\text{cell}) \cdot ([\text{IC3}] \cdot a_{11} - [\text{IC2}] \cdot b_{11}) \quad (22)$$

8.2 Reaction [reaction_0000002](#)

This is a reversible reaction of one reactant forming one product.

Name IC2=IF

Reaction equation



Reactant

Table 8: Properties of each reactant.

Id	Name	SBO
IC2		

Product

Table 9: Properties of each product.

Id	Name	SBO
IF		

Kinetic Law

Derived unit contains undeclared units

$$v_2 = \text{vol}(\text{cell}) \cdot ([\text{IC2}] \cdot a_{12} - [\text{IF}] \cdot b_{12}) \quad (24)$$

8.3 Reaction `reaction_0000003`

This is a reversible reaction of one reactant forming one product.

Name IF=IM1

Reaction equation



Reactant

Table 10: Properties of each reactant.

Id	Name	SBO
IF		

Product

Table 11: Properties of each product.

Id	Name	SBO
IM1		

Kinetic Law

Derived unit contains undeclared units

$$v_3 = \text{vol}(\text{cell}) \cdot ([\text{IF}] \cdot a_4 - [\text{IM1}] \cdot b_4) \quad (26)$$

8.4 Reaction `reaction_0000004`

This is a reversible reaction of one reactant forming one product.

Name IM1=IM2

Reaction equation



Reactant

Table 12: Properties of each reactant.

Id	Name	SBO
IM1		

Product

Table 13: Properties of each product.

Id	Name	SBO
IM2		

Kinetic Law

Derived unit contains undeclared units

$$v_4 = \text{vol}(\text{cell}) \cdot ([\text{IM1}] \cdot a_5 - [\text{IM2}] \cdot b_5) \quad (28)$$

8.5 Reaction `reaction_0000005`

This is a reversible reaction of one reactant forming one product.

Name C3=IC3

Reaction equation



Reactant

Table 14: Properties of each reactant.

Id	Name	SBO
C3		

Product

Table 15: Properties of each product.

Id	Name	SBO
IC3		

Kinetic Law

Derived unit contains undeclared units

$$v_5 = \text{vol}(\text{cell}) \cdot ([\text{C3}] \cdot b_3 - [\text{IC3}] \cdot a_3) \tag{30}$$

8.6 Reaction `reaction_0000006`

This is a reversible reaction of one reactant forming one product.

Name C2=C3

Reaction equation



Reactant

Table 16: Properties of each reactant.

Id	Name	SBO
C2		

Product

Table 17: Properties of each product.

Id	Name	SBO
C3		

Kinetic Law

Derived unit contains undeclared units

$$v_6 = \text{vol}(\text{cell}) \cdot ([\text{C2}] \cdot b_{11} - [\text{C3}] \cdot a_{11}) \quad (32)$$

8.7 Reaction `reaction_0000007`

This is a reversible reaction of one reactant forming one product.

Name C2=IC2

Reaction equation



Reactant

Table 18: Properties of each reactant.

Id	Name	SBO
C2		

Product

Table 19: Properties of each product.

Id	Name	SBO
IC2		

Kinetic Law

Derived unit contains undeclared units

$$v_7 = \text{vol}(\text{cell}) \cdot ([\text{C2}] \cdot b_3 - [\text{IC2}] \cdot a_3) \quad (34)$$

8.8 Reaction reaction_0000008

This is a reversible reaction of one reactant forming one product.

Name C1=C2

Reaction equation



Reactant

Table 20: Properties of each reactant.

Id	Name	SBO
C1		

Product

Table 21: Properties of each product.

Id	Name	SBO
C2		

Kinetic Law

Derived unit contains undeclared units

$$v_8 = \text{vol}(\text{cell}) \cdot ([\text{C1}] \cdot b_{12} - [\text{C2}] \cdot a_{12})$$

(36)

8.9 Reaction reaction_0000009

This is a reversible reaction of one reactant forming one product.

Name C1=IF

Reaction equation



Reactant

Table 22: Properties of each reactant.

Id	Name	SBO
C1		

Product

Table 23: Properties of each product.

Id	Name	SBO
IF		

Kinetic Law

Derived unit contains undeclared units

$$v_9 = \text{vol}(\text{cell}) \cdot ([\text{C1}] \cdot b3 - [\text{IF}] \cdot a3) \quad (38)$$

8.10 Reaction `reaction_0000010`

This is a reversible reaction of one reactant forming one product.

Name IF=O

Reaction equation



Reactant

Table 24: Properties of each reactant.

Id	Name	SBO
IF		

Product

Table 25: Properties of each product.

Id	Name	SBO
0	open states	

Kinetic Law

Derived unit contains undeclared units

$$v_{10} = \text{vol}(\text{cell}) \cdot ([\text{IF}] \cdot b2 - a2 \cdot [\text{O}]) \quad (40)$$

8.11 Reaction `reaction_0000011`

This is a reversible reaction of one reactant forming one product.

Name O=C1

Reaction equation



Reactant

Table 26: Properties of each reactant.

Id	Name	SBO
0	open states	

Product

Table 27: Properties of each product.

Id	Name	SBO
C1		

Kinetic Law

Derived unit contains undeclared units

$$v_{11} = \text{vol}(\text{cell}) \cdot ([\text{C1}] \cdot a13 + [\text{O}] \cdot b13) \quad (42)$$

9 Derived Rate Equations

When interpreted as an ordinary differential equation framework, this model implies the following set of equations for the rates of change of each species.

Identifiers for kinetic laws highlighted in gray cannot be verified to evaluate to units of SBML substance per time. As a result, some SBML interpreters may not be able to verify the consistency of the units on quantities in the model. Please check if

- parameters without an unit definition are involved or
- volume correction is necessary because the `hasOnlySubstanceUnits` flag may be set to `false` and `spacialDimensions > 0` for certain species.

9.1 Species C1

Initial concentration $0 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in three reactions (as a reactant in [reaction_0000008](#), [reaction_0000009](#) and as a product in [reaction_0000011](#)).

$$\frac{d}{dt}C1 = v_{11} - v_8 - v_9 \quad (43)$$

9.2 Species C2

Initial concentration $0 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in three reactions (as a reactant in [reaction_0000006](#), [reaction_0000007](#) and as a product in [reaction_0000008](#)).

$$\frac{d}{dt}C2 = v_8 - v_6 - v_7 \quad (44)$$

9.3 Species C3

Initial concentration $1 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in two reactions (as a reactant in [reaction_0000005](#) and as a product in [reaction_0000006](#)).

$$\frac{d}{dt}C3 = v_6 - v_5 \quad (45)$$

9.4 Species IC3

Initial concentration $0 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in two reactions (as a reactant in [reaction_0000001](#) and as a product in [reaction_0000005](#)).

$$\frac{d}{dt}IC3 = v_5 - v_1 \quad (46)$$

9.5 Species IC2

Initial concentration $0 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in three reactions (as a reactant in [reaction_0000002](#) and as a product in [reaction_0000001](#), [reaction_0000007](#)).

$$\frac{d}{dt}\text{IC2} = v_1 + v_7 - v_2 \quad (47)$$

9.6 Species IM1

Initial concentration $0 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in two reactions (as a reactant in [reaction_0000004](#) and as a product in [reaction_0000003](#)).

$$\frac{d}{dt}\text{IM1} = v_3 - v_4 \quad (48)$$

9.7 Species IM2

Initial concentration $0 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in one reaction (as a product in [reaction_0000004](#)).

$$\frac{d}{dt}\text{IM2} = v_4 \quad (49)$$

9.8 Species O

Name open states

Initial concentration $0 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in two reactions (as a reactant in [reaction_0000011](#) and as a product in [reaction_0000010](#)).

$$\frac{d}{dt}\text{O} = v_{10} - v_{11} \quad (50)$$

9.9 Species IF

Initial concentration $0 \text{ mol} \cdot \text{l}^{-1}$

This species takes part in four reactions (as a reactant in [reaction_0000003](#), [reaction_0000010](#) and as a product in [reaction_0000002](#), [reaction_0000009](#)).

$$\frac{d}{dt}\text{IF} = v_2 + v_9 - v_3 - v_{10} \quad (51)$$

SBML²TeX was developed by Andreas Dräger^a, Hannes Planatscher^a, Dieudonné M Wouamba^a, Adrian Schröder^a, Michael Hucka^b, Lukas Endler^c, Martin Golebiewski^d and Andreas Zell^a. Please see <http://www.ra.cs.uni-tuebingen.de/software/SBML2LaTeX> for more information.

^aCenter for Bioinformatics Tübingen (ZBIT), Germany

^bCalifornia Institute of Technology, Beckman Institute BNMC, Pasadena, United States

^cEuropean Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, United Kingdom

^dEML Research gGmbH, Heidelberg, Germany