

Simulation of Simpson's Paradox With Hospital Data

Andy Grogan-Kaylor

2 Nov 2020 12:36:46

Background

Simpson's paradox occurs when a bivariate association is reversed in a multivariate model. This example uses simulated data from hospitals (Wang et al., 2018).

Setup

```
. clear all

. cd "/Users/agrogan/Desktop/newstuff/categorical/simpsons-paradox-hospital-data"
/Users/agrogan/Desktop/newstuff/categorical/simpsons-paradox-hospital-data

. use "hospitaldata.dta"

. list
```

	hospital	severity	outcome	count
1.	better	less severe	success	18
2.	better	less severe	failure	2
3.	better	more severe	success	32
4.	better	more severe	failure	48
5.	normal	less severe	success	64
6.	normal	less severe	failure	16
7.	normal	more severe	success	4
8.	normal	more severe	failure	16

Outcome By Hospital Type

It appears as though patients do better at the *normal* hospital, as opposed to the *better* hospital.

```
. tabulate hospital outcome [fweight = count], row
```

Key
<i>frequency</i>
<i>row percentage</i>

hospital	outcome		Total
	failure	success	
better	50	50	100

	50.00	50.00	100.00
normal	32	68	100
	32.00	68.00	100.00
Total	82	118	200
	41.00	59.00	100.00

```
. graph bar (count) [fweight = count], over(outcome) over(hospital) blabel(bar) title("Hospital Ou
> tcomes") scheme(michigan)
```

```
. graph export bivariategraph.png, width(1000) replace
(file bivariategraph.png written in PNG format)
```




Figure 1: Bivariate Relationships

Outcome By Hospital Type by Severity

When we factor in the *severity* of the illness, we arrive at the reverse conclusion. Patients do better at the *better* hospital.

```
. generate severity_hospital = severity + " " + hospital // concatenate severity + hospital type
. tabulate severity_hospital outcome [fweight=count], row
```

Key
<i>frequency</i>
<i>row percentage</i>

severity_hospital	outcome		Total
	failure	success	
less severe better	2	18	20
	10.00	90.00	100.00
less severe normal	16	64	80
	20.00	80.00	100.00
more severe better	48	32	80
	60.00	40.00	100.00
more severe normal	16	4	20
	80.00	20.00	100.00
Total	82	118	200

```

|      41.00      59.00 |      100.00

. graph bar [fweight = count] if severity == "less severe", ///
> title(Less Severe) ///
> over(outcome) ///
> blabel(bar) ///
> by(hospital) ///
> scheme(michigan) ///
> name(lesssevere, replace)

. graph bar [fweight = count] if severity == "more severe", ///
> title(More Severe) ///
> over(outcome) ///
> blabel(bar) ///
> by(hospital) ///
> scheme(michigan) ///
> name(moresevere, replace)

. graph combine lesssevere moresevere, title(Hospital Outcomes) scheme(michigan)

. graph export multivariategraph.png, width(1000) replace
(file multivariategraph.png written in PNG format)

```




Figure 2: Multivariate Relationships

Reference

Wang, B., Wu, P., Kwan, B., Tu, X. M., & Feng, C. (2018). Simpson's Paradox: Examples. *Shanghai Archives of Psychiatry*, 30(2), 139–143. <https://doi.org/10.11919/j.issn.1002-0829.218026>