Dissertation Title

A DISSERTATION PRESENTED

BY

JOHN HARVARD

TO

THE DEPARTMENT OF PHYSICS

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE OF
DOCTOR OF PHYSICS
IN THE SUBJECT OF
PHYSICS

Harvard University Cambridge, Massachusetts March 2025 This work is licensed via $\underline{\text{CC BY 4.0}}$

Copyright © 2025 John Harvard

Dissertation Advisor: Melissa Franklin John Harvard

Dissertation Title

Abstract

While the search for ever heavier Beyond the Standard Model (BSM) particles is a popular excercise at the energy frontier, the search for XXX has been less explored. This thesis presents a search for YYY in a novel 140 fb⁻¹ dataset collected by the ATLAS experiment during Run 2 at the Large Hadron Collider (LHC) at $\sqrt{s} = 13$ TeV. The dataset is unique in that it is collected at the

CONTENTS

1	THE LHC AND THE ATLAS			
	1.1	Calorimeter · · · · · · · · · · · · · · · · · · ·	2	
		1.1.1 Electromagnetic Calorimetry (ECal) · · · · · · · · · · · · · · · · · · ·	2	
	1.2	Some equations · · · · · · · · · · · · · · · · · · ·	2	
A.2	Appendix			
	A.1	Appendix is hard · · · · · · · · · · · · · · · · · · ·	4	

1

THE LHC AND THE ATLAS

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magnam aliquam quaerat voluptatem. Ut enim aeque doleamus animo, cum corpore dolemus, fieri tamen permagna accessio potest, si aliquod aeternum et infinitum impendere malum nobis opinemur. Quod idem licet transferre in voluptatem, ut postea variari voluptas distinguique possit, augeri amplificarique non possit. At etiam Athenis, ut e patre audiebam

facete et urbane Stoicos irridente, statua est in quo a nobis philosophia defensa et.

t	1	2	3
у	0.3s	0.4s	0.8s

Table 1.1: Timing results

1.1 Calorimeter

1.1.1 Electromagnetic Calorimetry (ECal)

ATLAS uses Liquid Argon (LAr) calorimeter for electromagnetic energy measurements in both the central region¹ ($|\eta| < 1.475$) and end-caps regions² (1.375 $< |\eta| < 3.2$). Together, they provide three layers of calorimeter cells with varying granularities. Additionally, in the $|\eta| < 1.8$ region, a LAr presampler sits in front of the first layer of the LAr ECal and is used to correct the energy loss in the passive material between LAr ECal and the IP. Figure 1.1 shows the schematic of the EMB in regions with four layers.



Figure 1.1: Schematic of the EM Barrel Calorimeter, showing four layers including the presampler (PS) layer

1.2 Some equations

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magnam aliquam quaerat.

¹Electromagnetic Barrel Calorimeter, or EMB

²Electromagnetic Endcap Calorimeter, or EMEC

$$0.002(x + 89.6)^{-1.06\log(x)} \tag{1.1}$$



APPENDIX

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magnam aliquam quaerat. Figure A.1

$$a^2 + b^2 = c^2 (A.1)$$

A.1 Appendix is hard

$$a^3 + b^3 = c^3$$
 (A.2)



Figure A.1: Here's a figure in Appendix