

# Modeling of Bézier curves/surfaces

## Part 1

– a CAGD approach based on OpenGL and C++ –

Ágoston Róth

Department of Mathematics and Computer Science, Babeş-Bolyai University, Cluj-Napoca, Romania

(agoston.roth@gmail.com)

Lecture 9 – May 2, 2022



© Ágoston Róth, 2022

# Pierre Étienne Bézier

September 1, 1910 – November 25, 1999



- Born in Paris.
- **Mathematical engineering degree** (1930) at the École Nationale Supérieure d'Arts et Métiers.
- **Electrical engineering degree** (1931) at the École Supérieure d'Électricité.
- **Doctorate in Mathematics** (1977) at the University of Paris.
- He worked for **Renault** from 1933–1975, where he developed his **UNISURF CAD/CAM** system.
- **Professor of Production Engineering** (1968–1979) at the Conservatoire National des Arts et Métiers.
- In 1985 he was recognized by ACM SIGGRAPH with a "**Steven A. Coons**" award for his lifetime contribution to computer graphics and interactive techniques.
- Patentor (but **not the inventor**) of the Bézier curves and surfaces that are now used in CAD and CG systems.

© Ágoston Roth, 2022



# Pierre Étienne Bézier

September 1, 1910 – November 25, 1999



- Born in Paris.
- Mathematical engineering degree (1930) at the École Nationale Supérieure d'Arts et Métiers.
- Electrical engineering degree (1931) at the École Supérieure d'Électricité.
- Doctorate in Mathematics (1977) at the University of Paris.
- He worked for Renault from 1933–1975, where he developed his UNISURF CAD/CAM system.
- Professor of Production Engineering (1968–1979) at the Conservatoire National des Arts et Métiers.
- In 1985 he was recognized by ACM SIGGRAPH with a "Steven A. Coons" award for his lifetime contribution to computer graphics and interactive techniques.
- Patentor (but not the inventor) of the Bézier curves and surfaces that are now used in CAD and CG systems.

© Ágoston Roth, 2022



# Pierre Étienne Bézier

September 1, 1910 – November 25, 1999



- Born in Paris.
- **Mathematical engineering degree** (1930) at the École Nationale Supérieure d'Arts et Métiers.
- **Electrical engineering degree** (1931) at the École Supérieure d'Électricité.
- **Doctorate in Mathematics** (1977) at the University of Paris.
- He worked for **Renault** from 1933–1975, where he developed his **UNISURF CAD/CAM** system.
- **Professor of Production Engineering** (1968–1979) at the Conservatoire National des Arts et Métiers.
- In 1985 he was recognized by ACM SIGGRAPH with a "**Steven A. Coons**" award for his lifetime contribution to computer graphics and interactive techniques.
- Patentor (but **not the inventor**) of the Bézier curves and surfaces that are now used in CAD and CG systems.

© Ágoston Roth, 2022



# Pierre Étienne Bézier

September 1, 1910 – November 25, 1999



- Born in Paris.
- **Mathematical engineering degree** (1930) at the École Nationale Supérieure d'Arts et Métiers.
- **Electrical engineering degree** (1931) at the École Supérieure d'Électricité.
- **Doctorate in Mathematics** (1977) at the University of Paris.
- He worked for **Renault** from 1933–1975, where he developed his **UNISURF CAD/CAM** system.
- **Professor of Production Engineering** (1968–1979) at the Conservatoire National des Arts et Métiers.
- In 1985 he was recognized by ACM SIGGRAPH with a "**Steven A. Coons**" award for his lifetime contribution to computer graphics and interactive techniques.
- Patentor (but **not the inventor**) of the Bézier curves and surfaces that are now used in CAD and CG systems.



© Ágoston Roth, 2022

# Pierre Étienne Bézier

September 1, 1910 – November 25, 1999



- Born in Paris.
- **Mathematical engineering degree** (1930) at the École Nationale Supérieure d'Arts et Métiers.
- **Electrical engineering degree** (1931) at the École Supérieure d'Électricité.
- **Doctorate in Mathematics** (1977) at the University of Paris.
- He worked for **Renault** from 1933–1975, where he developed his **UNISURF CAD/CAM** system.
- **Professor of Production Engineering** (1968–1979) at the Conservatoire National des Arts et Métiers.
- In 1985 he was recognized by ACM SIGGRAPH with a "**Steven A. Coons**" award for his lifetime contribution to computer graphics and interactive techniques.
- Patentor (but **not the inventor**) of the Bézier curves and surfaces that are now used in CAD and CG systems.



© Ágoston Roth, 2022

# Pierre Étienne Bézier

September 1, 1910 – November 25, 1999



- Born in Paris.
- **Mathematical engineering degree** (1930) at the École Nationale Supérieure d'Arts et Métiers.
- **Electrical engineering degree** (1931) at the École Supérieure d'Électricité.
- **Doctorate in Mathematics** (1977) at the University of Paris.
- He worked for **Renault** from 1933–1975, where he developed his **UNISURF CAD/CAM** system.
- **Professor of Production Engineering** (1968–1979) at the Conservatoire National des Arts et Métiers.
- In 1985 he was recognized by ACM SIGGRAPH with a "**Steven A. Coons**" award for his lifetime contribution to computer graphics and interactive techniques.
- Patentor (but **not the inventor**) of the Bézier curves and surfaces that are now used in CAD and CG systems.



© Ágoston Roth, 2022

# Pierre Étienne Bézier

September 1, 1910 – November 25, 1999



- Born in Paris.
- **Mathematical engineering degree** (1930) at the École Nationale Supérieure d'Arts et Métiers.
- **Electrical engineering degree** (1931) at the École Supérieure d'Électricité.
- **Doctorate in Mathematics** (1977) at the University of Paris.
- He worked for **Renault** from 1933–1975, where he developed his **UNISURF CAD/CAM** system.
- **Professor of Production Engineering** (1968–1979) at the Conservatoire National des Arts et Métiers.
- In 1985 he was recognized by ACM SIGGRAPH with a "**Steven A. Coons**" award for his lifetime contribution to computer graphics and interactive techniques.
- Patentor (but **not the inventor**) of the Bézier curves and surfaces that are now used in CAD and CG systems.



© Ágoston Róth, 2022



# Pierre Étienne Bézier

September 1, 1910 – November 25, 1999



- Born in Paris.
- **Mathematical engineering degree** (1930) at the École Nationale Supérieure d'Arts et Métiers.
- **Electrical engineering degree** (1931) at the École Supérieure d'Électricité.
- **Doctorate in Mathematics** (1977) at the University of Paris.
- He worked for **Renault** from 1933–1975, where he developed his **UNISURF CAD/CAM** system.
- **Professor of Production Engineering** (1968–1979) at the Conservatoire National des Arts et Métiers.
- In 1985 he was recognized by ACM SIGGRAPH with a "**Steven A. Coons**" award for his lifetime contribution to computer graphics and interactive techniques.
- Patentor (but **not the inventor**) of the Bézier curves and surfaces that are now used in CAD and CG systems.



© Ágoston Roth, 2022

# Pierre Étienne Bézier

September 1, 1910 – November 25, 1999



- Born in Paris.
- **Mathematical engineering degree** (1930) at the École Nationale Supérieure d'Arts et Métiers.
- **Electrical engineering degree** (1931) at the École Supérieure d'Électricité.
- **Doctorate in Mathematics** (1977) at the University of Paris.
- He worked for **Renault** from 1933–1975, where he developed his **UNISURF CAD/CAM** system.
- **Professor of Production Engineering** (1968–1979) at the Conservatoire National des Arts et Métiers.
- In 1985 he was recognized by ACM SIGGRAPH with a "**Steven A. Coons**" award for his lifetime contribution to computer graphics and interactive techniques.
- Patentor (but **not the inventor**) of the Bézier curves and surfaces that are now used in most CAD and CG systems.



© Jonathan Roch, 2022

# Paul de Faget de Casteljaou

November 19, 1930 –



- Born in Besançon.
- Studies at École Normale Supérieure de Paris.
- Hired by **Citröen** (1959).
- **Invented (1959) the recursive definition of Bézier curves** (also called de Casteljaou-algorithm).
- Unfortunately, his discoveries were closely guarded **industrial secrets** and were not published until after Bézier.



© Ágoston Rósch, 2022

# Paul de Faget de Casteljau

November 19, 1930 –



- Born in Besançon.
- Studies at École Normale Supérieure de Paris.
- Hired by Citroën (1959).
- Invented (1959) the recursive definition of Bézier curves (also called de Casteljau-algorithm).
- Unfortunately, his discoveries were closely guarded industrial secrets and were not published until after Bézier.



© Agoston Roth, 2022

# Paul de Faget de Casteljau

November 19, 1930 –



- Born in Besançon.
- Studies at École Normale Supérieure de Paris.
- Hired by Citroën (1959).
- Invented (1959) the recursive definition of Bézier curves (also called de Casteljau-algorithm).
- Unfortunately, his discoveries were closely guarded industrial secrets and were not published until after Bézier.



© Agoston Roth, 2022

# Paul de Faget de Casteljau

November 19, 1930 –



- Born in Besançon.
- Studies at École Normale Supérieure de Paris.
- Hired by **Citröen** (1959).
- Invented (1959) the recursive definition of **Bézier curves** (also called de Casteljau-algorithm).
- Unfortunately, his discoveries were closely guarded **industrial secrets** and were not published until after Bézier.



© Ágoston Rósch, 2022

# Paul de Faget de Casteljau

November 19, 1930 –



- Born in Besançon.
- Studies at École Normale Supérieure de Paris.
- Hired by **Citröen** (1959).
- **Invented (1959) the recursive definition of Bézier curves** (also called de Casteljau-algorithm).
- Unfortunately, his discoveries were closely guarded **industrial secrets** and were not published until after Bézier.



© Ágoston Rósch, 2022

# Paul de Faget de Casteljau

November 19, 1930 –



- Born in Besançon.
- Studies at École Normale Supérieure de Paris.
- Hired by **Citröen** (1959).
- **Invented (1959) the recursive definition of Bézier curves** (also called de Casteljau-algorithm).
- Unfortunately, his discoveries were closely guarded **industrial secrets** and were not published until after Bézier.



© Agoston Roth, 2022



### A parabola construction




Fig. 1: A parabola construction

- Consider the non collinear points  $\mathbf{b}_0, \mathbf{b}_1, \mathbf{b}_2 \in \mathbb{R}^\delta, \delta \geq 2$ .
- Let  $u \in [0, 1]$  be an arbitrarily fixed real number.
- Define the inner points

$$\mathbf{b}_0^1(u) = (1-u)\mathbf{b}_0 + u\mathbf{b}_1,$$

$$\mathbf{b}_1^1(u) = (1-u)\mathbf{b}_1 + u\mathbf{b}_2$$

of the segments  $\mathbf{b}_0\mathbf{b}_1$  and  $\mathbf{b}_1\mathbf{b}_2$ .

- Let

$$\mathbf{b}_0^2(u) = (1-u)\mathbf{b}_0^1(u) + u\mathbf{b}_1^1(u)$$

be the inner point of the segment  $\mathbf{b}_0^1(u)\mathbf{b}_1^1(u)$ .

- If  $u$  varies in  $[0, 1]$ , then the orbit of the point  $\mathbf{b}_0^2(u)$  is an arc of a **parabola**:

$$\begin{aligned}\mathbf{b}_0^2(u) &= (1-u)[(1-u)\mathbf{b}_0 + u\mathbf{b}_1] \\ &\quad + u[(1-u)\mathbf{b}_1 + u\mathbf{b}_2] \\ &= (1-u)^2\mathbf{b}_0 + 2u(1-u)\mathbf{b}_1 + u^2\mathbf{b}_2\end{aligned}$$



### A parabola construction




Fig. 1: A parabola construction

- Consider the non collinear points  $\mathbf{b}_0, \mathbf{b}_1, \mathbf{b}_2 \in \mathbb{R}^\delta, \delta \geq 2$ .
- Let  $u \in [0, 1]$  be an arbitrarily fixed real number.
- Define the inner points

$$\mathbf{b}_0^1(u) = (1-u)\mathbf{b}_0 + u\mathbf{b}_1,$$

$$\mathbf{b}_1^1(u) = (1-u)\mathbf{b}_1 + u\mathbf{b}_2$$

of the segments  $\mathbf{b}_0\mathbf{b}_1$  and  $\mathbf{b}_1\mathbf{b}_2$ .

- Let

$$\mathbf{b}_0^2(u) = (1-u)\mathbf{b}_0^1(u) + u\mathbf{b}_1^1(u)$$

be the inner point of the segment  $\mathbf{b}_0^1(u)\mathbf{b}_1^1(u)$ .

- If  $u$  varies in  $[0, 1]$ , then the orbit of the point  $\mathbf{b}_0^2(u)$  is an arc of a **parabola**:

$$\begin{aligned}\mathbf{b}_0^2(u) &= (1-u)[(1-u)\mathbf{b}_0 + u\mathbf{b}_1] \\ &\quad + u[(1-u)\mathbf{b}_1 + u\mathbf{b}_2] \\ &= (1-u)^2\mathbf{b}_0 + 2u(1-u)\mathbf{b}_1 + u^2\mathbf{b}_2\end{aligned}$$



### A parabola construction




Fig. 1: A parabola construction

- Consider the non collinear points  $\mathbf{b}_0, \mathbf{b}_1, \mathbf{b}_2 \in \mathbb{R}^\delta, \delta \geq 2$ .
- Let  $u \in [0, 1]$  be an arbitrarily fixed real number.
- Define the inner points

$$\mathbf{b}_0^1(u) = (1-u)\mathbf{b}_0 + u\mathbf{b}_1,$$

$$\mathbf{b}_1^1(u) = (1-u)\mathbf{b}_1 + u\mathbf{b}_2$$

of the segments  $\mathbf{b}_0\mathbf{b}_1$  and  $\mathbf{b}_1\mathbf{b}_2$ .

- Let

$$\mathbf{b}_0^2(u) = (1-u)\mathbf{b}_0^1(u) + u\mathbf{b}_1^1(u)$$


be the inner point of the segment  $\mathbf{b}_0^1(u)\mathbf{b}_1^1(u)$ .

- If  $u$  varies in  $[0, 1]$ , then the orbit of the point  $\mathbf{b}_0^2(u)$  is an arc of a **parabola**:

$$\begin{aligned}\mathbf{b}_0^2(u) &= (1-u)[(1-u)\mathbf{b}_0 + u\mathbf{b}_1] \\ &\quad + u[(1-u)\mathbf{b}_1 + u\mathbf{b}_2] \\ &= (1-u)^2\mathbf{b}_0 + 2u(1-u)\mathbf{b}_1 + u^2\mathbf{b}_2\end{aligned}$$



### A parabola construction



**Fig. 1:** A parabola construction

- Consider the non collinear points  $\mathbf{b}_0, \mathbf{b}_1, \mathbf{b}_2 \in \mathbb{R}^\delta, \delta \geq 2$ .
- Let  $u \in [0, 1]$  be an arbitrarily fixed real number.
- Define the inner points

$$\mathbf{b}_0^1(u) = (1-u)\mathbf{b}_0 + u\mathbf{b}_1,$$

$$\mathbf{b}_1^1(u) = (1-u)\mathbf{b}_1 + u\mathbf{b}_2$$

of the segments  $\mathbf{b}_0\mathbf{b}_1$  and  $\mathbf{b}_1\mathbf{b}_2$ .

- Let

$$\mathbf{b}_0^2(u) = (1-u)\mathbf{b}_0^1(u) + u\mathbf{b}_1^1(u)$$

be the inner point of the segment  $\mathbf{b}_0^1(u)\mathbf{b}_1^1(u)$ .

- If  $u$  varies in  $[0, 1]$ , then the orbit of the point  $\mathbf{b}_0^2(u)$  is an arc of a parabola:

$$\begin{aligned} \mathbf{b}_0^2(u) &= (1-u)[(1-u)\mathbf{b}_0 + u\mathbf{b}_1] \\ &\quad + u[(1-u)\mathbf{b}_1 + u\mathbf{b}_2] \\ &= (1-u)^2\mathbf{b}_0 + 2u(1-u)\mathbf{b}_1 + u^2\mathbf{b}_2 \end{aligned}$$

### A parabola construction




Fig. 1: A parabola construction

- Consider the non collinear points  $\mathbf{b}_0, \mathbf{b}_1, \mathbf{b}_2 \in \mathbb{R}^\delta, \delta \geq 2$ .
- Let  $u \in [0, 1]$  be an arbitrarily fixed real number.
- Define the inner points

$$\mathbf{b}_0^1(u) = (1-u)\mathbf{b}_0 + u\mathbf{b}_1,$$

$$\mathbf{b}_1^1(u) = (1-u)\mathbf{b}_1 + u\mathbf{b}_2$$

of the segments  $\mathbf{b}_0\mathbf{b}_1$  and  $\mathbf{b}_1\mathbf{b}_2$ .

- Let

$$\mathbf{b}_0^2(u) = (1-u)\mathbf{b}_0^1(u) + u\mathbf{b}_1^1(u)$$

be the inner point of the segment  $\mathbf{b}_1^1(u)\mathbf{b}_1^1(u)$ .

- If  $u$  varies in  $[0, 1]$ , then the orbit of the point  $\mathbf{b}_0^2(u)$  is an arc of a **parabola**:

$$\begin{aligned}\mathbf{b}_0^2(u) &= (1-u)[(1-u)\mathbf{b}_0 + u\mathbf{b}_1] \\ &\quad + u[(1-u)\mathbf{b}_1 + u\mathbf{b}_2] \\ &= (1-u)^2\mathbf{b}_0 + 2u(1-u)\mathbf{b}_1 + u^2\mathbf{b}_2.\end{aligned}$$



# Introduction

## Generalization – the de Casteljau algorithm

### A recurrence relation




Fig. 2: The de Casteljau algorithm

- Consider the control polygon

$$[\mathbf{b}_i]_{i=0}^n \in \mathcal{M}_{1,n+1}(\mathbb{R}^\delta), \delta \geq 2, n \geq 1$$

and the recurrence relation

$$\mathbf{b}_i^r(u) = (1-u)\mathbf{b}_i^{r-1}(u) + u\mathbf{b}_{i+1}^{r-1}(u), u \in [0, 1],$$

where

$$r = 1, 2, \dots, n,$$


$$i = 0, 1, \dots, n-r,$$

$$\mathbf{b}_i^0 = \mathbf{b}_i, i = 0, 1, \dots, n.$$

- The last point  $\mathbf{b}_0^n(u)$ ,  $u \in [0, 1]$  of the recurrence defines the **Bézier curve of degree  $n$** .



### A recurrence relation



**Fig. 2:** The de Casteljau algorithm

- Consider the control polygon

$$[b_i]_{i=0}^n \in \mathcal{M}_{1,n+1}(\mathbb{R}^\delta), \delta \geq 2, n \geq 1$$

and the recurrence relation

$$\mathbf{b}_i^r(u) = (1-u)\mathbf{b}_i^{r-1}(u) + u\mathbf{b}_{i+1}^{r-1}(u), \quad u \in [0, 1],$$

where

$$r = 1, 2, \dots, n,$$


$$i = 0, 1, \dots, n-r,$$

$$\mathbf{b}_i^0 = \mathbf{b}_i, \quad i = 0, 1, \dots, n.$$

- The last point  $\mathbf{b}_0^n(u)$ ,  $u \in [0, 1]$  of the recurrence defines the **Bézier curve of degree  $n$** .



### A recurrence relation



**Fig. 2:** The de Casteljau algorithm

- Consider the control polygon

$$[\mathbf{b}_i]_{i=0}^n \in \mathcal{M}_{1,n+1}(\mathbb{R}^\delta), \delta \geq 2, n \geq 1$$

and the recurrence relation

$$\mathbf{b}_i^r(u) = (1-u)\mathbf{b}_i^{r-1}(u) + u\mathbf{b}_{i+1}^{r-1}(u), u \in [0, 1],$$

where

$$r = 1, 2, \dots, n,$$

$$i = 0, 1, \dots, n-r,$$


$$\mathbf{b}_i^0 = \mathbf{b}_i, i = 0, 1, \dots, n.$$

- The last point  $\mathbf{b}_0^n(u)$ ,  $u \in [0, 1]$  of the recurrence defines the **Bézier curve of degree  $n$** .





### A recurrence relation



**Fig. 2:** The de Casteljau algorithm

- Consider the control polygon

$$[\mathbf{b}_i]_{i=0}^n \in \mathcal{M}_{1,n+1}(\mathbb{R}^\delta), \delta \geq 2, n \geq 1$$

and the recurrence relation

$$\mathbf{b}_i^r(u) = (1-u)\mathbf{b}_i^{r-1}(u) + u\mathbf{b}_{i+1}^{r-1}(u), u \in [0, 1],$$

where

$$r = 1, 2, \dots, n,$$


$$i = 0, 1, \dots, n-r,$$

$$\mathbf{b}_i^0 = \mathbf{b}_i, i = 0, 1, \dots, n.$$

- The last point  $\mathbf{b}_0^n(u)$ ,  $u \in [0, 1]$  of the recurrence defines the **Bézier curve of degree  $n$** .



### A recurrence relation



**Fig. 2:** The de Casteljau algorithm

- Consider the control polygon

$$[\mathbf{b}_i]_{i=0}^n \in \mathcal{M}_{1,n+1}(\mathbb{R}^\delta), \delta \geq 2, n \geq 1$$

and the recurrence relation

$$\mathbf{b}_i^r(u) = (1 - u)\mathbf{b}_i^{r-1}(u) + u\mathbf{b}_{i+1}^{r-1}(u), u \in [0, 1],$$

where

$$r = 1, 2, \dots, n,$$

$$i = 0, 1, \dots, n - r,$$

$$\mathbf{b}_i^0 = \mathbf{b}_i, i = 0, 1, \dots, n.$$

- The last point  $\mathbf{b}_0^n(u)$ ,  $u \in [0, 1]$  of the recurrence defines the **Bézier curve of degree  $n$** .

# Introduction

## Generalization – the de Casteljau algorithm

### Triangular scheme of the de Casteljau corner cutting algorithm

$$\begin{array}{ccccccc}
 \mathbf{b}_0 & = & \mathbf{b}_0^0(u) & & & & \\
 & & & \mathbf{b}_0^1(u) & & & \\
 \mathbf{b}_1 & = & \mathbf{b}_1^0(u) & & \mathbf{b}_0^2(u) & & \\
 & & & \mathbf{b}_1^1(u) & & & \\
 \mathbf{b}_2 & = & \mathbf{b}_2^0(u) & & & & \\
 & & & & & & \mathbf{b}_0^{n-1} \\
 \vdots & & \vdots & & \vdots & & \vdots \\
 & & & & & & \mathbf{b}_0^n(u) \\
 & & & & & & \\
 & & & & & & \mathbf{b}_1^{n-1} \\
 \mathbf{b}_{n-2} & = & \mathbf{b}_{n-2}^0(u) & & & & \\
 & & & \mathbf{b}_{n-2}^1(u) & & & \\
 \mathbf{b}_{n-1} & = & \mathbf{b}_{n-1}^0(u) & & \mathbf{b}_{n-2}^2(u) & & \\
 & & & \mathbf{b}_{n-1}^1(u) & & & \\
 \mathbf{b}_n & = & \mathbf{b}_n^0(u) & & & & 
 \end{array}$$



# Introduction

## Generalization – the de Casteljau algorithm

### Triangular scheme of the de Casteljau corner cutting algorithm

$$\begin{array}{ccccccc} \mathbf{b}_0 & = & \mathbf{b}_0^0(u) & & & & \\ & & & \mathbf{b}_0^1(u) & & & \\ \mathbf{b}_1 & = & \mathbf{b}_1^0(u) & & \mathbf{b}_0^2(u) & & \\ & & & \mathbf{b}_1^1(u) & & & \\ \mathbf{b}_2 & = & \mathbf{b}_2^0(u) & & & & \\ & & & & & & \mathbf{b}_0^{n-1} \\ \vdots & & \vdots & \vdots & \vdots & \dots & \mathbf{b}_0^n(u) \\ & & & & & & \mathbf{b}_1^{n-1} \\ \mathbf{b}_{n-2} & = & \mathbf{b}_{n-2}^0(u) & & & & \\ & & & \mathbf{b}_{n-2}^1(u) & & & \\ \mathbf{b}_{n-1} & = & \mathbf{b}_{n-1}^0(u) & & \mathbf{b}_{n-2}^2(u) & & \\ & & & \mathbf{b}_{n-1}^1(u) & & & \\ \mathbf{b}_n & = & \mathbf{b}_n^0(u) & & & & \end{array}$$

# Introduction

Generalization – the de Casteljau algorithm

## Triangular scheme of the de Casteljau corner cutting algorithm

$$\begin{array}{ccccccc}
 \mathbf{b}_0 & = & \mathbf{b}_0^0(u) & & & & \\
 & & & \mathbf{b}_0^1(u) & & & \\
 \mathbf{b}_1 & = & \mathbf{b}_1^0(u) & & \mathbf{b}_0^2(u) & & \\
 & & & \mathbf{b}_1^1(u) & & & \\
 \mathbf{b}_2 & = & \mathbf{b}_2^0(u) & & & & \\
 & & & & & & \mathbf{b}_0^{n-1} \\
 \vdots & & \vdots & \vdots & \vdots & \dots & \mathbf{b}_0^n(u) \\
 & & & & & & \mathbf{b}_1^{n-1} \\
 \mathbf{b}_{n-2} & = & \mathbf{b}_{n-2}^0(u) & & & & \\
 & & & \mathbf{b}_{n-2}^1(u) & & & \\
 \mathbf{b}_{n-1} & = & \mathbf{b}_{n-1}^0(u) & & \mathbf{b}_{n-2}^2(u) & & \\
 & & & \mathbf{b}_{n-1}^1(u) & & & \\
 \mathbf{b}_n & = & \mathbf{b}_n^0(u) & & & & 
 \end{array}$$

# Introduction

## Generalization – the de Casteljau algorithm

### Triangular scheme of the de Casteljau corner cutting algorithm

$$\begin{array}{ccccccc}
 \mathbf{b}_0 & = & \mathbf{b}_0^0(u) & & & & \\
 & & & \mathbf{b}_0^1(u) & & & \\
 \mathbf{b}_1 & = & \mathbf{b}_1^0(u) & & \mathbf{b}_0^2(u) & & \\
 & & & \mathbf{b}_1^1(u) & & & \\
 \mathbf{b}_2 & = & \mathbf{b}_2^0(u) & & & & \\
 & & & & & & \mathbf{b}_0^{n-1} \\
 \vdots & & \vdots & & \vdots & & \vdots \\
 & & & & & & \mathbf{b}_0^n(u) \\
 & & & & & & \vdots \\
 & & & & & & \mathbf{b}_1^{n-1} \\
 \mathbf{b}_{n-2} & = & \mathbf{b}_{n-2}^0(u) & & & & \\
 & & & \mathbf{b}_{n-2}^1(u) & & & \\
 \mathbf{b}_{n-1} & = & \mathbf{b}_{n-1}^0(u) & & \mathbf{b}_{n-2}^2(u) & & \\
 & & & \mathbf{b}_{n-1}^1(u) & & & \\
 \mathbf{b}_n & = & \mathbf{b}_n^0(u) & & & & 
 \end{array}$$

# Introduction

Generalization – the de Casteljau algorithm

## Triangular scheme of the de Casteljau corner cutting algorithm

$$\begin{array}{ccccccc}
 \mathbf{b}_0 & = & \mathbf{b}_0^0(u) & & & & \\
 & & & \mathbf{b}_0^1(u) & & & \\
 \mathbf{b}_1 & = & \mathbf{b}_1^0(u) & & \mathbf{b}_0^2(u) & & \\
 & & & \mathbf{b}_1^1(u) & & & \\
 \mathbf{b}_2 & = & \mathbf{b}_2^0(u) & & & & \\
 & & & & & & \mathbf{b}_0^{n-1} \\
 \vdots & & \vdots & & \vdots & & \vdots \\
 & & & & & & \mathbf{b}_0^n(u) \\
 & & & & & & \vdots \\
 & & & & & & \mathbf{b}_1^{n-1} \\
 \mathbf{b}_{n-2} & = & \mathbf{b}_{n-2}^0(u) & & & & \\
 & & & \mathbf{b}_{n-2}^1(u) & & & \\
 \mathbf{b}_{n-1} & = & \mathbf{b}_{n-1}^0(u) & & \mathbf{b}_{n-2}^2(u) & & \\
 & & & \mathbf{b}_{n-1}^1(u) & & & \\
 \mathbf{b}_n & = & \mathbf{b}_n^0(u) & & & & 
 \end{array}$$

# Introduction

Generalization – the de Casteljau algorithm

## Triangular scheme of the de Casteljau corner cutting algorithm

$$\begin{array}{ccccccc}
 \mathbf{b}_0 & = & \mathbf{b}_0^0(u) & & & & \\
 & & & \mathbf{b}_0^1(u) & & & \\
 \mathbf{b}_1 & = & \mathbf{b}_1^0(u) & & \mathbf{b}_0^2(u) & & \\
 & & & \mathbf{b}_1^1(u) & & & \\
 \mathbf{b}_2 & = & \mathbf{b}_2^0(u) & & & & \\
 & & & & & & \mathbf{b}_0^{n-1} \\
 \vdots & & \vdots & \vdots & \vdots & \dots & \mathbf{b}_0^n(u) \\
 & & & & & & \mathbf{b}_1^{n-1} \\
 \mathbf{b}_{n-2} & = & \mathbf{b}_{n-2}^0(u) & & & & \\
 & & & \mathbf{b}_{n-2}^1(u) & & & \\
 \mathbf{b}_{n-1} & = & \mathbf{b}_{n-1}^0(u) & & \mathbf{b}_{n-2}^2(u) & & \\
 & & & \mathbf{b}_{n-1}^1(u) & & & \\
 \mathbf{b}_n & = & \mathbf{b}_n^0(u) & & & & 
 \end{array}$$



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Bernstein polynomials

Consider the function system

$$B = \left\{ B_i^n(u) = \binom{n}{i} u^i (1-u)^{n-i} : u \in [0, 1] \right\}_{i=0}^n,$$

where:

- the function  $B_i^n(u)$ ,  $u \in [0, 1]$  is the  $i$ th Bernstein polynomial of degree  $n$ ;
- the function system  $B$  is in fact a basis of the vector space of polynomials of degree at most  $n$ .



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Bernstein polynomials

Consider the function system

$$B = \left\{ B_i^n(u) = \binom{n}{i} u^i (1-u)^{n-i} : u \in [0, 1] \right\}_{i=0}^n,$$

where:

- the function  $B_i^n(u)$ ,  $u \in [0, 1]$  is the  $i$ th Bernstein polynomial of degree  $n$ ;
- the function system  $B$  is in fact a basis of the vector space of polynomials of degree at most  $n$ .



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Bernstein polynomials

Consider the function system

$$B = \left\{ B_i^n(u) = \binom{n}{i} u^i (1-u)^{n-i} : u \in [0, 1] \right\}_{i=0}^n,$$

where:

- the function  $B_i^n(u)$ ,  $u \in [0, 1]$  is the  $i$ th Bernstein polynomial of degree  $n$ ;
- the function system  $B$  is in fact **a basis of the vector space of polynomials of degree at most  $n$ .**



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Recurrence property of Bernstein polynomials

The Bernstein polynomials of degree  $n$  fulfill the **recurrence property**

$$B_i^n(u) = (1-u)B_i^{n-1}(u) + uB_{i-1}^{n-1}(u), \forall u \in [0, 1]$$

and the identities

$$B_0^n(u) \equiv 1, \forall u \in [0, 1], \text{ and } B_i^n(u) \equiv 0, \forall u \in [0, 1], \text{ if } i < 0 \text{ or } i > n.$$

### Proof.

- We will use the recurrence property  $\binom{n}{i} = \binom{n-1}{i} + \binom{n-1}{i-1}$  of the binomial coefficients.
- We can successively write:

$$\begin{aligned} B_i^n(u) &= \binom{n}{i} u^i (1-u)^{n-i} \\ &= \binom{n-1}{i} u^i (1-u)^{n-i} + \binom{n-1}{i-1} u^i (1-u)^{n-i} \\ &= (1-u) \binom{n-1}{i} u^i (1-u)^{n-1-i} + u \binom{n-1}{i-1} u^{i-1} (1-u)^{n-1-i} \\ &= (1-u) B_i^{n-1}(u) + u B_{i-1}^{n-1}(u), \forall u \in [0, 1]. \quad \square \end{aligned}$$



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Recurrence property of Bernstein polynomials

The Bernstein polynomials of degree  $n$  fulfill the **recurrence property**

$$B_i^n(u) = (1-u)B_i^{n-1}(u) + uB_{i-1}^{n-1}(u), \forall u \in [0, 1]$$

and the identities

$$B_0^n(u) \equiv 1, \forall u \in [0, 1], \text{ and } B_i^n(u) \equiv 0, \forall u \in [0, 1], \text{ if } i < 0 \text{ or } i > n.$$

### Proof.

- We will use the recurrence property  $\binom{n}{i} = \binom{n-1}{i} + \binom{n-1}{i-1}$  of the binomial coefficients.
- We can successively write:

$$\begin{aligned} B_i^n(u) &= \binom{n}{i} u^i (1-u)^{n-i} \\ &= \binom{n-1}{i} u^i (1-u)^{n-i} + \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} \\ &= (1-u) \binom{n-1}{i} u^i (1-u)^{n-1-i} + u \binom{n-1}{i-1} u^{i-1} (1-u)^{n-1-i} \\ &= (1-u) B_i^{n-1}(u) + u B_{i-1}^{n-1}(u), \forall u \in [0, 1]. \quad \square \end{aligned}$$



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Recurrence property of Bernstein polynomials

The Bernstein polynomials of degree  $n$  fulfill the **recurrence property**

$$B_i^n(u) = (1-u)B_i^{n-1}(u) + uB_{i-1}^{n-1}(u), \forall u \in [0, 1]$$

and the identities

$$B_0^n(u) \equiv 1, \forall u \in [0, 1], \text{ and } B_i^n(u) \equiv 0, \forall u \in [0, 1], \text{ if } i < 0 \text{ or } i > n.$$

### Proof.

- We will use the recurrence property  $\binom{n}{i} = \binom{n-1}{i} + \binom{n-1}{i-1}$  of the binomial coefficients.
- We can successively write:

$$\begin{aligned} B_i^n(u) &= \binom{n}{i} u^i (1-u)^{n-i} \\ &= \binom{n-1}{i} u^i (1-u)^{n-i} + \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} \\ &= (1-u) \binom{n-1}{i} u^i (1-u)^{n-1-i} + u \binom{n-1}{i-1} u^{i-1} (1-u)^{n-1-i} \\ &= (1-u) B_i^{n-1}(u) + u B_{i-1}^{n-1}(u), \forall u \in [0, 1]. \quad \square \end{aligned}$$



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Recurrence property of Bernstein polynomials

The Bernstein polynomials of degree  $n$  fulfill the **recurrence property**

$$B_i^n(u) = (1-u)B_i^{n-1}(u) + uB_{i-1}^{n-1}(u), \forall u \in [0, 1]$$

and the identities

$$B_0^n(u) \equiv 1, \forall u \in [0, 1], \text{ and } B_i^n(u) \equiv 0, \forall u \in [0, 1], \text{ if } i < 0 \text{ or } i > n.$$

### Proof.

- We will use the recurrence property  $\binom{n}{i} = \binom{n-1}{i} + \binom{n-1}{i-1}$  of the binomial coefficients.
- We can successively write:

$$\begin{aligned} B_i^n(u) &= \binom{n}{i} u^i (1-u)^{n-i} \\ &= \binom{n-1}{i} u^i (1-u)^{n-i} + \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} \\ &= (1-u) \binom{n-1}{i} u^i (1-u)^{n-1-i} + u \binom{n-1}{i-1} u^{i-1} (1-u)^{n-1-i} \\ &= (1-u) B_i^{n-1}(u) + u B_{i-1}^{n-1}(u), \forall u \in [0, 1]. \quad \square \end{aligned}$$



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Recurrence property of Bernstein polynomials

The Bernstein polynomials of degree  $n$  fulfill the **recurrence property**

$$B_i^n(u) = (1-u)B_i^{n-1}(u) + uB_{i-1}^{n-1}(u), \forall u \in [0, 1]$$

and the identities

$$B_0^n(u) \equiv 1, \forall u \in [0, 1], \text{ and } B_i^n(u) \equiv 0, \forall u \in [0, 1], \text{ if } i < 0 \text{ or } i > n.$$

### Proof.

- We will use the recurrence property  $\binom{n}{i} = \binom{n-1}{i} + \binom{n-1}{i-1}$  of the binomial coefficients.
- We can successively write:

$$\begin{aligned} B_i^n(u) &= \binom{n}{i} u^i (1-u)^{n-i} \\ &= \binom{n-1}{i} u^i (1-u)^{n-i} + \binom{n-1}{i-1} u^i (1-u)^{n-i} \\ &= (1-u) \binom{n-1}{i} u^i (1-u)^{n-1-i} + u \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} \\ &= (1-u) B_i^{n-1}(u) + u B_{i-1}^{n-1}(u), \forall u \in [0, 1]. \quad \square \end{aligned}$$



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Recurrence property of Bernstein polynomials

The Bernstein polynomials of degree  $n$  fulfill the **recurrence property**

$$B_i^n(u) = (1-u)B_i^{n-1}(u) + uB_{i-1}^{n-1}(u), \forall u \in [0, 1]$$

and the identities

$$B_0^n(u) \equiv 1, \forall u \in [0, 1], \text{ and } B_i^n(u) \equiv 0, \forall u \in [0, 1], \text{ if } i < 0 \text{ or } i > n.$$

### Proof.

- We will use the recurrence property  $\binom{n}{i} = \binom{n-1}{i} + \binom{n-1}{i-1}$  of the binomial coefficients.
- We can successively write:

$$\begin{aligned} B_i^n(u) &= \binom{n}{i} u^i (1-u)^{n-i} \\ &= \binom{n-1}{i} u^i (1-u)^{n-i} + \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} \\ &= (1-u) \binom{n-1}{i} u^i (1-u)^{n-1-i} + u \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} \\ &= (1-u) B_i^{n-1}(u) + u B_{i-1}^{n-1}(u), \forall u \in [0, 1]. \quad \square \end{aligned}$$

# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Partition of the unity

The system

$$B = \left\{ B_i^n(u) = \binom{n}{i} u^i (1-u)^{n-i} : u \in [0, 1] \right\}_{i=0}^n,$$

of Bernstein polynomials of degree  $n$  forms the partition of the unity, i.e.

$$\sum_{i=0}^n B_i^n(u) \equiv 1, \forall u \in [0, 1].$$

Proof.

- By the binomial theorem of Newton:

$$\sum_{i=0}^n B_i^n(u) = \sum_{i=0}^n \binom{n}{i} u^i (1-u)^{n-i} = [u + (1-u)]^n \equiv 1, \forall u \in [0, 1].$$

- Moreover,  $B_i^n(u) \geq 0, \forall u \in [0, 1]$ .



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Partition of the unity

The system

$$B = \left\{ B_i^n(u) = \binom{n}{i} u^i (1-u)^{n-i} : u \in [0, 1] \right\}_{i=0}^n,$$

of Bernstein polynomials of degree  $n$  forms the partition of the unity, i.e.

$$\sum_{i=0}^n B_i^n(u) \equiv 1, \forall u \in [0, 1].$$

Proof.

- By the binomial theorem of Newton:

$$\sum_{i=0}^n B_i^n(u) = \sum_{i=0}^n \binom{n}{i} u^i (1-u)^{n-i} = [u + (1-u)]^n \equiv 1, \forall u \in [0, 1].$$

- Moreover,  $B_i^n(u) \geq 0, \forall u \in [0, 1]$ .



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Partition of the unity

The system

$$B = \left\{ B_i^n(u) = \binom{n}{i} u^i (1-u)^{n-i} : u \in [0, 1] \right\}_{i=0}^n,$$

of Bernstein polynomials of degree  $n$  forms the partition of the unity, i.e.

$$\sum_{i=0}^n B_i^n(u) \equiv 1, \forall u \in [0, 1].$$

Proof.

- By the binomial theorem of Newton:

$$\sum_{i=0}^n B_i^n(u) = \sum_{i=0}^n \binom{n}{i} u^i (1-u)^{n-i} = [u + (1-u)]^n \equiv 1, \forall u \in [0, 1].$$

- Moreover,  $B_i^n(u) \geq 0, \forall u \in [0, 1]$ .



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Maximum points of Bernstein polynomials of degree $n$




Fig. 3: Bernstein polynomials of degree 3.

The  $i$ th ( $i = 0, 1, \dots, n$ ) Bernstein polynomial

$$B_i^n(u) = \binom{n}{i} u^i (1-u)^{n-i}, u \in [0, 1]$$

of degree  $n$  attains its maximum at the point

$$\bar{u}_i = \frac{i}{n}.$$

Moreover,  $B_i^n(u)$  is **strictly increasing** (resp. **decreasing**) on the interval  $[0, \bar{u}_i]$  (resp.  $[\bar{u}_i, 1]$ ).



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Maximum points of Bernstein polynomials of degree $n$




Fig. 3: Bernstein polynomials of degree 3.

The  $i$ th ( $i = 0, 1, \dots, n$ ) Bernstein polynomial

$$B_i^n(u) = \binom{n}{i} u^i (1-u)^{n-i}, u \in [0, 1]$$

of degree  $n$  attains its maximum at the point

$$\bar{u}_i = \frac{i}{n}.$$

Moreover,  $B_i^n(u)$  is **strictly increasing** (resp. **decreasing**) on the interval  $[0, \bar{u}_i]$  (resp.  $[\bar{u}_i, 1]$ ).



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Maximum points of Bernstein polynomials of degree $n$




Fig. 3: Bernstein polynomials of degree 3.

The  $i$ th ( $i = 0, 1, \dots, n$ ) Bernstein polynomial

$$B_i^n(u) = \binom{n}{i} u^i (1-u)^{n-i}, u \in [0, 1]$$

of degree  $n$  attains its maximum at the point

$$\bar{u}_i = \frac{i}{n}.$$

Moreover,  $B_i^n(u)$  is **strictly increasing** (resp. **decreasing**) on the interval  $[0, \bar{u}_i]$  (resp.  $[\bar{u}_i, 1]$ ).



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Maximum points of Bernstein polynomials of degree $n$




Fig. 3: Bernstein polynomials of degree 3.

The  $i$ th ( $i = 0, 1, \dots, n$ ) Bernstein polynomial

$$B_i^n(u) = \binom{n}{i} u^i (1-u)^{n-i}, u \in [0, 1]$$

of degree  $n$  attains its maximum at the point

$$\bar{u}_i = \frac{i}{n}.$$

Moreover,  $B_i^n(u)$  is **strictly increasing** (resp. **decreasing**) on the interval  $[0, \bar{u}_i]$  (resp.  $[\bar{u}_i, 1]$ ).






# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Maximum points of Bernstein polynomials of degree $n$



**Fig. 3:** Bernstein polynomials of degree 3.

The  $i$ th ( $i = 0, 1, \dots, n$ ) Bernstein polynomial

$$B_i^n(u) = \binom{n}{i} u^i (1-u)^{n-i}, u \in [0, 1]$$

of degree  $n$  attains its maximum at the point

$$\bar{u}_i = \frac{i}{n}.$$

Moreover,  $B_i^n(u)$  is **strictly increasing** (resp. **decreasing**) on the interval  $[0, \bar{u}_i]$  (resp.  $[\bar{u}_i, 1]$ ).



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof.

- When  $i = 0$  or  $i = n$  the property follows from the strict monotonicity of the functions  $(1 - u)^n$  and  $u^n$ .
- When  $1 \leq i \leq n - 1$ :
  - the derivative of the  $i$ th Bernstein polynomial of degree  $n$  is:

$$\begin{aligned}\frac{d}{du} B_i^n(u) &= \frac{d}{du} \left[ \binom{n}{i} u^i (1-u)^{n-i} \right] \\ &= i \binom{n}{i} u^{i-1} (1-u)^{n-i} - (n-i) \binom{n}{i} u^i (1-u)^{n-1-i} \\ &= n \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} - n \binom{n-1}{i} u^i (1-u)^{n-1-i} \\ &= n \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right); \end{aligned}$$

- we need to solve the system

$$\begin{cases} \frac{d}{du} B_i^n(u) = 0, \\ \frac{d^2}{du^2} B_i^n(u) < 0. \end{cases}$$



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof.

- When  $i = 0$  or  $i = n$  the property follows from the strict monotonicity of the functions  $(1 - u)^n$  and  $u^n$ .
- When  $1 \leq i \leq n - 1$ :
  - the derivative of the  $i$ th Bernstein polynomial of degree  $n$  is:

$$\begin{aligned}\frac{d}{du} B_i^n(u) &= \frac{d}{du} \left[ \binom{n}{i} u^i (1 - u)^{n-i} \right] \\ &= i \binom{n}{i} u^{i-1} (1 - u)^{n-i} - (n - i) \binom{n}{i} u^i (1 - u)^{n-1-i} \\ &= n \binom{n-1}{i-1} u^{i-1} (1 - u)^{n-i} - n \binom{n-1}{i} u^i (1 - u)^{n-1-i} \\ &= n \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right);\end{aligned}$$

- we need to solve the system

$$\begin{cases} \frac{d}{du} B_i^n(u) = 0, \\ \frac{d^2}{du^2} B_i^n(u) < 0. \end{cases}$$



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof.

- When  $i = 0$  or  $i = n$  the property follows from the strict monotonicity of the functions  $(1 - u)^n$  and  $u^n$ .
- When  $1 \leq i \leq n - 1$ :
  - the derivative of the  $i$ th Bernstein polynomial of degree  $n$  is:

$$\begin{aligned}\frac{d}{du} B_i^n(u) &= \frac{d}{du} \left[ \binom{n}{i} u^i (1 - u)^{n-i} \right] \\&= i \binom{n}{i} u^{i-1} (1 - u)^{n-i} - (n - i) \binom{n}{i} u^i (1 - u)^{n-1-i} \\&= n \binom{n-1}{i-1} u^{i-1} (1 - u)^{n-i} - n \binom{n-1}{i} u^i (1 - u)^{n-1-i} \\&= n \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right);\end{aligned}$$

- we need to solve the system

$$\begin{cases} \frac{d}{du} B_i^n(u) = 0, \\ \frac{d^2}{du^2} B_i^n(u) < 0. \end{cases}$$



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof.

- When  $i = 0$  or  $i = n$  the property follows from the strict monotonicity of the functions  $(1 - u)^n$  and  $u^n$ .
- When  $1 \leq i \leq n - 1$ :
  - the derivative of the  $i$ th Bernstein polynomial of degree  $n$  is:

$$\begin{aligned}\frac{d}{du} B_i^n(u) &= \frac{d}{du} \left[ \binom{n}{i} u^i (1 - u)^{n-i} \right] \\&= i \binom{n}{i} u^{i-1} (1 - u)^{n-i} - (n - i) \binom{n}{i} u^i (1 - u)^{n-1-i} \\&= n \binom{n-1}{i-1} u^{i-1} (1 - u)^{n-i} - n \binom{n-1}{i} u^i (1 - u)^{n-1-i} \\&= n \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right);\end{aligned}$$

- we need to solve the system

$$\begin{cases} \frac{d}{du} B_i^n(u) = 0, \\ \frac{d^2}{du^2} B_i^n(u) < 0. \end{cases}$$



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof.

- When  $i = 0$  or  $i = n$  the property follows from the strict monotonicity of the functions  $(1 - u)^n$  and  $u^n$ .
- When  $1 \leq i \leq n - 1$ :
  - the derivative of the  $i$ th Bernstein polynomial of degree  $n$  is:

$$\begin{aligned}\frac{d}{du} B_i^n(u) &= \frac{d}{du} \left[ \binom{n}{i} u^i (1 - u)^{n-i} \right] \\&= i \binom{n}{i} u^{i-1} (1 - u)^{n-i} - (n - i) \binom{n}{i} u^i (1 - u)^{n-1-i} \\&= n \binom{n-1}{i-1} u^{i-1} (1 - u)^{n-i} - n \binom{n-1}{i} u^i (1 - u)^{n-1-i} \\&= n \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right); \end{aligned}$$

- we need to solve the system

$$\begin{cases} \frac{d}{du} B_i^n(u) = 0, \\ \frac{d^2}{du^2} B_i^n(u) < 0. \end{cases}$$



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof.

- When  $i = 0$  or  $i = n$  the property follows from the strict monotonicity of the functions  $(1 - u)^n$  and  $u^n$ .
- When  $1 \leq i \leq n - 1$ :
  - the derivative of the  $i$ th Bernstein polynomial of degree  $n$  is:

$$\begin{aligned}\frac{d}{du} B_i^n(u) &= \frac{d}{du} \left[ \binom{n}{i} u^i (1 - u)^{n-i} \right] \\&= i \binom{n}{i} u^{i-1} (1 - u)^{n-i} - (n - i) \binom{n}{i} u^i (1 - u)^{n-1-i} \\&= n \binom{n-1}{i-1} u^{i-1} (1 - u)^{n-i} - n \binom{n-1}{i} u^i (1 - u)^{n-1-i} \\&= n \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right);\end{aligned}$$

- we need to solve the system

$$\begin{cases} \frac{d}{du} B_i^n(u) = 0, \\ \frac{d^2}{du^2} B_i^n(u) < 0. \end{cases}$$



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof.

- When  $i = 0$  or  $i = n$  the property follows from the strict monotonicity of the functions  $(1 - u)^n$  and  $u^n$ .
- When  $1 \leq i \leq n - 1$ :
  - the derivative of the  $i$ th Bernstein polynomial of degree  $n$  is:

$$\begin{aligned}\frac{d}{du} B_i^n(u) &= \frac{d}{du} \left[ \binom{n}{i} u^i (1 - u)^{n-i} \right] \\&= i \binom{n}{i} u^{i-1} (1 - u)^{n-i} - (n - i) \binom{n}{i} u^i (1 - u)^{n-1-i} \\&= n \binom{n-1}{i-1} u^{i-1} (1 - u)^{n-i} - n \binom{n-1}{i} u^i (1 - u)^{n-1-i} \\&= n \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right);\end{aligned}$$

- we need to solve the system

$$\begin{cases} \frac{d}{du} B_i^n(u) = 0, \\ \frac{d^2}{du^2} B_i^n(u) < 0. \end{cases}$$





# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof – continued.

- In order to simplify the equation

$$\frac{d}{du} B_i^n(u) = 0, \quad i = 1, 2, \dots, n-1,$$

we can successively write:

$$\begin{aligned} n \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} &= n \binom{n-1}{i} u^i (1-u)^{n-1-i} \\ \binom{n-1}{i-1} (1-u) &= \binom{n-1}{i} u \Big/ \cdot \frac{(i-1)!(n-1-i)!}{(n-1)!} \\ \frac{1-u}{n-i} &= \frac{u}{i}, \end{aligned}$$

the solution of which is

$$\bar{u}_i = \frac{i}{n}.$$

- It is easy to verify that  $\left. \frac{d^2}{du^2} B_i^n(u) \right|_{u=\bar{u}_i} < 0$ .



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof – continued.

- In order to simplify the equation

$$\frac{d}{du} B_i^n(u) = 0, \quad i = 1, 2, \dots, n-1,$$

we can successively write:

$$\begin{aligned} n \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} &= n \binom{n-1}{i} u^i (1-u)^{n-1-i} \\ \binom{n-1}{i-1} (1-u) &= \binom{n-1}{i} u \Big/ \cdot \frac{(i-1)!(n-1-i)!}{(n-1)!} \\ \frac{1-u}{n-i} &= \frac{u}{i}, \end{aligned}$$

the solution of which is

$$\bar{u}_i = \frac{i}{n}.$$

- It is easy to verify that  $\left. \frac{d^2}{du^2} B_i^n(u) \right|_{u=\bar{u}_i} < 0$ .



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof – continued.

- In order to simplify the equation

$$\frac{d}{du} B_i^n(u) = 0, \quad i = 1, 2, \dots, n-1,$$

we can successively write:

$$\begin{aligned} n \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} &= n \binom{n-1}{i} u^i (1-u)^{n-1-i} \\ \binom{n-1}{i-1} (1-u) &= \binom{n-1}{i} u \Big/ \cdot \frac{(i-1)!(n-1-i)!}{(n-1)!} \\ \frac{1-u}{n-i} &= \frac{u}{i}, \end{aligned}$$

the solution of which is

$$\bar{u}_i = \frac{i}{n}.$$

- It is easy to verify that  $\left. \frac{d^2}{du^2} B_i^n(u) \right|_{u=\bar{u}_i} < 0$ .



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof – continued.

- In order to simplify the equation

$$\frac{d}{du} B_i^n(u) = 0, \quad i = 1, 2, \dots, n-1,$$

we can successively write:

$$\begin{aligned} n \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} &= n \binom{n-1}{i} u^i (1-u)^{n-1-i} \\ \binom{n-1}{i-1} (1-u) &= \binom{n-1}{i} u \Big/ \cdot \frac{(i-1)!(n-1-i)!}{(n-1)!} \\ \frac{1-u}{n-i} &= \frac{u}{i}, \end{aligned}$$

the solution of which is

$$\bar{u}_i = \frac{i}{n}.$$

- It is easy to verify that  $\left. \frac{d^2}{du^2} B_i^n(u) \right|_{u=\bar{u}_i} < 0$ .



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof – continued.

- In order to simplify the equation

$$\frac{d}{du} B_i^n(u) = 0, \quad i = 1, 2, \dots, n-1,$$

we can successively write:

$$\begin{aligned} n \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} &= n \binom{n-1}{i} u^i (1-u)^{n-1-i} \\ \binom{n-1}{i-1} (1-u) &= \binom{n-1}{i} u \Big/ \cdot \frac{(i-1)!(n-1-i)!}{(n-1)!} \\ \frac{1-u}{n-i} &= \frac{u}{i}, \end{aligned}$$

the solution of which is

$$\bar{u}_i = \frac{i}{n}.$$

- It is easy to verify that  $\left. \frac{d^2}{du^2} B_i^n(u) \right|_{u=\bar{u}_i} < 0$ .



# Bernstein representation of Bézier curves

## Properties of Bernstein polynomials

### Proof – continued.

- In order to simplify the equation

$$\frac{d}{du} B_i^n(u) = 0, \quad i = 1, 2, \dots, n-1,$$

we can successively write:

$$\begin{aligned} n \binom{n-1}{i-1} u^{i-1} (1-u)^{n-i} &= n \binom{n-1}{i} u^i (1-u)^{n-1-i} \\ \binom{n-1}{i-1} (1-u) &= \binom{n-1}{i} u \Big/ \cdot \frac{(i-1)!(n-1-i)!}{(n-1)!} \\ \frac{1-u}{n-i} &= \frac{u}{i}, \end{aligned}$$

the solution of which is

$$\bar{u}_i = \frac{i}{n}.$$

- It is easy to verify that  $\left. \frac{d^2}{du^2} B_i^n(u) \right|_{u=\bar{u}_i} < 0$ .



# Bernstein representation of Bézier curves

## Parametric form of Bézier curves

Theorem (points of the de Casteljau algorithm by means of Bernstein polynomials)

The point  $\mathbf{b}_i^r(u)$  ( $r = 0, 1, \dots, n$ ;  $i = 0, 1, \dots, n - r$ ;  $u \in [0, 1]$ ) of the de Casteljau algorithm can be expressed as

$$\mathbf{b}_i^r(u) = \sum_{j=0}^r \mathbf{b}_{j+i} B_j^r(u)$$

by means of Bernstein polynomials of degree  $r$ .

Corollary (Bernstein/parametric representation of Bézier curves)

By substituting  $i = 0$  and  $r = n$  into the previous formula we get the

$$\mathbf{b}(u) := \mathbf{b}_0^n(u) = \sum_{j=0}^n \mathbf{b}_j B_j^n(u), \quad u \in [0, 1]$$

Bernstein representation of the Bézier curve of degree  $n$ .



# Bernstein representation of Bézier curves

## Parametric form of Bézier curves

Theorem (points of the de Casteljau algorithm by means of Bernstein polynomials)

The point  $\mathbf{b}_i^r(u)$  ( $r = 0, 1, \dots, n$ ;  $i = 0, 1, \dots, n - r$ ;  $u \in [0, 1]$ ) of the de Casteljau algorithm can be expressed as

$$\mathbf{b}_i^r(u) = \sum_{j=0}^r \mathbf{b}_{j+i} B_j^r(u)$$

by means of Bernstein polynomials of degree  $r$ .

Corollary (Bernstein/parametric representation of Bézier curves)

By substituting  $i = 0$  and  $r = n$  into the previous formula we get the

$$\mathbf{b}(u) := \mathbf{b}_0^n(u) = \sum_{j=0}^n \mathbf{b}_j B_j^n(u), \quad u \in [0, 1]$$

Bernstein representation of the Bézier curve of degree  $n$ .





# Bernstein representation of Bézier curves

## Parametric form of Bézier curves

### Proof.

- First, we perform the index transformation  $j \rightarrow j - i$ :

$$\sum_{j=0}^r \mathbf{b}_{j+i} B_j^r(u) = \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^r(u).$$

- Second, we prove the theorem by **mathematical induction with respect to  $r$** .
  - The statement trivially holds for  $r = 0$ :

$$\mathbf{b}_i^0(u) = \sum_{j=i}^{i+0} \mathbf{b}_j B_{j-i}^0(u) = \mathbf{b}_i B_0^0(u) = \mathbf{b}_i, \forall u \in [0, 1], i = 0, 1, \dots, n.$$

- We assume that the statement is valid for all  $0 \leq k \leq r - 1$  and we also prove it for  $k = r$  ( $1 \leq r \leq n$ ).
- Based on the recurrence relation of the de Casteljau algorithm, we have:

$$\mathbf{b}_i^r(u) = (1-u)\mathbf{b}_i^{r-1}(u) + u\mathbf{b}_{i+1}^{r-1}(u), \forall u \in [0, 1].$$

- By applying the induction hypothesis to the points  $\mathbf{b}_i^{r-1}(u)$  and  $\mathbf{b}_{i+1}^{r-1}(u)$  we can successively write:



# Bernstein representation of Bézier curves

## Parametric form of Bézier curves

### Proof.

- First, we perform the index transformation  $j \rightarrow j - i$ :

$$\sum_{j=0}^r \mathbf{b}_{j+i} B_j^r(u) = \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^r(u).$$

- Second, we prove the theorem by **mathematical induction with respect to  $r$** .
  - The statement trivially holds for  $r = 0$ :

$$\mathbf{b}_i^0(u) = \sum_{j=i}^{i+0} \mathbf{b}_j B_{j-i}^0(u) = \mathbf{b}_i B_0^0(u) = \mathbf{b}_i, \forall u \in [0, 1], i = 0, 1, \dots, n.$$

- We assume that the statement is valid for all  $0 \leq k \leq r - 1$  and we also prove it for  $k = r$  ( $1 \leq r \leq n$ ).
- Based on the recurrence relation of the de Casteljau algorithm, we have:

$$\mathbf{b}_i^r(u) = (1 - u) \mathbf{b}_i^{r-1}(u) + u \mathbf{b}_{i+1}^{r-1}(u), \forall u \in [0, 1].$$

- By applying the induction hypothesis to the points  $\mathbf{b}_i^{r-1}(u)$  and  $\mathbf{b}_{i+1}^{r-1}(u)$  we can successively write:



# Bernstein representation of Bézier curves

## Parametric form of Bézier curves

### Proof.

- First, we perform the index transformation  $j \rightarrow j - i$ :

$$\sum_{j=0}^r \mathbf{b}_{j+i} B_j^r(u) = \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^r(u).$$

- Second, we prove the theorem by **mathematical induction with respect to  $r$** .
- The statement trivially holds for  $r = 0$ :

$$\mathbf{b}_i^0(u) = \sum_{j=i}^{i+0} \mathbf{b}_j B_{j-i}^0(u) = \mathbf{b}_i B_0^0(u) = \mathbf{b}_i, \forall u \in [0, 1], i = 0, 1, \dots, n.$$

- We assume that the statement is valid for all  $0 \leq k \leq r - 1$  and we also prove it for  $k = r$  ( $1 \leq r \leq n$ ).
- Based on the recurrence relation of the de Casteljau algorithm, we have:

$$\mathbf{b}_i^r(u) = (1 - u) \mathbf{b}_i^{r-1}(u) + u \mathbf{b}_{i+1}^{r-1}(u), \forall u \in [0, 1].$$

- By applying the induction hypothesis to the points  $\mathbf{b}_i^{r-1}(u)$  and  $\mathbf{b}_{i+1}^{r-1}(u)$  we can successively write:



# Bernstein representation of Bézier curves

## Parametric form of Bézier curves

### Proof.

- First, we perform the index transformation  $j \rightarrow j - i$ :

$$\sum_{j=0}^r \mathbf{b}_{j+i} B_j^r(u) = \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^r(u).$$

- Second, we prove the theorem by **mathematical induction with respect to  $r$** .
- The statement trivially holds for  $r = 0$ :

$$\mathbf{b}_i^0(u) = \sum_{j=i}^{i+0} \mathbf{b}_j B_{j-i}^0(u) = \mathbf{b}_i B_0^0(u) = \mathbf{b}_i, \forall u \in [0, 1], i = 0, 1, \dots, n.$$

- We assume that the statement is valid for all  $0 \leq k \leq r - 1$  and we also prove it for  $k = r$  ( $1 \leq r \leq n$ ).
- Based on the recurrence relation of the de Casteljau algorithm, we have:

$$\mathbf{b}_i^r(u) = (1 - u) \mathbf{b}_i^{r-1}(u) + u \mathbf{b}_{i+1}^{r-1}(u), \forall u \in [0, 1].$$

- By applying the induction hypothesis to the points  $\mathbf{b}_i^{r-1}(u)$  and  $\mathbf{b}_{i+1}^{r-1}(u)$  we can successively write:



# Bernstein representation of Bézier curves

## Parametric form of Bézier curves

### Proof.

- First, we perform the index transformation  $j \rightarrow j - i$ :

$$\sum_{j=0}^r \mathbf{b}_{j+i} B_j^r(u) = \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^r(u).$$

- Second, we prove the theorem by **mathematical induction with respect to  $r$** .
- The statement trivially holds for  $r = 0$ :

$$\mathbf{b}_i^0(u) = \sum_{j=i}^{i+0} \mathbf{b}_j B_{j-i}^0(u) = \mathbf{b}_i B_0^0(u) = \mathbf{b}_i, \forall u \in [0, 1], i = 0, 1, \dots, n.$$

- We assume that the statement is valid for all  $0 \leq k \leq r - 1$  and we also prove it for  $k = r$  ( $1 \leq r \leq n$ ).
- Based on the recurrence relation of the de Casteljau algorithm, we have:

$$\mathbf{b}_i^r(u) = (1 - u) \mathbf{b}_i^{r-1}(u) + u \mathbf{b}_{i+1}^{r-1}(u), \forall u \in [0, 1].$$

- By applying the induction hypothesis to the points  $\mathbf{b}_i^{r-1}(u)$  and  $\mathbf{b}_{i+1}^{r-1}(u)$  we can successively write:



# Bernstein representation of Bézier curves

## Parametric form of Bézier curves

### Proof.

- First, we perform the index transformation  $j \rightarrow j - i$ :

$$\sum_{j=0}^r \mathbf{b}_{j+i} B_j^r(u) = \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^r(u).$$

- Second, we prove the theorem by **mathematical induction with respect to  $r$** .
- The statement trivially holds for  $r = 0$ :

$$\mathbf{b}_i^0(u) = \sum_{j=i}^{i+0} \mathbf{b}_j B_{j-i}^0(u) = \mathbf{b}_i B_0^0(u) = \mathbf{b}_i, \forall u \in [0, 1], i = 0, 1, \dots, n.$$

- We assume that the statement is valid for all  $0 \leq k \leq r - 1$  and we also prove it for  $k = r$  ( $1 \leq r \leq n$ ).
- Based on the recurrence relation of the de Casteljau algorithm, we have:

$$\mathbf{b}_i^r(u) = (1 - u) \mathbf{b}_i^{r-1}(u) + u \mathbf{b}_{i+1}^{r-1}(u), \forall u \in [0, 1].$$

- By applying the induction hypothesis to the points  $\mathbf{b}_i^{r-1}(u)$  and  $\mathbf{b}_{i+1}^{r-1}(u)$ , we can successively write:



# Bernstein representation of Bézier curves

## Parametric form of Bézier curves

Proof – continued.

$$\begin{aligned} \mathbf{b}_i^r(u) &= (1-u) \sum_{j=i}^{i+r-1} \mathbf{b}_j B_{j-i}^{r-1}(u) + u \sum_{j=i+1}^{i+r} \mathbf{b}_j B_{j-i-1}^{r-1}(u) \\ &= (1-u) \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^{r-1}(u) + u \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i-1}^{r-1}(u) \\ &= \sum_{j=i}^{i+r} \mathbf{b}_j \underbrace{\left( (1-u) B_{j-i}^{r-1}(u) + u B_{j-i-1}^{r-1}(u) \right)}_{B_{j-i}^r(u)} \\ &= \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^r(u), \forall u \in [0, 1], \end{aligned}$$

where we have also applied the recurrence property of the Bernstein polynomials of degree  $r$ .



# Bernstein representation of Bézier curves

## Parametric form of Bézier curves

Proof – continued.

$$\begin{aligned}\mathbf{b}_i^r(u) &= (1-u) \sum_{j=i}^{i+r-1} \mathbf{b}_j B_{j-i}^{r-1}(u) + u \sum_{j=i+1}^{i+r} \mathbf{b}_j B_{j-i-1}^{r-1}(u) \\&= (1-u) \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^{r-1}(u) + u \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i-1}^{r-1}(u) \\&= \sum_{j=i}^{i+r} \mathbf{b}_j \underbrace{\left( (1-u) B_{j-i}^{r-1}(u) + u B_{j-i-1}^{r-1}(u) \right)}_{B_{j-i}^r(u)} \\&= \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^r(u), \forall u \in [0, 1],\end{aligned}$$

where we have also applied the recurrence property of the Bernstein polynomials of degree  $r$ .





# Bernstein representation of Bézier curves

## Parametric form of Bézier curves

Proof – continued.

$$\begin{aligned}\mathbf{b}_i^r(u) &= (1-u) \sum_{j=i}^{i+r-1} \mathbf{b}_j B_{j-i}^{r-1}(u) + u \sum_{j=i+1}^{i+r} \mathbf{b}_j B_{j-i-1}^{r-1}(u) \\&= (1-u) \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^{r-1}(u) + u \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i-1}^{r-1}(u) \\&= \sum_{j=i}^{i+r} \mathbf{b}_j \underbrace{\left( (1-u) B_{j-i}^{r-1}(u) + u B_{j-i-1}^{r-1}(u) \right)}_{B_{j-i}^r(u)} \\&= \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^r(u), \forall u \in [0, 1],\end{aligned}$$

where we have also applied the recurrence property of the Bernstein polynomials of degree  $r$ .



# Bernstein representation of Bézier curves

## Parametric form of Bézier curves

Proof – continued.

$$\begin{aligned}\mathbf{b}_i^r(u) &= (1-u) \sum_{j=i}^{i+r-1} \mathbf{b}_j B_{j-i}^{r-1}(u) + u \sum_{j=i+1}^{i+r} \mathbf{b}_j B_{j-i-1}^{r-1}(u) \\&= (1-u) \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^{r-1}(u) + u \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i-1}^{r-1}(u) \\&= \sum_{j=i}^{i+r} \mathbf{b}_j \underbrace{\left( (1-u) B_{j-i}^{r-1}(u) + u B_{j-i-1}^{r-1}(u) \right)}_{B_{j-i}^r(u)} \\&= \sum_{j=i}^{i+r} \mathbf{b}_j B_{j-i}^r(u), \forall u \in [0, 1],\end{aligned}$$

where we have also applied the recurrence property of the Bernstein polynomials of degree  $r$ .



# Properties of Bézier curves

## Closure for the affine transformations

The shape of a Bézier curve of degree  $n \geq 1$  is invariant under the affine transformation of its control polygon.

### Proof 1.

The de Casteljau algorithm consists of proportional subdivision points that are invariant under affine transformations. □

### Proof 2 – cf. Lecture 1.

The Bernstein polynomials of degree  $n \geq 1$  form the partition of unity. □

## Convex hull property




Fig. 4: The convex hull of a plane Bézier curve of degree 3.

The shape of a Bézier curve of degree  $n \geq 1$  lies in the convex hull of its control polygon.

### Proof 1.

It follows automatically from the de Casteljau algorithm, since all subdivision points are on a segment of the convex hull of the control polygon. □

### Proof 2 – cf. Lecture 1.

The Bernstein polynomials of degree  $n \geq 1$  form a normalized system of functions (i.e.

$$B_i^n(u) \geq 0, \sum_{i=0}^n B_i^n(u) \equiv 1, \forall u \in [0, 1]).$$



## Properties of Bézier curves

### Closure for the affine transformations

The shape of a Bézier curve of degree  $n \geq 1$  is invariant under the affine transformation of its control polygon.

#### Proof 1.

The de Casteljau algorithm consists of proportional subdivision points that are invariant under affine transformations. □

#### Proof 2 – cf. Lecture 1.

The Bernstein polynomials of degree  $n \geq 1$  form the partition of unity. □

### Convex hull property




Fig. 4: The convex hull of a plane Bézier curve of degree 3.

The shape of a Bézier curve of degree  $n \geq 1$  lies in the convex hull of its control polygon.

#### Proof 1.

It follows automatically from the de Casteljau algorithm, since all subdivision points are on a segment of the convex hull of the control polygon. □

#### Proof 2 – cf. Lecture 1.

The Bernstein polynomials of degree  $n \geq 1$  form a normalized system of functions (i.e.

$$B_i^n(u) \geq 0, \sum_{i=0}^n B_i^n(u) \equiv 1, \forall u \in [0, 1]).$$



# Properties of Bézier curves

## Closure for the affine transformations

The shape of a Bézier curve of degree  $n \geq 1$  is invariant under the affine transformation of its control polygon.

### Proof 1.

The de Casteljau algorithm consists of proportional subdivision points that are invariant under affine transformations. □

### Proof 2 – cf. Lecture 1.

The Bernstein polynomials of degree  $n \geq 1$  form the partition of unity. □

## Convex hull property




Fig. 4: The convex hull of a plane Bézier curve of degree 3.

The shape of a Bézier curve of degree  $n \geq 1$  lies in the convex hull of its control polygon.

### Proof 1.

It follows automatically from the de Casteljau algorithm, since all subdivision points are on a segment of the convex hull of the control polygon. □

### Proof 2 – cf. Lecture 1.

The Bernstein polynomials of degree  $n \geq 1$  form a normalized system of functions (i.e.

$$B_i^n(u) \geq 0, \sum_{i=0}^n B_i^n(u) \equiv 1, \forall u \in [0, 1]).$$



## Properties of Bézier curves

### Closure for the affine transformations

The shape of a Bézier curve of degree  $n \geq 1$  is invariant under the affine transformation of its control polygon.


#### Proof 1.

The de Casteljau algorithm consists of proportional subdivision points that are invariant under affine transformations. □

#### Proof 2 – cf. Lecture 1.

The Bernstein polynomials of degree  $n \geq 1$  form the partition of unity. □

### Convex hull property



**Fig. 4:** The convex hull of a plane Bézier curve of degree 3.

The shape of a Bézier curve of degree  $n \geq 1$  lies in the convex hull of its control polygon.

#### Proof 1.

It follows automatically from the de Casteljau algorithm, since all subdivision points are on a segment of the convex hull of the control polygon. □

#### Proof 2 – cf. Lecture 1.

The Bernstein polynomials of degree  $n \geq 1$  form a normalized system of functions (i.e.

$$B_i^n(u) \geq 0, \sum_{i=0}^n B_i^n(u) \equiv 1, \forall u \in [0, 1]).$$

© Agoston Rósch, 2022



# Properties of Bézier curves

## Closure for the affine transformations

The shape of a Bézier curve of degree  $n \geq 1$  is invariant under the affine transformation of its control polygon.


### Proof 1.

The de Casteljau algorithm consists of proportional subdivision points that are invariant under affine transformations. □

### Proof 2 – cf. Lecture 1.

The Bernstein polynomials of degree  $n \geq 1$  form the partition of unity. □

## Convex hull property



**Fig. 4:** The convex hull of a plane Bézier curve of degree 3.

The shape of a Bézier curve of degree  $n \geq 1$  lies in the convex hull of its control polygon.

### Proof 1.

It follows automatically from the de Casteljau algorithm, since all subdivision points are on a segment of the convex hull of the control polygon. □

### Proof 2 – cf. Lecture 1.

The Bernstein polynomials of degree  $n \geq 1$  form a normalized system of functions (i.e.

$$B_i^n(u) \geq 0, \sum_{i=0}^n B_i^n(u) \equiv 1, \forall u \in [0, 1]).$$



# Properties of Bézier curves

## Closure for the affine transformations

The shape of a Bézier curve of degree  $n \geq 1$  is invariant under the affine transformation of its control polygon.


### Proof 1.

The de Casteljau algorithm consists of proportional subdivision points that are invariant under affine transformations. □

### Proof 2 – cf. Lecture 1.

The Bernstein polynomials of degree  $n \geq 1$  form the partition of unity. □

## Convex hull property



**Fig. 4:** The convex hull of a plane Bézier curve of degree 3.

The shape of a Bézier curve of degree  $n \geq 1$  lies in the convex hull of its control polygon.

### Proof 1.

It follows automatically from the de Casteljau algorithm, since all subdivision points are on a segment of the convex hull of the control polygon. □

### Proof 2 – cf. Lecture 1.

The Bernstein polynomials of degree  $n \geq 1$  form a normalized system of functions (i.e.

$$B_i^n(u) \geq 0, \sum_{i=0}^n B_i^n(u) \equiv 1, \forall u \in [0, 1]).$$





## Properties of Bézier curves

### Invariance under linear transformations of the parameter domain

The shape of the Bézier curve of degree  $n \geq 1$  is invariant under affine transformations of the parameter domain.

Proof.

The proportional subdivision points of the de Casteljau algorithm do not change under the linear parameter transformation

$$u(t) = \frac{t - a}{b - a}, \quad t \in [a, b].$$



### Endpoint interpolation

The Bézier curve of degree  $n \geq 1$  interpolates the first and last control point of its control polygon.

Proof.

$$\mathbf{b}(0) = \sum_{i=0}^n \mathbf{b}_i B_i^n(0) = \mathbf{b}_0 B_0^n(0) + \mathbf{0} = \mathbf{b}_0,$$

$$\mathbf{b}(1) = \sum_{i=0}^n \mathbf{b}_i B_i^n(1) = \mathbf{0} + \mathbf{b}_n B_n^n(1) = \mathbf{b}_n. \quad \square$$



## Properties of Bézier curves

### Invariance under linear transformations of the parameter domain

The shape of the Bézier curve of degree  $n \geq 1$  is invariant under affine transformations of the parameter domain.

**Proof.**

The proportional subdivision points of the de Casteljau algorithm do not change under the linear parameter transformation

$$u(t) = \frac{t - a}{b - a}, \quad t \in [a, b].$$



### Endpoint interpolation

The Bézier curve of degree  $n \geq 1$  interpolates the first and last control point of its control polygon.

**Proof.**

$$\mathbf{b}(0) = \sum_{i=0}^n \mathbf{b}_i B_i^n(0) = \mathbf{b}_0 B_0^n(0) + \mathbf{0} = \mathbf{b}_0,$$

$$\mathbf{b}(1) = \sum_{i=0}^n \mathbf{b}_i B_i^n(1) = \mathbf{0} + \mathbf{b}_n B_n^n(1) = \mathbf{b}_n. \quad \square$$



## Properties of Bézier curves

### Invariance under linear transformations of the parameter domain

The shape of the Bézier curve of degree  $n \geq 1$  is invariant under affine transformations of the parameter domain.

**Proof.**

The proportional subdivision points of the de Casteljau algorithm do not change under the linear parameter transformation

$$u(t) = \frac{t - a}{b - a}, \quad t \in [a, b].$$



### Endpoint interpolation

The Bézier curve of degree  $n \geq 1$  interpolates the first and last control point of its control polygon.

**Proof.**

$$\mathbf{b}(0) = \sum_{i=0}^n \mathbf{b}_i B_i^n(0) = \mathbf{b}_0 B_0^n(0) + \mathbf{0} = \mathbf{b}_0,$$

$$\mathbf{b}(1) = \sum_{i=0}^n \mathbf{b}_i B_i^n(1) = \mathbf{0} + \mathbf{b}_n B_n^n(1) = \mathbf{b}_n. \quad \square$$



## Properties of Bézier curves

### Invariance under linear transformations of the parameter domain

The shape of the Bézier curve of degree  $n \geq 1$  is invariant under affine transformations of the parameter domain.

**Proof.**

The proportional subdivision points of the de Casteljau algorithm do not change under the linear parameter transformation

$$u(t) = \frac{t - a}{b - a}, \quad t \in [a, b].$$

□

### Endpoint interpolation

The Bézier curve of degree  $n \geq 1$  interpolates the first and last control point of its control polygon.

**Proof.**

$$\begin{aligned} \mathbf{b}(0) &= \sum_{i=0}^n \mathbf{b}_i B_i^n(0) = \mathbf{b}_0 B_0^n(0) + \mathbf{0} = \mathbf{b}_0, \\ \mathbf{b}(1) &= \sum_{i=0}^n \mathbf{b}_i B_i^n(1) = \mathbf{0} + \mathbf{b}_n B_n^n(1) = \mathbf{b}_n. \quad \square \end{aligned}$$



# Properties of Bézier curves

## Symmetry

The Bézier curve of degree  $n \geq 1$  defined by the control polygon  $[\mathbf{b}_i]_{i=0}^n$  is symmetric, i.e. the control polygon  $[\mathbf{b}_{n-i}]_{i=0}^n$  generates the same Bézier curve (of course with different orientation).

### Proof.

Since

$$\begin{aligned} B_i^n(u) &= \binom{n}{i} u^i (1-u)^{n-i} \\ &= \binom{n}{n-i} (1-u)^{n-i} (1-(1-u))^{n-(n-i)} \\ &= B_{n-i}^n(1-u), \quad \forall u \in [0, 1], \end{aligned}$$

it follows that

$$\mathbf{b}(u) = \sum_{i=0}^n \mathbf{b}_i B_i^n(u) = \sum_{i=0}^n \mathbf{b}_{n-i} B_{n-i}^n(1-u), \quad \forall u \in [0, 1].$$

□



# Properties of Bézier curves

## Symmetry

The Bézier curve of degree  $n \geq 1$  defined by the control polygon  $[\mathbf{b}_i]_{i=0}^n$  is symmetric, i.e. the control polygon  $[\mathbf{b}_{n-i}]_{i=0}^n$  generates the same Bézier curve (of course with different orientation).

## Proof.

Since

$$\begin{aligned} B_i^n(u) &= \binom{n}{i} u^i (1-u)^{n-i} \\ &= \binom{n}{n-i} (1-u)^{n-i} (1-(1-u))^{n-(n-i)} \\ &= B_{n-i}^n(1-u), \quad \forall u \in [0, 1], \end{aligned}$$

it follows that

$$\mathbf{b}(u) = \sum_{i=0}^n \mathbf{b}_i B_i^n(u) = \sum_{i=0}^n \mathbf{b}_{n-i} B_{n-i}^n(1-u), \quad \forall u \in [0, 1].$$

□



# Properties of Bézier curves

## Symmetry

The Bézier curve of degree  $n \geq 1$  defined by the control polygon  $[\mathbf{b}_i]_{i=0}^n$  is symmetric, i.e. the control polygon  $[\mathbf{b}_{n-i}]_{i=0}^n$  generates the same Bézier curve (of course with different orientation).

### Proof.

Since

$$\begin{aligned} B_i^n(u) &= \binom{n}{i} u^i (1-u)^{n-i} \\ &= \binom{n}{n-i} (1-u)^{n-i} (1-(1-u))^{n-(n-i)} \\ &= B_{n-i}^n(1-u), \quad \forall u \in [0, 1], \end{aligned}$$

it follows that

$$\mathbf{b}(u) = \sum_{i=0}^n \mathbf{b}_i B_i^n(u) = \sum_{i=0}^n \mathbf{b}_{n-i} B_{n-i}^n(1-u), \quad \forall u \in [0, 1].$$

□



# Properties of Bézier curves

## Symmetry

The Bézier curve of degree  $n \geq 1$  defined by the control polygon  $[\mathbf{b}_i]_{i=0}^n$  is symmetric, i.e. the control polygon  $[\mathbf{b}_{n-i}]_{i=0}^n$  generates the same Bézier curve (of course with different orientation).

## Proof.

Since

$$\begin{aligned} B_i^n(u) &= \binom{n}{i} u^i (1-u)^{n-i} \\ &= \binom{n}{n-i} (1-u)^{n-i} (1-(1-u))^{n-(n-i)} \\ &= B_{n-i}^n(1-u), \quad \forall u \in [0, 1], \end{aligned}$$

it follows that

$$\mathbf{b}(u) = \sum_{i=0}^n \mathbf{b}_i B_i^n(u) = \sum_{i=0}^n \mathbf{b}_{n-i} B_{n-i}^n(1-u), \quad \forall u \in [0, 1].$$

□





## Properties of Bézier curves

### Symmetry

The Bézier curve of degree  $n \geq 1$  defined by the control polygon  $[\mathbf{b}_i]_{i=0}^n$  is symmetric, i.e. the control polygon  $[\mathbf{b}_{n-i}]_{i=0}^n$  generates the same Bézier curve (of course with different orientation).

### Proof.

Since

$$\begin{aligned} B_i^n(u) &= \binom{n}{i} u^i (1-u)^{n-i} \\ &= \binom{n}{n-i} (1-u)^{n-i} (1-(1-u))^{n-(n-i)} \\ &= B_{n-i}^n(1-u), \quad \forall u \in [0, 1], \end{aligned}$$

it follows that

$$\mathbf{b}(u) = \sum_{i=0}^n \mathbf{b}_i B_i^n(u) = \sum_{i=0}^n \mathbf{b}_{n-i} B_{n-i}^n(1-u), \quad \forall u \in [0, 1].$$



## Properties of Bézier curves

### Global/pseudo local controllability




Fig. 5: Effect of a translated control point

$$\begin{aligned}
 \mathbf{b}_{i,t}(u) &= \sum_{j=0, j \neq i}^n \mathbf{b}_j B_j^n(u) + (\mathbf{b}_i + \mathbf{t}) B_i^n(u) \\
 &= \sum_{j=0}^n \mathbf{b}_j B_j^n(u) + \mathbf{t} B_i^n(u) \\
 &= \mathbf{b}(u) + \mathbf{t} B_i^n(u), \forall u \in [0, 1].
 \end{aligned}$$

Modification of a single control point  $\mathbf{b}_i$  has a global effect on the shape of the Bézier curve

$$\mathbf{b}(u) = \sum_{j=0}^n \mathbf{b}_j B_j^n(u), \quad u \in [0, 1]$$

of degree  $n \geq 1$ .

Proof.

$$B_j^n(u) > 0, \forall u \in (0, 1), j = 0, \dots, n. \quad \square$$

However, we also say that the curve is **pseudo locally controllable**, because the maximal effect on the shape of the curve, generated by the modification of the control point  $\mathbf{b}_i$ , can be easily forecasted since the unique maximal weight of a control point is  $B_i^n(\bar{u}_i)$ , where  $\bar{u}_i = \frac{i}{n}$ .



## Properties of Bézier curves

### Global/pseudo local controllability




Fig. 5: Effect of a translated control point

$$\begin{aligned}
 \mathbf{b}_{i,t}(u) &= \sum_{j=0, j \neq i}^n \mathbf{b}_j B_j^n(u) + (\mathbf{b}_i + \mathbf{t}) B_i^n(u) \\
 &= \sum_{j=0}^n \mathbf{b}_j B_j^n(u) + \mathbf{t} B_i^n(u) \\
 &= \mathbf{b}(u) + \mathbf{t} B_i^n(u), \forall u \in [0, 1].
 \end{aligned}$$

Modification of a single control point  $\mathbf{b}_i$  has a global effect on the shape of the Bézier curve

$$\mathbf{b}(u) = \sum_{j=0}^n \mathbf{b}_j B_j^n(u), \quad u \in [0, 1]$$

of degree  $n \geq 1$ .

**Proof.**

$B_i^n(u) > 0, \forall u \in (0, 1), j = 0, \dots, n.$   $\square$

However, we also say that the curve is **pseudo locally controllable**, because the maximal effect on the shape of the curve, generated by the modification of the control point  $\mathbf{b}_i$ , can be easily forecasted, since the unique maximal weight of this control point is  $B_i^n(\bar{u}_i)$ , where  $\bar{u}_i = \frac{i}{n}$ .

## Properties of Bézier curves

### Global/pseudo local controllability




Fig. 5: Effect of a translated control point

$$\begin{aligned}
 \mathbf{b}_{i,t}(u) &= \sum_{j=0, j \neq i}^n \mathbf{b}_j B_j^n(u) + (\mathbf{b}_i + \mathbf{t}) B_i^n(u) \\
 &= \sum_{j=0}^n \mathbf{b}_j B_j^n(u) + \mathbf{t} B_i^n(u) \\
 &= \mathbf{b}(u) + \mathbf{t} B_i^n(u), \forall u \in [0, 1].
 \end{aligned}$$

Modification of a single control point  $\mathbf{b}_i$  has a global effect on the shape of the Bézier curve

$$\mathbf{b}(u) = \sum_{j=0}^n \mathbf{b}_j B_j^n(u), \quad u \in [0, 1]$$

of degree  $n \geq 1$ .

**Proof.**

$$B_i^n(u) > 0, \quad \forall u \in (0, 1), \quad j = 0, \dots, n. \quad \square$$

However, we also say that the curve is **pseudo locally controllable**, because the maximal effect on the shape of the curve, generated by the modification of the control point  $\mathbf{b}_i$ , can be easily forecasted, since the unique maximal weight of this control point is  $B_i^n(\bar{u}_i)$ , where  $\bar{u}_i = \frac{i}{n}$ .

## Properties of Bézier curves

### Global/pseudo local controllability




Fig. 5: Effect of a translated control point

$$\begin{aligned}
 \mathbf{b}_{i,t}(u) &= \sum_{j=0, j \neq i}^n \mathbf{b}_j B_j^n(u) + (\mathbf{b}_i + \mathbf{t}) B_i^n(u) \\
 &= \sum_{j=0}^n \mathbf{b}_j B_j^n(u) + \mathbf{t} B_i^n(u) \\
 &= \mathbf{b}(u) + \mathbf{t} B_i^n(u), \forall u \in [0, 1].
 \end{aligned}$$

Modification of a single control point  $\mathbf{b}_i$  has a global effect on the shape of the Bézier curve

$$\mathbf{b}(u) = \sum_{j=0}^n \mathbf{b}_j B_j^n(u), \quad u \in [0, 1]$$

of degree  $n \geq 1$ .

**Proof.**

$$B_i^n(u) > 0, \quad \forall u \in (0, 1), \quad j = 0, \dots, n. \quad \square$$

However, we also say that the curve is **pseudo locally controllable**, because the maximal effect on the shape of the curve, generated by the modification of the control point  $\mathbf{b}_i$ , can be easily forecasted, since the unique maximal weight of this control point is  $B_i^n(\bar{u}_i)$ , where  $\bar{u}_i = \frac{i}{n}$ .

## Properties of Bézier curves

### Global/pseudo local controllability




Fig. 5: Effect of a translated control point

$$\begin{aligned}
 \mathbf{b}_{i,t}(u) &= \sum_{j=0, j \neq i}^n \mathbf{b}_j B_j^n(u) + (\mathbf{b}_i + \mathbf{t}) B_i^n(u) \\
 &= \sum_{j=0}^n \mathbf{b}_j B_j^n(u) + \mathbf{t} B_i^n(u) \\
 &= \mathbf{b}(u) + \mathbf{t} B_i^n(u), \forall u \in [0, 1].
 \end{aligned}$$

Modification of a single control point  $\mathbf{b}_i$  has a global effect on the shape of the Bézier curve

$$\mathbf{b}(u) = \sum_{j=0}^n \mathbf{b}_j B_j^n(u), \quad u \in [0, 1]$$

of degree  $n \geq 1$ .

**Proof.**

$$B_i^n(u) > 0, \quad \forall u \in (0, 1), \quad j = 0, \dots, n. \quad \square$$

However, we also say that the curve is **pseudo locally controllable**, because the maximal effect on the shape of the curve, generated by the modification of the control point  $\mathbf{b}_i$ , can be easily forecasted, since the unique maximal weight of this control point is  $B_i^n(\bar{u}_i)$ , where  $\bar{u}_i = \frac{i}{n}$ .

## Properties of Bézier curves

### Global/pseudo local controllability




Fig. 5: Effect of a translated control point

$$\begin{aligned}
 \mathbf{b}_{i,t}(u) &= \sum_{j=0, j \neq i}^n \mathbf{b}_j B_j^n(u) + (\mathbf{b}_i + \mathbf{t}) B_i^n(u) \\
 &= \sum_{j=0}^n \mathbf{b}_j B_j^n(u) + \mathbf{t} B_i^n(u) \\
 &= \mathbf{b}(u) + \mathbf{t} B_i^n(u), \forall u \in [0, 1].
 \end{aligned}$$

Modification of a single control point  $\mathbf{b}_i$  has a global effect on the shape of the Bézier curve

$$\mathbf{b}(u) = \sum_{j=0}^n \mathbf{b}_j B_j^n(u), \quad u \in [0, 1]$$

of degree  $n \geq 1$ .

**Proof.**

$$B_i^n(u) > 0, \quad \forall u \in (0, 1), \quad j = 0, \dots, n. \quad \square$$

However, we also say that the curve is **pseudo locally controllable**, because the maximal effect on the shape of the curve, generated by the modification of the control point  $\mathbf{b}_i$ , can be easily forecasted, since the unique maximal weight of this control point is  $B_i^n(\bar{u}_i)$ , where  $\bar{u}_i = \frac{i}{n}$ .

# Properties of Bézier curves

## Representation of lines

The Bézier curve is a line if and only if its **control points are collinear**.

**Proof.**

Immediate corollary of the de Casteljau algorithm. □

## Linear precision

If the control points are uniformly distributed on a straight line, then the Bézier curve determined by them describes exactly the initial straight line. This property is equivalent to

$$\sum_{i=0}^n \frac{i}{n} B_i^n(u) \equiv u, \forall u \in [0, 1]$$

and it is called **linear precision property**.

**Proof.**

$$\begin{aligned} \sum_{i=0}^n \frac{i}{n} B_i^n(u) &= \sum_{i=1}^n \frac{i}{n} B_i^n(u) = u \sum_{i=1}^n \frac{(n-1)!}{(i-1)!(n-i)!} u^{i-1} (1-u)^{n-i} \\ &\stackrel{i-1 \rightarrow i}{=} u \sum_{i=0}^{n-1} \frac{(n-1)!}{i!(n-1-i)!} u^i (1-u)^{n-1-i} \equiv u, \forall u \in [0, 1]. \end{aligned}$$





# Properties of Bézier curves

## Representation of lines

The Bézier curve is a line if and only if its **control points are collinear**.

**Proof.**

Immediate corollary of the de Casteljau algorithm. □

## Linear precision

If the control points are uniformly distributed on a straight line, then the Bézier curve determined by them describes exactly the initial straight line. This property is equivalent to

$$\sum_{i=0}^n \frac{i}{n} B_i^n(u) \equiv u, \forall u \in [0, 1]$$

and it is called **linear precision property**.

**Proof.**

$$\begin{aligned} \sum_{i=0}^n \frac{i}{n} B_i^n(u) &= \sum_{i=1}^n \frac{i}{n} B_i^n(u) = u \sum_{i=1}^n \frac{(n-1)!}{(i-1)!(n-i)!} u^{i-1} (1-u)^{n-i} \\ &\stackrel{i-1 \rightarrow i}{=} u \sum_{i=0}^{n-1} \frac{(n-1)!}{i!(n-1-i)!} u^i (1-u)^{n-1-i} \equiv u, \forall u \in [0, 1]. \end{aligned}$$



# Properties of Bézier curves

## Representation of lines

The Bézier curve is a line if and only if its **control points are collinear**.

**Proof.**

Immediate corollary of the de Casteljau algorithm. □

## Linear precision

If the control points are uniformly distributed on a straight line, then the Bézier curve determined by them describes exactly the initial straight line. This property is equivalent to

$$\sum_{i=0}^n \frac{i}{n} B_i^n(u) \equiv u, \forall u \in [0, 1]$$

and it is called **linear precision property**.

**Proof.**

$$\begin{aligned} \sum_{i=0}^n \frac{i}{n} B_i^n(u) &= \sum_{i=1}^n \frac{i}{n} B_i^n(u) = u \sum_{i=1}^n \frac{(n-1)!}{(i-1)!(n-i)!} u^{i-1} (1-u)^{n-i} \\ &\stackrel{i-1 \rightarrow i}{=} u \sum_{i=0}^{n-1} \frac{(n-1)!}{i!(n-1-i)!} u^i (1-u)^{n-1-i} \equiv u, \forall u \in [0, 1]. \end{aligned}$$



# Properties of Bézier curves

## Representation of lines

The Bézier curve is a line if and only if its **control points are collinear**.

**Proof.**

Immediate corollary of the de Casteljau algorithm. □

## Linear precision

If the control points are uniformly distributed on a straight line, then the Bézier curve determined by them describes exactly the initial straight line. This property is equivalent to

$$\sum_{i=0}^n \frac{i}{n} B_i^n(u) \equiv u, \forall u \in [0, 1]$$

and it is called **linear precision property**.

**Proof.**

$$\begin{aligned} \sum_{i=0}^n \frac{i}{n} B_i^n(u) &= \sum_{i=1}^n \frac{i}{n} B_i^n(u) = u \sum_{i=1}^n \frac{(n-1)!}{(i-1)!(n-i)!} u^{i-1} (1-u)^{n-i} \\ &\stackrel{i-1 \rightarrow i}{=} u \sum_{i=0}^{n-1} \frac{(n-1)!}{i!(n-1-i)!} u^i (1-u)^{n-1-i} \equiv u, \forall u \in [0, 1]. \end{aligned}$$



## Properties of Bézier curves

### Hodograph of a Bézier curve of degree $n \geq 1$

We can successively write:

$$\begin{aligned}\frac{d}{du} \mathbf{b}(u) &= \frac{d}{du} \left( \sum_{i=0}^n \mathbf{b}_i B_i^n(u) \right) = \sum_{i=0}^n \mathbf{b}_i \frac{d}{du} B_i^n(u) \\&= n \sum_{i=0}^n \mathbf{b}_i \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right) \\&= n \sum_{i=0}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^n \mathbf{b}_i B_i^{n-1}(u) \\&= n \sum_{i=1}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u) \\&\stackrel{i-1 \rightarrow i}{=} n \sum_{i=0}^{n-1} \mathbf{b}_{i+1} B_i^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u) \\&= n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u) \\&= \sum_{i=0}^{n-1} \Delta \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1]; \Delta \mathbf{b}_i = n(\mathbf{b}_{i+1} - \mathbf{b}_i), i = 0, 1, \dots, n-1.\end{aligned}$$



## Properties of Bézier curves

### Hodograph of a Bézier curve of degree $n \geq 1$

We can successively write:

$$\frac{d}{du} \mathbf{b}(u) = \frac{d}{du} \left( \sum_{i=0}^n \mathbf{b}_i B_i^n(u) \right) = \sum_{i=0}^n \mathbf{b}_i \frac{d}{du} B_i^n(u)$$

$$= n \sum_{i=0}^n \mathbf{b}_i \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right)$$

$$= n \sum_{i=0}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^n \mathbf{b}_i B_i^{n-1}(u)$$

$$= n \sum_{i=1}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u)$$

$$\stackrel{i-1 \rightarrow i}{=} n \sum_{i=0}^{n-1} \mathbf{b}_{i+1} B_i^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u)$$

$$= n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u)$$

$$= \sum_{i=0}^{n-1} \Delta \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1]; \Delta \mathbf{b}_i = n(\mathbf{b}_{i+1} - \mathbf{b}_i), i = 0, 1, \dots$$



## Properties of Bézier curves

### Hodograph of a Bézier curve of degree $n \geq 1$

We can successively write:

$$\frac{d}{du} \mathbf{b}(u) = \frac{d}{du} \left( \sum_{i=0}^n \mathbf{b}_i B_i^n(u) \right) = \sum_{i=0}^n \mathbf{b}_i \frac{d}{du} B_i^n(u)$$

$$= n \sum_{i=0}^n \mathbf{b}_i \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right)$$

$$= n \sum_{i=0}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^n \mathbf{b}_i B_i^{n-1}(u)$$

$$= n \sum_{i=1}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u)$$

$$\stackrel{i-1 \rightarrow i}{=} n \sum_{i=0}^{n-1} \mathbf{b}_{i+1} B_i^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u)$$

$$= n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u)$$

$$= \sum_{i=0}^{n-1} \Delta \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1]; \Delta \mathbf{b}_i = n(\mathbf{b}_{i+1} - \mathbf{b}_i), i = 0, 1, \dots$$



## Properties of Bézier curves

### Hodograph of a Bézier curve of degree $n \geq 1$

We can successively write:

$$\frac{d}{du} \mathbf{b}(u) = \frac{d}{du} \left( \sum_{i=0}^n \mathbf{b}_i B_i^n(u) \right) = \sum_{i=0}^n \mathbf{b}_i \frac{d}{du} B_i^n(u)$$

$$= n \sum_{i=0}^n \mathbf{b}_i \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right)$$

$$= n \sum_{i=0}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^n \mathbf{b}_i B_i^{n-1}(u)$$

$$= n \sum_{i=1}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u)$$

$$\stackrel{i-1 \rightarrow i}{=} n \sum_{i=0}^{n-1} \mathbf{b}_{i+1} B_i^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u)$$

$$= n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u)$$

$$= \sum_{i=0}^{n-1} \Delta \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1]; \Delta \mathbf{b}_i = n(\mathbf{b}_{i+1} - \mathbf{b}_i), i = 0, 1, \dots$$



## Properties of Bézier curves

### Hodograph of a Bézier curve of degree $n \geq 1$

We can successively write:

$$\frac{d}{du} \mathbf{b}(u) = \frac{d}{du} \left( \sum_{i=0}^n \mathbf{b}_i B_i^n(u) \right) = \sum_{i=0}^n \mathbf{b}_i \frac{d}{du} B_i^n(u)$$

$$= n \sum_{i=0}^n \mathbf{b}_i \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right)$$

$$= n \sum_{i=0}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^n \mathbf{b}_i B_i^{n-1}(u)$$

$$= n \sum_{i=1}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u)$$

$$\stackrel{i-1 \rightarrow i}{=} n \sum_{i=0}^{n-1} \mathbf{b}_{i+1} B_i^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u)$$

$$= n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u)$$

$$= \sum_{i=0}^{n-1} \Delta \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1]; \Delta \mathbf{b}_i = n(\mathbf{b}_{i+1} - \mathbf{b}_i), i = 0, 1, \dots, n-1$$





## Properties of Bézier curves

### Hodograph of a Bézier curve of degree $n \geq 1$

We can successively write:

$$\frac{d}{du} \mathbf{b}(u) = \frac{d}{du} \left( \sum_{i=0}^n \mathbf{b}_i B_i^n(u) \right) = \sum_{i=0}^n \mathbf{b}_i \frac{d}{du} B_i^n(u)$$

$$= n \sum_{i=0}^n \mathbf{b}_i \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right)$$

$$= n \sum_{i=0}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^n \mathbf{b}_i B_i^{n-1}(u)$$

$$= n \sum_{i=1}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u)$$

$$\stackrel{i-1 \rightarrow i}{=} n \sum_{i=0}^{n-1} \mathbf{b}_{i+1} B_i^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u)$$

$$= n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u)$$

$$= \sum_{i=0}^{n-1} \Delta \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1]; \Delta \mathbf{b}_i = n(\mathbf{b}_{i+1} - \mathbf{b}_i), i = 0, 1, \dots$$



## Properties of Bézier curves

### Hodograph of a Bézier curve of degree $n \geq 1$

We can successively write:

$$\frac{d}{du} \mathbf{b}(u) = \frac{d}{du} \left( \sum_{i=0}^n \mathbf{b}_i B_i^n(u) \right) = \sum_{i=0}^n \mathbf{b}_i \frac{d}{du} B_i^n(u)$$

$$= n \sum_{i=0}^n \mathbf{b}_i \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right)$$

$$= n \sum_{i=0}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^n \mathbf{b}_i B_i^{n-1}(u)$$

$$= n \sum_{i=1}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u)$$

$$\stackrel{i-1 \rightarrow i}{=} n \sum_{i=0}^{n-1} \mathbf{b}_{i+1} B_i^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u)$$

$$= n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u)$$

$$= \sum_{i=0}^{n-1} \Delta \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1]; \Delta \mathbf{b}_i = n(\mathbf{b}_{i+1} - \mathbf{b}_i), i = 0, 1, \dots, n-1$$



## Properties of Bézier curves

### Hodograph of a Bézier curve of degree $n \geq 1$

We can successively write:

$$\begin{aligned}\frac{d}{du} \mathbf{b}(u) &= \frac{d}{du} \left( \sum_{i=0}^n \mathbf{b}_i B_i^n(u) \right) = \sum_{i=0}^n \mathbf{b}_i \frac{d}{du} B_i^n(u) \\&= n \sum_{i=0}^n \mathbf{b}_i \left( B_{i-1}^{n-1}(u) - B_i^{n-1}(u) \right) \\&= n \sum_{i=0}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^n \mathbf{b}_i B_i^{n-1}(u) \\&= n \sum_{i=1}^n \mathbf{b}_i B_{i-1}^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u) \\&\stackrel{i-1 \rightarrow i}{=} n \sum_{i=0}^{n-1} \mathbf{b}_{i+1} B_i^{n-1}(u) - n \sum_{i=0}^{n-1} \mathbf{b}_i B_i^{n-1}(u) \\&= n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u) \\&= \sum_{i=0}^{n-1} \Delta \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1]; \Delta \mathbf{b}_i = n(\mathbf{b}_{i+1} - \mathbf{b}_i), i = 0, 1, \dots, n-1.\end{aligned}$$



## Properties of Bézier curves

### Corollary (Bézier curves are closed under derivation)

The hodograph (i.e. first order derivative) of the Bézier curve

$$\mathbf{b}(u) = \sum_{i=0}^n \mathbf{b}_i B_i^n(u), \quad u \in [0, 1]$$

of degree  $n \geq 1$  is also a Bézier curve of degree  $n - 1$  that is determined by the control polygon

$$[\Delta \mathbf{b}_i]_{i=0}^{n-1} = [n(\mathbf{b}_{i+1} - \mathbf{b}_i)]_{i=0}^{n-1}.$$

### Corollary (Hodograph diminishing)




Fig. 6: Any Bézier curve fulfills the hodograph diminishing property

© Agoston Rósch, 2022



## Properties of Bézier curves

### Corollary (Bézier curves are closed under derivation)

The hodograph (i.e. first order derivative) of the Bézier curve

$$\mathbf{b}(u) = \sum_{i=0}^n \mathbf{b}_i B_i^n(u), \quad u \in [0, 1]$$

of degree  $n \geq 1$  is also a Bézier curve of degree  $n - 1$  that is determined by the control polygon

$$[\Delta \mathbf{b}_i]_{i=0}^{n-1} = [n(\mathbf{b}_{i+1} - \mathbf{b}_i)]_{i=0}^{n-1}.$$

### Corollary (Hodograph diminishing)





Fig. 6: Any Bézier curve fulfills the hodograph diminishing property



# Properties of Bézier curves


## Inflection points



**Fig. 7:** If the hodograph has a tangent line that goes through the origin and its touching point is a regular point of the hodograph, then the Bézier curve has an inflection point.

# Properties of Bézier curves


## Inflection points



**Fig. 7:** If the hodograph has a tangent line that goes through the origin and its touching point is a regular point of the hodograph, then the Bézier curve has an inflection point.

# Properties of Bézier curves

## Inflection points




**Fig. 7:** If the hodograph has a tangent line that goes through the origin and its touching point is a regular point of the hodograph, then the Bézier curve has an inflection point.



# Properties of Bézier curves

## Inflection points




**Fig. 7:** If the hodograph has a tangent line that goes through the origin and its touching point is a regular point of the hodograph, then the Bézier curve has an inflection point.



# Properties of Bézier curves


## Inflection points



**Fig. 7:** If the hodograph has a tangent line that goes through the origin and its touching point is a regular point of the hodograph, then the Bézier curve has an inflection point.

# Properties of Bézier curves


## Inflection points



**Fig. 7:** If the hodograph has a tangent line that goes through the origin and its touching point is a regular point of the hodograph, then the Bézier curve has an inflection point.

# Properties of Bézier curves


## Inflection points



**Fig. 7:** If the hodograph has a tangent line that goes through the origin and its touching point is a regular point of the hodograph, then the Bézier curve has an inflection point.

# Properties of Bézier curves

## Cusps




**Fig. 8:** If the hodograph goes through the origin, then the Bézier curve has a cusp.



# Properties of Bézier curves

## Cusps




**Fig. 8:** If the hodograph goes through the origin, then the Bézier curve has a cusp.



# Properties of Bézier curves

## Cusps




**Fig. 8:** If the hodograph goes through the origin, then the Bézier curve has a cusp.



# Properties of Bézier curves

## Cusps




**Fig. 8:** If the hodograph goes through the origin, then the Bézier curve has a cusp.





# Properties of Bézier curves

## Cusps




**Fig. 8:** If the hodograph goes through the origin, then the Bézier curve has a cusp.



# Properties of Bézier curves

## Cusps




**Fig. 8:** If the hodograph goes through the origin, then the Bézier curve has a cusp.



# Properties of Bézier curves


## Null curvature points



**Fig. 9:** If the hodograph has a tangent line that goes through the origin and its touching point is either a cusp or an inflection point of the hodograph, then the Bézier curve has a null curvature point.

# Properties of Bézier curves

## Null curvature points



**Fig. 9:** If the hodograph has a tangent line that goes through the origin and its touching point is either a cusp or an inflection point of the hodograph, then the Bézier curve has a null curvature point.




## Null curvature points

**Fig. 9:** If the hodograph has a tangent line that goes through the origin and its touching point is either a cusp or an inflection point of the hodograph, then the Bézier curve has a null curvature point.

# Properties of Bézier curves

## Null curvature points




**Fig. 9:** If the hodograph has a tangent line that goes through the origin and its touching point is either a cusp or an inflection point of the hodograph, then the Bézier curve has a null curvature point.



# Properties of Bézier curves


## Null curvature points



**Fig. 9:** If the hodograph has a tangent line that goes through the origin and its touching point is either a cusp or an inflection point of the hodograph, then the Bézier curve has a null curvature point.

# Properties of Bézier curves

## Null curvature points




**Fig. 9:** If the hodograph has a tangent line that goes through the origin and its touching point is either a cusp or an inflection point of the hodograph, then the Bézier curve has a null curvature point.



# Properties of Bézier curves


## Null curvature points



**Fig. 9:** If the hodograph has a tangent line that goes through the origin and its touching point is either a cusp or an inflection point of the hodograph, then the Bézier curve has a null curvature point.

# Properties of Bézier curves

## Null curvature points



**Fig. 9:** If the hodograph has a tangent line that goes through the origin and its touching point is either a cusp or an inflection point of the hodograph, then the Bézier curve has a null curvature point.

## Properties of Bézier curves

A differential operator of order  $r \geq 0$

$$\begin{cases} \Delta^0 \mathbf{b}_i = \mathbf{b}_i, \\ \Delta^r \mathbf{b}_i = \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i, \quad r \geq 1. \end{cases}$$

Example ( $r = 1, 2, 3$ )

$$\begin{aligned} \Delta^1 \mathbf{b}_i &= \Delta^0 \mathbf{b}_{i+1} - \Delta^0 \mathbf{b}_i \\ &= \mathbf{b}_{i+1} - \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^2 \mathbf{b}_i &= \Delta^1 \mathbf{b}_{i+1} - \Delta^1 \mathbf{b}_i \\ &= (\mathbf{b}_{i+2} - \mathbf{b}_{i+1}) - (\mathbf{b}_{i+1} - \mathbf{b}_i) \\ &= \mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^3 \mathbf{b}_i &= \Delta^2 \mathbf{b}_{i+1} - \Delta^2 \mathbf{b}_i \\ &= (\mathbf{b}_{i+3} - 2\mathbf{b}_{i+2} + \mathbf{b}_{i+1}) - (\mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i) \\ &= \mathbf{b}_{i+3} - 3\mathbf{b}_{i+2} + 3\mathbf{b}_{i+1} - \mathbf{b}_i. \end{aligned}$$



## Properties of Bézier curves

A differential operator of order  $r \geq 0$

$$\begin{cases} \Delta^0 \mathbf{b}_i = \mathbf{b}_i, \\ \Delta^r \mathbf{b}_i = \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i, \quad r \geq 1. \end{cases}$$

Example ( $r = 1, 2, 3$ )

$$\begin{aligned} \Delta^1 \mathbf{b}_i &= \Delta^0 \mathbf{b}_{i+1} - \Delta^0 \mathbf{b}_i \\ &= \mathbf{b}_{i+1} - \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^2 \mathbf{b}_i &= \Delta^1 \mathbf{b}_{i+1} - \Delta^1 \mathbf{b}_i \\ &= (\mathbf{b}_{i+2} - \mathbf{b}_{i+1}) - (\mathbf{b}_{i+1} - \mathbf{b}_i) \\ &= \mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^3 \mathbf{b}_i &= \Delta^2 \mathbf{b}_{i+1} - \Delta^2 \mathbf{b}_i \\ &= (\mathbf{b}_{i+3} - 2\mathbf{b}_{i+2} + \mathbf{b}_{i+1}) - (\mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i) \\ &= \mathbf{b}_{i+3} - 3\mathbf{b}_{i+2} + 3\mathbf{b}_{i+1} - \mathbf{b}_i. \end{aligned}$$



## Properties of Bézier curves

A differential operator of order  $r \geq 0$

$$\begin{cases} \Delta^0 \mathbf{b}_i = \mathbf{b}_i, \\ \Delta^r \mathbf{b}_i = \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i, \quad r \geq 1. \end{cases}$$

Example ( $r = 1, 2, 3$ )

$$\begin{aligned} \Delta^1 \mathbf{b}_i &= \Delta^0 \mathbf{b}_{i+1} - \Delta^0 \mathbf{b}_i \\ &= \mathbf{b}_{i+1} - \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^2 \mathbf{b}_i &= \Delta^1 \mathbf{b}_{i+1} - \Delta^1 \mathbf{b}_i \\ &= (\mathbf{b}_{i+2} - \mathbf{b}_{i+1}) - (\mathbf{b}_{i+1} - \mathbf{b}_i) \\ &= \mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^3 \mathbf{b}_i &= \Delta^2 \mathbf{b}_{i+1} - \Delta^2 \mathbf{b}_i \\ &= (\mathbf{b}_{i+3} - 2\mathbf{b}_{i+2} + \mathbf{b}_{i+1}) - (\mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i) \\ &= \mathbf{b}_{i+3} - 3\mathbf{b}_{i+2} + 3\mathbf{b}_{i+1} - \mathbf{b}_i. \end{aligned}$$



## Properties of Bézier curves

A differential operator of order  $r \geq 0$

$$\begin{cases} \Delta^0 \mathbf{b}_i = \mathbf{b}_i, \\ \Delta^r \mathbf{b}_i = \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i, \quad r \geq 1. \end{cases}$$

Example ( $r = 1, 2, 3$ )

$$\begin{aligned} \Delta^1 \mathbf{b}_i &= \Delta^0 \mathbf{b}_{i+1} - \Delta^0 \mathbf{b}_i \\ &= \mathbf{b}_{i+1} - \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^2 \mathbf{b}_i &= \Delta^1 \mathbf{b}_{i+1} - \Delta^1 \mathbf{b}_i \\ &= (\mathbf{b}_{i+2} - \mathbf{b}_{i+1}) - (\mathbf{b}_{i+1} - \mathbf{b}_i) \\ &= \mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^3 \mathbf{b}_i &= \Delta^2 \mathbf{b}_{i+1} - \Delta^2 \mathbf{b}_i \\ &= (\mathbf{b}_{i+3} - 2\mathbf{b}_{i+2} + \mathbf{b}_{i+1}) - (\mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i) \\ &= \mathbf{b}_{i+3} - 3\mathbf{b}_{i+2} + 3\mathbf{b}_{i+1} - \mathbf{b}_i. \end{aligned}$$



## Properties of Bézier curves

A differential operator of order  $r \geq 0$

$$\begin{cases} \Delta^0 \mathbf{b}_i = \mathbf{b}_i, \\ \Delta^r \mathbf{b}_i = \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i, \quad r \geq 1. \end{cases}$$

Example ( $r = 1, 2, 3$ )

$$\begin{aligned} \Delta^1 \mathbf{b}_i &= \Delta^0 \mathbf{b}_{i+1} - \Delta^0 \mathbf{b}_i \\ &= \mathbf{b}_{i+1} - \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^2 \mathbf{b}_i &= \Delta^1 \mathbf{b}_{i+1} - \Delta^1 \mathbf{b}_i \\ &= (\mathbf{b}_{i+2} - \mathbf{b}_{i+1}) - (\mathbf{b}_{i+1} - \mathbf{b}_i) \\ &= \mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^3 \mathbf{b}_i &= \Delta^2 \mathbf{b}_{i+1} - \Delta^2 \mathbf{b}_i \\ &= (\mathbf{b}_{i+3} - 2\mathbf{b}_{i+2} + \mathbf{b}_{i+1}) - (\mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i) \\ &= \mathbf{b}_{i+3} - 3\mathbf{b}_{i+2} + 3\mathbf{b}_{i+1} - \mathbf{b}_i. \end{aligned}$$



## Properties of Bézier curves

A differential operator of order  $r \geq 0$

$$\begin{cases} \Delta^0 \mathbf{b}_i = \mathbf{b}_i, \\ \Delta^r \mathbf{b}_i = \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i, \quad r \geq 1. \end{cases}$$

Example ( $r = 1, 2, 3$ )

$$\begin{aligned} \Delta^1 \mathbf{b}_i &= \Delta^0 \mathbf{b}_{i+1} - \Delta^0 \mathbf{b}_i \\ &= \mathbf{b}_{i+1} - \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^2 \mathbf{b}_i &= \Delta^1 \mathbf{b}_{i+1} - \Delta^1 \mathbf{b}_i \\ &= (\mathbf{b}_{i+2} - \mathbf{b}_{i+1}) - (\mathbf{b}_{i+1} - \mathbf{b}_i) \\ &= \mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^3 \mathbf{b}_i &= \Delta^2 \mathbf{b}_{i+1} - \Delta^2 \mathbf{b}_i \\ &= (\mathbf{b}_{i+3} - 2\mathbf{b}_{i+2} + \mathbf{b}_{i+1}) - (\mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i) \\ &= \mathbf{b}_{i+3} - 3\mathbf{b}_{i+2} + 3\mathbf{b}_{i+1} - \mathbf{b}_i. \end{aligned}$$





## Properties of Bézier curves

A differential operator of order  $r \geq 0$

$$\begin{cases} \Delta^0 \mathbf{b}_i = \mathbf{b}_i, \\ \Delta^r \mathbf{b}_i = \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i, \quad r \geq 1. \end{cases}$$

Example ( $r = 1, 2, 3$ )

$$\begin{aligned} \Delta^1 \mathbf{b}_i &= \Delta^0 \mathbf{b}_{i+1} - \Delta^0 \mathbf{b}_i \\ &= \mathbf{b}_{i+1} - \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^2 \mathbf{b}_i &= \Delta^1 \mathbf{b}_{i+1} - \Delta^1 \mathbf{b}_i \\ &= (\mathbf{b}_{i+2} - \mathbf{b}_{i+1}) - (\mathbf{b}_{i+1} - \mathbf{b}_i) \\ &= \mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^3 \mathbf{b}_i &= \Delta^2 \mathbf{b}_{i+1} - \Delta^2 \mathbf{b}_i \\ &= (\mathbf{b}_{i+3} - 2\mathbf{b}_{i+2} + \mathbf{b}_{i+1}) - (\mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i) \\ &= \mathbf{b}_{i+3} - 3\mathbf{b}_{i+2} + 3\mathbf{b}_{i+1} - \mathbf{b}_i. \end{aligned}$$



## Properties of Bézier curves

A differential operator of order  $r \geq 0$

$$\begin{cases} \Delta^0 \mathbf{b}_i = \mathbf{b}_i, \\ \Delta^r \mathbf{b}_i = \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i, \quad r \geq 1. \end{cases}$$

Example ( $r = 1, 2, 3$ )

$$\begin{aligned} \Delta^1 \mathbf{b}_i &= \Delta^0 \mathbf{b}_{i+1} - \Delta^0 \mathbf{b}_i \\ &= \mathbf{b}_{i+1} - \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^2 \mathbf{b}_i &= \Delta^1 \mathbf{b}_{i+1} - \Delta^1 \mathbf{b}_i \\ &= (\mathbf{b}_{i+2} - \mathbf{b}_{i+1}) - (\mathbf{b}_{i+1} - \mathbf{b}_i) \\ &= \mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^3 \mathbf{b}_i &= \Delta^2 \mathbf{b}_{i+1} - \Delta^2 \mathbf{b}_i \\ &= (\mathbf{b}_{i+3} - 2\mathbf{b}_{i+2} + \mathbf{b}_{i+1}) - (\mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i) \\ &= \mathbf{b}_{i+3} - 3\mathbf{b}_{i+2} + 3\mathbf{b}_{i+1} - \mathbf{b}_i. \end{aligned}$$



## Properties of Bézier curves

A differential operator of order  $r \geq 0$

$$\begin{cases} \Delta^0 \mathbf{b}_i = \mathbf{b}_i, \\ \Delta^r \mathbf{b}_i = \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i, \quad r \geq 1. \end{cases}$$

Example ( $r = 1, 2, 3$ )

$$\begin{aligned} \Delta^1 \mathbf{b}_i &= \Delta^0 \mathbf{b}_{i+1} - \Delta^0 \mathbf{b}_i \\ &= \mathbf{b}_{i+1} - \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^2 \mathbf{b}_i &= \Delta^1 \mathbf{b}_{i+1} - \Delta^1 \mathbf{b}_i \\ &= (\mathbf{b}_{i+2} - \mathbf{b}_{i+1}) - (\mathbf{b}_{i+1} - \mathbf{b}_i) \\ &= \mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^3 \mathbf{b}_i &= \Delta^2 \mathbf{b}_{i+1} - \Delta^2 \mathbf{b}_i \\ &= (\mathbf{b}_{i+3} - 2\mathbf{b}_{i+2} + \mathbf{b}_{i+1}) - (\mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i) \\ &= \mathbf{b}_{i+3} - 3\mathbf{b}_{i+2} + 3\mathbf{b}_{i+1} - \mathbf{b}_i. \end{aligned}$$



## Properties of Bézier curves

A differential operator of order  $r \geq 0$

$$\begin{cases} \Delta^0 \mathbf{b}_i = \mathbf{b}_i, \\ \Delta^r \mathbf{b}_i = \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i, \quad r \geq 1. \end{cases}$$

Example ( $r = 1, 2, 3$ )

$$\begin{aligned} \Delta^1 \mathbf{b}_i &= \Delta^0 \mathbf{b}_{i+1} - \Delta^0 \mathbf{b}_i \\ &= \mathbf{b}_{i+1} - \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^2 \mathbf{b}_i &= \Delta^1 \mathbf{b}_{i+1} - \Delta^1 \mathbf{b}_i \\ &= (\mathbf{b}_{i+2} - \mathbf{b}_{i+1}) - (\mathbf{b}_{i+1} - \mathbf{b}_i) \\ &= \mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i, \end{aligned}$$

$$\begin{aligned} \Delta^3 \mathbf{b}_i &= \Delta^2 \mathbf{b}_{i+1} - \Delta^2 \mathbf{b}_i \\ &= (\mathbf{b}_{i+3} - 2\mathbf{b}_{i+2} + \mathbf{b}_{i+1}) - (\mathbf{b}_{i+2} - 2\mathbf{b}_{i+1} + \mathbf{b}_i) \\ &= \mathbf{b}_{i+3} - 3\mathbf{b}_{i+2} + 3\mathbf{b}_{i+1} - \mathbf{b}_i. \end{aligned}$$



## Properties of Bézier curves

Theorem (Closed formula of the differential operator  $\Delta^r \mathbf{b}_i, r \geq 0$ )

$$\Delta^r \mathbf{b}_i = \sum_{k=0}^r \binom{r}{k} (-1)^{r-k} \mathbf{b}_{i+k}.$$

Proof.

- We prove the theorem by **mathematical induction with respect to  $r$** .
- The statement trivially holds for  $r = 0$  and  $r = 1$ .
- We assume that the statement is valid for all  $0 \leq j \leq r - 1$  and we also prove it for  $j = r \geq 1$ .
- We can successively write:

$$\begin{aligned} \Delta^r \mathbf{b}_i &= \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i \\ &= \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+1+k} - \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+k} \\ &\stackrel{k \rightarrow k-1}{=} \sum_{k=1}^r \binom{r-1}{k-1} (-1)^{r-k} \mathbf{b}_{i+k} + \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-k} \mathbf{b}_{i+k} \end{aligned}$$



## Properties of Bézier curves

Theorem (Closed formula of the differential operator  $\Delta^r \mathbf{b}_i, r \geq 0$ )

$$\Delta^r \mathbf{b}_i = \sum_{k=0}^r \binom{r}{k} (-1)^{r-k} \mathbf{b}_{i+k}.$$

Proof.

- We prove the theorem by **mathematical induction with respect to  $r$** .
- The statement trivially holds for  $r = 0$  and  $r = 1$ .
- We assume that the statement is valid for all  $0 \leq j \leq r - 1$  and we also prove it for  $j = r \geq 1$ .
- We can successively write:

$$\begin{aligned} \Delta^r \mathbf{b}_i &= \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i \\ &= \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+1+k} - \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+k} \\ &\stackrel{k \rightarrow k-1}{=} \sum_{k=1}^r \binom{r-1}{k-1} (-1)^{r-k} \mathbf{b}_{i+k} + \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-k} \mathbf{b}_{i+k} \end{aligned}$$



## Properties of Bézier curves

Theorem (Closed formula of the differential operator  $\Delta^r \mathbf{b}_i, r \geq 0$ )

$$\Delta^r \mathbf{b}_i = \sum_{k=0}^r \binom{r}{k} (-1)^{r-k} \mathbf{b}_{i+k}.$$

Proof.

- We prove the theorem by **mathematical induction with respect to  $r$** .
- The statement trivially holds for  $r = 0$  and  $r = 1$ .
- We assume that the statement is valid for all  $0 \leq j \leq r-1$  and we also prove it for  $j = r \geq 1$ .
- We can successively write:

$$\begin{aligned} \Delta^r \mathbf{b}_i &= \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i \\ &= \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+1+k} - \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+k} \\ &\stackrel{k \rightarrow k-1}{=} \sum_{k=1}^r \binom{r-1}{k-1} (-1)^{r-k} \mathbf{b}_{i+k} + \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-k} \mathbf{b}_{i+k} \end{aligned}$$



## Properties of Bézier curves

Theorem (Closed formula of the differential operator  $\Delta^r \mathbf{b}_i, r \geq 0$ )

$$\Delta^r \mathbf{b}_i = \sum_{k=0}^r \binom{r}{k} (-1)^{r-k} \mathbf{b}_{i+k}.$$

Proof.

- We prove the theorem by **mathematical induction with respect to  $r$** .
- The statement trivially holds for  $r = 0$  and  $r = 1$ .
- We assume that the statement is valid for all  $0 \leq j \leq r - 1$  and we also prove it for  $j = r \geq 1$ .
- We can successively write:

$$\begin{aligned} \Delta^r \mathbf{b}_i &= \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i \\ &= \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+1+k} - \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+k} \\ &\stackrel{k \rightarrow k-1}{=} \sum_{k=1}^r \binom{r-1}{k-1} (-1)^{r-k} \mathbf{b}_{i+k} + \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-k} \mathbf{b}_{i+k} \end{aligned}$$





## Properties of Bézier curves

Theorem (Closed formula of the differential operator  $\Delta^r \mathbf{b}_i, r \geq 0$ )

$$\Delta^r \mathbf{b}_i = \sum_{k=0}^r \binom{r}{k} (-1)^{r-k} \mathbf{b}_{i+k}.$$

Proof.

- We prove the theorem by **mathematical induction with respect to  $r$** .
- The statement trivially holds for  $r = 0$  and  $r = 1$ .
- We assume that the statement is valid for all  $0 \leq j \leq r - 1$  and we also prove it for  $j = r \geq 1$ .
- We can successively write:

$$\begin{aligned} \Delta^r \mathbf{b}_i &= \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i \\ &= \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+1+k} - \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+k} \\ &\stackrel{k \rightarrow k-1}{=} \sum_{k=1}^r \binom{r-1}{k-1} (-1)^{r-k} \mathbf{b}_{i+k} + \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-k} \mathbf{b}_{i+k} \end{aligned}$$

## Properties of Bézier curves

Theorem (Closed formula of the differential operator  $\Delta^r \mathbf{b}_i, r \geq 0$ )

$$\Delta^r \mathbf{b}_i = \sum_{k=0}^r \binom{r}{k} (-1)^{r-k} \mathbf{b}_{i+k}.$$

Proof.

- We prove the theorem by **mathematical induction with respect to  $r$** .
- The statement trivially holds for  $r = 0$  and  $r = 1$ .
- We assume that the statement is valid for all  $0 \leq j \leq r - 1$  and we also prove it for  $j = r \geq 1$ .
- We can successively write:

$$\begin{aligned} \Delta^r \mathbf{b}_i &= \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i \\ &= \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+1+k} - \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+k} \\ &\stackrel{k \rightarrow k-1}{=} \sum_{k=1}^r \binom{r-1}{k-1} (-1)^{r-k} \mathbf{b}_{i+k} + \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-k} \mathbf{b}_{i+k} \end{aligned}$$

## Properties of Bézier curves

Theorem (Closed formula of the differential operator  $\Delta^r \mathbf{b}_i, r \geq 0$ )

$$\Delta^r \mathbf{b}_i = \sum_{k=0}^r \binom{r}{k} (-1)^{r-k} \mathbf{b}_{i+k}.$$

Proof.

- We prove the theorem by **mathematical induction with respect to  $r$** .
- The statement trivially holds for  $r = 0$  and  $r = 1$ .
- We assume that the statement is valid for all  $0 \leq j \leq r - 1$  and we also prove it for  $j = r \geq 1$ .
- We can successively write:

$$\begin{aligned} \Delta^r \mathbf{b}_i &= \Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i \\ &= \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+1+k} - \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-1-k} \mathbf{b}_{i+k} \\ &\stackrel{k \rightarrow k-1}{=} \sum_{k=1}^r \binom{r-1}{k-1} (-1)^{r-k} \mathbf{b}_{i+k} + \sum_{k=0}^{r-1} \binom{r-1}{k} (-1)^{r-k} \mathbf{b}_{i+k}. \end{aligned}$$

## Properties of Bézier curves

Proof – continued.

- Since  $\binom{r-1}{-1} = \binom{r-1}{r} = 0$  and  $\binom{r}{k} = \binom{r-1}{k} + \binom{r-1}{k-1}$ , it follows that

$$\begin{aligned}\Delta^r \mathbf{b}_i &= \sum_{k=0}^r \binom{r-1}{k-1} (-1)^{r-k} \mathbf{b}_{i+k} + \sum_{k=0}^r \binom{r-1}{k} (-1)^{r-k} \mathbf{b}_{i+k} \\ &= \sum_{k=0}^r \binom{r}{k} (-1)^{r-k} \mathbf{b}_{i+k}. \quad \square\end{aligned}$$



## Properties of Bézier curves

Proof – continued.

- Since  $\binom{r-1}{-1} = \binom{r-1}{r} = 0$  and  $\binom{r}{k} = \binom{r-1}{k} + \binom{r-1}{k-1}$ , it follows that

$$\begin{aligned}\Delta^r \mathbf{b}_i &= \sum_{k=0}^r \binom{r-1}{k-1} (-1)^{r-k} \mathbf{b}_{i+k} + \sum_{k=0}^r \binom{r-1}{k} (-1)^{r-k} \mathbf{b}_{i+k} \\ &= \sum_{k=0}^r \binom{r}{k} (-1)^{r-k} \mathbf{b}_{i+k}. \quad \square\end{aligned}$$



## Properties of Bézier curves

Proof – continued.

- Since  $\binom{r-1}{-1} = \binom{r-1}{r} = 0$  and  $\binom{r}{k} = \binom{r-1}{k} + \binom{r-1}{k-1}$ , it follows that

$$\begin{aligned}\Delta^r \mathbf{b}_i &= \sum_{k=0}^r \binom{r-1}{k-1} (-1)^{r-k} \mathbf{b}_{i+k} + \sum_{k=0}^r \binom{r-1}{k} (-1)^{r-k} \mathbf{b}_{i+k} \\ &= \sum_{k=0}^r \binom{r}{k} (-1)^{r-k} \mathbf{b}_{i+k}. \quad \square\end{aligned}$$



## Properties of Bézier curves

### Theorem (Higher order derivatives of Bézier curves)

The  $r$ th ( $r \geq 0$ ) order derivative of a Bézier curve of degree  $n \geq 1$  is

$$\frac{d^r}{du^r} \mathbf{b}(u) = \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \Delta^r \mathbf{b}_i B_i^{n-r}(u), \forall u \in [0, 1].$$

Proof.

- The proof is based on **mathematical induction** with respect to  $r \geq 0$ .
- The statement trivially holds for  $r = 0$  and in case of  $r = 1$  we have seen that

$$\frac{d}{du} \mathbf{b}(u) = n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u)$$

$$= \frac{n!}{(n-1)!} \sum_{i=0}^{n-1} \Delta \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1].$$

- We assume that the statement is valid for all  $0 \leq j \leq r-1$  and we also prove it for  $j = r \geq 2$ .



## Properties of Bézier curves

### Theorem (Higher order derivatives of Bézier curves)

The  $r$ th ( $r \geq 0$ ) order derivative of a Bézier curve of degree  $n \geq 1$  is

$$\frac{d^r}{du^r} \mathbf{b}(u) = \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \Delta^r \mathbf{b}_i B_i^{n-r}(u), \forall u \in [0, 1].$$

Proof.

- The proof is based on **mathematical induction** with respect to  $r \geq 0$ .
- The statement trivially holds for  $r = 0$  and in case of  $r = 1$  we have seen that

$$\begin{aligned} \frac{d}{du} \mathbf{b}(u) &= n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u) \\ &= \frac{n!}{(n-1)!} \sum_{i=0}^{n-1} \Delta^1 \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1]. \end{aligned}$$

- We assume that the statement is valid for all  $0 \leq j \leq r-1$  and we also prove it for  $j = r \geq 2$ .





## Properties of Bézier curves

### Theorem (Higher order derivatives of Bézier curves)

The  $r$ th ( $r \geq 0$ ) order derivative of a Bézier curve of degree  $n \geq 1$  is

$$\frac{d^r}{du^r} \mathbf{b}(u) = \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \Delta^r \mathbf{b}_i B_i^{n-r}(u), \forall u \in [0, 1].$$

Proof.

- The proof is based on **mathematical induction** with respect to  $r \geq 0$ .
- The statement trivially holds for  $r = 0$  and in case of  $r = 1$  we have seen that

$$\begin{aligned} \frac{d}{du} \mathbf{b}(u) &= n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u) \\ &= \frac{n!}{(n-1)!} \sum_{i=0}^{n-1} \Delta^1 \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1]. \end{aligned}$$

- We assume that the statement is valid for all  $0 \leq j \leq r-1$  and we also prove it for  $j = r \geq 2$ .



## Properties of Bézier curves

### Theorem (Higher order derivatives of Bézier curves)

The  $r$ th ( $r \geq 0$ ) order derivative of a Bézier curve of degree  $n \geq 1$  is

$$\frac{d^r}{du^r} \mathbf{b}(u) = \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \Delta^r \mathbf{b}_i B_i^{n-r}(u), \forall u \in [0, 1].$$

Proof.

- The proof is based on **mathematical induction** with respect to  $r \geq 0$ .
- The statement trivially holds for  $r = 0$  and in case of  $r = 1$  we have seen that

$$\begin{aligned} \frac{d}{du} \mathbf{b}(u) &= n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u) \\ &= \frac{n!}{(n-1)!} \sum_{i=0}^{n-1} \Delta^1 \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1]. \end{aligned}$$

- We assume that the statement is valid for all  $0 \leq j \leq r-1$  and we also prove it for  $j = r \geq 2$ .



## Properties of Bézier curves

### Theorem (Higher order derivatives of Bézier curves)

The  $r$ th ( $r \geq 0$ ) order derivative of a Bézier curve of degree  $n \geq 1$  is

$$\frac{d^r}{du^r} \mathbf{b}(u) = \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \Delta^r \mathbf{b}_i B_i^{n-r}(u), \forall u \in [0, 1].$$

Proof.

- The proof is based on **mathematical induction** with respect to  $r \geq 0$ .
- The statement trivially holds for  $r = 0$  and in case of  $r = 1$  we have seen that

$$\begin{aligned} \frac{d}{du} \mathbf{b}(u) &= n \sum_{i=0}^{n-1} (\mathbf{b}_{i+1} - \mathbf{b}_i) B_i^{n-1}(u) \\ &= \frac{n!}{(n-1)!} \sum_{i=0}^{n-1} \Delta^1 \mathbf{b}_i B_i^{n-1}(u), \forall u \in [0, 1]. \end{aligned}$$

- We assume that the statement is valid for all  $0 \leq j \leq r-1$  and we also prove it for  $j = r \geq 2$ .



## Properties of Bézier curves

### Proof – continued.

- Based on the induction hypothesis:

$$\frac{d^{r-1}}{du^{r-1}} \mathbf{b}(u) = \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_i^{n-r+1}(u), \forall u \in [0, 1].$$

- We can successively write:

$$\begin{aligned} \frac{d^r}{du^r} \mathbf{b}(u) &= \frac{d}{du} \left( \frac{d^{r-1}}{du^{r-1}} \mathbf{b}(u) \right) \\ &= \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i \frac{d}{du} B_i^{n-r+1}(u) \\ &= \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i (n-r+1) (B_{i-1}^{n-r}(u) - B_i^{n-r}(u)) \\ &= \frac{n!}{(n-r)!} \left( \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_{i-1}^{n-r}(u) - \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \\ &= \frac{n!}{(n-r)!} \left( \sum_{i=1}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_{i-1}^{n-r}(u) - \sum_{i=0}^{n-r} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \end{aligned}$$



## Properties of Bézier curves

### Proof – continued.

- Based on the induction hypothesis:

$$\frac{d^{r-1}}{du^{r-1}} \mathbf{b}(u) = \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_i^{n-r+1}(u), \forall u \in [0, 1].$$

- We can successively write:

$$\begin{aligned} \frac{d^r}{du^r} \mathbf{b}(u) &= \frac{d}{du} \left( \frac{d^{r-1}}{du^{r-1}} \mathbf{b}(u) \right) \\ &= \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i \frac{d}{du} B_i^{n-r+1}(u) \\ &= \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i (n-r+1) (B_{i-1}^{n-r}(u) - B_i^{n-r}(u)) \\ &= \frac{n!}{(n-r)!} \left( \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_{i-1}^{n-r}(u) - \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \\ &= \frac{n!}{(n-r)!} \left( \sum_{i=1}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_{i-1}^{n-r}(u) - \sum_{i=0}^{n-r} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \end{aligned}$$



## Properties of Bézier curves

### Proof – continued.

- Based on the induction hypothesis:

$$\frac{d^{r-1}}{du^{r-1}} \mathbf{b}(u) = \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_i^{n-r+1}(u), \forall u \in [0, 1].$$

- We can successively write:

$$\begin{aligned} \frac{d^r}{du^r} \mathbf{b}(u) &= \frac{d}{du} \left( \frac{d^{r-1}}{du^{r-1}} \mathbf{b}(u) \right) \\ &= \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i \frac{d}{du} B_i^{n-r+1}(u) \\ &= \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i (n-r+1) (B_{i-1}^{n-r}(u) - B_i^{n-r}(u)) \\ &= \frac{n!}{(n-r)!} \left( \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_{i-1}^{n-r}(u) - \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \\ &= \frac{n!}{(n-r)!} \left( \sum_{i=1}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_{i-1}^{n-r}(u) - \sum_{i=0}^{n-r} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \end{aligned}$$



## Properties of Bézier curves

### Proof – continued.

- Based on the induction hypothesis:

$$\frac{d^{r-1}}{du^{r-1}} \mathbf{b}(u) = \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_i^{n-r+1}(u), \forall u \in [0, 1].$$

- We can successively write:

$$\begin{aligned} \frac{d^r}{du^r} \mathbf{b}(u) &= \frac{d}{du} \left( \frac{d^{r-1}}{du^{r-1}} \mathbf{b}(u) \right) \\ &= \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i \frac{d}{du} B_i^{n-r+1}(u) \\ &= \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i (n-r+1) (B_{i-1}^{n-r}(u) - B_i^{n-r}(u)) \\ &= \frac{n!}{(n-r)!} \left( \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_{i-1}^{n-r}(u) - \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \\ &= \frac{n!}{(n-r)!} \left( \sum_{i=1}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_{i-1}^{n-r}(u) - \sum_{i=0}^{n-r} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \end{aligned}$$



## Properties of Bézier curves

### Proof – continued.

- Based on the induction hypothesis:

$$\frac{d^{r-1}}{du^{r-1}} \mathbf{b}(u) = \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_i^{n-r+1}(u), \forall u \in [0, 1].$$

- We can successively write:

$$\begin{aligned} \frac{d^r}{du^r} \mathbf{b}(u) &= \frac{d}{du} \left( \frac{d^{r-1}}{du^{r-1}} \mathbf{b}(u) \right) \\ &= \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i \frac{d}{du} B_i^{n-r+1}(u) \\ &= \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i (n-r+1) (B_{i-1}^{n-r}(u) - B_i^{n-r}(u)) \\ &= \frac{n!}{(n-r)!} \left( \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_{i-1}^{n-r}(u) - \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \\ &= \frac{n!}{(n-r)!} \left( \sum_{i=1}^{n-r+1} \Delta^{r-1} \mathbf{b}_{i-1} B_{i-1}^{n-r}(u) - \sum_{i=0}^{n-r} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \end{aligned}$$





## Properties of Bézier curves

### Proof – continued.

- Based on the induction hypothesis:

$$\frac{d^{r-1}}{du^{r-1}} \mathbf{b}(u) = \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_i^{n-r+1}(u), \forall u \in [0, 1].$$

- We can successively write:

$$\begin{aligned} \frac{d^r}{du^r} \mathbf{b}(u) &= \frac{d}{du} \left( \frac{d^{r-1}}{du^{r-1}} \mathbf{b}(u) \right) \\ &= \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i \frac{d}{du} B_i^{n-r+1}(u) \\ &= \frac{n!}{(n-r+1)!} \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i (n-r+1) \left( B_{i-1}^{n-r}(u) - B_i^{n-r}(u) \right) \\ &= \frac{n!}{(n-r)!} \left( \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_{i-1}^{n-r}(u) - \sum_{i=0}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \\ &= \frac{n!}{(n-r)!} \left( \sum_{i=1}^{n-r+1} \Delta^{r-1} \mathbf{b}_i B_{i-1}^{n-r}(u) - \sum_{i=0}^{n-r} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \end{aligned}$$

## Properties of Bézier curves

Proof – continued.

$$\begin{aligned} & \stackrel{i-1 \rightarrow i}{=} \frac{n!}{(n-r)!} \left( \sum_{i=0}^{n-r} \Delta^{r-1} \mathbf{b}_{i+1} B_i^{n-r}(u) - \sum_{i=0}^{n-r} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \\ &= \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \underbrace{(\Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i)}_{\Delta^r \mathbf{b}_i} B_i^{n-r}(u) \\ &= \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \Delta^r \mathbf{b}_i B_i^{n-r}(u), \quad \forall u \in [0, 1]. \quad \square \end{aligned}$$



## Properties of Bézier curves

Proof – continued.

$$\begin{aligned} & \stackrel{i-1 \rightarrow i}{=} \frac{n!}{(n-r)!} \left( \sum_{i=0}^{n-r} \Delta^{r-1} \mathbf{b}_{i+1} B_i^{n-r}(u) - \sum_{i=0}^{n-r} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \\ &= \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \underbrace{(\Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i)}_{\Delta^r \mathbf{b}_i} B_i^{n-r}(u) \\ &= \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \Delta^r \mathbf{b}_i B_i^{n-r}(u), \quad \forall u \in [0, 1]. \quad \square \end{aligned}$$



## Properties of Bézier curves

Proof – continued.

$$\begin{aligned} & \stackrel{i-1 \rightarrow i}{=} \frac{n!}{(n-r)!} \left( \sum_{i=0}^{n-r} \Delta^{r-1} \mathbf{b}_{i+1} B_i^{n-r}(u) - \sum_{i=0}^{n-r} \Delta^{r-1} \mathbf{b}_i B_i^{n-r}(u) \right) \\ &= \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \underbrace{(\Delta^{r-1} \mathbf{b}_{i+1} - \Delta^{r-1} \mathbf{b}_i)}_{\Delta^r \mathbf{b}_i} B_i^{n-r}(u) \\ &= \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \Delta^r \mathbf{b}_i B_i^{n-r}(u), \forall u \in [0, 1]. \quad \square \end{aligned}$$



## Properties of Bézier curves




Fig. 10: The direction vectors of the first and second order derivatives at the beginning of a Bézier curve of degree 4.

### Derivatives at the endpoints of a Bézier curve of degree $n$

Since

$$\begin{aligned}\frac{d^r}{du^r} \mathbf{b}(0) &= \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \Delta^r \mathbf{b}_i B_i^{n-r}(0) \\ &= \frac{n!}{(n-r)!} \Delta^r \mathbf{b}_0\end{aligned}$$

and

$$\begin{aligned}\frac{d^r}{du^r} \mathbf{b}(1) &= \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \Delta^r \mathbf{b}_i B_i^{n-r}(1) \\ &= \frac{n!}{(n-r)!} \Delta^r \mathbf{b}_{n-r},\end{aligned}$$

the  $r$ th order derivatives at the endpoints of a Bézier curve of degree  $n$  depend only on the first

$$\mathbf{b}_0, \mathbf{b}_1, \dots, \mathbf{b}_r$$


and the last

$$\mathbf{b}_{n-r}, \mathbf{b}_{n-r+1}, \dots, \mathbf{b}_n$$

$r + 1$  control points.



## Properties of Bézier curves



**Fig. 10:** The direction vectors of the first and second order derivatives at the beginning of a Bézier curve of degree 4.

### Derivatives at the endpoints of a Bézier curve of degree $n$

Since

$$\begin{aligned}\frac{d^r}{du^r} \mathbf{b}(0) &= \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \Delta^r \mathbf{b}_i B_i^{n-r}(0) \\ &= \frac{n!}{(n-r)!} \Delta^r \mathbf{b}_0\end{aligned}$$

and

$$\begin{aligned}\frac{d^r}{du^r} \mathbf{b}(1) &= \frac{n!}{(n-r)!} \sum_{i=0}^{n-r} \Delta^r \mathbf{b}_i B_i^{n-r}(1) \\ &= \frac{n!}{(n-r)!} \Delta^r \mathbf{b}_{n-r},\end{aligned}$$

the  $r$ th order derivatives at the endpoints of a Bézier curve of degree  $n$  depend only on the first

$$\mathbf{b}_0, \mathbf{b}_1, \dots, \mathbf{b}_r$$

and the last

$$\mathbf{b}_{n-r}, \mathbf{b}_{n-r+1}, \dots, \mathbf{b}_n$$

$r + 1$  control points.

## Properties of Bézier curves

### Theorem (The relationship between derivatives and the de Casteljau-algorithm)

The  $r$ th ( $r \geq 0$ ) derivative of a Bézier curve of degree  $n \geq 1$  can also be represented as

$$\frac{d^r}{du^r} \mathbf{b}(u) = \frac{d^r}{du^r} \mathbf{b}_0^n(u) = \frac{n!}{(n-r)!} \Delta^r \mathbf{b}_0^{n-r}(u), \forall u \in [0, 1],$$

where  $\mathbf{b}_i^k$  are intermediate points of the de Casteljau-algorithm.

### Proof.

- We will use the fact that the summation and difference operators are commutable, e.g.

$$\sum_{j=0}^{n-1} \Delta^1 \mathbf{b}_j = \sum_{j=0}^{n-1} (\mathbf{b}_{j+1} - \mathbf{b}_j) = \sum_{j=1}^n \mathbf{b}_j - \sum_{j=0}^{n-1} \mathbf{b}_j = \Delta^1 \sum_{j=0}^{n-1} \mathbf{b}_j.$$

- Thus, we can write successively

$$\begin{aligned} \frac{d^r}{du^r} \mathbf{b}(u) &= \frac{d^r}{du^r} \mathbf{b}_0^n(u) = \frac{n!}{(n-r)!} \sum_{j=0}^{n-r} \Delta^r \mathbf{b}_j \mathcal{B}_j^{n-r}(u) = \frac{n!}{(n-r)!} \underbrace{\Delta^r \sum_{j=0}^{n-r} \mathbf{b}_j \mathcal{B}_j^{n-r}(u)}_{\mathbf{b}_0^{n-r}(u)} \\ &= \frac{n!}{(n-r)!} \Delta^r \mathbf{b}_0^{n-r}(u), \forall u \in [0, 1]. \quad \square \end{aligned}$$

## Properties of Bézier curves

### Theorem (The relationship between derivatives and the de Casteljau-algorithm)

The  $r$ th ( $r \geq 0$ ) derivative of a Bézier curve of degree  $n \geq 1$  can also be represented as

$$\frac{d^r}{du^r} \mathbf{b}(u) = \frac{d^r}{du^r} \mathbf{b}_0^n(u) = \frac{n!}{(n-r)!} \Delta^r \mathbf{b}_0^{n-r}(u), \forall u \in [0, 1],$$

where  $\mathbf{b}_i^k$  are intermediate points of the de Casteljau-algorithm.

### Proof.

- We will use the fact that the summation and difference operators are commutable, e.g.

$$\sum_{j=0}^{n-1} \Delta^1 \mathbf{b}_j = \sum_{j=0}^{n-1} (\mathbf{b}_{j+1} - \mathbf{b}_j) = \sum_{j=1}^n \mathbf{b}_j - \sum_{j=0}^{n-1} \mathbf{b}_j = \Delta^1 \sum_{j=0}^{n-1} \mathbf{b}_j.$$

- Thus, we can write successively

$$\begin{aligned} \frac{d^r}{du^r} \mathbf{b}(u) &= \frac{d^r}{du^r} \mathbf{b}_0^n(u) = \frac{n!}{(n-r)!} \sum_{j=0}^{n-r} \Delta^r \mathbf{b}_j B_j^{n-r}(u) = \frac{n!}{(n-r)!} \underbrace{\Delta^r \sum_{j=0}^{n-r} \mathbf{b}_j B_j^{n-r}(u)}_{\mathbf{b}_0^{n-r}(u)} \\ &= \frac{n!}{(n-r)!} \Delta^r \mathbf{b}_0^{n-r}(u), \forall u \in [0, 1]. \quad \square \end{aligned}$$



## Properties of Bézier curves

### Theorem (The relationship between derivatives and the de Casteljau-algorithm)

The  $r$ th ( $r \geq 0$ ) derivative of a Bézier curve of degree  $n \geq 1$  can also be represented as

$$\frac{d^r}{du^r} \mathbf{b}(u) = \frac{d^r}{du^r} \mathbf{b}_0^n(u) = \frac{n!}{(n-r)!} \Delta^r \mathbf{b}_0^{n-r}(u), \forall u \in [0, 1],$$

where  $\mathbf{b}_i^k$  are intermediate points of the de Casteljau-algorithm.

### Proof.

- We will use the fact that the summation and difference operators are commutable, e.g.

$$\sum_{j=0}^{n-1} \Delta^1 \mathbf{b}_j = \sum_{j=0}^{n-1} (\mathbf{b}_{j+1} - \mathbf{b}_j) = \sum_{j=1}^n \mathbf{b}_j - \sum_{j=0}^{n-1} \mathbf{b}_j = \Delta^1 \sum_{j=0}^{n-1} \mathbf{b}_j.$$

- Thus, we can write successively

$$\begin{aligned} \frac{d^r}{du^r} \mathbf{b}(u) &= \frac{d^r}{du^r} \mathbf{b}_0^n(u) = \frac{n!}{(n-r)!} \sum_{j=0}^{n-r} \Delta^r \mathbf{b}_j B_j^{n-r}(u) = \frac{n!}{(n-r)!} \underbrace{\Delta^r \sum_{j=0}^{n-r} \mathbf{b}_j B_j^{n-r}(u)}_{\mathbf{b}_0^{n-r}(u)} \\ &= \frac{n!}{(n-r)!} \Delta^r \mathbf{b}_0^{n-r}(u), \forall u \in [0, 1]. \quad \square \end{aligned}$$

## Properties of Bézier curves

### Bézier curves joined with $C^0$ continuity

Consider two Bézier curves

$$\mathbf{a}(u) = \sum_{i=0}^n \mathbf{a}_i B_i^n \left( \frac{u - u_0}{u_1 - u_0} \right), \quad u \in [u_0, u_1]$$

and

$$\mathbf{b}(u) = \sum_{i=0}^n \mathbf{b}_i B_i^n \left( \frac{u - u_1}{u_2 - u_1} \right), \quad u \in [u_1, u_2]$$

of degree  $n \geq 1$ . The condition of  $C^0$  continuity is

$$\mathbf{a}(u_1) = \mathbf{b}(u_1),$$

from which results that

$$\mathbf{a}_n = \mathbf{b}_0.$$



## Properties of Bézier curves

### Bézier curves joined with $C^0$ continuity

Consider two Bézier curves

$$\mathbf{a}(u) = \sum_{i=0}^n \mathbf{a}_i B_i^n \left( \frac{u - u_0}{u_1 - u_0} \right), \quad u \in [u_0, u_1]$$

and

$$\mathbf{b}(u) = \sum_{i=0}^n \mathbf{b}_i B_i^n \left( \frac{u - u_1}{u_2 - u_1} \right), \quad u \in [u_1, u_2]$$

of degree  $n \geq 1$ . The **condition of  $C^0$  continuity** is

$$\mathbf{a}(u_1) = \mathbf{b}(u_1),$$

from which results that

$$\mathbf{a}_n = \mathbf{b}_0.$$



## Properties of Bézier curves

### Bézier curves joined with $C^0$ continuity

Consider two Bézier curves

$$\mathbf{a}(u) = \sum_{i=0}^n \mathbf{a}_i B_i^n \left( \frac{u - u_0}{u_1 - u_0} \right), \quad u \in [u_0, u_1]$$

and

$$\mathbf{b}(u) = \sum_{i=0}^n \mathbf{b}_i B_i^n \left( \frac{u - u_1}{u_2 - u_1} \right), \quad u \in [u_1, u_2]$$

of degree  $n \geq 1$ . The **condition of  $C^0$  continuity** is

$$\mathbf{a}(u_1) = \mathbf{b}(u_1),$$

from which results that

$$\mathbf{a}_n = \mathbf{b}_0.$$



## Properties of Bézier curves

### Bézier curves joined with $C^1$ continuity

Using the same notations as in case of  $C^0$  continuity, Bézier curves

$$\mathbf{a}(u), u \in [u_0, u_1], \Delta u_0 = u_1 - u_0,$$

and

$$\mathbf{b}(u), u \in [u_1, u_2], \Delta u_1 = u_2 - u_1$$

of degree  $n \geq 1$  are joined with  $C^1$  continuity, if they are  $C^0$  continuous (i.e.  $\mathbf{a}_n = \mathbf{b}_0$ ) and

$$\frac{d}{du}\mathbf{a}(u_1) = \frac{d}{du}\mathbf{b}(u_1),$$

from which follows that

$$\frac{n!}{(n-1)!} \frac{1}{u_1 - u_0} \sum_{i=0}^{n-1} \Delta^1 \mathbf{a}_i B_i^{n-1}(1) = \frac{n!}{(n-1)!} \frac{1}{u_2 - u_1} \sum_{i=0}^{n-1} \Delta^1 \mathbf{b}_i B_i^{n-1}(0),$$

$$n \frac{1}{\Delta u_0} (\mathbf{a}_n - \mathbf{a}_{n-1}) = n \frac{1}{\Delta u_1} (\mathbf{b}_1 - \mathbf{b}_0),$$

$$\frac{\|\mathbf{a}_n - \mathbf{a}_{n-1}\|}{\Delta u_0} = \frac{\|\mathbf{b}_1 - \mathbf{b}_0\|}{\Delta u_1},$$

$$\|\Delta^1 \mathbf{a}_{n-1}\| : \|\Delta^1 \mathbf{b}_0\| = \Delta u_0 : \Delta u_1.$$



## Properties of Bézier curves

### Bézier curves joined with $C^1$ continuity

Using the same notations as in case of  $C^0$  continuity, Bézier curves

$$\mathbf{a}(u), u \in [u_0, u_1], \Delta u_0 = u_1 - u_0,$$

and

$$\mathbf{b}(u), u \in [u_1, u_2], \Delta u_1 = u_2 - u_1$$

of degree  $n \geq 1$  are joined with  $C^1$  continuity, if they are  $C^0$  continuous (i.e.  $\mathbf{a}_n = \mathbf{b}_0$ ) and

$$\frac{d}{du}\mathbf{a}(u_1) = \frac{d}{du}\mathbf{b}(u_1),$$

from which follows that

$$\begin{aligned} \frac{n!}{(n-1)!} \frac{1}{u_1 - u_0} \sum_{i=0}^{n-1} \Delta^1 \mathbf{a}_i B_i^{n-1}(1) &= \frac{n!}{(n-1)!} \frac{1}{u_2 - u_1} \sum_{i=0}^{n-1} \Delta^1 \mathbf{b}_i B_i^{n-1}(0), \\ n \frac{1}{\Delta u_0} (\mathbf{a}_n - \mathbf{a}_{n-1}) &= n \frac{1}{\Delta u_1} (\mathbf{b}_1 - \mathbf{b}_0), \\ \frac{\|\mathbf{a}_n - \mathbf{a}_{n-1}\|}{\Delta u_0} &= \frac{\|\mathbf{b}_1 - \mathbf{b}_0\|}{\Delta u_1}, \\ \|\Delta^1 \mathbf{a}_{n-1}\| : \|\Delta^1 \mathbf{b}_0\| &= \Delta u_0 : \Delta u_1. \end{aligned}$$



## Properties of Bézier curves

### Bézier curves joined with $C^1$ continuity

Using the same notations as in case of  $C^0$  continuity, Bézier curves

$$\mathbf{a}(u), u \in [u_0, u_1], \Delta u_0 = u_1 - u_0,$$

and

$$\mathbf{b}(u), u \in [u_1, u_2], \Delta u_1 = u_2 - u_1$$

of degree  $n \geq 1$  are joined with  $C^1$  continuity, if they are  $C^0$  continuous (i.e.  $\mathbf{a}_n = \mathbf{b}_0$ ) and

$$\frac{d}{du} \mathbf{a}(u_1) = \frac{d}{du} \mathbf{b}(u_1),$$

from which follows that

$$\begin{aligned} \frac{n!}{(n-1)!} \frac{1}{u_1 - u_0} \sum_{i=0}^{n-1} \Delta^1 \mathbf{a}_i B_i^{n-1}(1) &= \frac{n!}{(n-1)!} \frac{1}{u_2 - u_1} \sum_{i=0}^{n-1} \Delta^1 \mathbf{b}_i B_i^{n-1}(0), \\ n \frac{1}{\Delta u_0} (\mathbf{a}_n - \mathbf{a}_{n-1}) &= n \frac{1}{\Delta u_1} (\mathbf{b}_1 - \mathbf{b}_0), \\ \frac{\|\mathbf{a}_n - \mathbf{a}_{n-1}\|}{\Delta u_0} &= \frac{\|\mathbf{b}_1 - \mathbf{b}_0\|}{\Delta u_1}, \\ \|\Delta^1 \mathbf{a}_{n-1}\| : \|\Delta^1 \mathbf{b}_0\| &= \Delta u_0 : \Delta u_1. \end{aligned}$$



## Properties of Bézier curves

### Bézier curves joined with $C^1$ continuity

Using the same notations as in case of  $C^0$  continuity, Bézier curves

$$\mathbf{a}(u), u \in [u_0, u_1], \Delta u_0 = u_1 - u_0,$$

and

$$\mathbf{b}(u), u \in [u_1, u_2], \Delta u_1 = u_2 - u_1$$

of degree  $n \geq 1$  are joined with  $C^1$  continuity, if they are  $C^0$  continuous (i.e.  $\mathbf{a}_n = \mathbf{b}_0$ ) and

$$\frac{d}{du} \mathbf{a}(u_1) = \frac{d}{du} \mathbf{b}(u_1),$$

from which follows that

$$\begin{aligned} \frac{n!}{(n-1)!} \frac{1}{u_1 - u_0} \sum_{i=0}^{n-1} \Delta^1 \mathbf{a}_i B_i^{n-1}(1) &= \frac{n!}{(n-1)!} \frac{1}{u_2 - u_1} \sum_{i=0}^{n-1} \Delta^1 \mathbf{b}_i B_i^{n-1}(0), \\ n \frac{1}{\Delta u_0} (a_n - a_{n-1}) &= n \frac{1}{\Delta u_1} (b_1 - b_0), \\ \frac{\|a_n - a_{n-1}\|}{\Delta u_0} &= \frac{\|b_1 - b_0\|}{\Delta u_1}, \\ \|\Delta^1 \mathbf{a}_{n-1}\| : \|\Delta^1 \mathbf{b}_0\| &= \Delta u_0 : \Delta u_1. \end{aligned}$$





## Properties of Bézier curves

### Bézier curves joined with $C^1$ continuity

Using the same notations as in case of  $C^0$  continuity, Bézier curves

$$\mathbf{a}(u), u \in [u_0, u_1], \Delta u_0 = u_1 - u_0,$$

and

$$\mathbf{b}(u), u \in [u_1, u_2], \Delta u_1 = u_2 - u_1$$

of degree  $n \geq 1$  are joined with  $C^1$  continuity, if they are  $C^0$  continuous (i.e.  $\mathbf{a}_n = \mathbf{b}_0$ ) and

$$\frac{d}{du} \mathbf{a}(u_1) = \frac{d}{du} \mathbf{b}(u_1),$$

from which follows that

$$\frac{n!}{(n-1)!} \frac{1}{u_1 - u_0} \sum_{i=0}^{n-1} \Delta^1 \mathbf{a}_i B_i^{n-1}(1) = \frac{n!}{(n-1)!} \frac{1}{u_2 - u_1} \sum_{i=0}^{n-1} \Delta^1 \mathbf{b}_i B_i^{n-1}(0),$$

$$n \frac{1}{\Delta u_0} (\mathbf{a}_n - \mathbf{a}_{n-1}) = n \frac{1}{\Delta u_1} (\mathbf{b}_1 - \mathbf{b}_0),$$

$$\frac{\|\mathbf{a}_n - \mathbf{a}_{n-1}\|}{\Delta u_0} = \frac{\|\mathbf{b}_1 - \mathbf{b}_0\|}{\Delta u_1},$$

$$\|\Delta^1 \mathbf{a}_{n-1}\| : \|\Delta^1 \mathbf{b}_0\| = \Delta u_0 : \Delta u_1.$$



## Properties of Bézier curves

### Bézier curves joined with $C^1$ continuity

Using the same notations as in case of  $C^0$  continuity, Bézier curves

$$\mathbf{a}(u), u \in [u_0, u_1], \Delta u_0 = u_1 - u_0,$$

and

$$\mathbf{b}(u), u \in [u_1, u_2], \Delta u_1 = u_2 - u_1$$

of degree  $n \geq 1$  are joined with  $C^1$  continuity, if they are  $C^0$  continuous (i.e.  $\mathbf{a}_n = \mathbf{b}_0$ ) and

$$\frac{d}{du} \mathbf{a}(u_1) = \frac{d}{du} \mathbf{b}(u_1),$$

from which follows that

$$\begin{aligned} \frac{n!}{(n-1)!} \frac{1}{u_1 - u_0} \sum_{i=0}^{n-1} \Delta^1 \mathbf{a}_i B_i^{n-1}(1) &= \frac{n!}{(n-1)!} \frac{1}{u_2 - u_1} \sum_{i=0}^{n-1} \Delta^1 \mathbf{b}_i B_i^{n-1}(0), \\ n \frac{1}{\Delta u_0} (\mathbf{a}_n - \mathbf{a}_{n-1}) &= n \frac{1}{\Delta u_1} (\mathbf{b}_1 - \mathbf{b}_0), \\ \frac{\|\mathbf{a}_n - \mathbf{a}_{n-1}\|}{\Delta u_0} &= \frac{\|\mathbf{b}_1 - \mathbf{b}_0\|}{\Delta u_1}, \\ \|\Delta^1 \mathbf{a}_{n-1}\| : \|\Delta^1 \mathbf{b}_0\| &= \Delta u_0 : \Delta u_1. \end{aligned}$$



## Properties of Bézier curves

### Bézier curves joined with $C^1$ continuity

Using the same notations as in case of  $C^0$  continuity, Bézier curves

$$\mathbf{a}(u), u \in [u_0, u_1], \Delta u_0 = u_1 - u_0,$$

and

$$\mathbf{b}(u), u \in [u_1, u_2], \Delta u_1 = u_2 - u_1$$

of degree  $n \geq 1$  are joined with  $C^1$  continuity, if they are  $C^0$  continuous (i.e.  $\mathbf{a}_n = \mathbf{b}_0$ ) and

$$\frac{d}{du} \mathbf{a}(u_1) = \frac{d}{du} \mathbf{b}(u_1),$$

from which follows that

$$\frac{n!}{(n-1)!} \frac{1}{u_1 - u_0} \sum_{i=0}^{n-1} \Delta^1 \mathbf{a}_i B_i^{n-1}(1) = \frac{n!}{(n-1)!} \frac{1}{u_2 - u_1} \sum_{i=0}^{n-1} \Delta^1 \mathbf{b}_i B_i^{n-1}(0),$$

$$n \frac{1}{\Delta u_0} (\mathbf{a}_n - \mathbf{a}_{n-1}) = n \frac{1}{\Delta u_1} (\mathbf{b}_1 - \mathbf{b}_0),$$


$$\frac{\|\mathbf{a}_n - \mathbf{a}_{n-1}\|}{\Delta u_0} = \frac{\|\mathbf{b}_1 - \mathbf{b}_0\|}{\Delta u_1},$$

$$\|\Delta^1 \mathbf{a}_{n-1}\| : \|\Delta^1 \mathbf{b}_0\| = \Delta u_0 : \Delta u_1.$$



## Properties of Bézier curves

### Example (Bézier curves joined with $C^1$ continuity)



**Fig. 11:** Several cubic Bézier arcs are joined with  $C^1$  continuity. For the sake of simplicity all cubic arcs are defined on the interval  $[0, 1]$ , i.e. the joint and its two adjacent control points are collinear and the segments defined by them are congruent.

## Properties of Bézier curves

### Bézier curves joined with $C^r$ ( $r \geq 0$ ) continuity

Bézier curves

$$\mathbf{a}(u) = \sum_{i=0}^n \mathbf{a}_i B_i^n \left( \frac{u - u_0}{u_1 - u_0} \right), \quad u \in [u_0, u_1], \quad \Delta u_0 = u_1 - u_0,$$

and

$$\mathbf{b}(u) = \sum_{i=0}^n \mathbf{b}_i B_i^n \left( \frac{u - u_1}{u_2 - u_1} \right), \quad u \in [u_1, u_2], \quad \Delta u_1 = u_2 - u_1$$

of degree  $n \geq 1$  are joined with  $C^r$  ( $r \geq 0$ ) continuity, if the system of conditions

$$\left( \frac{1}{\Delta u_0} \right)^i \Delta^i \mathbf{a}_{n-i} = \left( \frac{1}{\Delta u_1} \right)^i \Delta^i \mathbf{b}_0, \quad i = 0, 1, \dots, r$$

is fulfilled.



## Properties of Bézier curves

### Bézier curves joined with $C^r$ ( $r \geq 0$ ) continuity

Bézier curves

$$\mathbf{a}(u) = \sum_{i=0}^n \mathbf{a}_i B_i^n \left( \frac{u - u_0}{u_1 - u_0} \right), \quad u \in [u_0, u_1], \quad \Delta u_0 = u_1 - u_0,$$

and

$$\mathbf{b}(u) = \sum_{i=0}^n \mathbf{b}_i B_i^n \left( \frac{u - u_1}{u_2 - u_1} \right), \quad u \in [u_1, u_2], \quad \Delta u_1 = u_2 - u_1$$

of degree  $n \geq 1$  are joined with  $C^r$  ( $r \geq 0$ ) continuity, if the system of conditions

$$\left( \frac{1}{\Delta u_0} \right)^i \Delta^i \mathbf{a}_{n-i} = \left( \frac{1}{\Delta u_1} \right)^i \Delta^i \mathbf{b}_0, \quad i = 0, 1, \dots, r$$

is fulfilled.

