

Figure 1 displays three sets of plots comparing the distribution of gene sets across different categories (not claimed by any selected set, all other selected sets) for three gene sets: WT_heat_up (202 genes), KO_heat_up (221 genes), and Interaction_fdrOnly_up (19 genes). The plots are categorized by 'fractional' and 'winner_take_all' methods. The x-axis represents the 'share of the arm' from 0% to 100%.

The legend indicates two types of gene sets: unpinned (white circle) and hypoxia_anchored (pink circle).

WT_heat_up (202 genes):

- fractional:**
 - not claimed by any selected set: 0.366 -> 0.366 the residue
 - all other selected sets: 0.608 -> 0.595 39 sets
- winner_take_all:**
 - not claimed by any selected set: 0.366 -> 0.366 the residue
 - all other selected sets: 0.614 -> 0.614 39 sets

KO_heat_up (221 genes):

- fractional:**
 - not claimed by any selected set: 0.452 -> 0.448 the residue
 - all other selected sets: 0.517 -> 0.507 35 sets
- winner_take_all:**
 - not claimed by any selected set: 0.452 -> 0.448 the residue
 - all other selected sets: 0.511 -> 0.511 35 sets

Interaction_fdrOnly_up (19 genes):

- fractional:**
 - not claimed by any selected set: 0.053 -> 0.053 the residue
 - all other selected sets: 0.947 -> 0.947 21 sets
- winner_take_all:**
 - not claimed by any selected set: 0.053 -> 0.053 the residue
 - all other selected sets: 0.947 -> 0.947 21 sets