

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84**Session Type:** Survival**Suggested Citation**

Software: Surveillance Research Program, National Cancer Institute SEER*Stat software (www.seer.cancer.gov/seerstat) version 8.3.8.
Data: Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Incidence - SEER Research Data, 9 Registries, Nov 2019 Sub (1975-2017) - Linked To County Attributes - Time Dependent (1990-2017) Income/Rurality, 1969-2018 Counties, National Cancer Institute, DCCPS, Surveillance Research Program, released April 2020, based on the November 2019 submission.

Data

Database: Incidence - SEER Research Data, 9 Registries, Nov 2019 Sub (1975-2017) - Linked To County Attributes - Time Dependent (1990-2017) Income/Rurality, 1969-2018 Counties

Statistic

Statistic: Crude Probability of Death Using Cause of Death
Method: Actuarial
Include: 95% Confidence Intervals
Expected Survival: U.S. 1970-2017 by individual year (White, Black, Other (AI/API), Ages 0-99, All races for Other Unspec 1991+ and Unknown) (Used for Expected Survival Exclusions Only)
Unknown Cause: Excluded
Cause of Death: Dead due to cancer using SEER cause-specific death classification
 Dead: {Cause of Death (COD) and Follow-up.SEER cause-specific death classification} = 'Dead (attributable to this cancer dx)'
 AND {Cause of Death (COD) and Follow-up.COD to site recode} = 'Prostate'

Selection

Select Only: Malignant Behavior, Known Age
Exclude All: Death Certificate and Autopsy, Alive with No Survival Time, Age Not in Expected Survival Table, Year Not in Expected Survival Table, Match Not Found for Other Expected Survival Variables
MP Selection: First Primary Only (Sequence Number 0 or 1)

Case: {User-Defined.PC age groups} = '50-54 years', '55-59 years', '60-64 years', '65-69 years', '70-74 years', '75-79 years', '80-84 years'
 AND {User-Defined.Period} = 'Pre-PSA (1980-1989)', 'Post-PSA (2000-2009)'
 AND {User-Defined.Race B&W} = 'White', 'Black'
 AND {Race, Sex, Year Dx.Sex} = 'Male'
 AND {Site and Morphology.Site recode ICD-O-3/WHO 2008} = 'Prostate'

Table

Row: PC age groups [Age recode with <1 year olds]
 Period [Year of diagnosis]
 Race B&W [Race recode (White, Black, Other)]

User Definitions

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84**User Defs. (cont'd)****PC age groups [Age recode with <1 year olds]**

50-54 years = 50-54 years
 55-59 years = 55-59 years
 60-64 years = 60-64 years
 65-69 years = 65-69 years
 70-74 years = 70-74 years
 75-79 years = 75-79 years
 80-84 years = 80-84 years

Period [Year of diagnosis]

Description: NAACCR Name=Date of Diagnosis–Year, Item #=390

Pre-PSA (1980-1989) = 1980, 1981, 1982, 1983, 1984, 1985, 1986, 1987, 1988, 1989

Post-PSA (2000-2009) = 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009

Race B&W [Race recode (White, Black, Other)]

Description: Caution should be exercised when using this variable. For more information, see https://seer.cancer.gov/seerstat/variables/seer/race_ethnicity.

White = White

Black = Black

Parameters

Calculate Survival: From Pre-calculated Duration

Duration: Survival months

Begin Date: Date of diagnosis recode

End Date: Date of follow-up recode

Study Cutoff: Dec 2017

Vital Status: Vital status recode (study cutoff used)

Censored: Attained Age Exceeds Max of Expected Rate Table

Intervals: 120 (1 month each)

Display: Cumulative Summary (Interval as row: 12,24,36,48,60,72,84,96,108,120) (Special Interval and Median Survival Included) (Stats: Cause-Specific; Crude; Standard Errors; Confidence Intervals)

Output

Title: Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Display as: Percents

Display with: 1 decimal place (default)

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Total number of cases read: 5,347,692

Case Exclusion Counts

Number Selected	Number Excluded	Statement

User-defined Exclusions/Selections:		
259,400	5,088,292	{User-Defined.PC age groups} = '50-54 years','55-59 years', '60-64 years','65-69 years','70-74 years','75-79 years', '80-84 years' AND {User-Defined.Period} = 'Pre-PSA (1980-1989)','Post-PSA (2000-2009)' ... (see Session Printout)
257,282	2,118	Exclude death certificate or autopsy cases
257,144	138	Select only malignant cancers
257,144	0	Select only known age
Expected Survival Table Exclusions:		
257,144	0	Exclude age values not in table
257,144	0	Exclude Race recode (White, Black, Other) values not in table
257,144	0	Exclude Sex values not in table
257,144	0	Exclude Invalid Age Year
257,144	0	Exclude Invalid Age Month
Multiple Primary Selection:		
233,729	23,415	First Primary Only (Sequence Number 0 or 1)
Survival Calculation Exclusions:		
233,729	0	Invalid vital status
233,729	0	Unknown survival duration
233,729	0	Calculation dates invalid
233,729	0	Coded survival duration conflicts with associated begin and end dates
233,729	0	Begin Date beyond Study Cutoff Date
233,729	0	Begin Date plus Survival Duration beyond Database Study Cutoff Date
231,969	1,760	Unknown/Missing Cause of Death
231,843	126	Alive with no survival time

Number of records censored due to attained age being greater than expected table max: 0

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Number of records for which a cohort table could not be located: 0

Total number of records used in the analysis: 231,843

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				N	Median Obs	Observed	SE Obs	Obs Cum CIs Lower	Obs Cum CIs Upper	Specified COD	Other CODs	SE Specified COD
50-54 years	Pre-PSA (1980-1989)	White	12 mo	1,184	+	96.7%	0.5%	95.5%	97.6%	2.4%	0.8%	0.4%
			24 mo	1,184	+	89.4%	0.9%	87.5%	91.1%	9.0%	1.6%	0.8%
			36 mo	1,184	+	83.4%	1.1%	81.1%	85.4%	13.6%	3.0%	1.0%
			48 mo	1,184	+	78.1%	1.2%	75.7%	80.4%	18.0%	3.9%	1.1%
			60 mo	1,184	+	73.9%	1.3%	71.3%	76.3%	20.9%	5.2%	1.2%
			72 mo	1,184	+	70.8%	1.3%	68.1%	73.3%	23.4%	5.8%	1.2%
			84 mo	1,184	+	67.5%	1.4%	64.7%	70.1%	25.4%	7.1%	1.3%
			96 mo	1,184	+	64.9%	1.4%	62.2%	67.6%	27.1%	7.9%	1.3%
			108 mo	1,184	+	62.7%	1.4%	59.8%	65.4%	28.5%	8.8%	1.3%
			120 mo	1,184	+	60.7%	1.4%	57.9%	63.4%	29.5%	9.8%	1.3%
		Black	12 mo	199	94.48	89.9%	2.1%	84.9%	93.4%	6.0%	4.0%	1.7%
			24 mo	199	94.48	76.9%	3.0%	70.4%	82.1%	17.1%	6.0%	2.7%
			36 mo	199	94.48	70.4%	3.2%	63.5%	76.2%	22.1%	7.5%	2.9%
			48 mo	199	94.48	63.8%	3.4%	56.7%	70.1%	28.1%	8.0%	3.2%
			60 mo	199	94.48	56.8%	3.5%	49.6%	63.3%	33.7%	9.5%	3.3%
			72 mo	199	94.48	54.3%	3.5%	47.1%	60.9%	34.7%	11.1%	3.4%
			84 mo	199	94.48	52.3%	3.5%	45.1%	58.9%	35.7%	12.1%	3.4%
			96 mo	199	94.48	49.7%	3.5%	42.6%	56.5%	37.2%	13.1%	3.4%
			108 mo	199	94.48	48.7%	3.5%	41.6%	55.4%	37.7%	13.6%	3.4%
			120 mo	199	94.48	46.2%	3.5%	39.1%	52.9%	38.7%	15.1%	3.5%
	Post-PSA (2000-20 09)	White	12 mo	10,505	+	99.3%	0.1%	99.1%	99.5%	0.4%	0.3%	0.1%
			24 mo	10,505	+	98.5%	0.1%	98.3%	98.7%	0.9%	0.6%	0.1%
			36 mo	10,505	+	97.4%	0.2%	97.1%	97.7%	1.5%	1.1%	0.1%
			48 mo	10,505	+	96.6%	0.2%	96.3%	97.0%	1.9%	1.5%	0.1%
			60 mo	10,505	+	95.7%	0.2%	95.3%	96.1%	2.4%	2.0%	0.1%
			72 mo	10,505	+	94.9%	0.2%	94.4%	95.3%	2.7%	2.4%	0.2%
			84 mo	10,505	+	94.0%	0.2%	93.5%	94.4%	3.2%	2.9%	0.2%
			96 mo	10,505	+	93.0%	0.2%	92.5%	93.5%	3.5%	3.5%	0.2%
			108 mo	10,505	+	92.0%	0.3%	91.5%	92.5%	3.9%	4.1%	0.2%
			120 mo	10,505	+	91.0%	0.3%	90.5%	91.6%	4.2%	4.7%	0.2%
		Black	12 mo	2,622	+	98.5%	0.2%	97.9%	98.9%	0.7%	0.8%	0.2%
			24 mo	2,622	+	96.2%	0.4%	95.4%	96.8%	2.1%	1.8%	0.3%
			36 mo	2,622	+	95.1%	0.4%	94.2%	95.9%	2.6%	2.3%	0.3%
			48 mo	2,622	+	94.0%	0.5%	93.0%	94.8%	3.0%	3.0%	0.3%
			60 mo	2,622	+	92.6%	0.5%	91.6%	93.6%	3.3%	4.0%	0.4%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				SE Other CODs	Specified COD Cum CI Lower	Specified COD Cum CI Upper	Other CODs Cum CI Lower	Other CODs Cum CI Upper
50-54 years	Pre-PSA (1980-1989)	White	12 mo	0.3%	1.7%	3.5%	0.4%	1.5%
			24 mo	0.4%	7.4%	10.7%	1.0%	2.4%
			36 mo	0.5%	11.7%	15.6%	2.2%	4.1%
			48 mo	0.6%	15.9%	20.2%	2.9%	5.1%
			60 mo	0.6%	18.7%	23.3%	4.0%	6.5%
			72 mo	0.7%	21.0%	25.8%	4.6%	7.3%
			84 mo	0.7%	23.0%	27.9%	5.7%	8.7%
			96 mo	0.8%	24.6%	29.7%	6.5%	9.6%
			108 mo	0.8%	26.0%	31.1%	7.3%	10.5%
			120 mo	0.9%	26.9%	32.1%	8.2%	11.6%
		Black	12 mo	1.4%	3.3%	9.9%	1.9%	7.4%
			24 mo	1.7%	12.2%	22.6%	3.3%	9.9%
			36 mo	1.9%	16.6%	28.1%	4.4%	11.7%
			48 mo	1.9%	22.1%	34.5%	4.8%	12.3%
			60 mo	2.1%	27.2%	40.2%	6.0%	14.1%
			72 mo	2.2%	28.1%	41.3%	7.2%	15.9%
			84 mo	2.3%	29.1%	42.3%	8.0%	17.0%
			96 mo	2.4%	30.5%	43.9%	8.8%	18.2%
			108 mo	2.4%	31.0%	44.4%	9.3%	18.7%
			120 mo	2.5%	31.9%	45.4%	10.5%	20.4%
	Post-PSA (2000-20 09)	White	12 mo	0.1%	0.3%	0.5%	0.2%	0.4%
			24 mo	0.1%	0.7%	1.1%	0.5%	0.8%
			36 mo	0.1%	1.3%	1.7%	0.9%	1.3%
			48 mo	0.1%	1.7%	2.2%	1.2%	1.7%
			60 mo	0.1%	2.1%	2.7%	1.7%	2.2%
			72 mo	0.1%	2.4%	3.1%	2.1%	2.7%
			84 mo	0.2%	2.8%	3.5%	2.6%	3.2%
			96 mo	0.2%	3.2%	3.9%	3.1%	3.8%
			108 mo	0.2%	3.5%	4.3%	3.7%	4.5%
			120 mo	0.2%	3.9%	4.6%	4.3%	5.2%
		Black	12 mo	0.2%	0.4%	1.1%	0.5%	1.3%
			24 mo	0.3%	1.6%	2.7%	1.3%	2.3%
			36 mo	0.3%	2.0%	3.2%	1.8%	2.9%
			48 mo	0.3%	2.4%	3.7%	2.4%	3.7%
			60 mo	0.4%	2.7%	4.1%	3.3%	4.8%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				N	Median Obs	Observed	SE Obs	Obs Cum CIs Lower	Obs Cum CIs Upper	Specified COD	Other CODs	SE Specified COD
50-54 years	Post-PSA (2000-2009)	Black	72 mo	2,622	+	90.9%	0.6%	89.7%	91.9%	4.0%	5.1%	0.4%
			84 mo	2,622	+	89.5%	0.6%	88.2%	90.6%	4.5%	6.0%	0.4%
			96 mo	2,622	+	88.1%	0.6%	86.8%	89.3%	5.0%	6.9%	0.4%
			108 mo	2,622	+	86.3%	0.7%	84.9%	87.6%	5.4%	8.3%	0.4%
			120 mo	2,622	+	84.4%	0.7%	82.9%	85.8%	5.8%	9.8%	0.5%
55-59 years	Pre-PSA (1980-1989)	White	12 mo	3,596	+	95.0%	0.4%	94.2%	95.6%	3.1%	1.9%	0.3%
			24 mo	3,596	+	88.7%	0.5%	87.6%	89.7%	7.8%	3.5%	0.4%
			36 mo	3,596	+	82.5%	0.6%	81.2%	83.7%	12.3%	5.2%	0.5%
			48 mo	3,596	+	77.5%	0.7%	76.1%	78.8%	15.8%	6.8%	0.6%
			60 mo	3,596	+	72.5%	0.7%	71.0%	73.9%	19.3%	8.2%	0.7%
			72 mo	3,596	+	68.8%	0.8%	67.2%	70.3%	21.7%	9.5%	0.7%
			84 mo	3,596	+	65.3%	0.8%	63.8%	66.9%	23.9%	10.8%	0.7%
			96 mo	3,596	+	62.3%	0.8%	60.7%	63.9%	25.5%	12.2%	0.7%
			108 mo	3,596	+	59.7%	0.8%	58.1%	61.3%	27.1%	13.2%	0.7%
			120 mo	3,596	+	56.9%	0.8%	55.3%	58.5%	28.5%	14.6%	0.8%
		Black	12 mo	629		93.17	1.0%	91.4%	95.3%	3.3%	3.0%	0.7%
			24 mo	629		93.17	1.5%	79.0%	85.0%	12.1%	5.7%	1.3%
			36 mo	629		93.17	1.7%	71.5%	78.2%	18.0%	7.0%	1.5%
			48 mo	629		93.17	1.9%	62.9%	70.3%	24.2%	9.1%	1.7%
			60 mo	629		93.17	2.0%	56.3%	64.0%	28.1%	11.6%	1.8%
			72 mo	629		93.17	2.0%	51.8%	59.6%	30.7%	13.5%	1.8%
			84 mo	629		93.17	2.0%	48.5%	56.3%	32.8%	14.8%	1.9%
			96 mo	629		93.17	2.0%	44.8%	52.6%	34.7%	16.5%	1.9%
			108 mo	629		93.17	2.0%	40.8%	48.5%	36.9%	18.4%	1.9%
			120 mo	629		93.17	2.0%	37.9%	45.6%	38.3%	19.9%	1.9%
	Post-PSA (2000-2009)	White	12 mo	19,927	+	99.2%	0.1%	99.1%	99.3%	0.4%	0.4%	0.0%
			24 mo	19,927	+	98.1%	0.1%	97.9%	98.2%	1.0%	1.0%	0.1%
			36 mo	19,927	+	97.0%	0.1%	96.7%	97.2%	1.5%	1.5%	0.1%
			48 mo	19,927	+	95.9%	0.1%	95.7%	96.2%	2.0%	2.1%	0.1%
			60 mo	19,927	+	94.9%	0.2%	94.6%	95.2%	2.3%	2.8%	0.1%
			72 mo	19,927	+	93.9%	0.2%	93.6%	94.2%	2.6%	3.4%	0.1%
			84 mo	19,927	+	92.8%	0.2%	92.4%	93.1%	3.0%	4.2%	0.1%
			96 mo	19,927	+	91.5%	0.2%	91.1%	91.9%	3.3%	5.1%	0.1%
			108 mo	19,927	+	90.3%	0.2%	89.9%	90.7%	3.7%	6.0%	0.1%
			120 mo	19,927	+	88.9%	0.2%	88.5%	89.4%	4.0%	7.1%	0.1%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				SE Other CODs	Specified COD Cum CI Lower	Specified COD Cum CI Upper	Other CODs Cum CI Lower	Other CODs Cum CI Upper
50-54 years	Post-PSA (2000-2009)	Black	72 mo	0.4%	3.3%	4.8%	4.3%	6.0%
			84 mo	0.5%	3.7%	5.3%	5.2%	7.0%
			96 mo	0.5%	4.2%	5.8%	6.0%	7.9%
			108 mo	0.5%	4.6%	6.3%	7.3%	9.4%
			120 mo	0.6%	4.9%	6.7%	8.7%	11.1%
55-59 years	Pre-PSA (1980-1989)	White	12 mo	0.2%	2.6%	3.7%	1.5%	2.4%
			24 mo	0.3%	7.0%	8.8%	2.9%	4.1%
			36 mo	0.4%	11.3%	13.4%	4.5%	5.9%
			48 mo	0.4%	14.6%	17.0%	6.0%	7.6%
			60 mo	0.5%	18.0%	20.6%	7.3%	9.1%
			72 mo	0.5%	20.4%	23.1%	8.6%	10.5%
			84 mo	0.5%	22.5%	25.3%	9.8%	11.8%
			96 mo	0.5%	24.1%	27.0%	11.1%	13.3%
			108 mo	0.6%	25.7%	28.6%	12.1%	14.3%
			120 mo	0.6%	27.1%	30.0%	13.4%	15.7%
		Black	12 mo	0.7%	2.1%	5.0%	1.9%	4.6%
			24 mo	0.9%	9.7%	14.8%	4.1%	7.7%
			36 mo	1.0%	15.1%	21.1%	5.2%	9.2%
			48 mo	1.1%	20.9%	27.6%	7.0%	11.5%
			60 mo	1.3%	24.7%	31.7%	9.3%	14.2%
			72 mo	1.4%	27.1%	34.3%	11.0%	16.3%
			84 mo	1.4%	29.1%	36.4%	12.1%	17.7%
			96 mo	1.5%	31.0%	38.4%	13.7%	19.5%
			108 mo	1.5%	33.1%	40.6%	15.5%	21.6%
			120 mo	1.6%	34.5%	42.1%	16.9%	23.1%
	Post-PSA (2000-2009)	White	12 mo	0.0%	0.3%	0.5%	0.3%	0.5%
			24 mo	0.1%	0.8%	1.1%	0.8%	1.1%
			36 mo	0.1%	1.3%	1.7%	1.4%	1.7%
			48 mo	0.1%	1.8%	2.2%	1.9%	2.3%
			60 mo	0.1%	2.1%	2.5%	2.5%	3.0%
			72 mo	0.1%	2.4%	2.9%	3.2%	3.7%
			84 mo	0.1%	2.7%	3.2%	4.0%	4.5%
			96 mo	0.2%	3.1%	3.6%	4.8%	5.4%
			108 mo	0.2%	3.4%	3.9%	5.7%	6.4%
			120 mo	0.2%	3.8%	4.3%	6.7%	7.4%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				N	Median Obs	Observed	SE Obs	Obs Cum CIs Lower	Obs Cum CIs Upper	Specified COD	Other CODs	SE Specified COD
55-59 years	Post-PSA (2000-2009)	Black	12 mo	4,352	+	98.5%	0.2%	98.0%	98.8%	0.7%	0.8%	0.1%
			24 mo	4,352	+	96.2%	0.3%	95.6%	96.7%	1.8%	2.0%	0.2%
			36 mo	4,352	+	94.2%	0.4%	93.5%	94.9%	2.4%	3.3%	0.2%
			48 mo	4,352	+	91.9%	0.4%	91.1%	92.7%	3.1%	5.0%	0.3%
			60 mo	4,352	+	90.0%	0.5%	89.1%	90.9%	3.7%	6.3%	0.3%
			72 mo	4,352	+	87.8%	0.5%	86.8%	88.8%	4.4%	7.8%	0.3%
			84 mo	4,352	+	86.2%	0.5%	85.1%	87.2%	4.7%	9.1%	0.3%
			96 mo	4,352	+	84.3%	0.6%	83.2%	85.4%	5.2%	10.5%	0.3%
			108 mo	4,352	+	82.0%	0.6%	80.8%	83.1%	5.8%	12.3%	0.4%
			120 mo	4,352	+	79.5%	0.6%	78.2%	80.7%	6.2%	14.3%	0.4%
60-64 years	Pre-PSA (1980-1989)	White	12 mo	8,152	+	95.6%	0.2%	95.1%	96.0%	2.3%	2.1%	0.2%
			24 mo	8,152	+	89.5%	0.3%	88.8%	90.1%	6.4%	4.1%	0.3%
			36 mo	8,152	+	83.4%	0.4%	82.6%	84.2%	10.4%	6.1%	0.3%
			48 mo	8,152	+	78.2%	0.5%	77.3%	79.1%	13.6%	8.2%	0.4%
			60 mo	8,152	+	73.3%	0.5%	72.3%	74.2%	16.3%	10.4%	0.4%
			72 mo	8,152	+	69.4%	0.5%	68.3%	70.3%	18.4%	12.2%	0.4%
			84 mo	8,152	+	65.4%	0.5%	64.3%	66.4%	20.3%	14.4%	0.4%
			96 mo	8,152	+	61.6%	0.5%	60.5%	62.6%	22.0%	16.4%	0.5%
			108 mo	8,152	+	58.2%	0.5%	57.1%	59.3%	23.3%	18.4%	0.5%
			120 mo	8,152	+	55.0%	0.6%	53.9%	56.0%	24.4%	20.6%	0.5%
		Black	12 mo	1,275	80.75	91.3%	0.8%	89.6%	92.7%	4.5%	4.2%	0.6%
			24 mo	1,275	80.75	80.2%	1.1%	77.9%	82.3%	11.9%	7.8%	0.9%
			36 mo	1,275	80.75	71.9%	1.3%	69.4%	74.3%	17.7%	10.4%	1.1%
			48 mo	1,275	80.75	63.5%	1.3%	60.7%	66.0%	23.1%	13.4%	1.2%
			60 mo	1,275	80.75	58.0%	1.4%	55.3%	60.7%	25.9%	16.1%	1.2%
			72 mo	1,275	80.75	53.1%	1.4%	50.3%	55.8%	28.2%	18.7%	1.3%
			84 mo	1,275	80.75	48.9%	1.4%	46.1%	51.6%	30.3%	20.9%	1.3%
			96 mo	1,275	80.75	44.6%	1.4%	41.9%	47.3%	31.8%	23.5%	1.3%
			108 mo	1,275	80.75	40.9%	1.4%	38.1%	43.5%	33.3%	25.9%	1.3%
			120 mo	1,275	80.75	36.4%	1.3%	33.7%	39.0%	35.5%	28.2%	1.3%
	Post-PSA (2000-2009)	White	12 mo	25,698	+	99.0%	0.1%	98.8%	99.1%	0.5%	0.6%	0.0%
			24 mo	25,698	+	97.6%	0.1%	97.5%	97.8%	1.0%	1.3%	0.1%
			36 mo	25,698	+	96.3%	0.1%	96.0%	96.5%	1.6%	2.2%	0.1%
			48 mo	25,698	+	94.9%	0.1%	94.6%	95.2%	2.1%	3.0%	0.1%
			60 mo	25,698	+	93.3%	0.2%	93.0%	93.6%	2.5%	4.2%	0.1%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				SE Other CODs	Specified COD Cum CI Lower	Specified COD Cum CI Upper	Other CODs Cum CI Lower	Other CODs Cum CI Upper
55-59 years	Post-PSA (2000-2009)	Black	12 mo	0.1%	0.5%	1.0%	0.6%	1.1%
			24 mo	0.2%	1.5%	2.3%	1.6%	2.4%
			36 mo	0.3%	2.0%	2.9%	2.8%	3.9%
			48 mo	0.3%	2.6%	3.6%	4.4%	5.7%
			60 mo	0.4%	3.2%	4.3%	5.6%	7.0%
			72 mo	0.4%	3.8%	5.0%	7.0%	8.6%
			84 mo	0.4%	4.1%	5.4%	8.3%	10.0%
			96 mo	0.5%	4.6%	5.9%	9.6%	11.4%
			108 mo	0.5%	5.1%	6.5%	11.3%	13.3%
			120 mo	0.5%	5.5%	7.0%	13.3%	15.4%
60-64 years	Pre-PSA (19 80-1989)	White	12 mo	0.2%	2.0%	2.6%	1.8%	2.5%
			24 mo	0.2%	5.9%	7.0%	3.7%	4.6%
			36 mo	0.3%	9.8%	11.1%	5.6%	6.7%
			48 mo	0.3%	12.8%	14.3%	7.7%	8.9%
			60 mo	0.3%	15.6%	17.2%	9.7%	11.0%
			72 mo	0.4%	17.6%	19.3%	11.5%	12.9%
			84 mo	0.4%	19.4%	21.1%	13.6%	15.2%
			96 mo	0.4%	21.1%	22.9%	15.6%	17.2%
			108 mo	0.4%	22.4%	24.3%	17.6%	19.3%
			120 mo	0.4%	23.5%	25.3%	19.8%	21.5%
		Black	12 mo	0.6%	3.5%	5.8%	3.2%	5.4%
			24 mo	0.8%	10.2%	13.8%	6.5%	9.4%
			36 mo	0.9%	15.7%	19.9%	8.8%	12.1%
			48 mo	1.0%	20.9%	25.5%	11.6%	15.3%
			60 mo	1.0%	23.5%	28.3%	14.1%	18.2%
			72 mo	1.1%	25.8%	30.7%	16.6%	20.9%
			84 mo	1.1%	27.8%	32.8%	18.7%	23.1%
			96 mo	1.2%	29.3%	34.4%	21.2%	25.9%
			108 mo	1.2%	30.7%	35.9%	23.5%	28.3%
			120 mo	1.3%	32.8%	38.1%	25.7%	30.7%
	Post-PSA (2000-2009)	White	12 mo	0.0%	0.4%	0.5%	0.5%	0.7%
			24 mo	0.1%	0.9%	1.2%	1.2%	1.5%
			36 mo	0.1%	1.4%	1.7%	2.0%	2.3%
			48 mo	0.1%	1.9%	2.2%	2.8%	3.2%
			60 mo	0.1%	2.4%	2.7%	3.9%	4.4%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				N	Median Obs	Observed	SE Obs	Obs Cum CIs Lower	Obs Cum CIs Upper	Specified COD	Other CODs	SE Specified COD
60-64 years	Post-PSA (2000-2009)	White	72 mo	25,698	+	91.7%	0.2%	91.4%	92.1%	2.9%	5.3%	0.1%
			84 mo	25,698	+	90.3%	0.2%	89.9%	90.6%	3.3%	6.4%	0.1%
			96 mo	25,698	+	88.7%	0.2%	88.3%	89.0%	3.7%	7.7%	0.1%
			108 mo	25,698	+	86.9%	0.2%	86.5%	87.3%	4.1%	9.0%	0.1%
			120 mo	25,698	+	84.9%	0.2%	84.5%	85.4%	4.5%	10.5%	0.1%
		Black	12 mo	4,768	+	97.7%	0.2%	97.2%	98.1%	1.0%	1.3%	0.1%
			24 mo	4,768	+	95.3%	0.3%	94.6%	95.8%	1.9%	2.9%	0.2%
			36 mo	4,768	+	92.8%	0.4%	92.0%	93.5%	2.7%	4.5%	0.2%
			48 mo	4,768	+	90.6%	0.4%	89.8%	91.4%	3.4%	5.9%	0.3%
			60 mo	4,768	+	88.2%	0.5%	87.2%	89.1%	4.1%	7.7%	0.3%
			72 mo	4,768	+	85.4%	0.5%	84.4%	86.4%	4.8%	9.8%	0.3%
			84 mo	4,768	+	82.6%	0.6%	81.5%	83.7%	5.4%	11.9%	0.3%
			96 mo	4,768	+	80.1%	0.6%	78.9%	81.2%	5.9%	14.0%	0.3%
			108 mo	4,768	+	77.1%	0.6%	75.9%	78.3%	6.5%	16.4%	0.4%
			120 mo	4,768	+	74.4%	0.6%	73.1%	75.7%	6.9%	18.7%	0.4%
65-69 years	Pre-PSA (1980-1989)	White	12 mo	13,151	109.81	94.2%	0.2%	93.8%	94.6%	2.4%	3.3%	0.1%
			24 mo	13,151	109.81	87.3%	0.3%	86.7%	87.8%	6.4%	6.3%	0.2%
			36 mo	13,151	109.81	80.5%	0.3%	79.9%	81.2%	10.2%	9.3%	0.3%
			48 mo	13,151	109.81	74.4%	0.4%	73.6%	75.1%	13.4%	12.3%	0.3%
			60 mo	13,151	109.81	69.1%	0.4%	68.3%	69.9%	16.1%	14.8%	0.3%
			72 mo	13,151	109.81	64.0%	0.4%	63.2%	64.8%	18.5%	17.5%	0.3%
			84 mo	13,151	109.81	59.2%	0.4%	58.4%	60.1%	20.4%	20.3%	0.4%
			96 mo	13,151	109.81	54.7%	0.4%	53.8%	55.5%	22.2%	23.1%	0.4%
			108 mo	13,151	109.81	50.6%	0.4%	49.7%	51.4%	23.7%	25.7%	0.4%
			120 mo	13,151	109.81	46.9%	0.4%	46.0%	47.7%	25.1%	28.1%	0.4%
		Black	12 mo	1,811	72.68	89.7%	0.7%	88.2%	91.0%	4.5%	5.9%	0.5%
			24 mo	1,811	72.68	78.0%	1.0%	76.0%	79.8%	11.4%	10.7%	0.7%
			36 mo	1,811	72.68	68.9%	1.1%	66.7%	71.0%	16.3%	14.8%	0.9%
			48 mo	1,811	72.68	62.3%	1.1%	60.0%	64.5%	19.4%	18.3%	0.9%
			60 mo	1,811	72.68	57.0%	1.2%	54.6%	59.2%	21.9%	21.2%	1.0%
			72 mo	1,811	72.68	50.3%	1.2%	47.9%	52.5%	24.7%	25.0%	1.0%
			84 mo	1,811	72.68	44.8%	1.2%	42.5%	47.1%	27.0%	28.2%	1.0%
			96 mo	1,811	72.68	40.1%	1.2%	37.8%	42.3%	28.7%	31.2%	1.1%
			108 mo	1,811	72.68	35.8%	1.1%	33.6%	38.0%	30.4%	33.8%	1.1%
			120 mo	1,811	72.68	32.8%	1.1%	30.6%	35.0%	31.7%	35.5%	1.1%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				SE Other CODs	Specified COD Cum CI Lower	Specified COD Cum CI Upper	Other CODs Cum CI Lower	Other CODs Cum CI Upper
60-64 years	Post-PSA (2000-2009)	White	72 mo	0.1%	2.7%	3.2%	5.1%	5.6%
			84 mo	0.2%	3.1%	3.5%	6.1%	6.7%
			96 mo	0.2%	3.5%	3.9%	7.3%	8.0%
			108 mo	0.2%	3.9%	4.3%	8.7%	9.4%
			120 mo	0.2%	4.3%	4.8%	10.2%	10.9%
		Black	12 mo	0.2%	0.7%	1.3%	1.0%	1.7%
			24 mo	0.2%	1.5%	2.3%	2.4%	3.4%
			36 mo	0.3%	2.3%	3.2%	3.9%	5.1%
			48 mo	0.3%	2.9%	4.0%	5.3%	6.6%
			60 mo	0.4%	3.6%	4.7%	7.0%	8.5%
			72 mo	0.4%	4.2%	5.4%	9.0%	10.6%
			84 mo	0.5%	4.8%	6.1%	11.0%	12.9%
			96 mo	0.5%	5.3%	6.6%	13.0%	15.0%
			108 mo	0.5%	5.8%	7.2%	15.3%	17.4%
			120 mo	0.6%	6.2%	7.7%	17.5%	19.8%
65-69 years	Pre-PSA (1980-1989)	White	12 mo	0.2%	2.2%	2.7%	3.0%	3.7%
			24 mo	0.2%	6.0%	6.8%	5.9%	6.7%
			36 mo	0.3%	9.7%	10.7%	8.8%	9.8%
			48 mo	0.3%	12.8%	13.9%	11.7%	12.8%
			60 mo	0.3%	15.5%	16.7%	14.2%	15.4%
			72 mo	0.3%	17.8%	19.1%	16.9%	18.2%
			84 mo	0.4%	19.8%	21.1%	19.6%	21.0%
			96 mo	0.4%	21.5%	23.0%	22.4%	23.8%
			108 mo	0.4%	23.0%	24.4%	25.0%	26.5%
			120 mo	0.4%	24.3%	25.8%	27.3%	28.8%
		Black	12 mo	0.6%	3.6%	5.5%	4.8%	7.0%
			24 mo	0.7%	10.0%	12.9%	9.3%	12.1%
			36 mo	0.8%	14.6%	18.0%	13.2%	16.5%
			48 mo	0.9%	17.7%	21.3%	16.5%	20.1%
			60 mo	1.0%	20.0%	23.8%	19.3%	23.1%
			72 mo	1.0%	22.7%	26.7%	23.1%	27.1%
			84 mo	1.1%	24.9%	29.0%	26.1%	30.3%
			96 mo	1.1%	26.7%	30.8%	29.1%	33.3%
			108 mo	1.1%	28.3%	32.5%	31.6%	36.0%
			120 mo	1.1%	29.5%	33.8%	33.3%	37.7%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				N	Median Obs	Observed	SE Obs	Obs Cum CIs Lower	Obs Cum CIs Upper	Specified COD	Other CODs	SE Specified COD
65-69 years	Post-PSA (2000-2009)	White	12 mo	27,807	+	98.4%	0.1%	98.2%	98.5%	0.6%	1.0%	0.0%
			24 mo	27,807	+	96.5%	0.1%	96.3%	96.7%	1.2%	2.3%	0.1%
			36 mo	27,807	+	94.7%	0.1%	94.4%	94.9%	1.9%	3.5%	0.1%
			48 mo	27,807	+	92.5%	0.2%	92.2%	92.8%	2.4%	5.0%	0.1%
			60 mo	27,807	+	90.4%	0.2%	90.1%	90.7%	2.9%	6.7%	0.1%
			72 mo	27,807	+	88.2%	0.2%	87.8%	88.6%	3.4%	8.4%	0.1%
			84 mo	27,807	+	85.8%	0.2%	85.3%	86.2%	3.9%	10.3%	0.1%
			96 mo	27,807	+	83.3%	0.2%	82.8%	83.7%	4.5%	12.2%	0.1%
			108 mo	27,807	+	80.4%	0.2%	80.0%	80.9%	4.9%	14.7%	0.1%
			120 mo	27,807	+	77.7%	0.3%	77.2%	78.2%	5.4%	17.0%	0.1%
		Black	12 mo	4,526	+	96.9%	0.3%	96.4%	97.4%	1.0%	2.0%	0.2%
			24 mo	4,526	+	93.9%	0.4%	93.1%	94.5%	2.0%	4.2%	0.2%
			36 mo	4,526	+	90.3%	0.4%	89.4%	91.1%	2.9%	6.8%	0.2%
			48 mo	4,526	+	86.7%	0.5%	85.7%	87.7%	4.1%	9.2%	0.3%
			60 mo	4,526	+	83.2%	0.6%	82.0%	84.2%	4.9%	11.9%	0.3%
			72 mo	4,526	+	79.9%	0.6%	78.7%	81.1%	5.7%	14.4%	0.3%
			84 mo	4,526	+	76.2%	0.6%	75.0%	77.4%	6.4%	17.4%	0.4%
			96 mo	4,526	+	72.7%	0.7%	71.4%	74.0%	7.0%	20.3%	0.4%
			108 mo	4,526	+	69.4%	0.7%	68.0%	70.7%	7.4%	23.2%	0.4%
			120 mo	4,526	+	65.2%	0.7%	63.8%	66.6%	8.3%	26.5%	0.4%
70-74 years	Pre-PSA (1980-1989)	White	12 mo	15,193	84.84	92.3%	0.2%	91.9%	92.8%	3.0%	4.6%	0.1%
			24 mo	15,193	84.84	84.2%	0.3%	83.7%	84.8%	7.0%	8.7%	0.2%
			36 mo	15,193	84.84	76.3%	0.3%	75.6%	77.0%	11.0%	12.7%	0.3%
			48 mo	15,193	84.84	69.4%	0.4%	68.6%	70.1%	14.0%	16.6%	0.3%
			60 mo	15,193	84.84	62.6%	0.4%	61.8%	63.3%	16.8%	20.6%	0.3%
			72 mo	15,193	84.84	56.5%	0.4%	55.7%	57.3%	19.1%	24.4%	0.3%
			84 mo	15,193	84.84	50.5%	0.4%	49.7%	51.3%	21.3%	28.1%	0.3%
			96 mo	15,193	84.84	45.4%	0.4%	44.6%	46.2%	22.9%	31.7%	0.3%
			108 mo	15,193	84.84	40.7%	0.4%	39.9%	41.4%	24.3%	35.0%	0.3%
			120 mo	15,193	84.84	36.0%	0.4%	35.3%	36.8%	25.7%	38.3%	0.4%
		Black	12 mo	1,737	57.75	87.1%	0.8%	85.4%	88.6%	6.6%	6.3%	0.6%
			24 mo	1,737	57.75	74.8%	1.0%	72.7%	76.8%	12.4%	12.8%	0.8%
			36 mo	1,737	57.75	65.3%	1.1%	63.1%	67.5%	17.6%	17.1%	0.9%
			48 mo	1,737	57.75	56.4%	1.2%	54.0%	58.7%	21.9%	21.7%	1.0%
			60 mo	1,737	57.75	49.3%	1.2%	46.9%	51.6%	25.1%	25.6%	1.0%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				SE Other CODs	Specified COD Cum CI Lower	Specified COD Cum CI Upper	Other CODs Cum CI Lower	Other CODs Cum CI Upper
65-69 years	Post-PSA (2000-2009)	White	12 mo	0.1%	0.5%	0.6%	0.9%	1.2%
			24 mo	0.1%	1.1%	1.4%	2.1%	2.4%
			36 mo	0.1%	1.7%	2.0%	3.3%	3.7%
			48 mo	0.1%	2.3%	2.6%	4.8%	5.3%
			60 mo	0.1%	2.7%	3.1%	6.4%	7.0%
			72 mo	0.2%	3.2%	3.6%	8.1%	8.7%
			84 mo	0.2%	3.7%	4.1%	10.0%	10.7%
			96 mo	0.2%	4.2%	4.7%	11.9%	12.6%
			108 mo	0.2%	4.6%	5.2%	14.3%	15.1%
			120 mo	0.2%	5.1%	5.6%	16.5%	17.4%
		Black	12 mo	0.2%	0.8%	1.4%	1.6%	2.5%
			24 mo	0.3%	1.6%	2.4%	3.6%	4.8%
			36 mo	0.4%	2.4%	3.4%	6.1%	7.6%
			48 mo	0.4%	3.5%	4.7%	8.4%	10.1%
			60 mo	0.5%	4.3%	5.6%	11.0%	12.9%
			72 mo	0.5%	5.1%	6.4%	13.3%	15.4%
			84 mo	0.6%	5.7%	7.1%	16.3%	18.5%
			96 mo	0.6%	6.3%	7.8%	19.1%	21.5%
			108 mo	0.6%	6.7%	8.2%	21.9%	24.4%
			120 mo	0.7%	7.5%	9.1%	25.2%	27.8%
70-74 years	Pre-PSA (1980-1989)	White	12 mo	0.2%	2.8%	3.3%	4.3%	5.0%
			24 mo	0.2%	6.6%	7.5%	8.3%	9.2%
			36 mo	0.3%	10.5%	11.5%	12.2%	13.3%
			48 mo	0.3%	13.5%	14.6%	16.0%	17.2%
			60 mo	0.3%	16.2%	17.4%	20.0%	21.3%
			72 mo	0.3%	18.5%	19.7%	23.7%	25.1%
			84 mo	0.4%	20.7%	22.0%	27.4%	28.9%
			96 mo	0.4%	22.2%	23.6%	31.0%	32.4%
			108 mo	0.4%	23.7%	25.0%	34.2%	35.8%
			120 mo	0.4%	25.0%	26.4%	37.5%	39.1%
		Black	12 mo	0.6%	5.5%	7.8%	5.3%	7.5%
			24 mo	0.8%	10.9%	14.0%	11.3%	14.4%
			36 mo	0.9%	15.8%	19.4%	15.4%	18.9%
			48 mo	1.0%	20.0%	23.9%	19.8%	23.7%
			60 mo	1.0%	23.1%	27.2%	23.6%	27.7%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				N	Median Obs	Observed	SE Obs	Obs Cum CIs Lower	Obs Cum CIs Upper	Specified COD	Other CODs	SE Specified COD
70-74 years	Pre-PSA (1980-1989)	Black	72 mo	1,737	57.75	43.2%	1.2%	40.8%	45.5%	27.9%	28.9%	1.1%
			84 mo	1,737	57.75	37.2%	1.2%	34.9%	39.5%	30.6%	32.2%	1.1%
			96 mo	1,737	57.75	32.0%	1.1%	29.8%	34.2%	32.6%	35.3%	1.1%
			108 mo	1,737	57.75	27.5%	1.1%	25.4%	29.6%	34.1%	38.3%	1.1%
			120 mo	1,737	57.75	23.5%	1.0%	21.5%	25.5%	35.9%	40.6%	1.2%
	Post-PSA (2000-2009)	White	12 mo	24,904	+	97.5%	0.1%	97.3%	97.7%	0.9%	1.6%	0.1%
			24 mo	24,904	+	94.6%	0.1%	94.3%	94.9%	1.8%	3.6%	0.1%
			36 mo	24,904	+	91.5%	0.2%	91.2%	91.9%	2.7%	5.8%	0.1%
			48 mo	24,904	+	88.2%	0.2%	87.8%	88.6%	3.4%	8.4%	0.1%
			60 mo	24,904	+	84.9%	0.2%	84.4%	85.3%	4.1%	11.0%	0.1%
			72 mo	24,904	+	81.4%	0.2%	80.9%	81.9%	4.8%	13.8%	0.1%
			84 mo	24,904	+	77.8%	0.3%	77.3%	78.3%	5.4%	16.8%	0.1%
			96 mo	24,904	+	73.9%	0.3%	73.4%	74.5%	6.0%	20.1%	0.2%
			108 mo	24,904	+	69.7%	0.3%	69.1%	70.3%	6.6%	23.7%	0.2%
			120 mo	24,904	+	65.3%	0.3%	64.7%	65.9%	7.1%	27.5%	0.2%
		Black	12 mo	3,318	+	95.0%	0.4%	94.2%	95.7%	1.8%	3.2%	0.2%
			24 mo	3,318	+	90.4%	0.5%	89.4%	91.4%	3.1%	6.5%	0.3%
			36 mo	3,318	+	86.3%	0.6%	85.1%	87.4%	4.2%	9.5%	0.3%
			48 mo	3,318	+	81.4%	0.7%	80.1%	82.7%	5.4%	13.2%	0.4%
			60 mo	3,318	+	76.9%	0.7%	75.5%	78.3%	6.2%	16.8%	0.4%
			72 mo	3,318	+	72.4%	0.8%	70.8%	73.9%	7.3%	20.3%	0.5%
			84 mo	3,318	+	67.9%	0.8%	66.2%	69.4%	8.1%	24.0%	0.5%
			96 mo	3,318	+	63.7%	0.8%	62.0%	65.3%	8.6%	27.7%	0.5%
			108 mo	3,318	+	59.4%	0.9%	57.7%	61.1%	9.2%	31.4%	0.5%
			120 mo	3,318	+	54.2%	0.9%	52.5%	56.0%	9.9%	35.9%	0.5%
75-79 years	Pre-PSA (1980-1989)	White	12 mo	13,322	62.71	88.4%	0.3%	87.9%	89.0%	4.4%	7.2%	0.2%
			24 mo	13,322	62.71	78.1%	0.4%	77.3%	78.8%	9.0%	12.9%	0.2%
			36 mo	13,322	62.71	68.5%	0.4%	67.7%	69.3%	12.8%	18.7%	0.3%
			48 mo	13,322	62.71	59.8%	0.4%	59.0%	60.6%	15.9%	24.3%	0.3%
			60 mo	13,322	62.71	51.6%	0.4%	50.8%	52.5%	18.7%	29.7%	0.3%
			72 mo	13,322	62.71	45.0%	0.4%	44.2%	45.8%	20.8%	34.2%	0.4%
			84 mo	13,322	62.71	38.6%	0.4%	37.8%	39.5%	22.5%	38.8%	0.4%
			96 mo	13,322	62.71	33.1%	0.4%	32.3%	33.9%	24.0%	42.9%	0.4%
			108 mo	13,322	62.71	28.2%	0.4%	27.4%	28.9%	25.4%	46.4%	0.4%
			120 mo	13,322	62.71	23.4%	0.4%	22.7%	24.2%	26.5%	50.0%	0.4%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				SE Other CODs	Specified COD Cum CI Lower	Specified COD Cum CI Upper	Other CODs Cum CI Lower	Other CODs Cum CI Upper
70-74 years	Pre-PSA (1980-1989)	Black	72 mo	1.1%	25.8%	30.0%	26.8%	31.0%
			84 mo	1.1%	28.4%	32.7%	30.1%	34.4%
			96 mo	1.1%	30.4%	34.9%	33.1%	37.6%
			108 mo	1.2%	31.9%	36.4%	36.1%	40.6%
			120 mo	1.2%	33.7%	38.2%	38.3%	42.9%
	Post-PSA (2000-20 09)	White	12 mo	0.1%	0.8%	1.0%	1.5%	1.8%
			24 mo	0.1%	1.7%	2.0%	3.4%	3.8%
			36 mo	0.1%	2.5%	2.9%	5.5%	6.1%
			48 mo	0.2%	3.2%	3.6%	8.0%	8.7%
			60 mo	0.2%	3.9%	4.4%	10.6%	11.4%
			72 mo	0.2%	4.5%	5.1%	13.4%	14.2%
			84 mo	0.2%	5.1%	5.7%	16.4%	17.3%
			96 mo	0.3%	5.7%	6.3%	19.6%	20.6%
			108 mo	0.3%	6.3%	6.9%	23.2%	24.3%
			120 mo	0.3%	6.8%	7.5%	27.0%	28.1%
		Black	12 mo	0.3%	1.4%	2.3%	2.6%	3.8%
			24 mo	0.4%	2.5%	3.7%	5.7%	7.4%
			36 mo	0.5%	3.6%	4.9%	8.5%	10.5%
			48 mo	0.6%	4.6%	6.2%	12.1%	14.4%
			60 mo	0.7%	5.5%	7.1%	15.6%	18.1%
			72 mo	0.7%	6.5%	8.2%	19.0%	21.7%
			84 mo	0.7%	7.2%	9.1%	22.6%	25.5%
			96 mo	0.8%	7.7%	9.6%	26.2%	29.2%
			108 mo	0.8%	8.2%	10.2%	29.8%	33.0%
			120 mo	0.9%	8.9%	11.0%	34.2%	37.5%
75-79 years	Pre-PSA (1980-1989)	White	12 mo	0.2%	4.0%	4.7%	6.8%	7.7%
			24 mo	0.3%	8.5%	9.5%	12.4%	13.5%
			36 mo	0.3%	12.2%	13.4%	18.0%	19.3%
			48 mo	0.4%	15.3%	16.5%	23.6%	25.1%
			60 mo	0.4%	18.0%	19.3%	28.9%	30.5%
			72 mo	0.4%	20.1%	21.5%	33.4%	35.0%
			84 mo	0.4%	21.8%	23.3%	38.0%	39.6%
			96 mo	0.4%	23.3%	24.8%	42.1%	43.7%
			108 mo	0.4%	24.7%	26.2%	45.6%	47.3%
			120 mo	0.4%	25.8%	27.3%	49.2%	50.9%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				N	Median Obs	Observed	SE Obs	Obs Cum CIs Lower	Obs Cum CIs Upper	Specified COD	Other CODs	SE Specified COD
75-79 years	Pre-PSA (1980-1989)	Black	12 mo	1,380	43.41	81.4%	1.0%	79.3%	83.4%	7.5%	11.0%	0.7%
			24 mo	1,380	43.41	68.1%	1.3%	65.6%	70.5%	14.9%	17.0%	1.0%
			36 mo	1,380	43.41	56.9%	1.3%	54.2%	59.4%	19.8%	23.3%	1.1%
			48 mo	1,380	43.41	46.0%	1.3%	43.3%	48.6%	24.5%	29.5%	1.2%
			60 mo	1,380	43.41	38.3%	1.3%	35.7%	40.8%	27.3%	34.5%	1.2%
			72 mo	1,380	43.41	31.6%	1.3%	29.2%	34.1%	29.5%	39.0%	1.2%
			84 mo	1,380	43.41	25.6%	1.2%	23.3%	27.9%	31.3%	43.1%	1.2%
			96 mo	1,380	43.41	21.2%	1.1%	19.1%	23.4%	33.4%	45.4%	1.3%
			108 mo	1,380	43.41	17.4%	1.0%	15.4%	19.4%	34.7%	48.0%	1.3%
			120 mo	1,380	43.41	14.2%	0.9%	12.4%	16.1%	35.3%	50.5%	1.3%
	Post-PSA (2000-20 09)	White	12 mo	18,732	116.48	95.9%	0.1%	95.6%	96.2%	1.6%	2.5%	0.1%
			24 mo	18,732	116.48	91.1%	0.2%	90.7%	91.5%	3.1%	5.8%	0.1%
			36 mo	18,732	116.48	86.0%	0.3%	85.5%	86.5%	4.6%	9.4%	0.2%
			48 mo	18,732	116.48	81.0%	0.3%	80.5%	81.6%	5.8%	13.2%	0.2%
			60 mo	18,732	116.48	75.9%	0.3%	75.3%	76.5%	6.7%	17.4%	0.2%
			72 mo	18,732	116.48	70.7%	0.3%	70.1%	71.4%	7.6%	21.6%	0.2%
			84 mo	18,732	116.48	65.2%	0.3%	64.5%	65.9%	8.5%	26.3%	0.2%
			96 mo	18,732	116.48	59.6%	0.4%	58.9%	60.3%	9.3%	31.1%	0.2%
			108 mo	18,732	116.48	53.8%	0.4%	53.1%	54.5%	10.0%	36.2%	0.2%
			120 mo	18,732	116.48	48.2%	0.4%	47.5%	49.0%	10.6%	41.1%	0.2%
		Black	12 mo	2,188	103.24	93.2%	0.5%	92.0%	94.2%	2.7%	4.2%	0.3%
			24 mo	2,188	103.24	87.3%	0.7%	85.8%	88.6%	4.4%	8.3%	0.4%
			36 mo	2,188	103.24	81.4%	0.8%	79.7%	82.9%	6.0%	12.6%	0.5%
			48 mo	2,188	103.24	75.8%	0.9%	74.0%	77.6%	7.4%	16.8%	0.6%
			60 mo	2,188	103.24	70.9%	1.0%	69.0%	72.8%	8.4%	20.7%	0.6%
			72 mo	2,188	103.24	65.3%	1.0%	63.2%	67.2%	9.5%	25.2%	0.6%
			84 mo	2,188	103.24	59.9%	1.1%	57.8%	61.9%	10.7%	29.4%	0.7%
			96 mo	2,188	103.24	53.8%	1.1%	51.7%	55.9%	11.6%	34.6%	0.7%
			108 mo	2,188	103.24	47.9%	1.1%	45.8%	50.0%	12.7%	39.4%	0.7%
			120 mo	2,188	103.24	43.2%	1.1%	41.1%	45.4%	13.6%	43.2%	0.7%
80-84 years	Pre-PSA (1980-1989)	White	12 mo	8,753	43.54	83.1%	0.4%	82.3%	83.8%	5.8%	11.1%	0.3%
			24 mo	8,753	43.54	69.3%	0.5%	68.3%	70.2%	11.5%	19.3%	0.3%
			36 mo	8,753	43.54	56.8%	0.5%	55.7%	57.8%	15.9%	27.3%	0.4%
			48 mo	8,753	43.54	46.7%	0.5%	45.6%	47.7%	19.2%	34.1%	0.4%
			60 mo	8,753	43.54	38.1%	0.5%	37.1%	39.1%	21.9%	40.0%	0.4%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				SE Other CODs	Specified COD Cum CI Lower	Specified COD Cum CI Upper	Other CODs Cum CI Lower	Other CODs Cum CI Upper
75-79 years	Pre-PSA (1980-1989)	Black	12 mo	0.8%	6.2%	9.0%	9.4%	12.7%
			24 mo	1.0%	13.0%	16.8%	15.1%	19.1%
			36 mo	1.1%	17.7%	21.9%	21.2%	25.6%
			48 mo	1.2%	22.3%	26.8%	27.1%	31.9%
			60 mo	1.3%	24.9%	29.6%	32.0%	37.0%
			72 mo	1.3%	27.1%	31.9%	36.4%	41.5%
			84 mo	1.3%	28.9%	33.8%	40.5%	45.7%
			96 mo	1.3%	30.9%	35.9%	42.8%	48.0%
			108 mo	1.3%	32.2%	37.2%	45.3%	50.6%
			120 mo	1.3%	32.8%	37.9%	47.8%	53.1%
	Post-PSA (2000-20 09)	White	12 mo	0.1%	1.4%	1.8%	2.3%	2.8%
			24 mo	0.2%	2.9%	3.4%	5.5%	6.2%
			36 mo	0.2%	4.3%	4.9%	9.0%	9.8%
			48 mo	0.2%	5.4%	6.1%	12.7%	13.7%
			60 mo	0.3%	6.4%	7.1%	16.8%	17.9%
			72 mo	0.3%	7.3%	8.0%	21.1%	22.2%
			84 mo	0.3%	8.1%	8.9%	25.7%	27.0%
			96 mo	0.3%	8.9%	9.7%	30.4%	31.7%
			108 mo	0.4%	9.5%	10.4%	35.5%	36.9%
			120 mo	0.4%	10.2%	11.1%	40.4%	41.8%
		Black	12 mo	0.4%	2.0%	3.4%	3.4%	5.1%
			24 mo	0.6%	3.6%	5.4%	7.1%	9.5%
			36 mo	0.7%	5.1%	7.1%	11.3%	14.1%
			48 mo	0.8%	6.3%	8.5%	15.3%	18.4%
			60 mo	0.9%	7.3%	9.6%	19.0%	22.4%
			72 mo	0.9%	8.3%	10.8%	23.4%	27.1%
			84 mo	1.0%	9.4%	12.0%	27.5%	31.4%
			96 mo	1.0%	10.3%	13.0%	32.6%	36.6%
			108 mo	1.1%	11.3%	14.1%	37.3%	41.4%
			120 mo	1.1%	12.1%	15.0%	41.1%	45.3%
80-84 years	Pre-PSA (1980-1989)	White	12 mo	0.3%	5.4%	6.4%	10.4%	11.8%
			24 mo	0.4%	10.8%	12.1%	18.4%	20.1%
			36 mo	0.5%	15.1%	16.7%	26.4%	28.3%
			48 mo	0.5%	18.4%	20.0%	33.1%	35.1%
			60 mo	0.5%	21.1%	22.8%	39.0%	41.0%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				N	Median Obs	Observed	SE Obs	Obs Cum CIs Lower	Obs Cum CIs Upper	Specified COD	Other CODs	SE Specified COD
80-84 years	Pre-PSA (1980-1989)	White	72 mo	8,753	43.54	31.2%	0.5%	30.2%	32.1%	23.8%	45.0%	0.5%
			84 mo	8,753	43.54	24.9%	0.5%	24.0%	25.8%	25.3%	49.8%	0.5%
			96 mo	8,753	43.54	19.7%	0.4%	18.9%	20.6%	26.5%	53.8%	0.5%
			108 mo	8,753	43.54	14.8%	0.4%	14.1%	15.6%	27.5%	57.6%	0.5%
			120 mo	8,753	43.54	11.6%	0.3%	10.9%	12.2%	28.2%	60.3%	0.5%
		Black	12 mo	796	37.00	78.1%	1.5%	75.1%	80.9%	10.1%	11.8%	1.1%
			24 mo	796	37.00	63.2%	1.7%	59.7%	66.4%	17.0%	19.8%	1.3%
			36 mo	796	37.00	51.1%	1.8%	47.6%	54.5%	22.1%	26.8%	1.5%
			48 mo	796	37.00	41.8%	1.7%	38.4%	45.2%	25.5%	32.7%	1.5%
			60 mo	796	37.00	30.0%	1.6%	26.9%	33.2%	29.9%	40.1%	1.6%
			72 mo	796	37.00	24.4%	1.5%	21.4%	27.4%	31.8%	43.8%	1.7%
			84 mo	796	37.00	19.7%	1.4%	17.0%	22.6%	33.5%	46.7%	1.7%
			96 mo	796	37.00	14.6%	1.3%	12.2%	17.1%	35.1%	50.4%	1.7%
			108 mo	796	37.00	10.8%	1.1%	8.8%	13.1%	36.4%	52.8%	1.7%
			120 mo	796	37.00	7.3%	0.9%	5.6%	9.2%	37.2%	55.5%	1.7%
	Post-PSA (2000-2009)	White	12 mo	10,236	74.79	91.3%	0.3%	90.7%	91.8%	3.5%	5.2%	0.2%
			24 mo	10,236	74.79	83.2%	0.4%	82.5%	84.0%	6.8%	10.0%	0.2%
			36 mo	10,236	74.79	75.3%	0.4%	74.4%	76.1%	9.0%	15.7%	0.3%
			48 mo	10,236	74.79	67.1%	0.5%	66.1%	68.0%	11.0%	21.9%	0.3%
			60 mo	10,236	74.79	59.1%	0.5%	58.1%	60.0%	12.6%	28.3%	0.3%
			72 mo	10,236	74.79	51.7%	0.5%	50.7%	52.6%	14.1%	34.2%	0.3%
			84 mo	10,236	74.79	44.8%	0.5%	43.9%	45.8%	15.4%	39.8%	0.4%
			96 mo	10,236	74.79	38.1%	0.5%	37.1%	39.0%	16.4%	45.5%	0.4%
			108 mo	10,236	74.79	31.3%	0.5%	30.4%	32.2%	17.6%	51.2%	0.4%
			120 mo	10,236	74.79	25.2%	0.4%	24.4%	26.1%	18.5%	56.3%	0.4%
		Black	12 mo	1,082	61.69	85.4%	1.1%	83.1%	87.3%	6.3%	8.3%	0.7%
			24 mo	1,082	61.69	75.5%	1.3%	72.9%	78.0%	9.5%	14.9%	0.9%
			36 mo	1,082	61.69	65.6%	1.4%	62.7%	68.4%	12.2%	22.1%	1.0%
			48 mo	1,082	61.69	58.0%	1.5%	55.0%	60.9%	14.5%	27.4%	1.1%
			60 mo	1,082	61.69	50.7%	1.5%	47.7%	53.6%	15.9%	33.4%	1.1%
			72 mo	1,082	61.69	44.0%	1.5%	41.0%	46.9%	17.7%	38.3%	1.2%
			84 mo	1,082	61.69	37.5%	1.5%	34.6%	40.4%	19.1%	43.4%	1.2%
			96 mo	1,082	61.69	32.4%	1.4%	29.6%	35.2%	19.3%	48.3%	1.2%
			108 mo	1,082	61.69	26.6%	1.4%	23.9%	29.3%	19.9%	53.5%	1.2%
			120 mo	1,082	61.69	21.6%	1.3%	19.2%	24.2%	20.7%	57.7%	1.2%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.

Crude cause-specific, age-specific, 10-survival of PC among men aged 50-84

Cumulative Summary

				SE Other CODs	Specified COD Cum CI Lower	Specified COD Cum CI Upper	Other CODs Cum CI Lower	Other CODs Cum CI Upper
80-84 years	Pre-PSA (1980-1989)	White	72 mo	0.5%	22.9%	24.7%	44.0%	46.1%
			84 mo	0.5%	24.4%	26.2%	48.7%	50.8%
			96 mo	0.5%	25.6%	27.4%	52.7%	54.8%
			108 mo	0.5%	26.6%	28.5%	56.6%	58.7%
			120 mo	0.5%	27.2%	29.1%	59.2%	61.3%
		Black	12 mo	1.1%	8.1%	12.3%	9.7%	14.2%
			24 mo	1.4%	14.4%	19.7%	17.2%	22.7%
			36 mo	1.6%	19.3%	25.1%	23.7%	29.9%
			48 mo	1.7%	22.5%	28.6%	29.4%	35.9%
			60 mo	1.7%	26.8%	33.1%	36.7%	43.5%
			72 mo	1.8%	28.6%	35.0%	40.4%	47.3%
			84 mo	1.8%	30.3%	36.8%	43.2%	50.2%
			96 mo	1.8%	31.7%	38.4%	46.9%	53.8%
			108 mo	1.8%	33.1%	39.8%	49.2%	56.2%
			120 mo	1.8%	33.8%	40.5%	52.0%	58.9%
	Post-PSA (2000-2009)	White	12 mo	0.2%	3.2%	3.9%	4.8%	5.6%
			24 mo	0.3%	6.3%	7.3%	9.4%	10.6%
			36 mo	0.4%	8.5%	9.6%	15.0%	16.4%
			48 mo	0.4%	10.4%	11.6%	21.1%	22.7%
			60 mo	0.4%	12.0%	13.3%	27.4%	29.2%
			72 mo	0.5%	13.4%	14.8%	33.3%	35.2%
			84 mo	0.5%	14.7%	16.1%	38.8%	40.7%
			96 mo	0.5%	15.7%	17.2%	44.5%	46.5%
			108 mo	0.5%	16.8%	18.3%	50.2%	52.1%
			120 mo	0.5%	17.7%	19.2%	55.3%	57.3%
		Black	12 mo	0.8%	5.0%	7.8%	6.8%	10.1%
			24 mo	1.1%	7.9%	11.4%	12.9%	17.1%
			36 mo	1.3%	10.4%	14.3%	19.7%	24.7%
			48 mo	1.4%	12.5%	16.7%	24.8%	30.1%
			60 mo	1.4%	13.8%	18.2%	30.6%	36.2%
			72 mo	1.5%	15.5%	20.0%	35.4%	41.2%
			84 mo	1.5%	16.8%	21.5%	40.4%	46.4%
			96 mo	1.5%	17.0%	21.7%	45.3%	51.2%
			108 mo	1.5%	17.6%	22.3%	50.5%	56.5%
			120 mo	1.5%	18.3%	23.1%	54.6%	60.6%

Actuarial method.

Confidence interval: Log(-Log()) Transformation. The level is 95%.

+ The statistic could not be calculated.