

Least Squares and Quadratic Programming

Computational Intelligence, Lecture 5

by Sergei Savin

Fall 2020

- Some problems with exact analytical solutions
- Problems with inequality constraints
- Quadratic programming
- Homework

Some problems with exact analytical solutions

Problem 1. minimize $\|\mathbf{x}\|$. Solution $\mathbf{x} = \mathbf{0}$.

Problem 2. minimize $\|\mathbf{Ax}\|$. Solution $\mathbf{x} = \mathbf{0}$.

Problem 3. minimize $\|\mathbf{Ax} + \mathbf{b}\|$.

We observe that minimum of $\|\mathbf{Ax} + \mathbf{b}\|$ coincides with the minimum of $(\mathbf{Ax} + \mathbf{b})^2 = (\mathbf{Ax} + \mathbf{b})^\top (\mathbf{Ax} + \mathbf{b}) = \mathbf{b}^\top \mathbf{b} + \mathbf{b}^\top \mathbf{Ax} + \mathbf{x}^\top \mathbf{A}^\top \mathbf{b} + \mathbf{x}^\top \mathbf{A}^\top \mathbf{Ax}$, whose minimum coincides with the minimum of $\mathbf{b}^\top \mathbf{Ax} + \mathbf{x}^\top \mathbf{A}^\top \mathbf{b} + \mathbf{x}^\top \mathbf{A}^\top \mathbf{Ax}$. Since $\mathbf{b}^\top \mathbf{Ax}$ is a scalar, it is equal to its own transpose $(\mathbf{b}^\top \mathbf{Ax})^\top = \mathbf{x}^\top \mathbf{A}^\top \mathbf{b}$. Taking derivative with respect to \mathbf{x} we find gradient of the obtained *cost function*, and set it to zero, since that is the condition for the extreme point:

$$\frac{\partial}{\partial \mathbf{x}} (2\mathbf{b}^\top \mathbf{Ax} + \mathbf{x}^\top \mathbf{A}^\top \mathbf{Ax}) = 2\mathbf{b}^\top \mathbf{A} + 2\mathbf{x}^\top \mathbf{A}^\top \mathbf{A} = \mathbf{0} \quad (1)$$

Some problems with exact analytical solutions

Part 2

Thus we get expression: $2\mathbf{x}^\top \mathbf{A}^\top \mathbf{A} = -2\mathbf{b}^\top \mathbf{A}$, which we transpose and find value of \mathbf{x} :

$$\mathbf{x} = -(\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top \mathbf{b} \quad (2)$$

The shorthand for this formula is $\mathbf{x} = -\mathbf{A}^+ \mathbf{b}$, and it is called Moore-Penrose pseudoinverse.

Remember that a projector is often defined as $\mathbf{P} = \mathbf{A} \mathbf{A}^+$. Now we can provide an explicit formula for a projector:

$$\mathbf{P} = \mathbf{A}(\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top \quad (3)$$

However, neither (2) not (3) should be directly used in as algorithms, since usually there are better ways to compute those quantities (via SVD decomposition, for example).

Some problems with exact analytical solutions

Part 3

Problem 4.

$$\begin{aligned} & \underset{\mathbf{x}}{\text{minimize}} \quad \|\mathbf{x}\|, \\ & \text{subject to} \quad \mathbf{Ax} + \mathbf{By} = \mathbf{c}. \end{aligned} \tag{4}$$

We already solved this problem before, during lecture 2, using projection. Solution is: $\mathbf{x} = -((\mathbf{I} - \mathbf{BB}^+)\mathbf{A})^+(\mathbf{I} - \mathbf{BB}^+)\mathbf{c}$.

Problem 5.

$$\begin{aligned} & \underset{\mathbf{x}}{\text{minimize}} \quad \|\mathbf{x}\|, \\ & \text{subject to} \quad \mathbf{Ax} = \mathbf{c}. \end{aligned} \tag{5}$$

We can treat it as a special case of the previous problem, where $\mathbf{B} = \mathbf{0}$, hence $(\mathbf{I} - \mathbf{BB}^+) = \mathbf{I}$, and the solution is: $\mathbf{x} = -\mathbf{A}^+\mathbf{c}$.

Notice that solution for the problem 5 and problem 3 are written identically, even though problem 3 asks us to minimize residual of the linear system, while problem 5 - find minimum norm solution.

Some problems with exact analytical solutions

Part 4

Problem 6.

$$\begin{aligned} & \underset{\mathbf{x}}{\text{minimize}} && ||\mathbf{D}\mathbf{x}||, \\ & \text{subject to} && \mathbf{A}\mathbf{x} = \mathbf{b}. \end{aligned} \tag{6}$$

One way to think about it is to first find all solution to the constraint equation $\mathbf{A}\mathbf{x} = \mathbf{b}$ and then find optimal one among them. As we know, all solutions are given as: $\mathbf{x} = \mathbf{A}^+\mathbf{b} + \mathbf{N}\mathbf{z}$, where $\mathbf{N} = \text{null}(\mathbf{A})$. Then our cost function becomes: $||\mathbf{D}\mathbf{A}^+\mathbf{b} + \mathbf{D}\mathbf{N}\mathbf{z}||$, which is equivalent to the problem 3. Thus, we can write solution as: $\mathbf{z}^* = -(\mathbf{D}\mathbf{N})^+\mathbf{D}\mathbf{A}^+\mathbf{b}$. In terms of \mathbf{x} solution is:

$$\mathbf{x}^* = \mathbf{A}^+\mathbf{b} - \mathbf{N}(\mathbf{D}\mathbf{N})^+\mathbf{D}\mathbf{A}^+\mathbf{b} \tag{7}$$

Some problems with exact analytical solutions

Part 5

Problem 7.

$$\begin{aligned} & \underset{\mathbf{x}}{\text{minimize}} && \|\mathbf{D}\mathbf{x} + \mathbf{f}\|, \\ & \text{subject to} && \mathbf{A}\mathbf{x} = \mathbf{b}. \end{aligned} \tag{8}$$

After the same initial step, we arrive at the cost function $\|\mathbf{D}\mathbf{N}\mathbf{z} + \mathbf{D}\mathbf{A}^+\mathbf{b} + \mathbf{f}\|$. It is only different in the constant term, and the solution is found as follows:

$$\mathbf{z}^* = -(\mathbf{D}\mathbf{N})^+(\mathbf{D}\mathbf{A}^+\mathbf{b} + \mathbf{f}) \tag{9}$$

$$\mathbf{x}^* = \mathbf{A}^+\mathbf{b} - \mathbf{N}(\mathbf{D}\mathbf{N})^+(\mathbf{D}\mathbf{A}^+\mathbf{b} + \mathbf{f}) \tag{10}$$

Some problems with exact analytical solutions

Part 6

Problem 8.

$$\begin{aligned} & \underset{\mathbf{x}}{\text{minimize}} && \mathbf{x}^\top \mathbf{H} \mathbf{x} + \mathbf{c}^\top \mathbf{x}, \\ & \text{subject to} && \mathbf{A} \mathbf{x} = \mathbf{b}. \end{aligned} \tag{11}$$

where \mathbf{H} is positive-definite.

Assume that we found a decomposition $\mathbf{H} = \mathbf{D}^\top \mathbf{D}$. We can also find such \mathbf{f} that $2\mathbf{f}^\top \mathbf{D} = \mathbf{c}^\top$. Then our cost function becomes $\mathbf{x}^\top \mathbf{D}^\top \mathbf{D} \mathbf{x} + 2\mathbf{f}^\top \mathbf{D} \mathbf{x}$, which as we saw before has coinciding minimum with the cost function $\|\mathbf{D} \mathbf{x} + \mathbf{f}\|$.

Therefore the problem has the same solution as Problem 6, after the mentioned above change in constants.

Problems with inequality constraints

Problem 9.

$$\begin{array}{ll}\underset{\mathbf{x}}{\text{minimize}} & \|\mathbf{D}\mathbf{x} + \mathbf{f}\|, \\ \text{subject to} & \mathbf{x} \leq \mathbf{b}.\end{array}\tag{12}$$

Problem 10.

$$\begin{array}{ll}\underset{\mathbf{x}}{\text{minimize}} & \|\mathbf{x}\|, \\ \text{subject to} & \mathbf{A}\mathbf{x} \leq \mathbf{b}.\end{array}\tag{13}$$

Problem 11.

$$\begin{array}{ll}\underset{\mathbf{x}}{\text{minimize}} & \|\mathbf{D}\mathbf{x} + \mathbf{f}\|, \\ \text{subject to} & \begin{cases} \mathbf{A}\mathbf{x} \leq \mathbf{b}, \\ \mathbf{C}\mathbf{x} = \mathbf{d}. \end{cases}\end{array}\tag{14}$$

Quadratic programming

Mentioned problems can be described together as quadratic programs. The name is due to the cost function being quadratic (or equivalent). They are allowed to have linear equality or inequality constraints.

General form of a quadratic program is given below:

$$\begin{aligned} & \underset{\mathbf{x}}{\text{minimize}} && \mathbf{x}^\top \mathbf{H} \mathbf{x} + \mathbf{f}^\top \mathbf{x}, \\ & \text{subject to} && \begin{cases} \mathbf{A} \mathbf{x} \leq \mathbf{b}, \\ \mathbf{C} \mathbf{x} = \mathbf{d}. \end{cases} \end{aligned} \tag{15}$$

where \mathbf{H} is positive-definite and $\mathbf{A} \mathbf{x} \leq \mathbf{b}$ describe a *convex region*.

Homework

- Solve all examples of problems with and without inequalities using `quadprog` function in the language of your choice.

Lecture slides are available via Moodle.

You can help improve these slides at:

github.com/SergeiSa/Computational-Intelligence-Slides-Fall-2020

Check Moodle for additional links, videos, textbook suggestions.