

# Coral Reef Monitoring Program — 2025 Field Season Results

*Prepared by Institute of Marine Science*

## Methods

Restoration nurseries propagated 18,000 coral fragments across the study sites. Monitoring transects were surveyed twice yearly at fixed depth contours. Restoration nurseries propagated 18,000 coral fragments across the study sites. Ocean acidification reduces calcification rates in most reef-building species studied. Herbivorous fish control macroalgae that would otherwise outcompete coral recruits.

Larval connectivity models suggest recovery depends on upstream source reefs remaining healthy. Reef fish biomass on protected reefs was 2.4 times higher than on adjacent fished reefs. Restoration nurseries propagated 18,000 coral fragments across the study sites. Restoration nurseries propagated 18,000 coral fragments across the study sites. Reef fish biomass on protected reefs was 2.4 times higher than on adjacent fished reefs.

## Results

Bleaching occurs when thermal stress causes corals to expel their symbiotic algae. Bleaching occurs when thermal stress causes corals to expel their symbiotic algae. Monitoring transects were surveyed twice yearly at fixed depth contours. Herbivorous fish control macroalgae that would otherwise outcompete coral recruits. Herbivorous fish control macroalgae that would otherwise outcompete coral recruits.

Bleaching occurs when thermal stress causes corals to expel their symbiotic algae. Restoration nurseries propagated 18,000 coral fragments across the study sites. Herbivorous fish control macroalgae that would otherwise outcompete coral recruits. Reef fish biomass on protected reefs was 2.4 times higher than on adjacent fished reefs. Ocean acidification reduces calcification rates in most reef-building species studied.

Monitoring transects were surveyed twice yearly at fixed depth contours. Reef fish biomass on protected reefs was 2.4 times higher than on adjacent fished reefs. Monitoring transects were surveyed twice yearly at fixed depth contours. Monitoring transects were surveyed twice yearly at fixed depth contours. Reef fish biomass on protected reefs was 2.4 times higher than on adjacent fished reefs.

Bleaching occurs when thermal stress causes corals to expel their symbiotic algae. Larval connectivity models suggest recovery depends on upstream source reefs remaining healthy. Reef fish biomass on protected reefs was 2.4 times higher than on adjacent fished reefs. Larval connectivity models suggest recovery depends on upstream source reefs remaining healthy. Restoration nurseries propagated 18,000 coral fragments across the study sites.

## Restoration Activities

Reef fish biomass on protected reefs was 2.4 times higher than on adjacent fished reefs. Monitoring transects were surveyed twice yearly at fixed depth contours. Restoration nurseries propagated 18,000 coral fragments across the study sites. Larval connectivity models suggest recovery depends on upstream source reefs remaining healthy. Larval connectivity models suggest recovery depends on upstream source reefs remaining healthy.

Restoration nurseries propagated 18,000 coral fragments across the study sites. Ocean acidification reduces calcification rates in most reef-building species studied. Monitoring transects were surveyed twice yearly at fixed depth contours. Reef fish biomass on protected reefs was 2.4 times higher than on adjacent fished reefs. Bleaching occurs when thermal stress causes corals to expel their symbiotic algae.