

HackerRank







40/115 challenges solved Rank: **74552** | Points: **545** (!)



X Your Introduction to Sets submission got 10.00 points. Try the next challenge | Try a Random Challenge Problem Leaderboard Editorial A Submissions A set is an unordered collection of elements without duplicate entries.

When printed, iterated or converted into a sequence, its elements will appear in an arbitrary order.

Example

```
>>> print set()
set([])
>>> print set('HackerRank')
set(['a', 'c', 'e', 'H', 'k', 'n', 'r', 'R'])
>>> print set([1,2,1,2,3,4,5,6,0,9,12,22,3])
set([0, 1, 2, 3, 4, 5, 6, 9, 12, 22])
>>> print set((1,2,3,4,5,5))
set([1, 2, 3, 4, 5])
>>> print set(set(['H','a','c','k','e','r','r','a','n','k']))
set(['a', 'c', 'r', 'e', 'H', 'k', 'n'])
>>> print set({'Hacker' : 'DOSHI', 'Rank' : 616 })
set(['Hacker', 'Rank'])
>>> print set(enumerate(['H','a','c','k','e','r','r','a','n','k']))
set([(6, 'r'), (7, 'a'), (3, 'k'), (4, 'e'), (5, 'r'), (9, 'k'), (2, 'c'), (0, 'H'), (1, 'a'), (8, 'n')])
```

Basically, sets are used for membership testing and eliminating duplicate entries.

Task

Now, let's use our knowledge of sets and help Mickey.

Ms. Gabriel Williams is a botany professor at District College. One day, she asked her student Mickey to compute the average of all the plants with distinct heights in her greenhouse.

Formula used:

$$Average = rac{Sum \ of \ Distinct \ Heights}{Total \ Number \ of \ Distinct \ Heights}$$

Function Description

Complete the average function in the editor below.

average has the following parameters:


• int arr: an array of integers


Returns

• float: the resulting float value rounded to 3 places after the decimal

Input Format

```
The first line contains the integer, N, the size of arr.
The second line contains the m{N} space-separated integers, m{arr}[i] .
Constraints
0 < N \le 100
Sample Input
   STDIN
                                                        Function
   10
                                                        arr[] size N = 10
   161 182 161 154 176 170 167 171 170 174
                                                        arr = [161, 181, ..., 174]
Sample Output
   169.375
Explanation
Here, set([154, 161, 167, 170, 171, 174, 176, 182]) is the set containing the distinct heights. Using the sum() and len() functions, we can compute the average.
                                                           Average = \frac{1355}{8} = 169.375
```





Contest Calendar | Blog | Scoring | Environment | FAQ | About Us | Support | Careers | Terms Of Service | Privacy Policy | Request a Feature