



Set .difference() Operation ★

36/115 challenges solved

Rank: 84185 | Points: 505



Your Set .difference() Operation submission got 10.00 points.

مشاركة

Tweet


[Try the next challenge](#) | [Try a Random Challenge](#)

Problem

Submissions

Leaderboard

Editorial

**A.difference(B) or A - B**

By DOSHI

.difference()

The tool .difference() returns a set with all the elements from the set that are not in an iterable.

Sometimes the - operator is used in place of the .difference() tool, but it only operates on the set of elements in set.

Set is immutable to the .difference() operation (or the - operation).

```
>>> s = set("Hacker")
>>> print s.difference("Rank")
set(['c', 'r', 'e', 'H'])

>>> print s.difference(set(['R', 'a', 'n', 'k']))
set(['c', 'r', 'e', 'H'])

>>> print s.difference(['R', 'a', 'n', 'k'])
set(['c', 'r', 'e', 'H'])

>>> print s.difference(enumerate(['R', 'a', 'n', 'k']))
set(['a', 'c', 'r', 'e', 'H', 'k'])

>>> print s.difference({"Rank":1})
set(['a', 'c', 'e', 'H', 'k', 'r'])

>>> s - set("Rank")
set(['H', 'c', 'r', 'e'])
```

Task



Students of District College have a subscription to English and French newspapers. Some students have subscribed to only the English newspaper, some have subscribed to only the French newspaper, and some have subscribed to both newspapers.

You are given two sets of student roll numbers. One set has subscribed to the English newspaper, and one set has subscribed to the French newspaper. Your task is to find the total number of students who have subscribed to only English newspapers.

Input Format

The first line contains the number of students who have subscribed to the English newspaper.

The second line contains the space separated list of student roll numbers who have subscribed to the English newspaper.

The third line contains the number of students who have subscribed to the French newspaper.

The fourth line contains the space separated list of student roll numbers who have subscribed to the French newspaper.

Constraints

$0 < \text{Total number of students in college} < 1000$

Output Format

Output the total number of students who are subscribed to the English newspaper only.

Sample Input

```
9
1 2 3 4 5 6 7 8 9
9
10 1 2 3 11 21 55 6 8
```

Sample Output

```
4
```

Explanation

The roll numbers of students who only have English newspaper subscriptions are:

4, 5, 7 and 9.

Hence, the total is **4** students.

[Change Theme](#)

Language

Python 3



```
1 n=int(input())
2 l1=set(map(int,input().split()))
3 m=int(input())
4 l2=set(map(int,input().split()))
5 print(len(l1.difference(l2)))
```

Line: 5 Col: 30

Upload Code as File

☐ Test against custom input


Run Code

Submit Code

You have earned 10.00 points!

36/115 challenges solved.




31%



Congratulations

You solved this challenge. Would you like to challenge your friends?

Next Challenge



Earn a certificate in Python

Kudos on your progress! Take the HackerRank Skills Certification test and enrich your profile

Get Certified

Test case 0

Test case 1

Test case 2

Test case 3

Test case 4

Test case 5

Compiler Message

Success

Input (stdin)

Download

19

21 2 3 4 5 6 7 8 9

39

410 1 2 3 11 21 55 6 8

Expected Output

Download

14

Contest Calendar

Blog

Scoring

Environment

FAQ

About Us

Support

Careers

Terms Of Service

Privacy Policy

Request a Feature

https://www.hackerrank.com/challenges/py-set-difference-operation/problem?utm_campaign=challenge-recommendation&utm_medium=email&utm_s... 3/3