

From the Sun to Earth

Exploring the Multifacets of Solar

Eruptive Events

Mohamed Nedal

A thesis submitted for the degree of
Doctor of Philosophy

Institute of Astronomy and National Astronomical Observatory,
Bulgarian Academy of Sciences
Solar and Space Weather Research
January 22, 2024

Supervisor: Assoc. Prof. Kamen Asenov Kozarev

Acknowledgments

To my family, professors, and friends ...

Contents

List of Tables

List of Figures

Chapter 1

Introduction

1.1 Background and Motivation

The Sun, an ordinary main-sequence star situated at the center of our Solar System, exhibits various forms of activity and variability on multiple spatial and temporal scales. One of the main manifestations of solar activity relevant to space weather research are transient energetic eruptive phenomena such as flares, Coronal Mass Ejections (CME), and wide-ranging emissions of electromagnetic radiation and energetic particles (??). These eruptive events originate due to the sudden release of free magnetic energy stored in complex, twisted or sheared magnetic field structures in the solar atmosphere (????). The energetic phenomena are driven by the rapid dissipation of magnetic energy via magnetic reconnection which can accelerate large numbers of electrons to relativistic energies and heat plasma to tens of million Kelvin (??).

The eruptive solar events drive major disturbances in the near-Earth space environment and planetary environments across the heliosphere, collectively termed space weather (??). Enhanced fluxes of solar energetic particles (SEPs), plasma ejecta, and electromagnetic radiation emitted during solar eruptions can impact the geomagnetic field, radiation belts, ionosphere, thermosphere, and upper atmosphere surrounding the Earth (??). Adverse effects range from disruption of radio communications to damage of satellites, power grid failures, aviation hazards due to radiation risks for airline crew and passengers, and increased radiation exposure for astronauts (?). The societal dependence on space-based infrastructure has increased exponentially, escalating the vulnerability to space weather disturbances. Recent studies estimate a severe space weather event could lead to trillion-dollar economic damages in the US alone (?). Besides the near-Earth space environment, solar eruptive transients also drive adverse space weather effects across the Solar System impacting activities such as deep space exploration and astronomy (?).

Therefore, advancing our understanding of the origins and propagation characteristics of solar eruptive phenomena, as well as quantifying their impacts on geospace and planetary environments, has become an extremely important pursuit for nations worldwide. Fundamental research seeks to uncover the physical processes involved using observations coupled with theory and modeling. Concurrently, significant efforts are underway to develop next-generation space environment modeling and forecasting capabilities for predicting the impacts of solar variability. The field combining these research and predictive aspects related to Sun-Earth connections is broadly termed heliophysics (?). It encompasses understanding the fundamental solar, heliospheric and geospace plasma processes; coupling across multiple spatial and temporal scales; quantifying the impacts on humanity's technological systems and space-borne assets; and utilizing this knowledge to prevent/mitigate adverse effects (??). NASA's Living With a Star program and the National Science Foundation's Space Weather activities exemplify strategic efforts to advance scientific understanding and predictive capabilities across the interconnected domains of heliophysics (?).

The present thesis focuses on studying several important phenomena related to solar eruptive activity and its impacts from the perspective of heliophysics research and space weather. The specific topics investigated include: (1) The propagation and evolution characteristics of large-scale coronal disturbances termed EUV waves that are triggered by solar flares and CMEs; (2) The generation, propagation and plasma characteristics of solar radio bursts emitted by accelerated electron beams traveling along open magnetic field lines in the corona; (3) The forecasting of gradual solar energetic particle (SEP) events which constitute one of the major components of space radiation hazards at Earth.

These diverse topics are united by the common theme of seeking to uncover the origins and propagation mechanisms of key transient phenomena resulting from solar eruptions, utilizing observational data,

analytical theory and modeling, and data science techniques. The phenomena have been studied for several decades using observations from multiple space missions, but gaps persist in our understanding of their underlying physics and space weather impacts. The thesis aims to provide new insights that help address some of the outstanding questions, guided by the overarching goals and framework of heliophysics research. The following sub-sections elaborate on the background, significance, observational challenges and knowledge gaps pertaining to each of the research topics investigated. The following subsection provide a concise overview of key literature related to the research topics investigated in the thesis. A detailed review is presented in each chapter specific to the respective phenomenon.

1.1.1 Coronal Waves

Coronal waves, or Coronal Bright Fronts (CBFs), also known as Extreme Ultra-Violet (EUV) waves, are large-scale arc-shaped bright fronts or disturbances observed propagating across significant portions of the solar corona following the eruption of CMEs and flares (??????). They are best observed in EUV and white-light coronal emission, as well as in radio wavelengths, spanning distances of up to several 100 Mm with speeds ranging from 100-1000 $km\ s^{-1}$, faster than the local characteristic speed in the solar corona, transforming into shock waves (????). These structures consist of piled-up plasma with higher density, making them appear brighter in white-light images.

The discovery of coronal waves dates back to observations obtained with the EIT instrument on SOHO launched in 1995, appearing as bright propagating fronts in 19.5 nm wavelength imaging of Fe XII emission lines formed at ~ 1.5 MK plasma (?). Subsequent studies based on SOHO/EIT and TRACE imaging found correlations between waves and CMEs, favoring an interpretation as fast-mode MHD waves driven by CME lateral expansions (?).

Since 2010, the initiation and evolution of coronal waves are being exquisitely observed with unprecedented resolution by the Atmospheric Imaging Assembly (AIA) on the Solar Dynamics Observatory (SDO) instrument (?) across multiple EUV passbands sensitive to a wide temperature range (?). Alternatively, shock waves can be indirectly observed through the detection of type II radio bursts, which are commonly associated with shock waves in the solar corona ?. The AIA instrument has provided valuable insights into the dynamics of the low solar corona over the past decade, thanks to its exceptional spatial and temporal resolution. Equipped with telescopes observing the solar disk in bands 193 and 211 Å, the AIA instrument has demonstrated its ability to distinguish compressive waves in the lower corona ??. These observations offer valuable information about the kinematics and geometric structure of CBFs. To accurately study the evolution of the wave's leading front, observations off the solar limb are preferred to mitigate projection effects, which may introduce ambiguities in estimating time-dependent positions and the global structure of the wave ?.

In situ observations of shock waves have revealed their classification into quasi-parallel, quasi-perpendicular, sub-critical, and super-critical shocks based on the angle between the wavefront normal vector and the upstream magnetic field lines ?. Quasi-parallel shocks have an angle (θ_{BN}) smaller than 45° , while quasi-perpendicular shocks have θ_{BN} greater than 45° . Supercritical shocks, often associated with accelerated particles, are promising candidates for generating type II radio bursts ?. However, obtaining accurate estimates of shock strength and obliquity solely from remote observations is challenging.

Coronal waves exhibit diverse morphology and kinematics ranging from circular fronts to narrow jets or expanding dome-like structures (?). A taxonomy of wave properties based on extensive observational surveys can be found in papers by ? and ?. However, despite being observed for over two decades since their serendipitous discovery, fundamental questions remain regarding the physical nature and drivers of coronal waves (??). The debate centers around two competing interpretations - the wave versus pseudo-wave (or non-wave) models. The wave models envisage coronal waves as fast-mode MHD waves or shocks that propagate freely after being launched by a CME lateral over-expansion or an initial flare pressure pulse (??). The pseudo-wave models interpret them as bright fronts produced by magnetic field restructuring related to the CME lift-off process rather than a true wave disturbance (??). Extensive observational and modeling studies have been undertaken to evaluate the two paradigms (??), but a consensus remains elusive. Addressing these outstanding questions related to the nature and origin of coronal waves is imperative, since they are being incorporated into models as a primary agent producing SEP events and geomagnetic storms during CMEs (??). Their use as a diagnostic tool for CME and shock kinematics predictions in these models requires discriminating between the different physical mechanisms proposed for their origin.

The present thesis undertakes an extensive statistical analysis of coronal EUV wave events observed by SDO to provide new insights into their kinematical properties and relationship to CMEs. We focus

on analyzing their large-scale evolution as a function of distance and direction from the source region, leveraging the extensive EUV full-disk imaging capabilities of SDO spanning nearly a decade. Statistical surveys to date have mostly focused on initial speeds and morphological classifications rather than large-scale propagation characteristics. Our study aims to uncover systematic trends in their propagation kinematics using a significantly larger sample compared to previous works. We also comprehensively evaluate associations with CME and flare parameters in order to discriminate between wave and pseudo-wave origins. The results have important implications for incorporating coronal waves into predictive models of CMEs and SEP events for future space weather forecasting.

1.1.2 Solar Radio Bursts

Solar radio emissions have been the subject of extensive study and research due to their connection with solar activity and their potential impact on Earth's atmosphere and technology. One area of particular interest is solar radio bursts, which are intense bursts of electromagnetic radiation originating from the Sun. These bursts can be classified into different types based on their characteristics and associated phenomena. Solar radio bursts, including Type III bursts, serve as remote diagnostics for the study of energetic electrons within the solar corona. These bursts result from transient energetic electron beams injected into the corona, which then propagate along interplanetary magnetic field (IMF) lines (??). As these electron beams traverse the corona, they induce plasma waves, also known as Langmuir waves, which subsequently transform into radio emission at the local plasma frequency or its harmonic components (?). The frequency of the radio emission is directly linked to the plasma density, making Type III bursts a valuable tool for investigating the inner heliosphere and understanding the underlying processes that drive solar active phenomena, such as solar flares and coronal mass ejections (??). These bursts offer insights into the acceleration of energetic electrons in the corona and their transport along magnetic field lines (?). The generation of electromagnetic emission at radio frequencies through plasma emission mechanisms is a key aspect of solar radio bursts, shedding light on the dynamic interplay between non-thermal electron distributions and the ambient plasma (?).

In radio spectrograms, type III bursts manifest as intense enhancements of radio flux over background levels, exhibiting rapid frequency drifts over timescales ranging from seconds to minutes, signifying plasma dynamics (?). These bursts are observable across a wide range of frequencies, spanning from GHz to kHz, and wavelengths extending from metric to decametric (??). This phenomenon is detectable by ground-based instruments on Earth as well as various spacecraft within the heliosphere, underscoring the significance of plasma dynamics in their manifestation.

Pioneering observations of solar radio bursts were made in the 1940s leading to their classifications (?). Subsequent spectrographic studies uncovered emission mechanisms, source regions and particle diagnostics (?). Magnetic reconnection models of flares provided theoretical explanations for particle acceleration generating radio bursts (?). Radio imaging spectroscopy using interferometric imaging arrays coupled with high time/frequency resolution spectrometers enables tracking radio sources as a function of frequency and position on the Sun, yielding particle acceleration locations and trajectories through the corona into interplanetary space (??). This provides a unique diagnostic of energetic particle transport from the Sun to the Earth which is crucial for improving SEP forecasting models.

Different types of bursts are observed, classified based on their spectral characteristics as documented in radio burst catalogs (?). The present thesis focuses on detailed analysis of solar type III radio bursts and their associated phenomena (?). Type III bursts appear as intense rapidly drifting emissions from high to low frequencies over seconds, corresponding to the propagation of energetic electron beams from the low corona to beyond 1 AU along open field lines. They signify the initial escape of flare-accelerated electrons into interplanetary space, making them an important precursor signature of SEP activity (??). Investigating their source locations, plasma environments, and beam kinematics based on multi-wavelength observations coupled with plasma emission theory is therefore vital for improved understanding of coronal particle acceleration and transport processes relevant for SEP forecasting models.

While type III bursts have been studied for over 50 years since their initial discovery by ???, gaps persist in our understanding of their exciter beams and emission mechanisms. Key outstanding questions pertain to the detailed electron acceleration and injection sites, beam configurations and energy spectra, drivers of burst onset and duration, and the role of density fluctuations in propagating beams (??). Recent work combines imaging and spectral data with modeling to constrain radio burst excitors in unprecedented detail (??). Key challenges remain in reconciling emission models with observations and predicting radio diagnostics. Advancing our knowledge of these aspects through coordinated observations and modeling can help constrain the predictions of energetic electron properties based on

radio diagnostics. The present work undertakes detailed investigation of a solar type III burst combining imaging and radio spectral data to derive electron beam trajectories and coronal densities, and models the emission sources. The results provide insights into the corona plasma environment and energetic electron transport relevant for SEP forecasting applications.

1.1.3 Solar Energetic Particle (SEP) Forecasting

Solar Energetic Protons (SEP) are high-energy particles that are believed to be originated from the acceleration of particles in the solar corona during coronal mass ejections (CMEs) and solar flares (?????). They are typically characterized by their high energy levels - with some particles having energies in the relativistic GeV/nucleon range - and their ability to penetrate through spacecraft shielding, causing radiation damage (??). The fluence and energy spectrum of SEP are influenced by several factors, including the strength of the solar flare or CME that produced them, and the conditions of the interplanetary environment (????????).

SEP exhibit a strong association with the solar cycle, with the frequency and flux of SEP events peaking during the maximum phase of the solar cycle (?). This is thought to be due to the increased activity of the Sun during this phase, which leads to more frequent and powerful flares and CMEs. Previous studies have shown a relationship between the occurrence frequency of SEP and the sunspot number (SN; ?, ?, ?, ?, ?). However, the exact relationship between the solar cycle and SEP is complex and not fully understood. Hence, more work is needed to better understand this connection, as previous studies have reported intense SEP events during relatively weak solar activity (??).

SEP have been a subject of interest and research in heliophysics for decades. It is hypothesized that shock waves generated in the corona can lead to an early acceleration of particles. However, SEP have sufficient energy to propagate themselves by *surfing* the interplanetary magnetic fields (IMF), and therefore, the expanding CME is not necessary for their transport (????). While this theory has gained acceptance, there is an ongoing debate among scientists over the specific mechanisms and conditions responsible for SEP production and acceleration.

The creation, acceleration, and transport mechanisms of SEP are complex and involve a combination of magnetic reconnection, shock acceleration, and wave-particle interactions (???). The specific mechanisms responsible for SEP production and acceleration can vary depending on the type and strength of the solar event that triggered them. Further research is imperative to better understand the processes involved in the production and transport of SEP in the heliosphere. This will facilitate the development of more precise models that assist in minimizing the impact of SEP on astronauts and space-based assets.

The arrival of solar energetic particles (SEPs) in the near-Earth space environment constitutes one of the major components of adverse space weather (??). SEPs consist primarily of protons (and some heavy ions), accelerated to very high energies by CME-driven shock waves during large solar eruptive events. The gradual SEP events, so called due to their long durations from several hours to a few days, involve protons accelerated to energies above ~ 10 MeV which can penetrate Earth's magnetic field and atmosphere posing radiation hazards to humans and equipment in space and at polar regions (?).

Initial SEP forecasting models were based on empirical correlations between proton intensity profiles and CME or flare properties (?). The complex physics of CME shock acceleration combined with modeling the transport of SEPs through turbulent interplanetary magnetic fields presents major challenges for first-principles based SEP forecasting models (??). As an alternative approach, empirical and data-driven models based on statistical/machine learning techniques applied to historical SEP event data have shown considerable promise for operational forecasting over the past decade (???). This motivates detailed investigation of data-driven SEP forecasting models using state-of-the-art machine learning algorithms which can outperform conventional empirical methods.

The emergence of deep learning techniques has enabled application of sophisticated machine learning models to SEP forecasting, yielding improved predictions since they can capture complex nonlinear relationships between parameters which has been leveraged for diverse space weather applications recently (??). Opportunities exist for novel forecasting approaches utilizing deep learning algorithms and expanded input parameters. However, applications to SEP forecasting problems are still limited, presenting an important research gap which this thesis aims to address. In the present work, we develop a deep neural network model for predicting the intensity profile of >10 MeV gradual SEP proton events utilizing near real-time solar wind plasma measurements as model inputs. The developed model is trained and tested on a database of historical SEP events spanning solar cycles 23 and 24, with the goal of producing SEP flux forecasts over an hour in advance of particle arrivals near Earth. Such capability can provide actionable information for mitigating radiation effects from extreme SEP events. The study demonstrates

the potential of state-of-the-art machine learning algorithms to achieve significant enhancement of SEP forecasting capabilities building upon conventional empirical methods.

1.2 Objectives and Scope

The primary objectives and research questions addressed through the investigations carried out in this thesis include:

1. Characterize the large-scale propagation kinematics of coronal EUV waves over distances of hundreds of Mm from the eruption source location. Compare observed spatial and temporal variations in speeds with analytical CME-driven wave/shock models.
2. Conduct a comprehensive statistical analysis correlating properties of EUV waves with associated CME and flare parameters utilizing a large event sample. Discriminate between wave and pseudo-wave models based on observational evidence.
3. Analyze coordinated observations of a solar type III radio burst across imaging and radio spectral domains to derive coronal density profiles, electron beam kinematics and emission source models.
4. Develop a deep neural network model for forecasting the intensity profile of >10 MeV gradual SEP proton events using real-time solar wind data as inputs. Evaluate model performance and forecast accuracy over different lead times.

The scope of the thesis encompasses key phenomena related to solar eruptions and their space weather impacts that align with the outstanding questions and challenges highlighted in the background discussion. While expansive in scope, some limitations exist that bound the present work:

- The studies rely primarily on remote sensing observations of the Sun and heliosphere, limited by measurement capabilities and resolution.
- Analytical modeling utilizes simplified theory and assumptions which cannot account for all complexities.
- Machine learning models have dependencies on data coverage and uncertainties in input parameters.
- Findings are constrained by the event samples studied and applicability to the broader population.

These factors imply appropriate care and diligence in interpretation of results and their generalizability. Nevertheless, the present work establishes an important foundation for future advances that can build upon these limitations.

1.3 Methodology Overview

The research presented in this thesis employs a synergistic methodology combining analytical theory, numerical modeling, and data science techniques. Both observational case studies and statistical analysis approaches are utilized for gaining new insights from application of these tools. The data sources, models, and algorithms employed in each of the investigations are concisely summarized below.

Coronal waves: The study utilizes an event database of ~ 60 coronal EUV waves observed by SDO/AIA since 2010, tracking kinematics to $30 R_{\odot}$ distances. Evolution trends are compared with analytical CME-driven wave propagation models. Statistical associations with CME and flare parameters provide corroboration for physical interpretation.

Solar radio bursts: Multi-wavelength observations of type III bursts from the LOFAR stations and Parker Solar Probe (PSP) are analyzed. Beam trajectories, densities, and emission sources are characterized by combining imaging data, plasma emission theory and coronal density models.

SEP forecasting: A database of >10 , >30 , and >60 MeV SEP integral fluxes during the last four solar cycles is generated using the GOES database. Deep neural network models are developed using solar wind data time-series as inputs. Model training, testing and validation is performed to evaluate forecast accuracy over different lead times.

This triangulation between data analysis, physics-based modeling and data-driven modeling provides confidence in the results obtained. Details of the methodological approaches are elaborated in their respective chapters.

1.4 Main Contributions

The primary contributions arising from the research presented in this thesis include:

- New large-scale kinematical characterization of coronal EUV waves propagating to distances over 100 Mm. Derived velocity and acceleration trends challenge steady-wave behavior assumed in models.
- Statistical analysis correlating EUV wave and CME/flare properties using a significantly larger event sample compared to prior studies. This enables stronger discrimination between competing initiation models.
- Novel methodology combining radio and EUV observations with analytical modeling to reconstruct plasma environments and electron beam trajectories for a solar type III radio burst.
- Deep learning forecasting model for intense SEP events using an expanded input parameter space based on solar wind data. This demonstrates cutting-edge artificial intelligence capabilities for space weather applications.
- Synergistic approach leveraging analytical theory, numerical modeling and data science techniques to gain new insights on long-standing problems in heliophysics research related to solar eruptions and their space weather impacts.

These contributions provide advances over prior state-of-the-art in the respective areas. They have implications for improving models used in operational space weather monitoring and forecasting systems, besides progressing fundamental physics understanding of solar and heliospheric phenomena. The results validate the merit of cross-disciplinary studies combining traditional analytical techniques with modern statistical and machine learning methods to enable discoveries from application of these synergies.

1.5 Outline

This thesis is divided into the following five chapters:

Chapter 1 - Introduction: Provides a background to the research topics, motivation and context of the work, summary of literature, overview of methodology, and the structure of the thesis.

Chapter 2 – Propagation and Drivers of Coronal EUV Waves: Presents a statistical analysis of the kinematics and physical interpretation of coronal waves using EUV imaging observations and analytical models.

Chapter 3 – Plasma Environment and Energetics of a Solar Type III Radio Burst: Details a multi-wavelength observational case study of a type III burst combining data analysis and modeling to probe the radio emission physics.

Chapter 4 – Deep Learning Approach for Forecasting Intense SEP Events: Describes the development and evaluation of a neural network model for predicting SEP properties using solar wind data.

Chapter 5 – Conclusions and Future Outlook: Summarizes the key findings, implications, and limitations of the research studies. Discusses future extensions building on the present work.

The core chapters 2 through 4 present the major research investigations carried out. The multi-faceted phenomena are studied by tailoring the methodology to leverage their key observational signatures. Together they provide new insights on different aspects of solar eruptions and space weather. Each chapter is structured to be reasonably self-contained, with relevant background and literature specific to the phenomenon under study. The findings are synergistic and united by the common thread of employing heliophysics principles to address outstanding questions using cutting-edge analytics.

Chapter 2

Characterization of Coronal Bright Fronts

2.1 Introduction

2.2 Observations

2.3 Data Analysis and Methods

2.4 Results and Discussion

2.5 Conclusions

2.6 Wavetrack: A Flexible Framework for Automated Recognition and Tracking of Wave-Like Events in Solar Imagery

2.6.1 Overview

Solar eruptive events encompass a myriad of intricate phenomena, including flares, filament eruptions, coronal mass ejections (CMEs), and CME-driven shock waves. The principal drivers of solar energetic particles (SEPs) are acknowledged to be CME-driven shocks within the corona and interplanetary space. This acceleration primarily occurs through the diffusive shock acceleration (DSA) and shock drift acceleration (SDA) processes (?). Efforts have been devoted to characterizing and modeling SEP acceleration under ideal conditions (???). Recent endeavors have shifted towards the development of more realistic CME-shock and SEP acceleration models, incorporating observational data (????).

Understanding the efficiency and spatial distribution of SEP acceleration necessitates a comprehensive grasp of CME-shock interactions with three-dimensional coronal fields ?. The modeling of DSA requires the deduction of shock front shapes from observations, given the profound impact of the local magnetic field-shock angle on acceleration ?. Moreover, the lateral over-expansion of CMEs early in their evolution modifies the compressive front ??, necessitating improvements to the idealized shock surface descriptions used in modeling (???).

The utilization of extreme ultraviolet (EUV) imaging, facilitated by instruments such as STEREO/EUVI (?) and SDO/AIA (?), enables the characterization of shocks. The high resolution and multi-wavelength coverage of these instruments permit the time-dependent modeling of SEP acceleration (???). With the burgeoning demand for Big Data, solar feature detection has become more automated, employing fundamental techniques reviewed by ? and applied to features at various heights (?). Some of these techniques are tailored to detect specific solar features, such as sunspots and active regions (?).

EUV wave recognition and tracking pose challenges due to their weak intensity in proximity to other features. While algorithms for this purpose exist, their complexity is notable (????). Specifically, CorPITA (?) and AWARE (?) fit shapes to flare-centered wave sectors but are constrained to the solar disk.

This study, led by ?, introduces Wavetrack, a flexible, object-oriented Python library designed for general solar feature detection and tracking. Wavetrack integrates multiscale representation (?), Δ trous wavelets (?), and filtering at its core. Notably, Wavetrack has the capability to generate training sets for machine learning-based recognition.

Machine learning applications are gaining prominence in solar physics, spanning areas such as EUV irradiance mapping (?), solar flare prediction (?), and magnetic flux generation (?). However, the potential of data-driven CME tracking is hindered by limited training sets. Wavetrack addresses this limitation by facilitating the conversion of results into annotated sets.

Central to Wavetrack is a method for general feature recognition and tracking, implemented as an accessible, open-source Python library. Its modular structure allows the configuration of applications by adjusting a few threshold parameters. The calculation scheme integrates multiscale data representation (?) and the Δ trous wavelet transform (?), supplemented by image filtering techniques for noise removal and edge enhancement.

Wavetrack distinguishes itself from existing algorithms by offering a flexible framework for detecting various solar features in EUV, white light, and other observations. Its modular design facilitates the generation of training sets for machine learning approaches, supporting automated analysis for deriving new insights from the extensive solar data. The Δ trous wavelet transform' employed by Wavetrack enables the isolation of features at different scales, effectively separating noise from the signal and enhancing edges. Wavelet coefficients encode the multiscale structure efficiently, with thresholding suppressing noise by zeroing insignificant values. Further refinement of the signal is achieved through filtering techniques. Morphological operations, such as opening, eliminate artifacts and smooth edges, while contrast enhancement sharpens edges and boundaries. The resulting output highlights features while suppressing noise.

Wavetrack strategically applies these techniques to recognize faint features like EUV waves and tracks them by identifying significant coefficients across wavelet scales. This approach avoids pre-processing steps, such as difference images, that may introduce false signals. The adaptability of Wavetrack is a key feature, allowing users to build specific applications by adjusting parameters such as thresholds and filters. Importantly, the original data remains unaltered, with only processed copies utilized for feature detection to prevent information loss.

Wavetrack, offered as an open-source tool, invites community involvement, enabling users to contribute modules, optimizations, and new techniques. This collaborative effort is poised to enhance solar data analysis capabilities in the era of Big Data, facilitating the discovery of new knowledge to deepen our understanding of the dynamic Sun.

In the realm of solar image processing and analysis, diverse techniques contribute significantly to understanding solar phenomena. This section delineates the methods employed, encompassing wavelet transform, image filtering, feature detection algorithms, machine learning, and solar feature tracking.

- **Wavelet Transform:** The application of the Δ trous wavelet transform' constitutes a fundamental multi-scale data representation concept widely employed in solar image analysis. This method facilitates the extraction of features at various decomposition and intensity levels, contributing to the comprehensive understanding of solar events.
- **Image Filtering:** Various image filtering techniques play a crucial role in enhancing and extracting specific features within solar images. These techniques, adept at detecting and tracking phenomena like CME shock waves and filaments, contribute significantly to the analysis of solar dynamics.
- **Feature Detection Algorithms:** Automated algorithms, leveraging image processing techniques, are instrumental in detecting and identifying diverse solar features, including sunspot groups, active regions, and eruptive fronts. These algorithms enhance the efficiency of solar feature identification across different types of observations.
- **Machine Learning:** The integration of machine learning methods, such as Convolutional Neural Networks (CNNs) and Generative Adversarial Networks (GANs), signifies a contemporary approach in solar image analysis. These methods find application in tasks ranging from predicting solar flares to generating magnetic flux distributions, enriching the analytical capabilities of solar researchers.
- **Solar Feature Tracking:** An essential facet of image analysis involves tracking the evolution of solar features over time. Tracking algorithms, instrumental in monitoring the movement and changes in features like CMEs, filaments, and EUV waves, contribute significantly to understanding dynamic solar processes.

The present study leveraged observations from the Atmospheric Imaging Assembly (AIA) instrument on the Solar Dynamics Observatory (SDO). The AIA instrument, functioning as an extreme ultraviolet (EUV) imager, offers high-resolution and high-temporal observations, particularly useful for detecting and characterizing large-scale shocks known as EUV waves or Coronal Bright Fronts (CBFs). Focusing specifically on the AIA channels centered on 193Å and 211Å wavelengths, the study utilized a standard AIA pipeline with the SunPy package to process the 193Å data.

The processing involved the creation of base images through the averaging of a series of images preceding the eruption. Subsequent construction of base difference images, achieved by subtracting the base images from the current raw image sequence, facilitated the enhancement of intensity changes caused by CBFs while mitigating static details and reducing noise. Given the typically dim nature of CBFs in EUV images, the study pre-selected a segment of the dynamic range where the shock wave is revealed from the base difference image of the sequence. The observations obtained from the SDO/AIA telescope were instrumental in studying and tracking the CBFs and their temporal evolution, providing essential data for analyzing the spatial and temporal relationships of CBFs and other solar dynamic features.

2.6.2 Image Filtering Techniques

Specific image filtering techniques, including thresholding, wavelet decomposition, and segmentation, formed the cornerstone of the method. Initial thresholding applied to the absolute values of pixels in the base difference images narrowed the dynamic range, concentrating on the segment containing the object of interest.

Subsequent to thresholding, the base difference images underwent Δ trous wavelet decomposition into multiple scales. Each wavelet coefficient underwent relative thresholding based on the statistical distribution of pixel intensities at each decomposition level. The segmentation process yielded object masks for each time step, subsequently multiplied by the original data to reveal the intensity of different parts of the object.

Given the variations in statistical distributions of pixel intensities in base and running difference images, tailored approaches were applied. In base difference images, pixel intensities were constrained to a specific segment, focusing on the object of interest, such as a CBF. Running difference images, employed to identify eruption initiation, depended on the specific features and objects under observation.

The Δ trous wavelet transform, constituting a multi-scale data representation, significantly contributed to the recognition and tracking of solar images. This hierarchical decomposition enabled the extraction of specific objects and their masks from imaging observations, facilitating their tracking and temporal evolution. The Δ trous wavelet decomposition enhanced the clarity and intensity of faint features like EUV waves, offering a flexible framework applicable to various solar phenomena across different wavelengths. This method emerges as a valuable tool for comprehensively analyzing and understanding the dynamics of solar features, including CME shock waves and filaments.

The study introduced the Wavetrack package, designed for automated detection and tracking of dynamic coronal features in solar observations. Utilizing wavelet decomposition, feature enhancement and filtering, and object recomposition, Wavetrack identified and tracked features such as CBFs and eruptive filaments. This object-oriented Python framework, adaptable to both on-disk and off-limb features, enables tracking of features that bifurcate over time.

2.6.3 Wavetrack for Coronal Wave Tracking

In tracking coronal waves, Wavetrack utilized wavelet decomposition, feature enhancement, and filtering techniques. The Δ trous wavelet decomposition method enhanced faint features like EUV waves, capturing different scales of features through convolutions. Automated feature recognition, incorporating intensity thresholding, image posterization, median filtering, segmentation, and intensity level splitting, identified and isolated coronal wave features. The output comprised time-dependent pixel masks, representing the tracked coronal wave, applicable to generating final feature maps for both on-disk and off-limb features.

2.6.4 Wavetrack for Filament Tracking

For filament tracking, Wavetrack followed a process involving wavelet decomposition, feature recognition, object mask generation, and image recomposition. The Δ trous wavelet decomposition identified different scales of features, and subsequent processing enhanced the features for ease of tracking. Object masks generated for each time step isolated filament features, and image recomposition from weighted wavelet

scales produced final feature maps. The choice of images, such as inverted AIA 193Å images, depended on source data and filament velocity.

2.6.5 Fourier Local Correlation Tracking (FLCT) Model

To determine the plane-of-sky velocity and speed of solar features, the study employed the FLCT model, utilizing the Fourier Local Correlation Tracking (FLCT) method. Through the analysis of consecutive solar images and the tracking of feature motion over time, the FLCT algorithm was applied to object masks derived from the Wavetrack methodology. The resultant output furnished detailed maps of instantaneous velocity, contributing to a deeper understanding of the expansion and evolution of solar features such as Coronal Bright Fronts (CBFs) and erupting filaments.

The overall methodology of the study integrated advanced image processing techniques, incorporating wavelet transform, image filtering, feature detection algorithms, and machine learning. This comprehensive approach facilitated an in-depth analysis of solar observations. Notably, the Wavetrack package emerged as a versatile tool, specifically valuable in the tracking of dynamic coronal features. Its application provided significant insights into the intricate dynamics governing the solar environment.

2.6.6 Results

The study presents several examples and case studies that demonstrate the application of the Wavetrack package. Here are some of the examples mentioned:

1. **May 11, 2011 event:** The Wavetrack algorithm was used to track both an erupting filament and the coronal bright front it drives. The relationship between the driver and wave was studied using the segmented features from AIA 193Å observations.
2. **September 29, 2013 event:** Wavetrack was applied to a large-scale filament in a slow eruption. The segmented filament feature was overlaid on the solar corona, and the on-disk feature was consistently tracked during the eruption.
3. **December 12, 2013 event:** Wavetrack was used to track the evolution of driven and non-driven regions of the CBF. The method revealed the relation between the CBF and eruptive features. These examples demonstrate the versatility of Wavetrack in recognizing and tracking various solar features, both on the solar disk and off the solar limb.
4. **June 07, 2011 event:** For this event, the variation in speed between the geometric center and center of mass is up to 300 km s^{-1} in the radial direction. The angle between the geometric center and center of mass changes quite a lot as different parts of the compressive front brighten and dim. However, the angle remains relatively stable, changing only slightly during this event.

Here, I will just focus on the first event since it was accompanied with a coronal bright front.

The Wavetrack methodology was applied to analyze three previously investigated eruptive events observed by the AIA telescope (??). In the first event on May 11, 2011, an erupting filament induced a relatively bright Coronal Bright Front (CBF) on the northwest region of the solar disk near the limb. The second event, occurring on June 07, 2011, originated close to the southwest limb and featured a large and luminous CBF. On December 12, 2013, the third event, also originating near the southwest limb, was examined.

Figure 4 illustrates the successfully extracted Wavetrack CBF objects for the three events, observed at four time points, each separated by 3 minutes. Wavetrack object masks, when applied to base difference images, were visualized using a blue-green color map overlaid on the original AIA 193Å observations in grayscale to emphasize the CBFs. Across all three events, Wavetrack adeptly delineated the extent of the CBF in consecutive time steps, demonstrating its ability to track the evolution of CBFs both on the solar disk and off the limb. This capability was maintained despite the distinct pixel distributions and intensities in these two regions. The application of Wavetrack facilitated the detailed study of the time-dependent shapes and changing intensity distributions of CBFs, separate from the broader corona. Moreover, Wavetrack effectively selected CBF objects under different coronal activity states. For instance, despite increased coronal activity on December 12, 2013, Wavetrack successfully segmented the CBF. Importantly, the segmentation of the CBF in the last event highlighted Wavetrack's capability to track multiple separate parts of the same feature.




Figure 2.1: Wavetrack images showing progression of a CBF and filament on May 11, 2011. Three snapshots captured ~ 2 minutes apart track the CBF and filament over time. Credit goes to ?.

The Wavetrack code was further applied to isolate and study the kinematics of CBFs and filaments in detail. The Fourier Local Correlation Tracking (FLCT) method, extensively utilized for estimating horizontal flows in photospheric magnetograms, was employed for this purpose. Initially proposed as a cross-correlation technique for precise motion measurement, FLCT calculates a 2D flow field that best resembles the scalar field between two consecutive 2D maps.

Figure 7 displays four pairs of consecutive AIA images from the May 11, 2011 event, used as input for the FLCT algorithm. Each pair, separated by 24 seconds, is strategically chosen to observe the CBF progression over 6 minutes. The corresponding FLCT results, presented in Figure 8, show the instantaneous plane-of-sky direction of motion and speed of each CBF region. Notably, the results reveal the uneven expansion of the CBF from the central source. In this event, the thinnest part of the CBF, strongly driven by the erupting filament, exhibited the fastest motion compared to other regions on the disk and off the limb. This nuanced information, not discernible from intensity observations alone, underscores the value of our combined Wavetrack and FLCT approach in elucidating the dynamic behavior of coronal features.

2.6.7 Conclusions

In this study, led by ?, we introduce Wavetrack, an innovative approach designed for the automated identification and monitoring of dynamic coronal phenomena. Employing wavelet decomposition, feature enhancement, filtering, and ultimate object recombination, Wavetrack generates time-dependent masks for feature pixels. These masks can be applied to integral or base-difference images, yielding comprehensive feature maps. Notably, Wavetrack excels in tracing pixels associated with faint, large-scale features, such as coronal bright fronts/EUV waves in AIA observations, and has demonstrated efficacy in tracking eruptive filaments. Operable for both on-disk and off-limb features, Wavetrack adeptly follows features that evolve into distinct components over time. Implemented as a versatile, object-oriented framework in Python, Wavetrack is freely accessible for download and utilization.

The application of Wavetrack to four distinct events, with emphasis on three – the CBFs occurring on May 11 and June 07, 2011, and December 12, 2013 – reveals its proficiency in tracking complete CBF pixel maps. These model results, when integrated with the FLCT method for calculating plane-of-sky speeds, unveil the dynamic evolution of driven and non-driven regions within CBFs, as well as their correlation with eruptive filament drivers. Our findings indicate that drivers induce a compression effect, causing CBF thinning and increased speed, aligning with theoretical models. However, the brightness of CBFs in the driven regions, as observed in AIA data, does not necessarily exhibit a significant increase.

Furthermore, Wavetrack facilitates the tracking of temporal changes in feature regions by computing the time-dependent vector between the pixel geometric center and the center of mass, determined by weighing the observed pixel intensities. This proves particularly valuable for large-scale features like CBFs, providing a straightforward metric in the form of a one-dimensional time series for characterizing feature evolution.

While the method demonstrates widespread applicability to various solar dynamic features and observational data, it currently relies on human input for the segmentation of specific feature types, particularly dim ones. Manual setting of object criteria, including threshold intervals and recombination weight coefficients, is necessary. In instances of base difference imaging, precise selection of the base image is crucial to prevent contamination of input data with spurious features. Additionally, parameter fine-tuning through trial and error may be required for specific events. These limitations are acknowledged, and future work will address them to enhance the model's versatility.




Figure 2.2: Wavetrack images of a May 11, 2011 eruption, with CBF mask applied. Four image pairs shown, separated by 2 minutes, to track CBF over time. Credit goes to ?.




Figure 2.3: FLCT model output for four image pairs from Fig. 3. Left: the plane-of-sky velocity vectors. Right: plane-of-sky speed. Credit goes to ?.

The methodology holds promise for extensive application across diverse solar dynamic features and observational datasets. Subsequent research endeavors will extend its use to in-depth studies of filament evolution and coronagraph data. The objective is to refine our comprehension of how large-scale eruptive fronts manifest variations across distinct observational contexts, spanning the low and middle corona.

2.7 Geomagnetic Storms: CME Speed De-Projection vs. In Situ Analysis

This study, led by ?, examines the relationship between the intensity of geomagnetic storms (GS) and parameters of solar and interplanetary phenomena. We utilize the recently developed PyThea framework to reconstruct the 3D geometry of geo-effective CMEs and compare on-sky and de-projected values, focusing on the reliability of the de-projection capabilities. Spheroid, ellipsoid, and graduated cylindrical shell models are used. We collected parameters of the GS-associated phenomena. Considerable variation in 3D de-projections of CME speeds was obtained depending on the reconstruction model chosen and subjective observations. Fast CME speeds combined with frontal magnetic structure orientation when reaching Earth's magnetosphere proved the best indicator of GS strength. More accurate estimations of geometry, direction, and de-projected speeds are critical for operational GS forecasting in space weather prediction schemes.

2.7.1 Overview

Solar eruptive events like coronal mass ejections (CMEs) and solar flares (SFs) can generate disturbances that propagate through interplanetary space and impact Earth's magnetosphere, leading to space weather effects (????). The electromagnetic radiation from SFs arrives at Earth first, followed by energetic particles. The plasma clouds of CMEs take longer, from tens of hours up to a few days, to impact Earth's environment (??). The temporary disruption of Earth's magnetosphere and atmosphere due to these solar events are known as geomagnetic storms (GSs) (????).

The coupling between solar wind plasma and Earth's magnetosphere occurs through magnetic reconnection, enabled when the interplanetary (IP) magnetic field turns southward (negative B_z component) and impacts Earth at high speed, as during CMEs (???). This leads to increased particle injection into the magnetosphere and atmosphere, causing bright auroral displays. Drifting electrons and protons also drive the westward ring current responsible for decreases in the horizontal magnetic field measured by the disturbance storm time (Dst) index (??).

Fast CMEs propagating through IP space (ICMEs) cause the most intense GSs, with sudden Dst decreases, compared to gradual storms from corotating interaction regions (CIRs) (????). ICME shock waves and magnetic ejecta produce cascading effects in near-Earth space that can disrupt technology (?).

Earth-directed fast ejecta with strong southward magnetic fields inside are the most geoeffective. However, single spacecraft observations are limited by projection effects, leading to uncertain CME speed estimations (??). Previous studies found no clear relationship between GS intensity and solar flare parameters or CME properties like projected speed and angular width (?). Furthermore, CME magnetic structure cannot be derived from remote sensing data. Reliable solar or near-Sun measurements that provide early warnings for potential GS strength remain lacking.

Accurately predicting when an incoming disturbance will impact Earth requires determining the arrival time and speed of CMEs. Different portions of these large structures, such as the apex or flanks, can hit Earth upon arrival at 1 AU. Flank hits may only involve the CME sheath while apex hits include both sheath and ejecta, leading to different magnetospheric effects (?). Therefore, deducing CME directionality and 3D geometry is important. To maximize forecast lead time, these parameters should be estimated as early as possible when the CME emerges in coronagraph fields of view. In images, CMEs appear as line-of-sight integrated brightness enhancements projected onto the observing plane (??).

Further developing reconstruction techniques to correct for projection effects can improve CME propagation understanding and impact forecasting (????). Several CME propagation models exist (????). A study found 2D CME speeds underestimate 3D speeds by $\sim 20\%$ while 2D widths overestimate 3D widths (?). Another study showed observer bias in 3D modeling using graduated cylindrical shells, even for experienced observers (?). CME structure interpretation differs, and line-of-sight integration leads to non-unique solutions.

This study focuses on deducing CME directionality and near-Sun speeds using new tools like PyThea (?). We analyze geo-effective cycle 24 CMEs using multiple reconstruction techniques by two team members. Comparisons are made between derived parameters and storm intensity (Dst). CME speed correlations with ICME and IP shock speeds evaluate 3D de-projection value for arrival forecasting. Other IP parameters are also examined, including shock speed, plasma jumps, and magnetic fields near L1. We examine the correlation between the intensity of geomagnetic storms and parameters of solar and interplanetary phenomena. More specifically, we focus on the speed and geometry of CMEs, as well as interplanetary shocks, as these factors are known to play a significant role in the occurrence and intensity of geomagnetic storms. The objective of the study is to analyze the correlations between the intensity of geomagnetic storms (GS) and parameters of solar and interplanetary (IP) phenomena. The study also aims to perform 3D geometry reconstructions of geo-effective coronal mass ejections (CMEs) using the PyThea framework and compare the de-projected CME speeds with in situ data.

The event selection process for this investigation commenced with the identification of significant Geomagnetic Storms (GSs) within Solar Cycle 24 (SC24). These storms were characterized by a Dst index exceeding 100 nT, as per the classification outlined in ?. A total of 25 GSs were identified, with Dst indices ranging from 101 to 234 nT. Previous studies have explored the solar and Interplanetary (IP) origins of these GSs in SC24 (?????). However, the comprehensive review of all pertinent literature falls beyond the purview of this study. It is noteworthy that SC24 exhibited a reduced number of GSs in comparison to preceding solar cycles (?).

In our investigation, distinct from prior analyses, our objective was to establish a causal connection between the identified GSs and potential IP and/or solar phenomena. This approach mirrors methodologies employed by other researchers (?????). To delineate the solar and IP drivers, we employed an association procedure widely acknowledged in the field of Space Weather (SW) research. The methodology involves searching for the IP and solar origins of a GS storm within a specific timeframe preceding the reported GS occurrence at Earth. The sequential steps of our approach are delineated below:

1. **Temporal Association with IP and ICME Events:** We initiated the analysis with a temporal association between the GS and recorded IP shocks in the vicinity of Earth. This association was established within a 1-day period preceding the reported minimum Dst of the GS. A parallel procedure was employed for the association with Interplanetary Coronal Mass Ejections (ICMEs) reported in proximity to Earth. Additionally, animations from the *helioweather* archive¹ (accessed on 5 April 2023) were utilized to validate potential ICME and IP shock candidates.
2. **Association with Coronal Mass Ejections (CMEs):** Subsequently, we extended the association to include Coronal Mass Ejections (CMEs) within a 3-to-5 day window prior to the IP (or GS) timing. Information from available solar and IP event catalogs, as well as animations from the *helioweather* archive¹ (accessed on 5 April 2023), was employed for this purpose.
3. **Completion of the Association with Solar Flares (SFs):** The final step involved associating the GS with the identification of a Solar Flare (SF) linked to the previously associated CME. This association was based on timing (within one hour between the SF onset and CME timing) and location constraints (the SF location had to align with the solar quadrant reported in the measurement position angle, MPA, of the CME).

All utilized databases, catalogs, and publicly available lists pertinent to the analysis are summarized as follows:

- GS database (Kyoto)² (accessed on 5 April 2023);
- SF database (GOES)³ (accessed on 5 April 2023);
- CME catalog (SOHO-LASCO)⁴ (accessed on 5 April 2023);
- ICME Wind database⁵ (accessed on 5 April 2023);
- ICME ACE database⁶ (accessed on 5 April 2023);

¹<http://helioweather.net/archive/>

²<https://wdc.kugi.kyoto-u.ac.jp/dstdir/index.html>

³<http://ftp.swpc.noaa.gov/pub/warehouse/>

⁴CMEcatalog(SOHO-LASCO)

⁵https://wind.nasa.gov/ICME_catalog/ICME_catalog_viewer.php

⁶<https://izw1.caltech.edu/ACE/ASC/DATA/level3/icmetable2.htm>

- IP shock database (Wind)^{7,8} (accessed on 5 April 2023).

2.7.2 GSs and IP Phenomena

The findings pertaining to Geomagnetic Storms (GSs) and their associated Interplanetary Coronal Mass Ejections (ICMEs) and Interplanetary (IP) shocks are succinctly presented in Table 1 in ?. The initial column designates the event number (#), a consistent reference utilized throughout this paper. Columns (2) and (3) detail the GS date, hour (mm-dd/hr), and the corresponding Dst index (in nT). Subsequently, columns (4)–(6) expound upon the ICME parameters (?), drawing from the Wind database accessible at this website⁹. The sheath duration (D, in hours), representing the temporal span between the initiation times of the ICME and the magnetic structure, is computed from the available timings within the plots offered by the aforementioned Wind database, and is presented in column (7). The ICME in situ measured speed is extracted from both the Wind and ACE databases, with the exception of E11 where no ICME is reported.

Column (8) integrates the minimum B_z component throughout the Interplanetary Coronal Mass Ejection (ICME) duration, as ascertained from the data available at this website¹⁰, thereby enhancing the comprehensiveness of the dataset. In Column (9), we conduct a qualitative assessment of the ICME arrival orientation. Specifically, we visually inspect the intersection point between the ICME structure and Earth using ecliptic plane animations accessible at this website¹¹. The orientations are denoted as either *hit* (nose) or *f* (flank) arrivals. Instances of discrepancies among diverse data sources, such as the presence of solar wind streams or Corotating Interaction Regions (CIRs) visible in animations contradicting ICME arrivals identified through in situ data, are marked as ‘u’ (uncertain) in the same column. This classification signifies situations where a distinct ICME structure propagating through the IP space could not be conclusively observed. Noteworthy is the occurrence of swift solar wind flows and/or CIRs recorded near Earth during ICME and/or shock wave events. For instance, in the cases of E11 and E18, a CIR was identified as their IP origin by ?; however, in contrast to these findings, our methodology does not differentiate between ICME and sheath origins.

The final columns, (10)–(13), enumerate the properties of the IP shock, incorporating timing, speed, magnetic field, density, temperature jump at the shock interface, and the Mach number (Mms). These details are sourced from Wind satellite data, available at this website¹², with the exception of E24, where the median shock speed is derived from this website¹³. Notably, for E17, E18, and E25, no IP shocks are reported in either database.

2.7.3 GSs and Solar Phenomena

In the context of five cases denoted as E05, E11, E17, E18, and E22, our attempts to identify Solar Flares (SFs) or Coronal Mass Ejections (CMEs) were unsuccessful. Furthermore, in an additional six cases, the specification of SFs proved unattainable. The ensuing presentation provides details on the parameters of the remaining cases, specifically focusing on the solar origins associated with Geomagnetic Storms (GSs), as outlined in Table ??.

Columns (2)–(5) of Table ?? elucidate the properties of the GS-associated SFs, while columns (6) to (9) furnish the parameters of the GS-associated CMEs. The identified SFs exhibit a range of magnitudes from C1.2 to X5.4 and are predominantly positioned proximate to the solar disk center, with the exception of E02 and E03. The CMEs associated with these events possess on-sky projected speeds, denoted as 2D, spanning from 126 to 2684 $km\ s^{-1}$, extracted from the SOHO-LASCO CDAW catalog. Notably, the majority of the GS-associated CMEs (15 out of 20) exhibit a halo configuration, while three others are in close proximity to halo.

It is imperative to note that events with uncertain CME origins have been excluded from the subsequent 3D analyses. Specifically, in the case of E07, the unique orientation of the double CME rendered the de-projection procedure unfeasible for the same CME structure, leading to its exclusion from the 3D analyses. Additionally, for seven other cases (E12, E14–E16, E19, E23, E25), the online tool utilized for

⁷<http://www.ipshocks.fi/database>

⁸https://lweb.cfa.harvard.edu/shocks/wi_data/

⁹https://wind.nasa.gov/ICME_catalog/ICME_catalog_viewer.php

¹⁰<https://cdaweb.gsfc.nasa.gov/>

¹¹<http://helioweather.net/archive/>

¹²<http://www.ipshocks.fi/database>

¹³https://lweb.cfa.harvard.edu/shocks/wi_data/

analyses failed to retrieve data simultaneously from both spacecraft. Consequently, these cases have also been omitted from the 3D analyses.

For the remaining 12 cases, successful 3D CME speed reconstructions were achieved from each model. The mean values, based on 2 or 4 available fits (as detailed in the subsequent subsection), are presented in the concluding columns (10)–(12) of Table ???. This comprehensive overview provides a foundation for the subsequent analytical discussions, offering a detailed characterization of the solar events associated with the studied GSs.

2.7.4 PyThea 3D De-Projection Tool

The de-projection methodology employed in this investigation relies on the innovative PyThea online tool designed for the 3D reconstruction of CMEs and shock waves (?). All three available models within PyThea, namely spheroid, ellipsoid, and the graduated cylindrical shell (GCS), are applied in this study. The fitting procedure is conducted independently by two observers within our team. Figure ?? presents an illustrative example of the fits for event E03.




Figure 2.4: 3D reconstructions of a CME (E03) using the spheroid, ellipsoid, and GCS model from the PyThea tool performed by observers 1 and 2 (top and bottom row, respectively). Credit goes to ?.

Upon scrutinizing the fitting outcomes for this particular example, it becomes evident that the reconstructions exhibit a discernible bias. The observer's subjective *choice* of structures to align with the model introduces a level of variability. In the top row of Figure ??, distinct shock-related structures (manifested as the bending of streamers) are observed, upon which the idealized GCS flux rope geometry is fitted. Consequently, there is a likelihood of overestimating the CME width. Despite this bias, it is noteworthy that the overall results, including directivity and speed for event E03, are minimally impacted. However, it is acknowledged that the complexity of structure choices can lead to larger discrepancies among different observers.

This study places emphasis on deriving de-projected CME speeds based on fits conducted at two distinct time steps. For each of the three models, the initial CME longitude and latitude are manually specified, utilizing the provided locations of the CME-accompanied SFs. It is important to note that these values exhibit minimal to no substantial changes post-finalization of the fitting procedure. Consequently, the final CME directivity provided by PyThea is considered to be a relatively crude estimate. The ultimate orientations of the CME in IP space and at Earth are derived solely from qualitative information extracted from animations available at this website¹⁴. This approach ensures a rigorous and consistent

¹⁴<http://helioweb.net/archive/>

basis for evaluating the de-projected CME speeds in our analyses.

2.7.5 Results

Projection Effects

In our investigation a fitting analysis of approximately 10 CMEs was undertaken by two designated observers within our research team, I was one of the observers. This analysis involved the utilization of all three models available in the PyThea framework, each selected based on the evaluators' individual considerations.

To elucidate, the fitting process was executed at two distinct time steps to derive a velocity parameter based on the height-time estimation. For each specific event, both observers iteratively conducted the 3D de-projection procedure twice, resulting in averaged values for the CME speeds. These values are meticulously presented in Table ??, rounded to the nearest tenth. The disparity between these two fitting instances is denoted as an error (or uncertainty), spanning from 10 km s^{-1} to twice the estimated speed. Additionally, Figure 2 illustrates the correlation between 3D speeds and the corresponding estimated errors for each observer. Notably, considerable variability is observed in both plots, particularly concerning the GCS model; nonetheless, a discernible positive trend emerges between the estimated error magnitude and the CME speed.

The inherent subjectivity and diverse levels of experience among observers play a crucial role in the visual fitting procedure. Noteworthy distinctions in evaluated speeds are evident not only between individual observers using the same model (e.g., spheroid fit for E06) but also between a single observer applying different models (e.g., spheroid and GCS for E13). Furthermore, variations in operating system software further contributed to differences in results. Notably, events E04, E08, and E21 were not completed by both observers due to either PyThea computing resource failures or the substantial uncertainty associated with the visual assessment of CME structures.

Our findings underscore the well-established subjectivity inherent in procedures relying on personal judgment for fit quality, where the alignment of structures with the model is subjective. A more comprehensive explanation of this human-in-the-loop bias is detailed in (?). The specific values of CME speeds are meticulously outlined in Table ??, while their averaged values, categorized by model, as determined by both observers, are presented in Table ???. These averaged values will serve as the basis for subsequent correlation studies.

Table 2.1: Parameters of the solar origin, SFs, and CMEs for GSs from Table 1 in ?. All times are in UT, speeds in km s^{-1} , AW and MPA in degrees. Credit goes to ?.

# (1)	Event (2)	SF (3)	Onset (4)	Location (5)	Time (6)	Speed (7)	AW (8)	MPA (9)	Spheroid (10)	Elliptical (11)	GCS (12)
2011											
E01	08-04	M9.3	03:41	N19W36	04:12	1315	360	298	1990	1920	1780
E02	09-24	M7.1	12:33	N10S56	12:48	1915	360	78	1570	1590	1720
E03	10-22	M1.3	10:00	N25W77	10:24	1005	360	311	760	690	840
2012											
E04	03-07	X5.4	00:02	N17S27	00:24	2684	360	57	2150	2460	2530
E05				uncertain origin					-	-	-
E06	07-12	X1.4	15:37	S15W01	16:48	885	360	158	1060	1780	1520
E07	09-28	C3.7	23:36	N06W34	24:12	947	360	251		multiple CMEs	
E08	10-05		uncertain		02:48	612	284	202	350	360	350
E09	11-09		uncertain		15:12	559	276	157	660	570	720
2013											
E10	03-15	X1.1	05:46	N11S11	07:12	1063	360	112	720	1040	1110
E11				uncertain origin					-	-	-
E12	06-28		uncertain		02:00	1037	360	214		no SOHO images	
2014											
E13	02-16	M1.1	09:20	S11E01	10:00	634	360	227	340	690	890
2015											
E14	01-03	C1.2	03:06	S05E21	03:12	163	153	144		no STEREO images	
E15	03-15	C9.1	01:15	S22W25	01:48	719	360	240		no STEREO images	
E16	06-21	M2.6	02:06	N12E13	02:36	1366	360	72		no STEREO images	
E17				uncertain origin					-	-	-
E18				uncertain origin					-	-	-
E19	12-16	C6.6	08:34	S13W04	09:36	579	360	334		no STEREO images	
2016											
E20	12-28	M1.8	11:20	S23W11	12:12	1212	360	163	820	680	1080
E21	01-14		uncertain		23:24	191	360	234	620	440	280
E22				uncertain origin					-	-	-
2017											
E23	05-23		uncertain		05:00	259	243	281		no SOHO images	
E24	09-04	M5.5	20:28	S11W16	20:36	1418	360	184	1020	1290	990
2018				uncertain		21:24	126	120	266		no STEREO images
E25	08-20										

Table 2.2: CME Speeds ($km\ s^{-1}$) for Observers 1 and 2. Credit goes to ?.

#	Spheroid		Ellipsoid		GCS	
	obs1	obs2	obs1	obs2	obs1	obs2
E01	2170 \pm 870	1800 \pm 270	2130 \pm 200	1710 \pm 450	1590 \pm 100	1760 \pm 10
E02	1780 \pm 140	1350 \pm 50	1880 \pm 580	1310 \pm 90	1780 \pm 260	1630 \pm 130
E03	770 \pm 40	740 \pm 10	640 \pm 180	740 \pm 180	1020 \pm 170	700 \pm 270
E04	-	2150 \pm 140	-	2460 \pm 70	-	2530 \pm 630
E06	1410 \pm 420	710 \pm 70	1870 \pm 50	1700 \pm 300	1680 \pm 870	1560 \pm 470
E08	350 \pm 90	-	360 \pm 150	-	350 \pm 70	-
E09	690 \pm 280	630 \pm 150	550 \pm 170	590 \pm 60	670 \pm 610	710 \pm 220
E10	840 \pm 380	610 \pm 1040	1120 \pm 360	960 \pm 90	1160 \pm 650	1310 \pm 80
E13	320 \pm 90	350 \pm 50	620 \pm 140	750 \pm 160	780 \pm 80	1310 \pm 700
E20	830 \pm 190	800 \pm 600	790 \pm 90	570 \pm 20	1240 \pm 280	1130 \pm 230
E21	620 \pm 230	-	440 \pm 40	-	280 \pm 180	-
E24	750 \pm 270	1310 \pm 220	880 \pm 350	2020 \pm 960	950 \pm 120	1560 \pm 540

Correlation between GSs, Coronal and Near-Sun Parameters

In Figure 3, we present a scatter plot depicting the relationship between the modulus of the GS Dst index and the CME speed, as derived from the data in Table ???. To enhance clarity, the averaged results of the three model fits are collectively illustrated and labeled as '3D-mean' in Table 4. These values are juxtaposed with the 2D SOHO-LASCO CME speed. Horizontal lines, representing the error estimates from the 3D de-projections, are also included for completeness, despite the substantial overlap. For demonstrative purposes, we highlight the largest error value among the two observers, as outlined in Table 3.

The analysis conducted reveals no discernible trend between the Dst index and the CME speed, irrespective of whether the 3D de-projection or the 2D CME speeds are considered. It is important to note that, due to data constraints, 3D speed de-projections were not feasible for the most robust GSs. This limitation results in a skewed distribution of the 3D speeds, impacting the overall findings. Despite the modest sample size (comprising between 10 and 20 event pairs), the quality of the fit is assessed through Pearson correlations. The correlation coefficients, reflecting the relationship between all CME speed estimations and the GS Dst index, are systematically documented in Table 4. These coefficients range from negligible (e.g., 0.04 for the 2D LASCO speeds) to moderate (with the highest value reaching 0.55, observed with the GCS model). Importantly, no significant correlations are identified between the Dst index and other coronal parameters, such as SF class, location, and CME AW, as inclusively presented in the same table for comprehensive evaluation.

Correlation between GSs and IP Parameters

Here, we explore the correlations between Geospace Storms (GSs) and various parameters associated with pre-selected Interplanetary (IP) phenomena. To visually represent these relationships, scatter plots are employed for specific parameter pairs, as follows: the Dst index versus Interplanetary Coronal Mass Ejection (ICME) speed and IP shock speed in Figure 4; Dst versus Mach number and sheath duration in Figure 5; Dst versus jBzj and Bd/Bu in Figure 6; and Dst versus Td/Tu and Nd/Nu in Figure 7. The corresponding Pearson correlation values elucidating these trends are documented in Table 5.

For the limited sample of GS storms utilized in our analyses, we observe a moderately positive trend between the Dst index and the plasma compression parameters at the shock interface (downstream to upstream ratio). Interestingly, these correlations are comparable to, or slightly larger than, those obtained when considering ICME speeds from the Wind and ACE spacecraft. Notably, the trend with jBzj is weaker (0.37 for our dataset), in contrast to the robust trend reported in previous studies. Furthermore, the correlations involving Dst and IP shock speed, Mach number, or sheath duration are relatively smaller.

Recent reports have highlighted strong correlations with different components of the electric and magnetic fields [12], although this aspect exceeds the scope of our current analyses. It is essential to exercise caution in interpreting these results, given the absence of uncertainty estimates for the correlation coefficients.

The ICME and IP shock catalogs employed in our study offer additional parameters, including averaged magnetic field B , plasma speed V inside the magnetic structure, and upstream plasma beta β_u (not presented in Table 1). However, no robust correlations emerge between these parameters and the Dst index, with all correlation coefficients found being smaller than 0.2.

On the GS Strength Forecasting Based on Solar and IP Parameters

We examine the collective impact of magnetic obstacle type and orientation upon Earth arrival (columns 5 and 9 from Table 1), along with the 3D reconstructed Coronal Mass Ejection (CME) speeds (columns 10–12 from Table 2), on the intensity of Geospace Storms (GSs), approximated in this study using the Dst index.

The most potent GSs in our compilation, arranged in descending order based on their Dst values (nT), exhibit distinctive magnetic structure parameters, including complexity, orientation of arrival, and speed at Earth (refer to Tables 1 and 2):

- E15 (-234): Magnetic obstacle type undefined, fast speed (f), no 3D speed estimation - E16 (-198): Complex (Cx) structure, nose-like (n) orientation, no 3D speed estimation - E25 (-175): Flux-rope (Fr) structure, nose-like (n) orientation, no 3D speed estimation - E19 (-166): Flux-rope (Fr) structure, nose-like (n) orientation, no 3D speed estimation - E03 (-147): Complex (Cx) structure, fast speed (f), reduced 3D speed compared to 2D - E04 (-145): Complex (Cx) structure, nose-like (n) orientation, similar 3D speed compared to 2D - E06 (-139): Flux-rope (Fr) structure, nose-like (n) orientation, larger 3D speed compared to 2D - E10 (-132): Flux-rope (Fr) structure, nose-like (n) orientation, similar 3D speed compared to 2D

Upon scrutiny of these cases, it becomes apparent that the most intense GSs are linked to magnetic obstacles characterized by a nose-like (n) orientation upon arrival, coupled with either a complex (Cx) or flux-rope (Fr) structure. Exceptions include instances of fast speed flank hits or flank hits in combination with a complex (Cx) structure. It has been established that sheath duration does not serve as a decisive ordering parameter. Other GSs of lesser intensity (refer to Table 1) predominantly result from flank hits or exhibit uncertain configurations, possibly influenced by fast solar wind streams or Corotating Interaction Regions (CIRs). Notably, E20 is an exception, resulting from a nose hit; however, the Interplanetary (IP) structure associated with it has a notably low speed, as observed in the examined animations.

It is crucial to acknowledge that the IP shock speed provided by the Wind and ACE satellites represents a single point sample within the entire structure. In contrast, animations offer a global speed distribution. Therefore, our interpretation considers information from both sources — speed reconstructions and in situ measurements — to enhance the robustness of our analysis.

2.7.6 Discussion

In this study, we conduct post-event analyses of all geo-effective storms observed during solar cycle 24 with the aim of identifying distinct and reliable predictors for GS intensity. The overarching objective is to derive dependable solar or near-Sun parameters from remote sensing image data that can be effectively utilized for early warnings regarding potential GS onsets and their strength. Our approach involves the integration of solar, near-Sun, and interplanetary parameters, primarily sourced from catalogs but also subject to our analysis using space-related databases. Notably, we incorporate the results obtained from a novel tool for CME speed de-projection, termed PyThea, which is employed for the first time alongside well-established parameters in space weather and geophysics research.

Among the solar and near-Sun phenomena considered, certain selected parameters exhibit a positive correlation with the Dst index. Notably, the correlation coefficient, when comparing the observed projected CME speed, experiences an improvement from 0.04 (as obtained from LASCO) to a range of 0.34–0.55, achieved through various geometrical models provided by the PyThea software package, combining LASCO and STEREO data. However, our exploration of different CME geometry reconstruction techniques reveals a susceptibility to large speed errors, particularly in the case of fast CMEs. This aligns with findings from prior studies focusing on CME arrival time and speed forecasts for Earth, suggesting a potential overestimation of CME launch speed for fast events, likely attributable to the increased complexity in discerning coronal structures due to the rapid expansion of the magnetic structure associated with the CME.

Moreover, our examination indicates that fast halo CMEs may exhibit significant deviations, stemming from the overlap in shock and magnetic structure components, thereby strongly impacting the reconstruction quality. Consequently, we assert that the derived near-Sun 3D parameters continue to

possess limited forecasting potential for predicting GS strength. In contrast, most selected IP parameters derived from in situ measurements display moderate positive correlations with GS strength, in line with expectations. However, the B_z parameter (southward component of the magnetic field) shows a relatively low correlation coefficient of 0.37, possibly influenced by the limited event sample used in our study.

Other IP parameters, such as ICME and IP shock speeds, along with their derivative parameters (e.g., Mach number), exhibit a positive trend with the Dst index, featuring correlation coefficients of 0.35–0.45. Nevertheless, none of these parameters emerge as predominant, and their reliance on single-point in situ observation raises considerations about their comprehensive predictive capability. Discrepancies in ICME speed values between ACE and Wind measurements prompt further investigation, with slightly stronger correlation coefficients (0.4–0.5) observed when utilizing different shock parameters. However, averaged values of magnetic field and speed within the magnetic structure, plasma beta in the upstream region, or duration of the sheath region show no correlation with GS strength.

Among the multitude of considered solar, near-Sun, and IP parameters, only the combination of speed and orientation (nose-like) of the magnetic obstacle appears to exert a positive influence on GS strength, as indicated by qualitative results obtained from animations accessible at *helioweather*. Consistent with earlier studies, de-projected CME speeds emerge as imperative for enhancing modeling accuracy when predicting CME propagation through the IP space. However, a notable observation is the apparent lack of direct impact of 3D de-projected CME speed on GS intensity. Consequently, an accurate estimation of the ICME speed distribution over the entire ICME structure upon arrival at Earth assumes significant importance. We emphasize the imperative need for permanent stereoscopic observations, exemplified by the upcoming ESA Vigil mission situated at Lagrange point L5. Future research endeavors should aim for a more comprehensive disentanglement of distinct CME structures, thereby enabling more reliable 3D reconstructions of CME geometries for a nuanced estimation of 3D speed and directivity.

Chapter 3

Solar Radio Observations: Integrating Space-based and Ground-based Data for Coronal Diagnostics

In this chapter, I focus on multi-wavelength observations of solar type III radio bursts, as well as modeling studies of plasma parameters and coronal magnetic fields, to gain insights into the mechanism of solar radio emission during quiet times and the coronal conditions influencing the propagation and detection of these bursts. First, I briefly introduce the characteristics of solar type III radio bursts. I then present the data I have worked with, detailing the multiple processing stages I undertook. Finally, I present my outputs and interpret their meaning.

3.1 Introduction

Type III radio bursts are manifestations of transient energetic electron beams injected into the solar corona, propagating along the interplanetary magnetic field (IMF) lines ????. As these beams traverse the corona, they trigger plasma waves (also known as Langmuir waves) that are then transformed into radio emission at the local plasma frequency or its harmonic components ?. In the radio spectrograms, type III bursts are usually observed as intense emissions that drift in frequency over timescales of several seconds to minutes and over a wide range of frequencies, from metric to decametric wavelengths ???, making them detectable by ground-based instruments on Earth and various spacecraft within the heliosphere. The frequency of the radio emission is directly related to the plasma density, making type III bursts a valuable diagnostic tool for examining the inner heliosphere and the processes that drive solar active phenomena, such as solar flares and coronal mass ejections ??.

The electron beams follow open magnetic field lines and can persist well beyond 1 astronomical unit (AU) (e.g., ??), offering in situ insights into the burst and ambient conditions of the heliosphere, including electron density, radio frequency drift, speed of the electron beams and even potential direct detection of Langmuir waves (see ?? and ? and references within). In addition, tracing the path of type III bursts provides a map of the density structure of the heliosphere, serving as a foundation for developing and testing density models. Since radio observations below ~ 10 MHz cannot be accomplished from the ground, it is important to combine high- and low-frequency observations from ground-based and space-borne instruments. In this work, I perform a study of several type III radio bursts that occurred in close succession on April 3, 2019. I use remote observations of type III radio bursts detected by the Low-Frequency Array (? , LOFAR) ground-based radio telescope and the Parker Solar Probe (? , PSP) spacecraft during Encounter 2 to study the sources of these radio emissions and to investigate the physical conditions responsible for their generation. Additionally, I incorporate results of two steady-state models of the solar corona: the potential field source surface (PFSS) model ?? and the magnetohydrodynamic algorithm outside a sphere (MAS) model ?, to gain a better understanding of the coronal magnetic environment and its role in the acceleration of electrons. The ground-based LOFAR imaging observations provide valuable insight into the actual location of the burst sources. This research aims to expand upon

current knowledge of the electron beams responsible for triggering type III radio bursts and the coronal conditions they experience. Gaining a deeper insight into this aspect is vital in comprehending other solar phenomena, such as solar energetic particles and solar wind, and how they influence the near-Earth space environment.

A number of recent studies investigate the physical mechanisms responsible for the generation of solar type III radio bursts. For example, ? investigated the association of type III bursts with flaring activities in February 2011, via combined multi-wavelength observation from the Solar Dynamic Observatory (SDO) instruments, as well as Wind/WAVE and ground-based instruments. They found that the SDO measurements indicated that type III emission was correlated with a hot plasma (7 MK) at the extreme ultraviolet (EUV) jet's footpoint. By using a triangulation method with the Wind and the twin STEREO spacecraft, ? reported the first measurements of the beaming characteristics for two type III bursts between 2007-2008, assuming the source was located near the ecliptic plane (see also ?). They concluded that the individual type III bursts have a broad beaming pattern that is roughly parallel to the Parker spiral magnetic field line at the source. ? conducted a study on almost 10,000 type III bursts observed by the Nancay Radioheliograph between 1998 and 2008. Their analysis revealed discrepancies in the location of type III sources that may have been caused by a tilted magnetic field. Additionally, they found that the average energy released during type III bursts throughout a solar cycle could be comparable to the energy produced by non-thermal bremsstrahlung mechanisms in nano-flares. ? utilized LOFAR data to investigate the statistical characteristics of over 800 type III radio bursts within an eight-hour period on July 9, 2013. They discovered that the drift rates of type III bursts were twice that of type S bursts and plasma emission was the primary emission mechanism for both types.

? introduced a statistical overview of type III radio bursts during the first two PSP solar encounters. While the first encounter in November 2018 revealed a small number of bursts, the second encounter in April 2019 exhibited frequent type III bursts, including continuous occurrences during noise storms. They reported the characteristics of type III bursts with spectral and polarization analysis.

? performed a statistical survey of 30 type III radio bursts detected by PSP during the second encounter in April 2019 and estimated their decay times, which were used to estimate the relative electron density fluctuations in the solar wind. They localized radio sources using a polarization-based-radio triangulation technique, which placed the sources near the modeled Parker spiral rooted in the active region AR12738 behind the plane of the sky as seen from Earth.

? explored correlations between type III radio bursts and EUV emission in the solar corona. Using coordinated observations from PSP, SDO, and Nuclear Spectroscopic Telescope Array (NuSTAR) on April 12, 2019, they identified periodicities in EUV emission correlated with type III burst rates. The findings suggested impulsive events causing heating and cooling in the corona, possibly nano-flares, despite the absence of observable flares in X-ray and EUV data, which implies periodic non-thermal electron acceleration processes associated with small-scale impulsive events.

? explored the origin of the type III radio bursts I am tackling in this paper and found that electron beams that triggered radio bursts may have emanated from the periphery of an active region that showed significant blue-shifted plasma. More recently, ? observed a distinct type III radio burst using the PSP and LOFAR between 0.1 and 80 MHz on April 9, 2019, around 12:40 UT, six days after the occurrence of the event analyzed in our study. While no detectable flare activity was linked with the event, a type III noise storm was ongoing during the PSP encounter 2. The authors determined the type III trajectory and reconstructed its source using observations from Wind and STEREO spacecraft, as well as measuring related electron enhancement in situ.

In the last few years, I have witnessed the emergence of modern instruments, such as LOFAR and PSP, that have allowed for the observation of solar radio emissions with higher sensitivity from a better vantage point. Although type III bursts have been extensively studied ?, there are still some unresolved issues regarding the exact mechanism of type III emissions. For example, it is not yet clear how the electrons are accelerated to the high energies required to generate type III radio bursts or what role the coronal magnetic field plays in this process. Furthermore, there are inconsistencies between the observations and the models, which need to be resolved in order to gain a more complete understanding of the dynamics of the solar corona. Examples of these inconsistencies are the origin of the type III radio bursts and the discrepancy between the estimated plasma densities from the models and the observations. This paper aims to address these unresolved challenges by using new observations from LOFAR and PSP and models of the solar corona to study the physical mechanisms responsible for the generation of type III bursts. The data analysis includes a combination of radio spectroscopy and imaging techniques to study the frequency, temporal and spatial variations of the radio bursts.

This chapter is organized as follows. In Section ??, I describe the observations of type III radio bursts

made with LOFAR and PSP. In Section ?? I explain the data analysis and modeling techniques used to study these events. In Section ??, I present the results of our analysis, including an investigation of the potential physical mechanisms responsible for the generation of type III radio bursts and a comparison of the observations with models of the solar corona. Finally, in Section ??, I summarize our findings and discuss their implications.

3.2 Observations

A number of studies focused on observing the solar radio emissions during the second encounter of the PSP in late 2019 ?????. In this study, our primary emphasis is directed towards investigating a set of type III radio bursts that took place on April 3, 2019, during the time interval spanning from \sim 12:10 to 12:50 UT. This period coincided with the presence of two distinct active regions (ARs) on the Sun, denoted as AR12737 and AR12738. AR12737 was situated on the Sun’s near side at coordinates E12°N06°. Notably, this region had eight sunspots and exhibited a β magnetic configuration according to the Hale magnetic classification ?. On the other hand, AR12738 was positioned on the solar far side at coordinates E140°N02°. Due to its remote location, detailed observations of the magnetic configuration and activity within AR12738 were unattainable in this time frame.

I observed a group of intense type III radio bursts by four instruments (Wind/WAVES, PSP/FIELDS, STEREO-A/SWAVES, and LOFAR/LBA) via a regular survey. In Figure ??, I show the first type III burst within the time of this study as observed by the four instruments. By taking the second derivative of the light curve at a specific frequency channels, I determined the start time of the burst, which is denoted by the vertical red dashed line. The frequency bands used for obtaining the start time at each instrument are as follows: 6.97 MHz (Wind), 7.03 MHz (STEREO), 5.03 MHz (PSP), and 40.16 MHz (LOFAR).

I checked the relative orientations of the instruments with respect to Earth (Fig. ??). Since the PSP and STEREO spacecraft were almost aligned (close in an angular sense) with the Sun, the STEREO/EUVI image could be taken as what PSP would see (Fig. ??). Figure ?? shows how the solar disk looks like from the Earth perspective (using the SDO/AIA instrument) and from the eastern side where the PSP and STEREO were located at that time (using the STEREO/EUVI instrument). The right panel shows a closer view of AR12737 with the contours of the photospheric magnetic field obtained from the Helioseismic and Magnetic Imager (HMI) on board SDO. From the GOES-15/XRS and SDO/EVE instruments in the panels below, they also confirm that there is no flaring activity at that time.

The solar disk was quiet, including only one AR that is visible with no X-rays and no EUV transient emissions over this period. Nevertheless, the very sensitive LOFAR telescope detected a number of bursts close to noon. I checked PSP data, and I found bursts there as well. Meanwhile, from the EUVI and AIA images, I see that there are numerous small localized regions of relatively higher intensity (i.e., likely small-scale coronal brightenings spots or campfires; see ???). In the next subsections, I introduce the PSP and LOFAR instruments and their observations of the radio bursts.

3.2.1 PSP Observations

Parker Solar Probe (PSP) is a pioneering spacecraft with cutting-edge technologies, launched on August 12, 2018, aimed at helping to resolve key questions about solar corona and solar wind (?). To study the radio bursts, I used the level-2 data of the radio dynamic spectrum obtained from the FIELDS instrument suite (??), which can be downloaded from this website¹. The data file is in CDF format and the unit of the data values is converted from V^2/Hz to dB units using the formula

$$I_{dB} = 10 \times \log_{10}(I/10^{-16}) \quad (3.1)$$

The minimum power spectral density (PSD) of $10^{-16} V^2/Hz$ is used as a threshold for radio bursts according to ? for converting to decibels. Then, both the High-Frequency Receiver (HFR: 1.3 – 19.2 MHz) and the Low-Frequency Receive (LFR: 10.5 kHz – 1.7 MHz) data are combined into a single dynamic spectrum as shown in Figure ?? with a full frequency range between 10.5 kHz - 19.2 MHz. The mean intensity value at each timestep over the full frequency range is subtracted from each frequency channel to clean the spectrum and minimize the noise level.

¹PSP FIELDS data products: <http://research.ssl.berkeley.edu/data/psp/data/sci/fields/>




Figure 3.1: Radio dynamic spectra for a single burst obtained from multiple instruments. The top-left panel is from the LOFAR/LBA instrument, the top-right is from the PSP/FIELDS instrument, the bottom-left is from the STEREO/SWAVES instrument, and the bottom-right is from the Wind/WAVES. The vertical red dashed line denotes the start time of the burst.




Figure 3.2: Top view of the spacecraft positions in the ecliptic plane at 12:15 UT on April 3, 2019, with the Sun-Earth line as the reference point for longitude. The Earth's location is representative of the positions of LOFAR, Wind/WAVES, and GOES-15/XRS instruments. The spacecraft were connected back to the Sun by a 400 km/s reference Parker Spiral. The black arrow represents the longitude of AR12737 and the blue arrow represents the longitude of the AR12738. The gray dotted lines are the background Parker spiral field lines. The black dashed spiral shows the field line connected to the AR12737, and the blue dashed spiral is connected to the AR12738. The figure is generated using the Solar MAgnetic Connection Haus (Solar-MACH) tool (?).

3.2.2 LOFAR Observations

The LOw Frequency ARray (LOFAR) radio telescope (?) is a powerful tool for studying the Sun at low radio frequencies ranging between 10 and 240 MHz. Its high sensitivity and high time resolution have enabled the detection of various solar phenomena, including radio bursts and CMEs, and the study of dynamic processes in the solar atmosphere on timescales of milliseconds. The LOFAR dynamic spectrum from the beamformed radio observations is obtained by the Low-Band Antenna (LBA: 10 – 90 MHz) and can be downloaded from the LOFAR long-term archive (LTA)². The High-Band Antenna (HBA: 110 – 190 MHz) data are not available for that timeframe. For this day under study, the LOFAR data are available between 11:42 – 13:27 UT. To clean the spectrum, background subtraction is performed, which flattens the sensitivity (response) with the frequency of the LBA antennas. Basically, the mean spectrum along each frequency band is calculated and subtracted from the whole frequency band, the same applied to the PSP spectrum. This operation effectively removes the constant background from the spectrum. Then a Gaussian smoothing filter is applied to the spectrum using the `scipy.ndimage.gaussian_filter` function with a sigma value of 1.5, which helps to reduce noise and variations in the data. After that, the PSP and LOFAR spectra are combined together in a single plot within the same time interval. The bursts' signals observed by the PSP occur earlier than those at LOFAR. This is due to the fact that the PSP spacecraft is much closer to the Sun and hence it detects the radio emissions earlier than LOFAR because of the shorter travel time of radio signals from the Sun. Therefore, the PSP dynamic spectrum must be shifted with respect to the LOFAR observations based on a calculation of the relative time travel of the radio emission from the Sun to PSP and to LOFAR. In addition, the time cadence of the PSP observations changes according to its distance from the Sun. On that day, the PSP data cadence was 7 seconds, while LOFAR's is 1 second. Therefore, the LOFAR dynamic spectrum was down-sampled to 7 seconds to match the time resolution of the PSP. Figure ?? shows the resulting combined LOFAR-PSP spectrum on a logarithmic y-axis. The LOFAR LBA frequency ranges between 19.82 – 80.16 MHz and for the PSP is 10.55 kHz – 19.17 MHz.

In order to detect the type III radio bursts automatically from the combined dynamic spectrum, I applied the ? algorithm based on the probabilistic Hough transformation that detects vertical bright edges in images, within a certain degree of deviation from the vertical direction.

3.3 Methods

3.3.1 Imaging of radio sources

As part of our task, I developed an automated pipeline consisting of several modules that not only preprocessed and calibrated the LOFAR interferometric data to produce cleaned images of the Sun in the radio band (?), but also utilized the resulting data to find the trajectory of the radio sources and sample the magnetic field and plasma parameters at their respective locations through modeling and simulations in subsequent modules.

First, I ran the burst detection algorithm (?) ³ on the combined dynamic radio spectrum of LOFAR and PSP (Fig. ??) to find the characteristics of each type III burst. I converted the spectrum into a binary map to isolate the bursts from the background. Then I applied the Hough transformation to get line segments of the features. For each type III burst, the line segments are grouped together into one group. To account for the interplanetary component within radio dynamic spectra, I employed the Parker electron-density model (?) assuming a fundamental emission. This model enabled mapping between the time and frequency indices for each type III burst and subsequently converted electron densities into radial distances. Finally, a least-squares fitting method was applied to derive both the frequency drifts and the speed of the electron beams.

After this step, I did the same for the LOFAR dynamic spectrum only (Fig. ??) to find the (f, t) pairs for every type III burst. Then I took snapshot frequencies for each burst defined by a list of 60 central frequencies between ~ 20 – 80 MHz from LOFAR LTA for the interferometric imaging. I obtained the interferometric data from LOFAR core and remote stations at the snapshot frequencies for all type III bursts. I used the concurrent observations of the radio source Tau-A to calibrate the interferometric observations. For that, I used the default preprocessing pipeline (?), DP3 for preliminary processing and calibrating the measurement sets (MS). Finally I obtained the cleaned images of the radio sources

²LOFAR LTA: <https://lta.lofar.eu/>

³Detection algorithm repository: <https://github.com/peijin94/type3detect>

by using *w*-stacking clean (WSClean) algorithm (?) only at the time indices in the MS files that are equivalent to the snapshot frequencies.

Table 3.1: Characteristics of the type III bursts detected via the automatic algorithm from the combined spectrum.

Burst ID	Start Time (UT)	End Time (UT)	Start Frequency (MHz)	End Frequency (MHz)	Frequency Drift (MHz s ⁻¹)	Beam Speed (c)
1	12:18:45	12:22:42	76.44	1.57	0.892	0.044
2	12:34:05	12:36:31	41.24	0.86	0.241	0.119
3	12:34:40	12:34:56	54.44	26.54	3.992	0.046
4	12:37:14	12:38:09	66.03	10.02	4.006	0.046
5	12:38:17	12:40:54	76.92	1.57	0.77	0.066
6	12:39:34	12:40:11	78.86	11.93	3.192	0.062
7	12:40:28	12:40:40	45.34	22.9	3.21	0.067
8	12:41:39	12:43:06	78.21	2.13	1.555	0.093
9	12:43:53	12:44:15	59.07	42.13	2.424	0.013

After processing and cleaning the interferometric measurements of LOFAR, I explored the observations of each burst individually. Out of the 60 frequency bands in the LOFAR LTA, I chose 54 frequency bands that have unique integer numeric, between 19.92 - 80.08 MHz. For each burst, at each timestamp, the nearest frequency of the fit model to the list of chosen frequencies is picked as the snapshot frequency at that particular timestamp. This process was repeated for all the 16 type III bursts detected in the LOFAR dynamic spectrum in order to obtain snapshot images for each type III burst (Fig. ??). For each type III burst, I applied persistence imaging in order to create a continuous display of the radio emissions (?).

Persistence imaging enables the creation of a clearer and more informative image. In the context of a time-ordered series of images, a method of persisting pixel values can be employed as follows: for each image, compare the value of each pixel to its corresponding value in the previous persistence image in the series. If the pixel value in the current image is brighter than its corresponding pixel in the previous image, replace the previous value with the current one; otherwise, retain the previous value. This process generates a new image, referred to as the current persistence image, which serves as the basis for the subsequent evaluation of the next image in the series. This evaluation involves a pixel-by-pixel comparison between the current image and its associated persistence image, allowing for the identification of any changes or patterns that may have occurred over time. The mathematical background is explained in Appendix ??.

In order to estimate the locations of the type III sources in 3D space, I combined observations with modeling. I used magnetogram data from the Global Oscillation Network Group project (GONG) (?). I constructed a grid of footpoints on the GONG map over two longitudinal belts around the two active regions AR12737 and AR12738, which are the two potential candidates source regions for the group of type III bursts under study. These points are used as the seed points for tracing the coronal magnetic field lines using pfsspy python package⁴, which is a robust implementation in python of the PFSS model developed by ?. Using the major and minor axes of the beam size, I estimated the radius of the radio source using Equation ??, which was used to approximate the source size. Since I already obtained the (*x*, *y*) positions of the type III sources in the plane of the sky (POS) through LOFAR observations, now it is necessary to determine their corresponding *z* position to have an overall understanding of their spatial distribution. Therefore, I employed ?'s approach here, assuming that the type III bursts were from harmonic emission. Firstly, I found the radial distance of the radio source from the Sun in the POS (Eq. ??). Secondly, I calculated the sources' radial distance (*r*_{model}) using the 2.5×Newkirk electron-density model (??). The 2.5 fold factor is taken to incorporate the effects of scattering and overdensity (streamers) beyond the nominal Newkirk quiet Sun model. The MAS model results (Fig. ??) show streamers above the eastern limb, supporting the inclusion of such a factor. Lastly, I estimated the *z* location of the type III sources (Eq. ??). I proceeded with the +*z* solution because the theory precludes emission behind POS in this region of high-density gradients (i.e., the emission would be absorbed by passing through the high-density regions of the corona). More details are explained in Appendix ??.

$$r_{\text{source}} = \sqrt{(b_{\text{major}}^2 + b_{\text{minor}}^2)}, \quad (3.2)$$

$$r_{\text{pos}} = \sqrt{(x^2 + y^2)}, \quad (3.3)$$

⁴Pfsspy tool: <https://pfsspy.readthedocs.io/>

$$z = \sqrt{(r_{model}^2 - r_{pos}^2)}. \quad (3.4)$$

The result of the deprojection of the type III sources for the sixth burst are shown in Figure ?? with 70% contours made for ten frequencies on the extrapolated magnetic field lines. The red dashed line is a spline fitting curve that represents the trajectory of the centroids of the radio sources. The black arrow points toward the Earth's line of sight (LOS). It is worth to mention that the axes direction in the POS of LOFAR images are different in the 3D space. The (x, y) coordinates in the POS are translated into (y, z) in the 3D space, and z in the POS is translated into x in the 3D space.

3.3.2 Modeling

To explore the characteristics of the coronal plasma environment during the studied events, I used the Predictive Science Inc. (PSI) standard coronal solutions from magnetohydrodynamic (MHD) simulations originating from the MAS code (?). The data are available on the PSI's data archive⁵. I obtained the PSI MAS coronal solution (a thermodynamic-with-heating MHD model) on April 3, 2019, at 12:00 UT, with the following simulation result ID⁶. Initially, I calculated the angle between the burst's source radial vector and the LOS. Moreover, I calculated the complement angle, which is the separation angle between the burst's radial vector and POS from the Earth's perspective. Subsequently, I utilized the complement angle to derive the Carrington longitude (?), facilitating the extraction of a longitudinal segment from the MAS datacube, as if it were in the POS. Following this, the selected data slice was fed into the FORWARD model, a toolset responsible for generating synthetic coronal maps of observable quantities describing the plasma state. For extracting the longitudinal slices from the MAS data, I utilized the psipy python package⁷. The MAS datacube is specifically defined on a spherical grid and represents a steady-state MHD model. Owing to the inherent attributes of this datacube, the utilization of the FORWARD toolset proves more practical and advantageous for our objective. In Figure ?? I show the first radio contour of the sixth type III burst on top of the equivalent 2D maps for six plasma parameters, as an example. The plasma parameters are, from left to right and from the top to bottom: plasma density, plasma temperature, magnetic field strength, plasma beta parameter, the total plasma pressure, and the Alfvén speed. By taking the value of these physical plasma quantities at the centroids' coordinates of the type III sources at each frequency band, I obtained estimates of local plasma conditions shown in Figure ?? for the sixth type III burst, as an example.

3.4 Results and discussion

3.4.1 Detection and characterization of type III radio bursts

I found that the radio waves arrived at STEREO one minute before they arrived at Wind (Fig. ??). However, the difference between the $+z$ and $-z$ positions of the burst this close to the Sun in terms of light travel time is ~ 10 seconds ($\sim 4 R_\odot$), which is within the time resolution of the observations (1 min time resolution). Thus, I cannot confidently conclude whether the emission arrived at one spacecraft first and the other second.

Figure ?? shows the combined dynamic spectrum from both LOFAR and PSP. The free parameters of the auto-detection algorithm do not have the same values as for detection the type III bursts in the LOFAR spectrum alone. Upon visual examination, I observed that the detection algorithm effectively identified type III bursts in the LOFAR dynamic spectrum (Fig. ??), but it had limitations in terms of detecting type III bursts in the combined spectrum of the LOFAR and PSP, as well as missing segments of the detected bursts and a few bursts entirely. This could be due to the increased frequency drift and dispersion of the radio bursts at lower frequencies, which made it a challenging task for the detection algorithm. I captured nine type III bursts from the combined dynamic spectrum and their characteristics are reported in Table ???. However, the detection algorithm performed better on the LOFAR dynamic spectrum only and I traced 16 type III bursts.

3.4.2 Imaging of radio emission sources

Figure ?? shows the persistence imaging for the 16 type III bursts in the LOFAR dynamic spectrum (Fig. ??). The observation frequencies and timestamps of the snapshot images used to produce the

⁵Predictive Science Inc.: <https://www.predsci.com/mhdweb/home.php>

⁶Simulation result ID: hmi_med-cor-thermo2-std01_med-hel-poly-std01

⁷PsiPy repository: <https://github.com/predsci/PsiPy>

persistence image are shown at the top-right corner of each image. From a visual inspection of Figure ??, it seems that all the type III emissions originated from the same quadrant in the images (south-east direction on the solar disk), although there was no active region presented at that location except for a single active region nearby the central meridian (Fig. ??). Based on the imaging data presented in Figure ??, I chose one representative type III burst (No. 6) for a single-burst analysis in this paper, as it shares similarities in extent and location with other bursts.

To determine the spatial connection between the sources of radio emissions and the coronal magnetic field, a three-dimensional (3D) projection of the radio source contours onto the extrapolated coronal magnetic field via the PFSS model was employed (Fig. ??). The result indicates a discernible south-eastward propagation of the radio sources relative to the Earth's perspective, with no open field line crossing the radio sources. In Figure ??, I performed an extrapolation only over the two active regions presented on the solar surface at that time. However, when I extrapolated the magnetic field over the entire solar surface, I noticed that the radio sources are aligned with the lower part of large-scale closed field lines and are placed onto the open field lines emanating from the southern coronal hole. No open field lines crossing the radio sources are observed. I note that the PFSS modeling is limited by the fact that AR12738 is behind the limb on April 3 as observed from Earth. Consequently, the magnetic data available to us could be around two weeks old or more. This might limit the reliability of PFSS extrapolation for that region during that specific timeframe.

From Figure ??, the results suggest several potential origins of these type III radio emissions: 1) they could be triggered in a closed-field lines structure such as large-scale coronal loops, given that the radio sources are aligned to closed-field lines geometry in the southern hemisphere; 2) they could be triggered by electron beams that are accelerated from an open-field active region (?). However, from the PFSS model, I found no evidence for magnetic connectivity from both ARs on the Sun at that time; 3) they may result from electron beams that are accelerated in the corona due to expanding magnetic fields from plasma upflows in the active region (??). Our findings indicate a notable inverse relationship between imaging quality and the level of solar radio emission brightness (e.g., for type III bursts 10 and 13, for instance). This observation is due to the leakage of solar radio emission into the side lobes of the calibrator beam, which disrupts the accuracy of calibration solutions.

3.4.3 Plasma diagnostics and magnetic field analysis

Considering the observed alignment of radio sources in Figure ?? and the case depicted in Figure ??, it becomes evident that radio sources at higher frequencies (indicating proximity to the Sun) align with a streamer-like structure near the equator within the coronal model. This structure is characterized by elevated plasma beta, reduced coronal temperature, and diminished Alfvén speed. The coronal plasma density was relatively homogeneous with no prominent structures, probably due to the model resolution.

The location of radio sources of all the bursts were in the same quadrant as seen from Earth. Therefore, I assumed that the former description applies for all bursts. I also found that the radio sources were confined between the equatorial sheet and the southern coronal hole and moving along that boundary. Figure ?? shows the variability of the coronal plasma quantities at the radio sources' centroids, taken from FORWARD maps in Figures ??, at different frequencies for the sixth burst. To estimate the error bars, I initialized random centroids, within the limits of the 70%-contours of the radio emissions, to sample the plasma quantities at those locations. Then the standard error (SE) is calculated using Equation ??, where σ is the standard deviation, and n is the number of points.

$$SE = \frac{\sigma}{\sqrt{n}} \quad (3.5)$$

The coronal temperature was increasing with radial distance, which implies there may have been some heating locally. The behavior of the coronal magnetic field, the plasma total dynamic pressure, and the Alfvén speed were decreasing over distance as expected. Finally the value of plasma beta parameter started increasing sharply around 40 MHz, which implies that the plasma pressure became more dominant than the magnetic pressure around that distance from the Sun (for a $2.5 \times$ Newkirk model, it is $1.89 R_{\odot}$ assuming a fundamental emission, or $2.57 R_{\odot}$ assuming a harmonic emission).

The top-left panel of Figure ?? shows a comparison between the density profiles of the MAS model, the $2.5 \times$ Newkirk model and the theoretical expected density profiles under the fundamental and harmonic assumptions. Although the Newkirk density model provided a useful approximation for determining the height of radio sources in the corona, it is not entirely accurate due to a number of its underlying assumptions; for instance, the assumption of a steady state and the spherical symmetry of the corona,

which do not always apply. Therefore, I tried to use the MAS density values to estimate the depth along the LOS of the radio source, since it is supposed to give a more realistic result. I found that the plasma density obtained from the MAS and FORWARD modeling results were significantly lower compared with the $2.5 \times$ Newkirk density model and the theoretical expected density obtained from the classical relation in Equation ??, where f_p is the plasma frequency (in MHz) and n_e is the electron density (in cm^{-3}):

$$n_e = \left(\frac{f_p}{8.98 \times 10^{-3}} \right)^2 \quad (3.6)$$

The required density from the fitted Newkirk model is much higher (about ten times) than what is obtained from the MAS model, even after accounting for the $2.5 \times$ enhancement already applied to the standard Newkirk model. This implies the discrepancy cannot be fully explained by the density enhancement factor alone. Furthermore, the imaging places the radio sources near a streamer which is an overdense region in the MAS model, so it seems unlikely the source's apparent location in the model is wrongly attached to a less dense feature, as there are not denser options available. The apparent source positions from the imaging are likely too high, possibly due to scattering effects (??), which could lead to fitting an overly dense Newkirk model. Another potential explanation is that there could be a stealth CME that pushed the coronal magnetic field outward, allowing the plasma to follow it to be perceived as having a higher density than expected, and there was not enough time for the magnetic field relaxation to occur (private communication with J. Magdalenić). However, scattering alone does not seem to fully explain the large density discrepancy. While further investigation is certainly needed regarding scattering and propagation effects on the radio waves, it is interesting to report this significant discrepancy between the model and observations, as it highlights limitations in the current modeling and suggests the need for additional physics to properly characterize the density distribution. Resolving this discrepancy could lead to important insights into the true nature of the corona.

3.5 Summary and conclusions

In this work I analyse the characteristics of a series of type III bursts that occurred on April 3, 2019, during the second near-Sun encounter period of PSP. The bursts were observed in dynamic spectra taken with the PSP/FIELDS (2.6 kHz – 19 MHz) instrument, as well as in interferometric imaging with the LOFAR (20 – 80 MHz) ground-based telescope, as part of a coordinated observing campaign. The series of 16 separate weak bursts were observed over the span of ~ 20 minutes, during an otherwise relatively quiet period. The solar disk as observed from Earth was dominated by a single active region near its center. I combined the dynamic spectra for the LOFAR frequency range and the PSP frequency range to study the solar radio emissions within the frequency range of 2.6 kHz – 80 MHz.

For the study, I developed a semi-automated pipeline, which allowed us to obtain the exact times and frequencies of the bursts. These I used to align the PSP to the LOFAR observations and to generate interferometric images between 20 and 80 MHz. I performed data pre-processing of the PSP and LOFAR dynamic spectra to resample and shift the data based on the relative location of the spacecraft with respect to the Sun and Earth, and found an excellent temporal match between the two sets of observations. Thus I automatically traced the type III bursts in the dynamic spectra algorithmically and estimated frequency drift and the electron beam speeds. I found that frequency drifts remained relatively uniform between the high-frequency (LOFAR) and low-frequency (PSP) observations, as well as among the bursts, suggesting that they are related.

In addition, I imaged the type III emission at multiple frequency bands using the interferometric observations from LOFAR to determine the locations of the sources in the solar corona. The type III emissions observed were all found to occur in the same general region off the southeast limb of the Sun, leading us to conclude that they shared a single source of electron beams low in the corona. The potential origins of these emissions are varied and include: 1) small-scale impulsive events such as nanoflares (??); 2) plasma upflows from the active region (?); 3) coronal closed-loop structures (?); 4) electron beams accelerated from interchange reconnection (?); 5) high-frequency Alfvén waves and/or magnetic reconnection in the outer corona (??).

Our magnetic extrapolation shows that there is no open potential field to either AR12737 or AR12738, which is consistent with ?. Our findings are in line with the conclusions of ?, who proposed that the likely origin of these type III bursts is the AR12737 region. The type III radio bursts in ? occurred between April 1 and 4, are temporally aligned with the emergence of AR12737 near the eastern limb of the solar disk.

While potential field source surface models provide valuable insight into the large-scale magnetic topology, their reliability decreases near active regions where the field can deviate significantly from a potential configuration. Therefore, the lack of open field connectivity directly to AR12737 suggested by the PFSS model should be viewed with some caution.

This work (?) complements those results by precisely locating the burst sources in the middle of the corona. I used the Newkirk density model to estimate the height of the radio sources from the Sun of one of the type III bursts as representative of all. Combining this with PFSS magnetic modeling, I found good agreement between the centroids of the radio sources and the location of the southern open field lines in the corona, which would be required to produce radio emissions at interplanetary wavelengths in general. On the other hand, this location does not seem to be well connected to the AR itself, according to the PFSS model.

I attempted to correct the radial distance of the radio sources from the Sun by replacing the Newkirk model with more realistic MHD results from the MAS model, but I found that there is a significant discrepancy between the Newkirk model profile fitted to the observations and the MAS density. This could result from scattering lensing the apparent burst location to a higher altitude, thus, overestimating the height of radio sources in the corona. The presence of type III radio sources at relatively high distances in the corona, with plasma density higher than expected from the MAS model, suggests that there may be missing information in the modeling. One possibility is the existence of a stealth CME that pushed the coronal magnetic field outward, causing the plasma to appear denser than expected (see ?) — or other non-obvious changes in large-scale coronal magnetic topology. These findings demonstrate that scattering and propagation effects play a significant role in determining the location and directionality of solar radio bursts (???). Therefore, the discrepancy between the observed and modeled density profiles could potentially be attributed to scattering and lensing effects that make the radio sources appear higher in the corona than their true location. Further investigation is required to disentangle these effects from limitations in the density models themselves. Overall, accounting for scattering and refraction will likely lead to improved modeling of the corona and solar radio bursts. In future work, I will also employ the time delay of arrival (TDoA) technique (?) to estimate the radio burst source positions from multi-instrument observations and compare that with the current methodology in this paper. Solar Orbiter observations shall also be included.

High-fidelity interferometric radio imaging in metric-decametric wavelengths provides a powerful method to characterize solar eruptive events. It is also becoming increasingly important for studying relatively quiet periods, during which there may be elevated levels of in situ particle fluxes. The ability to observe and image faint radio bursts such as those presented in this work, which may be related to episodes of reconnection on the solar surface, and potentially to episodes of solar wind release, is a testament to LOFAR’s power as a space weather instrument. In a future work, I will automate and use our method for studying hundreds of faint bursts observed with LOFAR and will investigate their relation to small-scale activity on the solar surface.

Through a novel combination between the LOFAR imaging and MAS model results, I found that the type III radio bursts experienced a weakening background magnetic field, decreasing solar wind dynamic pressure and Alfvén speed, increasing plasma beta and coronal temperature, and plasma rarefaction. The radio sources appeared at larger radial distances than the models predicted, which suggests scattering and density fluctuations are important when attempting to interpret the actual burst trajectory. The discrepancies between the observed and modeled radial distances of the radio sources suggest refinements are needed in the models to fully explain the radio imaging and modeling results. Overall, comparing the LOFAR imaging and MAS modeling for these type III bursts motivates further analysis on additional radio bursts to improve our understanding of the physical conditions that influence the propagation of radio emissions in the corona.




Figure 3.3: Exploring the X-ray and extreme ultraviolet (EUV) emissions from the Sun. The top panel showcases a cutout region of the SDO/AIA 193 Å image of the solar disk along with the STEREO-A EUVI 195 Å point of view. The white curve is the limb of the solar disk as seen by AIA from the right side. The red and blue colors are the contours of the line-of-sight magnetogram from the SDO/HMI instrument. The levels are (50, 100, 150, 300, 500, 1000) Gauss. The middle panel shows the X-ray flux from the GOES-14 spacecraft shows minimum activity. The bottom panel shows the time series of the ESP Quad band from the SDO/EVE instrument, which shows the solar irradiance in the extreme ultraviolet (EUV) band.




Figure 3.4: Automatic detection of type III radio bursts from the combined radio dynamic spectrum of LOFAR and PSP instruments. The dashed horizontal lines separates the LOFAR frequency range (top) and the PSP frequency range (bottom).




Figure 3.5: Automatic detection of type III bursts observed by LOFAR. The red symbols along the fit lines are the (f, t) coordinates of the image snapshots shown in Figure ??.




Figure 3.6: Persistence imaging for the 16 type III bursts detected in the LOFAR dynamic spectrum. The label shows the observation frequencies in MHz and times in (minutes:seconds from 12:00:00 UT). Here, the color coding is not absolute, but rather each panel has its own color code.




Figure 3.7: Different viewing angles for the deprojection of the radio sources of the sixth burst using the $2.5 \times$ Newkirk electron-density model on the PFSS solution. The black arrow points toward the Earth's LOS. The yz plane is the POS as seen from the Earth. The red dashed line is a spline curve fit for the sources' centroids. The red, black, and blue curves are the open northern, closed, and open southern field lines, respectively. The opacity of the closed field lines is decreased for better visualization.




Figure 3.8: Synthesized maps of plasma parameters obtained using the FORWARD toolset, with the 70%-contour of radio emission of the sixth burst at the first timestamp (12:34:06.8 UT) at the frequency of 72.26 MHz depicted on top of the 2D POS cuts. The left column represents, from top to bottom, plasma density, magnetic field, and the total plasma dynamic pressure. The right column represents, from top to bottom, the temperature, plasma beta, and the Alfvén speed.




Figure 3.9: Coronal plasma parameters sampled from the 2D maps by the source centroids. The top panel shows (from left to right) the plasma density profiles from the MAS model, $2.5 \times$ Newkirk model, and the theoretical densities under the fundamental and harmonic assumptions, plasma temperature, and magnetic field. The bottom panel shows, from left to right, the total plasma dynamic pressure, Alfvén speed, and plasma beta. The x-axis is inverted to demonstrate a progression of increasing radial distance from the Sun as the observer moves towards the right.

Chapter 4

Modeling and Forecasting of Solar Energetic Protons (SEP)

This chapter has two parts. In the first part, I describe a modeling study that we conducted on energetic proton acceleration and propagation from the solar corona to 1 AU. For this modeling, we employ the physics-based approach utilized in the first chapter, including the 3D coronal models and a 3D MAS-MHD model run. With these models, we use the EPREM model to find the fluxes and spectra of energetic protons at 1 AU. We then compare the modeling results with in-situ measurements. In the second part, I describe a deep learning approach that I developed to forecast the integral flux of energetic protons in three energy channels across three different forecasting horizons.

4.1 Introduction

Coronal Mass Ejections (CME) are a major source of accelerated solar energetic particles (SEPs), exceeding the thermal plasma in ions and electrons (?). While flares contribute SEPs, CME-driven shocks play a key role through compression and acceleration above 100 MeV/nucleon, even in early stages below $10 R_\odot$ (??). Consequently, research has focused on CME and shock dynamics in the corona using multi-wavelength observations. Recent studies of CMEs have utilized Extreme Ultra-Violet (EUV) and radio wavelength observations to uncover nuances beyond the prominence of these solar eruptions in visible light (???). Observation of the solar corona in EUV wavelengths allows the initiation of CMEs to be imaged, through the use of instrumentation such as that on the Solar Dynamics Observatory (??, SDO). CME propagation can generate shock waves visible as EUV waves or bright fronts in the corona (??).

Extending prior work, ? explored early SEP production using coronal bright front (CBF) events and a model of Diffusive Shock Acceleration (DSA) of ?. This revealed event-to-event variations and temporal evolution in acceleration efficiency strongly influenced by diverse coronal environments through which shock waves propagate. This approach demonstrates the use of computationally complex physics-based models to predict SEP flux at multiple energies.

Several models are available, or under development, for forecasting SEP, which use diverse approaches and serve different objectives. These models comprise computationally complex physics-based models, quick and simple empirical models, Machine Learning (ML)-based models, and hybrid models that combine different approaches and produce different types of outputs, including deterministic, probabilistic, categorical, and binary. Deterministic models always generate the same output without any randomness or stochastic components, such as predicting the SEP flux at a specific moment or the arrival time of SEP. On the other hand, probabilistic models provide a probability value that reflects the likelihood of an SEP event occurring. However, replicating SEP fluxes at a specific time is still a significant challenge for current models.

An excellent review on SEP models and predictive efforts was recently published by ?, which summarizes the majority of the existing models. For instance, ? introduced the Probabilistic Solar Particle Event Forecasting (PROSPER) model, which is incorporated into the Advanced Solar Particle Event Casting System (ASPECS)¹. The PROSPER model utilizes a Bayesian approach and data-driven methodology to probabilistically predict SEP events for 3 integral energy channels >10 , >30 , and >100 MeV. The model's validation results indicate that the solar flare and CME modules have hit rates of 90% and

¹ASPECS: <http://phobos-srv.space.noa.gr/>

100%, respectively, while the combined flare and CME module has a hit rate of 100%. ? developed an empirical model to predict the peak intensity and spectra of SEP at 1 AU between 10 and 130 MeV, using data from multiple spacecraft. The model is tested on 20 SEP events and shows good agreement with observed values. The spatial distribution of SEP intensities was reconstructed successfully, and they found a correlation between SEP intensities and CME speed.

? extended the Particle Acceleration and Transport in the Heliosphere (PATH) model to study particle acceleration and transport at CME-driven shocks. They showed that the model can be used to obtain simultaneous calculations of SEP characteristics such as time-intensity profiles, instantaneous particle spectra, and particle pitch angle distributions at multiple heliospheric locations. Overall, results resemble closely those observed in situ near the Earth but also yield results at other places of interest, such as Mars, making it of particular interest to Mars missions. SPREAdFAST (??) is a physics-based, data-driven framework that utilizes EUV observations and models to simulate SEP fluxes at 1 AU and to estimate energetic particle acceleration and transport to various locations in the inner heliosphere. It generates time-dependent histograms and movies distributing them through an online catalog. The accuracy and efficiency of the model were encouraging, but the highest energy fluxes showed disagreement with in situ observations by the SOHO/ERNE instrument. However, the framework has great potential for space weather science and forecasting.

In ?, they used neural networks to provide probabilities for the occurrence of SEP based on soft X-rays data from 1988 to 2013. They obtained >85% for correct SEP occurrence predictions and >92% for correct no-SEP predictions. ? described a consistent approach to making a binary prediction of SEP events using ML and conventional statistical techniques. The study evaluated various ML models and concluded that random forests could be the best approach for an optimal sample comprising both flares and CMEs. The most important features for identifying SEP were found to be the CME speed, width, and flare soft X-ray fluence. ? employed ML techniques to anticipate the occurrence of a SEP event in an active region that generates flares. They utilized the Space-Weather MDI Active Region Patches (SMARP) dataset, which comprises observations of solar magnetograms between June 1996 and August 2010. The SMARP dataset had a success rate of 72% in accurately predicting whether an active region that produces a flare would result in a SEP event. Moreover, it provided a competitive lead time of 55.3 min in forecasting SEP events.

? introduced the Space Radiation Intelligence System (SPRINTS), a technology that uses pre- and post-event data to forecast solar-driven events such as SEP. It integrates automatic detections and ML to produce forecasts. Results show that SPRINTS can predict SEP with a 56% probability of detection and 34% false alarm rate. Nevertheless, the HESPERIA REleASE tools provide real-time predictions of the proton flux at L1 by using near-relativistic electrons as a warning for the later arrival of protons and have been set to operation (?). Historical data analysis indicates high prediction accuracy, with a low false alarm rate of approximately 30% and a high probability of detection of 63% (?).

Forecasting SEP is a critical task that serves operational needs and provides insight into the broader field of space weather science and heliophysics. As emphasized in previous works, a high precision forecasting model is urgently required to predict SEP flux within a period of time, given the risks associated with these events. This highlights the critical requirement for a dependable forecasting system that can mitigate the risks associated with SEP.

Scientists have been using physics-based and empirical models for decades to forecast SEP. However, these models have certain limitations. Physics-based models require accurate input data and underlying physical assumptions. In addition, the complexity of the physics involved and incorrect parameters may introduce uncertainties that can lead to inaccurate predictions. On the other hand, empirical models rely on historical data to make predictions. While they can be accurate sometimes, they may be unable to account for changes in physical conditions related to the acceleration and propagation of SEP, which can influence prediction accuracy. ML models, however, provide a different approach to SEP forecasting. These models can analyze vast amounts of data, learning patterns from the data that are used, and connections that may not be obvious to experts. Additionally, ML models can adapt to changes in underlying physical conditions, resulting in more accurate predictions as more data is collected; they also provide relatively rapid forecasts, which allows for incorporation into a real-time forecasting workflow.

In the upcoming sections, I will explore the limitations in accuracy that arise from dealing with an imbalanced dataset and low-resolution data. Specifically, the presence of intrinsic outliers in the time series data pertaining to SEP flux poses a significant challenge in modeling. These outliers correspond to occurrences of SEP events and, consequently, have an impact on the accuracy of predictions. Notably, they often lead to an underestimation of the SEP fluxes, primarily due to the predominance of relatively low values throughout the majority of the time interval.

In the first part, we extend the work in Chapter 2 on the kinematics of CBFs and expand on previous relevant investigations by modeling CBF-related shocks and particle acceleration up to 10 solar radii. Our modeling approach incorporates coupling to a numerical model of particle transport throughout the heliosphere, with validation against in-situ spacecraft measurements. Our study implements, for the first time, an extensive physics-based model linking CME-driven shock acceleration with the propagation of SEPs from the Sun to Earth. In the second part, I present advanced deep learning models to forecast the daily integral flux of SEP over a 3-day forecasting window by using bi-directional long short-term memory (BiLSTM) neural networks, for 3 energy channels (>10 , >30 , and >60 MeV). Our models can forecast the time-dependent development of SEP events in different energy domains, which can be used to model the space radiation profiles using frameworks such as BRYNTRN [?](#) and GEANT4 [\(?\)](#).

4.2 Early-Stage SEP Acceleration by CME-Driven Shocks

4.2.1 Overview

The purpose of the SPREAdFAST (Solar Particle Radiation Environment Analysis and Forecasting-Acceleration and Scattering Transport) project is to model and analyze the particle fluxes from the Sun to Earth, specifically focusing on solar energetic particle (SEP) events. The project combines detailed observations of coronal bright fronts (CBFs) with modeling of the coronal plasma and the resulting SEP production and interplanetary transport. The SPREAdFAST framework utilizes physics-based modeling to simulate the evolution of the plasma upstream of the coronal shock associated with CBFs and the subsequent acceleration and transport of protons from the Sun to 1 AU. It incorporates various components such as EUV observations, shock dynamics, particle acceleration, and interplanetary transport. The project aims to provide a better understanding of the processes involved in SEP events and improve forecasting capabilities for these events. By modeling a large number of events and comparing the model results with in situ observations, the SPREAdFAST project contributes to the advancement of Sun-to-Earth physics-based modeling of SEP acceleration and transport. Overall, the SPREAdFAST project is a comprehensive effort to study and simulate the complex phenomena associated with SEP events, with the goal of enhancing our knowledge and predictive capabilities in this field.

The SPREAdFAST framework includes the following components:

1. CBF Kinematics and Geometric Modeling: This component characterizes the kinematics of Coronal Bright Fronts (CBFs) using observations from the Atmospheric Imaging Assembly (AIA) instrument. It estimates the CBF kinematics, including the front, peak, and back edge positions over time, as well as the mean intensity and thickness of the CBFs.
2. Coronal Shock and Particle Acceleration Modeling: This component models the evolution of the plasma immediately upstream of the coronal shock associated with CBFs. It incorporates the physics of coronal shock waves and the process of particle acceleration through diffusive shock acceleration.
3. Interplanetary Particle Transport Modeling: This component simulates the transport of accelerated particles from the corona to 1 astronomical unit (1 au), which is the distance between the Sun and Earth. It takes into account the interplanetary magnetic field and other factors that influence particle propagation.
4. Comparison with Observations: The framework compares the modeled particle fluxes and fluences at 1 au with observations from instruments like the Solar and Heliospheric Observatory (SOHO) and the Energetic and Relativistic Nuclei and Electron (ERNE) instrument. It evaluates the accuracy of the model predictions by calculating metrics such as the Mean Squared Logarithmic Error (MSLE).

Overall, the SPREAdFAST framework combines detailed observations, physics-based modeling of coronal shocks and particle acceleration, and interplanetary transport modeling to analyze and forecast Solar Energetic Particle (SEP) events from the Sun to Earth.

4.2.2 Data Analysis and Methodology

We used a combination of telescopic observations and dynamic physical models to simulate the acceleration of solar energetic particles (SEPs) in global coronal shock events. We first observed off-limb coronal

bright fronts (CBF) and studied their interaction with the coronal plasma using synoptic magnetohydrodynamic (MHD) simulations. Based on these observations and simulations, we then employed an analytical diffusive shock acceleration (DSA) model to simulate the SEP acceleration. The simulated fluxes obtained from the DSA model were used as time-dependent inner boundary conditions for modeling the particle transport to 1 AU. This approach allowed us to study the early-stage acceleration and transport of SEPs from the Sun to 1 AU.

The criteria used to select the events for analysis in the study of the SPREAdFAST framework were as follows:

1. Proton events in the energy range of 17-22 MeV observed by the SOHO/ERNE instrument from 2010 to 2017 were initially identified.
2. Events without identified flares and coronal mass ejections (CMEs) and without EUV waves preceding the SEP event were excluded.
3. Events without EUV waves or no EUV data, even if they had identified flares/CMEs, were also excluded.
4. Uncertain EUV waves that were not relevant to the specific solar eruption were dropped.
5. Events with measurable near-limb or off-limb Coronal Bright Fronts (CBFs) that could be analyzed with the SPREAdFAST framework were selected.

In total, 62 events met the selection criteria and were included in the analysis.

The kinematics of Coronal Bright Fronts (CBFs) are characterized using the methodology of the Coronal Analysis of SHocks and Waves (CASHeW) framework. This framework estimates the CBF kinematics by following the leading edge of the front on consecutive images. It calculates the kinematics of the front, peak, and back edge of the CBFs over time, allowing for the estimation of their time-dependent mean intensity and thickness. The kinematics are determined using time-height maps (J-maps) generated with the CASHeW code for each event. The radial and lateral wave front positions are measured in these J-maps, providing information on the radial and lateral positions, speeds, accelerations, mean wave intensities, and wave thickness of the CBFs.

The methodology used to characterize the kinematics of Coronal Bright Fronts (CBFs) is based on the Coronal Analysis of SHocks and Waves (CASHeW) framework. This framework involves analyzing observations from the Atmospheric Imaging Assembly (AIA) instrument on board the Solar Dynamics Observatory (SDO). The kinematics of CBFs are determined by tracking the leading edge of the front on consecutive images. Time-height maps, also known as J-maps, are created by stacking columns of pixels in a desired direction from a solar image. The shape of the track on these J-maps depends on the direction and speed of the CBF. The CASHeW code identifies the radial and lateral wave front positions over time in the J-maps, allowing for the estimation of the CBF kinematics, including speeds, accelerations, mean wave intensities, and wave thickness. A three-dimensional geometric model, known as the Synthetic Shock Model (S2M), is then created based on the measured front positions, which describes the shock surface at regular intervals. This model is propagated through the solar corona using a synoptic coronal magnetohydrodynamic (MHD) model, providing information on the relevant parameters for coronal shock acceleration of solar energetic particles (SEPs).

In the SPREAdFAST DSA model, the shock-crossing field lines are modeled by dividing the shock surface into three regions: the "nose" of the shock model, which consists of model points on the spheroidal cap; and two flanks or zones, divided by a plane parallel to the Sun-Earth line. The plasma parameters at the points on these three surfaces are examined separately. The model calculates the proton acceleration along these shock-crossing field lines based on time-dependent estimates of shock speed, density jump ratio, magnetic field strength, and shock angle. The model solves for the coronal charged particle acceleration by large-scale CME-driven shocks and provides time-dependent distribution function spectra or fluxes as output.

The method used to compare the modeled and observed proton fluences is by analyzing scatter plots of the fitted power indices of the proton fluences from the EPREM model and the ERNE observations. The power law indices are compared between the two sets, and the onset hours for the proton events are also compared. The comparison helps evaluate the performance of the modeling framework in predicting the proton fluxes. Additionally, histograms and Mean Squared Logarithmic Error (MSLE) are used to assess the agreement between the modeled and observed fluence spectra and onset times.

4.2.3 Results and Discussions

The study's findings have important implications for understanding and predicting solar particle radiation. By modeling the dynamics of shock waves and particle acceleration in the solar corona, the study provides valuable insights into the factors that influence the efficiency of particle acceleration. The results highlight the significant role of the coronal environment in shaping the acceleration and transport of solar energetic particles (SEPs) from the Sun to Earth. One key implication is that the overlying coronal structure and the particle energy play a crucial role in determining where SEPs are produced during CME-driven shock and compressive waves. The study shows that the large gradients in plasma parameters between neighboring streamers, quiet-Sun areas, and coronal holes lead to continuous changes in the acceleration process. This knowledge can help improve our understanding of the spatial distribution of SEPs and their energy dependence. Furthermore, the study's findings contribute to the development of physics-based models for forecasting SEP events. The SPREADFAST framework used in the study demonstrates the potential for accurately simulating the evolution of SEPs from the Sun to 1 AU. This framework can be further refined and utilized for early-stage forecasting of SEP events, providing valuable information for space weather prediction and mitigation efforts. Overall, the study enhances our understanding of the complex processes involved in solar particle radiation and provides a foundation for improving our ability to predict and mitigate the impacts of these events on space weather.

The main discrepancies between the modeled and observed fluxes in the study are primarily seen at higher energies. Above 15 MeV, there is a discrepancy in the time profile, with the observed proton fluxes rising approximately 1 hour before the simulation. Additionally, the fluxes at the highest energies show the most disagreement, mainly due to the slope of the increase and the onset times. These discrepancies indicate the need for further improvements and refinements in the modeling framework to better match the observations.

4.3 Solar Proton Flux Forecasting with Deep Learning Models

4.3.1 Data preparation

In this section, I describe the physical quantities, the types of inputs and their sources, as well as the outputs I am forecasting. Some of the technical terms used in this study are explained further in the appendices.

In order to capture the variability of solar activity, which modulates the SEP flux, I selected input physical quantities that describe both the interplanetary medium and solar activity. These input features can be categorized into two groups: remote signatures and in-situ measurements. The remote signatures consist of the F10.7 index, as well as the long-wavelength (X_L) and short-wavelength (X_S) x-ray fluxes. The F10.7 index represents the flux of solar radio emission at a wavelength of 10.7 cm, measured in solar flux units (sfu). To obtain the x-ray fluxes, I utilized 1- and 5-minute averaged data from the Geostationary Operational Environmental Satellite (GOES) database², specifically at long wavelengths (1 - 8 Å) and short wavelengths (0.5 - 4.0 Å).

The in-situ measurements encompass the near-Earth solar wind magnetic field and plasma parameters. These include the solar wind speed (in km s⁻¹), average interplanetary magnetic field (IMF) strength (in nT), and the integral SEP fluxes at three energy channels: >10, >30, and >60 MeV, which correspond to the GOES channels (in 1/cm² sec ster). These SEP fluxes were obtained from multiple spacecraft stationed at the first Lagrange point (L1) throughout the study period. In particular, the IMF and plasma data in the OMNI database are obtained from the IMP, Wind, and ACE missions, while the energetic particle fluxes are obtained from the IMP and GOES spacecraft³.

To ensure a comprehensive dataset, I acquired hourly-averaged data covering a timeframe from December 1976 to July 2019, which spans the past four solar cycles. These data were sourced from the Space Physics Data Facility (SPDF) OMNIWeb database⁴, hosted by the Goddard Space Flight Center. This database provides a wealth of information, including integral proton fluxes, as well as an extensive range of solar wind plasma and magnetic field parameters. Lastly, the daily data on sunspot numbers were obtained from the Sunspot Index and Long-term Solar Observations (SILSO) archive⁵, maintained by the World Data Center.

²GOES SXR Database: <https://satdat.ngdc.noaa.gov/sem/goes/data/avg/>

³OMNIWeb Data Documentation: https://omniweb.gsfc.nasa.gov/html/ow_data.html

⁴OMNI Database: <https://omniweb.gsfc.nasa.gov>

⁵Sunspot Number Dataset: <https://www.sidc.be/silso/home>

Figure ?? shows a plot for the timeseries data of all features. The top 3 panels are the logarithms of the SEP integral flux at the 3 energy channels ($\log_{\text{PF}10}$, $\log_{\text{PF}30}$, and $\log_{\text{PF}60}$), then the sunspot number, the F10.7 index ($\text{F}10_{\text{idx}}$), the logarithms of the x-ray fluxes (\log_{Xs} and \log_{XI}), the solar wind speed (V_{sw}), and the average magnitude of the IMF (avg_{IMF}). Throughout this paper, I adopt the convention that "log" refers to the common logarithm with a base of 10. The gray shades refer to the timespan of solar cycles. The blue, orange, and gold colors refer to the training, validation, and test sets, respectively. The data split method will be explained shortly.

Since the input SEP data have been compiled from various spacecraft, it may have artifacts even after processing. In particular, there are occasional jumps in the background level. There are also several day-long gaps in the OMNI solar wind parameters from the early 1980s to mid-1990s where only IMP 8 data are available and this spacecraft spent part of each orbit in the magnetosphere. I am reasonably confident that these issues do not influence the overall analysis significantly.

In deep learning applications, the dataset is split into 3 sets; namely the training set, the validation set, and the test set. The training set is usually the largest chunk of data that is used to fit the model. The validation set is a smaller chunk of data used to fine-tune the model and evaluate its accuracy to ensure it is unbiased. The test set is the out-of-sample data exclusively used to assess the final model when performing on unseen data (?).

After inspecting the correlation between the solar wind indices and the SEP integral fluxes in the OMNIWeb database, I chose the top-correlated features with the SEP flux. The correlations were made between the SEP fluxes and the individual parameters. Hence I took only timeseries of logarithms of the protons' integral flux at 3 energy channels (>10 , >30 , and >60 MeV), the timeseries of logarithm of the X-ray fluxes, the F10.7 index, the sunspot number, the solar wind speed, and the average strength of the IMF as input parameters to our model. The log of the SEP flux was used across the whole study. The correlation matrices for the training, validation, and test sets are shown in Figure ???. The X-ray and proton fluxes were converted into the logarithmic form because it was more convenient than the original form of data since the time series data were mostly quiet and had numerous sharp spikes, which correspond to solar events. Based on a previous experience with NNs (?), I found that training separate models for each target (output) feature can lead to better results. This is because a dedicated model for each output feature can more easily learn the interrelationships between input features and make more accurate predictions. Therefore, in our current study, I trained 3 separate models, each one targeting the logarithm of the protons integral flux at a specific energy channel.

In order to ensure consistency across all features, all durations of the time series data of the physical quantities were matched to be within the same time range. Subsequently, the dataset was resampled to obtain daily averaged data, resulting in a significant reduction of the dataset size by a factor of 24. This reduction facilitated expeditious training and yielded prompt results.

There were missing data values in the original dataset; for the B_{avg} ($\sim 10.7\%$), V_{sw} ($\sim 10.5\%$), F10.7-index ($\sim 0.08\%$), short-band x-ray flux ($\sim 8\%$), long-band x-ray flux ($\sim 9.8\%$), and proton fluxes ($\sim 4.3\%$). The data gaps were linearly interpolated.

In timeseries forecasting, it is a common practice to take a continuous set of data points from the main dataset to be the validation set and another smaller chunk of data to be the test set, for instance in ??. From our experiments, I got descent results when I applied the same data split method, but the results were a bit biased toward the end of the solar cycle 24 and the testing set was biased towards a quiet period. So, I adopted the 9-2-1 strategy, that is taking from each year 9 months to be added in the training set, 2 months to be added in the validation set, and 1 month to be added in the test set. This is applied over the ~ 43 years of data (Fig. ??), which yields 74.29% of the data for the training set, 16.2% for the validation set, and 9.51% for the testing set. By doing so, I eliminated the need to do cross-validation and hence, made the training more efficient. It is worth to mention that the timeseries data must not be shuffled as that will break temporal and logical order of measurements, which must be maintained.

4.3.2 Method

In this section, I introduce the data analysis methods used in this work. I start with explaining the model selection phase, followed by a discussion of the bidirectional long-short-term memory (BiLSTM) neural network architecture. The technical terminologies are described in the appendices.




Figure 4.1: Data splitting for all input features, showing the training, validation, and testing sets. Daily data from 1976-12-25 00:00 to 2019-07-30 00:00. The gray shading labels the solar cycles from SC21 to SC24.

The Bi-LSTM Model

Recurrent neural networks (RNNs) that support processing input sequences both forward and backward are known as Bidirectional Long Short-Term Memory (BiLSTM) neural networks (?). Regular RNNs (??) depend on the prior hidden state and the current input to determine the output at a given time. The output of a BiLSTM network, on the other hand, is dependent on the input at a given moment as well as the previous and future hidden states. As a result, the network is able to make predictions using contexts from the past as well as the future. Hence, accuracy is improving. Each BiLSTM layer consists of two LSTM layers; a forward layer that processes the input sequences from the past to future, and a backward layer that processes the input sequences from the future to the past, as illustrated in




Figure 4.2: Correlation matrices show the correlation between the features in the training, validation, and test sets.

Figure ??, to capture information from both past and future contexts. The output from each layer is concatenated and fed to the next layer, which can be another BiLSTM layer or a fully connected layer for final prediction.

BiLSTM networks are advantageous than traditional LSTM networks in a variety of aspects (???). First, as I demonstrate in this study, they are excellent for tasks like timeseries forecasting, as well as speech recognition and language translation (?????) because they can capture long-term dependencies in the input sequence in both forward and backward directions. Second, unlike feedforward networks, BiLSTM networks do not demand fixed-length input sequences, thus being able to handle variable-length sequences better. Furthermore, by taking into account both past and future contexts, BiLSTM networks can handle noisy data. However, BiLSTM networks are computationally more expensive than regular LSTM networks due to the need for processing the input sequence in both directions. They also have a higher number of parameters and require more training data to achieve good performance.




Figure 4.3: Architecture of a single BiLSTM layer. The blue circles at the bottom labeled by $(x_0, x_1, x_2, \dots, x_i)$ are the input data values at multiple time steps. The purple circles, on the other hand, are the output data values at multiple time steps labeled by $(y_0, y_1, y_2, \dots, y_i)$. The dark green and light green boxes are the activation units of the forward layer and the backward layer, respectively. The orange and yellow circles are the hidden states at the forward layer and the backward layer, respectively. Both the forward and backward layers compose a single hidden BiLSTM layer. The figure is adopted from ?

The final dataset has 7 features, including the target feature, from December 25th 1976 to July 30th 2019, with a total of 15,558 samples (number of days). The training set has 11,558 samples, the validation set has 2,520 samples, and the test set has 1,480 samples.

The input horizon of 270 steps (30 days \times 9 months) was used. A data batch size of 30 was used, which is the number of samples processed that result in one update to the model's weights (Appendix ??). The model consists of 4 BiLSTM layers with 64 neurons each, and an output dense layer with 3 neurons, representing the output forecasting horizon. The total number of trainable parameters is 333,699. The number of training epochs was set to 50 because from experiments, the model stopped improving remarkably after almost 50 epochs. Thus, there was no need to waste time and computational resources to train the model for more than 50 epochs.

The *ModelCheckpoint* callback function was used to register the model version with the minimal

validation loss. The *EarlyStopping* callback function was used to halt the model run when detecting overfitting, with a *patience* parameter of 7. *ReduceLROnPlateau* callback function was used to reduce the learning rate when the validation loss stops improving, with a *patience* parameter of 5, a reduction factor of 0.1 and minimal learning rate of $1e^{-6}$.

Model Selection

To determine the most suitable model for our objective and provide justifiable reasons, I conducted the following analysis. First I examined the naive (persistence) model, which is very simplistic and assumes that the timeseries values will remain constant in the future. In other words, it assumes that the future value will be the same as the most recent historical value. That was the baseline. Next I examined the moving-average model, which calculates the future values based on the average value of historical data within a specific time widow. This gives a little bit lower error.



Figure 4.4: Illustration of the sliding window technique for a sample of 10 timesteps, where each number denotes a distinct time step. As an example here, the input horizon (blue color) length is 4 timesteps and the output horizon length is 3 timesteps. The input window slides 1 time step at a time across the entire data sequence to generate 4 distinct input and forecast horizon pairs. The purple, orange, and green colors of the output horizon represent 1-day, 2-day, and 3-day ahead forecasting, respectively. The timesteps of 1-day ahead forecasting across the data sequences are then concatenated into a single timeseries list that is called 1-day ahead prediction. The same for 2-day and 3-day ahead.

After that, I went towards the machine learning (ML)-based models. For all the ML models, I chose the Adaptive moment estimation (Adam) optimizer (?) as the optimization algorithm due to its minimal memory requirements and high computational efficiency as it is well-suited for applications that involve large number of parameters or large datasets. As a rule of thumb, I set the optimizer's learning rate to be 0.001 as it is usually recommended (?).

In order to prepare the data in a readable format to the ML models, I created a windowed dataset with an input horizon of 365 steps representing 1 year of data and an output horizon of 3 steps representing the forecast window of three days. I call this windowing method as Multi-Input Multiple Output (MIMO) strategy, in which the entire output sequence is predicted in one shot. The MIMO strategy adopts the sliding window method that was mentioned in ? in which each sequence is shifted by one step with respect to the previous sequence until reaching the end of the available data (Fig. ??). This approach minimized the imbalance of active days, with high SEP fluxes, and quiet days.

After experiments with different loss functions and evaluate their performance on our dataset, I chose the Huber function ?? as the loss function and the Mean Absolute Error (MAE) is used as the metric function to monitor the model performance. I used the Huber function because it is robust and combines the advantages of both Mean Squared Error (MSE) and MAE loss functions. It is less sensitive to outliers than MSE, while still being differentiable and providing gradients, unlike MAE. Since our data is noisy and contains outliers that may negatively impact the model's performance, the Huber loss function is a good choice.

I examined various neural network models to determine the optimal architecture for our task. Initially, I started with a simple linear model comprising of a single layer with a single neuron. However, this model did not yield satisfactory results. I then explored a dense ML model consisting of two hidden layers, each with 32 neurons and a *ReLU* activation function. Next, I experimented with a simple RNN model with the same number of hidden layers and neurons. To find the optimal learning rate, I utilized the *LearningRateScheduler* callback function and discovered that a rate of $1.58e^{-4}$ under the basic settings minimized the loss. I proceeded to examine stateful versions of RNN, LSTM, and BiLSTM models with three hidden layers, each with 32 neurons and a learning rate of $1.58e^{-4}$. In addition, I explored




Figure 4.5: Benchmarking of 10 models, shows the Huber loss for the validation and test sets.

a hybrid model that consisted of a 1-dimensional convolutional layer with 32 filters, a kernel size of 5, and a *ReLU* activation function. I combined this with a two-hidden layer LSTM network with 32 neurons each and a learning rate of $1.58e^{-4}$. I experimented with *Dropout* layers but did not observe any significant improvement in the results. Finally, I evaluated a BiLSTM model with five hidden layers, 64 neurons each, and a learning rate of 0.001. Based on the evaluation of all the models on both the validation and test sets (Fig. ?? and Table ??), I selected the BiLSTM model for further refinement. More details on the final model architecture and hyperparameters are explained in the Appendix ???. Figure ?? presents a comparative analysis of the Huber loss within the validation and testing sets across the ten aforementioned models. I used several evaluation measures to assess our models since each metric provides valuable insights into the accuracy and performance of the forecasts (Appendix ??), helping to identify areas for improvement and adjust the forecasting models accordingly.

4.3.3 Results and discussion

The benchmarking in Figure ?? showed that, in general, the ML-based methods were not much different. On the other hand, the persistence model and moving average model resulted in the highest errors compared with the ML-based models, and their results were close to some extent. As I see, the BiLSTM model performed the best over both the validation and test sets compared with the other models.

I developed and trained 3 BiLSTM models to forecast the integral flux of SEP, one model per energy channels. After the training was completed, I evaluated the performance of the models from the loss curve (Fig. ??) using the Huber loss (the left panel) and the metric MAE (the middle panel). During the training, the learning rate was reduced multiple times via the *LearningRateScheduler* callback function (the right panel). The left panel quantifies the discrepancy between the model's predictions and the true values over time. It shows how the Huber loss function changes during the training iterations (Epochs) for the training and validation sets for the three energy channels so that each channel has one color. The middle panel shows how the model's metric MAE changes with training epochs. It is used to evaluate the performance of the trained model by measuring the average absolute difference between the model's predictions and the true values, providing a single numerical value that indicates the model's error at a given epoch. The right panel shows how the learning rate of the model's optimizer changes with epochs via the *LearningRateScheduler* callback function, which changes the learning rate based on a predefined schedule to improve training efficiency and convergence. The learning rate refers to the rate at which the model's parameters are updated during the training process. I noticed that at the epochs where the learning rate has changed, there were bumps in the loss curves across all the energy channels, which is expected. This highlights the boundaries within which the learning rate yields better performance.

From experimentation, I found that the batch size and the optimizer learning rate are the most important hyperparameters that have a strong influence on the overall model's performance (?). In addition, adding *dropout* layers as well as varying the number of hidden layers and hidden neurons resulted in only marginal improvements to the final model performance, while substantially increasing training time and requiring greater computational resources.




Figure 4.6: *Left Panel* - The Huber loss vs. the number of training epochs for the BiLSTM model for the validation and test sets, for the 3 energy channels. *Middle Panel* - The mean absolute error (MAE); the model's metric vs. the number of training epochs. *Right Panel* - Shows how the learning rate of the Adam optimizer changes over the number of epochs.

The term *batch size* refers to the number of data sequences processed in one iteration during the training of a ML model (?). Initially, a batch size of 64 was selected, however, I observed that the model produced better results when a batch size of 30 was used instead. This could be related to the Carrington rotation, which lasts for ~ 27 days. There were ~ 570 Carrington rotations between December 25th 1976 and July 30th 2019. Therefore, updating the model's weights after every Carrington rotation could be a reasonable choice for improving its performance. Figure ?? shows how good the model predictions are (on the y-axis) compared with the observations of the validation set (on the x-axis). The blue, orange, and gold colors refer to 1-day, 2-day, and 3-day ahead predictions, respectively. The top panel is for the >10 MeV channel, the middle panel is for the >30 MeV channel, and the bottom panel is for the >60 MeV channel. The left column is for the entire validation set, while the right column is for the observations points ≥ 10 proton flux units (pfu). That is the threshold value of proton flux as measured by the National Oceanic and Atmospheric Administration (NOAA) GOES spacecraft to indicate severity of space weather events caused by SEP.

I found that, overall, the models performed very well. The R correlation was >0.9 for all points of the validation set across the forecasting windows for the 3 energy channels. The R correlation was >0.7 for the observations points ≥ 10 pfu as well. However, the correlation between the modeled data and the observations exhibited a decline as the forecast horizon increased, in accordance with the anticipated result. To confirm the validity of the models, I performed the same correlation analysis between the modeled data and the observations of the out-of-sample test set (Fig. ??), which was not given to the model. Again, I found a high correlation across the forecasting windows for the 3 energy channels. The points were more dispersed between 1 and 1.5 on the x-axis, which reflected in a bit lower correlation. This might be a limitation in the current version of the model between that range of SEP fluxes since the models underestimated the flux values within that range across all energy channels, possibly due to the relatively smaller training samples with fluxes above 10 pfu compared with the majority of the data.

In order to see the temporal variation of the correlation between the modeled data and the observations, I applied a rolling window of 90 steps (3 months \times 30 days/month = 1 season) that shows the seasonal variation of the correlation, as shown in Figure ???. Here, I show only the 1-day ahead predictions for the test set, for the 3 energy channels. I observe drops in the correlation factor synchronized with the transition between solar cycles (e.g., particularly between ~ 1995 - 2000, which represents the declining phase of the solar cycle 22 and the rising phase of the solar cycle 23). This could be related to the fact that the low SEP fluxes during quiet times are more random and thus more difficult to forecast (????).

During periods of low solar activity, the forecasting of low SEP fluxes becomes more challenging due to their increased randomness. This difficulty arises from the reduced occurrence of conventional SEP drivers, such as solar flares and CMEs. Studies have suggested that the most significant solar eruptions tend to happen shortly before or after the solar cycle reaches its maximum (?). Additionally, sporadic increases in solar activity have been observed (?), which might contribute to the diminished correlations observed in our research. There is clearly some factor that is influencing the correlation during certain periods where there are no or only weak SEP events. However, it is not obvious which physical phenomena are the cause rather than, for instance, some artifact of the data. Understanding the interplay between these factors and their influence on SEP fluxes during periods of reduced solar activity




Figure 4.7: Correlation between the model predictions and observations for 1-day, 2-day, and 3-day ahead for >10 MeV (top panel), >30 MeV (middle panel), and >60 MeV (bottom panel). The panels in the left column represent all the points of the validation set, those in the right column represent all the observations points with daily mean flux ≥ 10 pfu.




Figure 4.8: Same as Figure ?? but for the test set.

remains a critical area of research. It would be interesting to find what is reducing the correlations, thus more investigation is needed.

Overall, the modeled data was correlated the most with observations at >60 MeV, then the second rank was for the >10 MeV channel, and the third rank was for the >30 MeV channel. That could be related to the relatively larger extent of drops in correlation at the >30 MeV channel. The decline in correlation at the >30 MeV channel is consistent with the findings of ?. A summary of the performance results of the models for both the validation set and test set is presented in Table ??.




Figure 4.9: Comparison between the model outputs and observations of the test set for the 3 energy channels. In addition to the rolling-mean window correlation for 1-day ahead predictions.

From the visual inspection of the test set examples (Fig. ??, ??, and ??), I found that the predicted onset time, the peak time, and end times of SEP events were highly correlated with those of the observations, which implies that the model captured the temporal variations, as well as the trends in SEP




Figure 4.10: The model's forecasts for the out-of-sample testing set for the >10 MeV channel are shown at forecast horizons of 1 day, 2 days, and 3 days ahead, using samples of data from December in selected years mentioned in the top-left side of the plots.




Figure 4.11: The model's forecasts for the out-of-sample testing set for the >30 MeV channel are shown at forecast horizons of 1 day, 2 days, and 3 days ahead, using samples of data from December in selected years mentioned in the top-left side of the plots.




Figure 4.12: The model's forecasts for the out-of-sample testing set for the >60 MeV channel are shown at forecast horizons of 1 day, 2 days, and 3 days ahead, using samples of data from December in selected years mentioned in the top-left side of the plots.

flux.

Table 4.1: Summary of the performance results of the models for the validation and test sets.

Validation Set								
	log PF >10 MeV			log PF >30 MeV			log PF >60 MeV	
Model Loss	0.0016			0.0010			0.0009	
Model Metric	0.0329			0.0232			0.0218	
	1-Day	2-Day	3-Day	1-Day	2-Day	3-Day	1-Day	2-Day
MAE	0.061	0.091	0.125	0.053	0.079	0.098	0.052	0.069
MSE	0.013	0.028	0.054	0.010	0.031	0.055	0.009	0.027
RMSE	0.114	0.168	0.233	0.098	0.176	0.234	0.097	0.164
MAPE	22.156	28.104	34.721	13.039	18.590	22.735	10.036	13.994
Test Set								
	log PF >10 MeV			log PF >30 MeV			log PF >60 MeV	
Model Loss	0.0014			0.0011			0.0010	
Model Metric	0.0333			0.0283			0.0250	
	1-Day	2-Day	3-Day	1-Day	2-Day	3-Day	1-Day	2-Day
MAE	0.072	0.099	0.125	0.053	0.088	0.107	0.045	0.066
MSE	0.015	0.030	0.050	0.009	0.029	0.048	0.007	0.020
RMSE	0.121	0.172	0.224	0.094	0.170	0.218	0.082	0.141
MAPE	30.135	37.498	48.139	20.599	34.300	40.803	12.358	20.504

To get further insight into the model’s performance, I conducted an assessment of various skill scores, including True Positive (TP), True Negative (TN), False Positive (FP), and False Negative (FN). Additionally, skill score ratios such as Probability of Detection (POD), Probability of False Detection (POFD), False Alarm Rate (FAR), Critical Success Index (CSI), True Skill Statistic (TSS), and Heidke Skill Score (HSS). Detailed descriptions of these skill scores can be found in Appendix ???. To extract individual SEP events from the test dataset, I implemented a threshold-based clustering algorithm. This algorithm uses the NOAA/SWPC warning threshold value of 10 pfu for the $E \geq 10$ MeV channel. Upon analysis, I identified the number of detected SEP events for each output forecasting window and calculated the skill scores (Table ??). In the true test set, I identified 12 SEP events.

The evaluation of the model revealed notable trends as the length of the output forecasting window increased. The POD and CSI exhibited a declining pattern, indicating a reduced ability of the model to accurately detect and capture positive events (SEP occurrences) as the forecasting horizon extended further into the future. This suggests that the model’s performance in identifying and capturing true positive instances diminishes with longer forecasting windows. Moreover, the POFD demonstrated an increasing trend, indicating an elevated rate of false positive predictions as the forecasting horizon lengthened. The model’s propensity to generate false alarms rose with the lengthening forecasting window, leading to incorrect identification of non-events as positive events. Consequently, the TSS and HSS exhibited decreasing values, signifying a deterioration in the model’s overall skill in accurately capturing and distinguishing between positive and negative instances. Overall, our skill scores are comparable with those reported by previous studies (Table ??). Although the UMASEP model does better than ours (i.e., has a higher POD), our FAR is much lower, thus, making fewer false alarms than the UMASEP model.

Table 4.2: Confusion matrix for the energy channel ≥ 10 MeV predictions in the test set.

E >10 MeV	No. events	TP	TN	FP	FN
1-day ahead	15	21	1441	2	13
2-day ahead	13	14	1441	2	20
3-day ahead	5	5	1443	0	29

4.4 Conclusions

Forecasting the SEP flux is a crucial task in heliophysics since it affects satellite operations, astronaut safety, and ground-based communication systems. It is a challenging task due to its non-linear, non-stationary, and complex nature. Machine learning techniques, particularly neural networks, have shown

Table 4.3: Comparing the skill scores with previous models. The dashed entries mean the data is unavailable (?) for more details).

Model		POD	FAR	TSS	HSS	POFD	CSI	Accuracy	Precision
Our BiLSTM model	1-Day	0.618	0.087	0.531	0.732	0.001	0.583	0.99	0.913
	2-Day	0.412	0.125	0.287	0.553	0.001	0.389	0.985	0.875
	3-Day	0.147	0	0.147	0.252	0	0.147	0.980	1
UMASEP-10 (?)		0.822	0.219	—	—	—	—	—	—
PCA (?)		0.587	0.245	—	0.65	—	—	—	—
SPARX (?)		0.5	0.57	—	—	0.32	0.3	—	—
SPRINTS (?)		0.56	0.34	—	0.58	—	—	—	—
REleASE (?)		0.63	0.3	—	—	—	—	—	—

promising results in predicting SEP flux. In this study, I developed and trained BiLSTM neural network models to predict the daily-averaged integral flux of SEP at 1-day, 2-day, and 3-day ahead, for the energy channels >10 MeV, >30 MeV, and >60 MeV. I used a combination of solar and interplanetary magnetic field indices from the OMNIWeb database for the past 4 solar cycles as input to the model. I compared the models with baseline models and evaluated them using the Huber loss and the error metrics in Appendix ??.

The data windowing method I used, based on the MIMO strategy, eliminates the need to feed the output forecast as input back into the model and that allows to do forecasting relatively far into the future while maintaining decent results (e.g., the MSE is ranging between 0.007 and 0.015 for 1-day forecasting in the test set, compared to an MSE of 0.236 for a persistence model. See Table ??). The results show that the model can make reasonably accurate predictions given the difficulty and complexity of the problem. The MSE was ranged between 0.009 and 0.055 for the validation set, and between 0.007 and 0.05 for the test set. The correlations between the observations and predictions were >0.9 for the validation and test sets (Fig. ?? and Fig. ??). Nevertheless, the mean temporal correlation was ~ 0.8 for the test set (Fig. ??). Although our models performed well, I observed a relatively large discrepancy between the predictions and the observations in the >30 MeV energy band.

The findings of this study underscore the challenges encountered by the forecasting model in accurately predicting SEP data over longer time periods. As the length of the output forecasting window increased, the model’s ability to detect true positives and its overall skill in differentiating positive and negative instances diminished. Additionally, the model displayed an elevated rate of false negative predictions, indicating an increased tendency to generate misses as the forecasting horizon extended. These results highlight the importance of carefully considering the appropriate forecasting window length for SEP data to ensure the model’s optimal performance. Our skill scores generally align with those from previous works (Table ??). There are variations in the metrics’ values across different studies, highlighting the complexities and nuances associated with each study. Nevertheless, it is important to acknowledge that the statistical significance of the results in this study is limited due to data averaging. Future studies should consider incorporating hourly data, as this is likely to result in a greater number of identified events. The model can provide short-term predictions, which can be used to anticipate the behavior of the near-Earth space environment. These predictions have important implications for space weather forecasting, which is essential for protecting satellites, spacecraft, and astronauts from the adverse effects of solar storms.

Multiple techniques exist for identifying the optimal combination of hidden layers and neurons for a given task such as empirical methods, parametric methods, and the grid search cross-validation method, which I will explore in future work. The observed reduction in correlation necessitates further investigation to determine its origin, whether stemming from tangible causal factors or potential aberrations within the model or data. I plan to expand upon this work by performing short-term forecasting using hourly-averaged data. This extension will involve integrating additional relevant features such as the location and area of active regions and coronal holes on the Sun.

BiLSTM networks are particularly useful for tasks involving sequential data such as timeseries forecasting. Given their capacity to handle input sequences in both directions in time and capture long-term dependencies, they are valuable in a broad range of applications. Nonetheless, one should carefully consider their data requirements and computational complexity before adopting them. Our results emphasize that the use of deep learning models in forecasting tasks in heliophysics are promising and encouraging, as pointed out by ?.

This work is a stepping stone towards real-time forecasting of SEP flux based on the public-available

datasets. As an extension, I am currently working on developing a set of models that deliver near-real time prediction of SEP fluxes at multiple energy bands, multiple forecasting windows, with hourly-averaged data resolution, with a more sophisticated model architecture, as well as more features that address the state of solar activity more comprehensively. I plan to extend the analysis to include more recent data from solar cycle 25, in order to improve the accuracy of the models. In conclusion, our study highlights the potential of using BiLSTM neural networks for forecasting SEP integral fluxes. Our models provide a promising approach for predicting the near-Earth space environment too, which is crucial for space weather forecasting and ensuring the safety of our space assets. Our findings contribute to the growing body of literature on the applications of deep learning techniques in heliophysics and space weather forecasting.

Chapter 5

Summary and Future Work

5.1

5.2

5.3

5.4

5.5 Future Work

The research presented in this thesis establishes an important foundation and provides a precursor for future advances that can build upon the present work. Some open questions and promising areas for future investigations include:

- Additional coronal wave statistical studies using expanded event samples and new imaging datasets from Solar Orbiter and ground observatories to improve generalizability of findings.
- Incorporating 3D analytical and numerical coronal wave propagation models for more physics-based forecasting approaches.
- Modeling mechanisms for type III radio burst onset and time profiles using particle-in-cell and MHD models.
- Ensemble forecasting models for SEP events combining multiple machine learning algorithms trained on multi-mission data.
- Validation of data-driven models for other solar wind driven geospace extremes such as radiation belt enhancements and ionospheric storms.
- Leveraging new solar observatory missions and assimilative models within operational prediction systems for real-time space weather forecasts.
- Exploring applications of deep learning and physics-informed machine learning to additional outstanding problems in heliophysics and astrophysics.

In summary, the present work opens exciting avenues for more cross-disciplinary studies synthesizing heliophysics domain knowledge with cutting-edge data science and artificial intelligence methods. The new generation of solar, heliospheric and geospace missions will yield transformative observations to continue advancing both science understanding and predictive capabilities.

5.6 Definitions and Acronyms

Some of the key technical terms and acronyms used in this thesis are listed below:

- SEP - Solar Energetic Particle
- CME - Coronal Mass Ejection
- EUV - Extreme Ultraviolet
- SOHO - Solar and Heliospheric Observatory
- SDO - Solar Dynamics Observatory
- AIA - Atmospheric Imaging Assembly

EIT - Extreme ultraviolet Imaging Telescope

TRACE - Transition Region and Coronal Explorer

LASCO - Large Angle and Spectrometric COronagraph

GOES - Geostationary Operational Environmental Satellite

AI - Artificial Intelligence

MHD - Magnetohydrodynamics

AU - Astronomical Unit

This provides definitions of the major domain-specific terms and measurement concepts used. Additional terminology is introduced as required in the respective chapters.

Appendix A

Appendix

A.1 Appendix for Radio Observation-related Work

A.1.1 Persistent imaging technique

Persistent imaging is a technique used in medical imaging, particularly ultrasound imaging, to create a continuous, real-time display of the anatomy being imaged (see ??, and references within). The core idea of persistent imaging is to use persistence, or the ability of the human eye to retain an image for a brief moment after it has disappeared to create a more informative and visually clear image ??.

At every image in a time-ordered series, the technique keeps the old pixel value if it is brighter than the current pixel value, else it takes the current pixel's value. The result is saved as the current persistence image. Then, the next image in the series is evaluated by comparing it pixel by pixel with respect to the previous persistence image. The resulting image emphasizes the changes between the current image and the previous persistent image, making them more visible to the human eye.

The persistent imaging technique can be described mathematically by a set of equations. If we let $I(t, x, y)$ be the intensity at time t and pixel coordinates (x, y) , and let $P(t, x, y)$ be the persistence image at time t and pixel coordinates (x, y) , then the persistence image at time t is computed as:

$$P(t, x, y) = \max\{I(t, x, y), P(t - 1, x, y)\}, \quad (\text{A.1})$$

where **max** represents the maximum of its two arguments. The current image at time t is then evaluated with respect to the previous persistence image as follows:

$$I'(t, x, y) = \max\{I(t, x, y) - P(t - 1, x, y), 0\}, \quad (\text{A.2})$$

The resulting image $I'(t, x, y)$ is a modified version of the current image that emphasizes the differences from the previous persistence image.

The persistent imaging technique has been shown to improve the visual quality of ultrasound images and other medical imaging modalities, and is commonly used in clinical practice. In this paper, we utilize the persistent imaging technique to improve the visualization of the solar radio sources of type III emissions (Fig. ??).

A.1.2 Resolving the radio emission location ambiguity

In this part, we show that the $-Z$ solution of Equation ?? is highly unlikely in our case. Figure ?? shows the positive and negative solutions of Equation ???. We take the innermost and outermost coronal radio sources at R_1 and R_2 , respectively, as an example. r_1 and r_2 are the projections of R_1 and R_2 on the POS, respectively. Harmonic radio emission from R_1 will theoretically be absorbed by a region along the LOS with plasma frequency (and corresponding density) equal to or higher than the harmonic emission frequency at R_1 . In the case of the spherically symmetric Newkirk model, the highest density location the emission from R_1 could pass through is r_1 on the POS. Thus, for harmonic radio emission from behind the POS ($-Z$, where $Z = 0$ is defined at the center of the Sun and positive Z is towards the observer) to be observed at the Earth, it must satisfy the following condition:

$$2f_{R_1} > f_{r_1}, \quad (\text{A.3})$$

where f_{R_1} is the plasma frequency of radio emission that occurred behind the POS, and f_{r_1} is the plasma frequency at the projected location of r_1 on the POS. The relation between the local plasma frequency and the electron density is defined by the equation:

$$f[\text{MHz}] = 8.93 \times 10^{-3} \sqrt{n[\text{cm}^{-3}]} \quad (\text{A.4})$$

The Newkirk electron-density model (??) describes the typical densities in the outer part of the corona according to the following equation:

$$n[\text{cm}^{-3}] = \alpha 4.2 \times 10^4 10^{4.32 \frac{R_\odot}{r}} \quad (\text{A.5})$$

where α is the fold number (i.e., a multiplicative factor that accounts for the density variations based on the degree of solar activity), and r is the radial distance from the Sun in solar radii. By substituting Equations ?? and ?? into Equation ??, we obtain

$$\frac{n_{r_1}}{n_{R_1}} = \frac{10^{4.32 \frac{R_\odot}{r_1}}}{10^{4.32 \frac{R_\odot}{R_1}}} < 4. \quad (\text{A.6})$$

After reduction we obtain the final formula that must be satisfied under these assumptions in order for radio emission behind the POS to pass through the corona and reach the Earth:

$$\frac{r_1}{R_\odot} < \left(\frac{\log 2}{2.16} + \frac{R_\odot}{R_1} \right)^{-1}. \quad (\text{A.7})$$




Figure A.1: Schematic shows the locations of the radio sources for the +Z and -Z solutions of Equation ???. The Sun is located in the middle as an orange circle, with a horizontal dashed black line representing the POS. The vertical dashed green line represents the Sun-Earth LOS. The dashed blue and red circles represent the plasma spheres of density equivalent to the observation frequencies of the innermost and outermost radio sources at R_1 and R_2 , respectively, under the Newkirk model assumption of spherically-symmetric density distribution. The impact parameters r_1 and r_2 are the projection of R_1 and R_2 on the POS. The dot-dashed blue and red circles are the circles passing through the impact parameters r_1 and r_2 , respectively.




Figure A.2: Cut of the flux density at 700 kHz observed by Wind (left panel) and STEREO-A (right panel). Note: for STEREO-A, there is no exact frequency channel at 700 kHz; therefore we selected the nearest one (675 kHz).

From Figure ??, r_1 and r_2 will always be smaller than R_1 and R_2 , respectively. The Newkirk model requires that the density at r_1 and r_2 be significantly higher than the density at R_1 and R_2 , respectively (Table ??). Additionally, from the geometric representation in Figure ??, we find that the electron density at r_1 is higher than at R_1 , hence the radio emission cannot reach the Earth from that point behind the POS (?).

From Table ??, the assumption of Equation ?? is not satisfied. Thus, the $-Z$ solution is invalid in our case. This implies that the harmonic emission from behind the POS will not reach the Earth. Thus, the $+Z$ assumption is the valid solution.

Table A.1: Radial distances and densities at the first (R_1) and last (R_2) radio sources were obtained from the $2.5 \times$ Newkirk model, as well as their impact parameters r_1 and r_2 , respectively.

Point	Radial distance (R_\odot)	Density (cm^{-3})	Ratio (n_r/n_R)
r_1	1.58	5.69×10^7	
R_1	1.81	4.82×10^6	11.81
r_2	2.6	2.59×10^7	
R_2	3.49	1.82×10^6	14.23

Furthermore, we analyzed the time difference of arrival of the radio emission at interplanetary wavelengths in Figure ???. Specifically, we compared the timing of peak signals at a low frequency between two spacecraft, Wind and STEREO. This analysis was conducted under the assumption of two possible scenarios:

- one in which the radio emission source follows a trajectory that is roughly equidistant between Wind and STEREO – if the $+Z$ assumption is true.
- the trajectory implies significantly longer travel times from the source to Wind compared to STEREO – if the $-Z$ assumption is true.

Examining the data, we selected the frequency channel 700 kHz observed by Wind and its nearest counterpart 675 kHz for STEREO. Interestingly, the difference in the arrival times of these signals was merely one minute, which is within the bounds of the time resolution of the instrument. This negligible difference in arrival times supports the $+Z$ assumption for the beam trajectory, meaning it travels approximately at an equal distance between the two spacecraft.

A.2 Appendix for SEP-related Work

A.2.1 Terminologies

In this section, we introduce the main concepts related to machine learning which are presented in this paper.

- **Cross-validation:** A technique used to evaluate the performance of a machine learning model by dividing the data into subsets and assessing the model on different combinations of these subsets.

- **Input Horizon:** The number of previous time steps considered as input to a model for time series forecasting. It represents the length of the historical sequence used for predictions.
- **Batch Size:** The number of samples processed together in a single iteration of the training algorithm. It affects training speed and memory requirements.
- **Updating the Model's Weights:** The process of adjusting the parameters of a neural network based on training data to minimize the difference between predicted and true outputs. The model's weights represent the parameters that are learned during the training process.
- **Loss:** A function that quantifies the difference between predicted and actual outputs. It guides the optimization process during training.
- **Minimum Validation Loss:** The lowest value achieved by the loss function on a validation dataset during training. It indicates the most accurate predictions on unseen data.
- **Overfitting:** When a model performs well on training data but fails to generalize to unseen data due to memorizing training examples instead of learning underlying patterns.
- **Learning Rate:** A hyperparameter that determines the step size at each iteration of the optimization algorithm during training. It affects learning speed and convergence. A high learning rate can cause the training process to converge quickly, but it may also result in overshooting the optimal solution or getting stuck in a suboptimal solution. On the other hand, a very low learning rate can make the training process slow, and may struggle to find the optimal solution.
- **Reducing the learning rate when the validation loss stops improving:** This concept involves adjusting the learning rate dynamically during the training process. When the validation loss reaches a plateau or stops improving, it indicates a suboptimal point. By reducing the learning rate, the model can take smaller steps in weight space, potentially finding a better solution. This technique, known as learning rate scheduling or learning rate decay, is commonly used to fine-tune the model's performance.
- **Patience:** A parameter used in training to determine the number of epochs to wait for an improvement in validation loss before stopping the training process.
- **Patience Parameter of 7:** In the context of early stopping, training will be stopped if the validation loss does not improve for 7 consecutive epochs.
- **Adam Optimizer:** A popular optimization algorithm in deep learning that combines Adaptive Gradient Algorithm (AdaGrad) and Root Mean Square Propagation (RMSprop) to achieve efficient optimization.
- **Optimal Architecture:** The best configuration of a neural network, including the number of layers, neurons, and other choices, for optimal performance on a specific task.
- **Hyperparameters:** Parameters set before training a model that control the learning algorithm's behavior, such as learning rate, batch size, and activation functions.
- **Layer:** A building block of a neural network that performs specific operations on input data. Includes input, hidden, output, fully connected, convolutional, recurrent, activation, and dropout layers. Here is a description for each layer:
 - **Input Layer:** The first layer of a neural network that receives raw input data. It passes the input to subsequent layers for further processing. The number of nodes in the input layer is determined by the dimensionality of the input data.
 - **Hidden Layers:** Intermediate layers between the input and output layers. They perform computations on the input data and capture higher-level representations or abstractions. Hidden layers are not directly exposed to the input or output.
 - **Output Layer:** The final layer of a neural network that produces model predictions or outputs based on computations from preceding layers. The number of neurons in the output layer depends on the problem being solved, such as regression or classification.

- **Fully Connected Layer (Dense Layer):** Each neuron in this layer is connected to every neuron in the previous layer. It allows information flow between all neurons, enabling complex relationships to be learned.
- **Convolutional Layer:** Commonly used in Convolutional Neural Networks (CNNs) for analyzing grid-like data, such as images. It applies convolution operations using filters or kernels to learn spatial patterns or features.
- **Recurrent Layer:** Used in Recurrent Neural Networks (RNNs) to process sequential data. These layers have feedback connections that allow information to be passed from one step to the next, capturing temporal dependencies and maintaining memory of past inputs.
- **Activation Layer:** Applies a non-linear function to the output of a layer, introducing non-linearity into the neural network. Activation functions like Sigmoid, Hyperbolic Tangent (\tanh), or Rectified Linear Unit (ReLU) determine neuron outputs based on weighted inputs.
- **Dropout Layer:** A regularization technique commonly used in deep learning models. It randomly sets a fraction of outputs from the previous layer to zero during training, preventing overfitting and improving generalization.

Layers play a crucial role in the information processing and learning capabilities of neural networks. The arrangement and combination of different layers determine the network's architecture and ultimately its ability to solve specific tasks.

- **Stateful:** A property of Recurrent Neural Networks (RNNs) where the hidden state is preserved between consecutive inputs, allowing the network to have memory.
- **Neuron:** A computational unit in a neural network that receives input, applies weights, and passes the result through an activation function to produce an output.
- **Hidden Neuron:** A neuron in a hidden layer of a neural network that performs intermediate computations.
- **Callback Function:** A function used during model training to perform specific actions at certain points or conditions, such as saving the best model, adjusting learning rates, or early stopping.
- **LearningRateScheduler Callback Function:** A function used in training to dynamically adjust the learning rate at specific points based on a predefined schedule or function. It improves training efficiency and convergence by allowing the model to make finer adjustments as it approaches the optimal solution.

A.2.2 Mathematical Representation of the LSTM NN Model

The computations inside one LSTM cell can be described by the following formulas (?):

$$f_t = \sigma(W_f x_t + U_f h_{t-1} + b_f) \quad (\text{A.8a})$$

$$i_t = \sigma(W_i x_t + U_i h_{t-1} + b_i) \quad (\text{A.8b})$$

$$\tilde{C}_t = \tanh(W_c x_t + U_c h_{t-1} + b_c) \quad (\text{A.8c})$$

$$C_t = f_t \odot C_{t-1} + i_t \odot \tilde{C}_t \quad (\text{A.8d})$$

$$o_t = \sigma(W_o x_t + U_o h_{t-1} + b_o) \quad (\text{A.8e})$$

$$h_t = o_t \odot \tanh(C_t) \quad (\text{A.8f})$$

where x_t is input data at time t . The input gate i_t determines which values from the updated cell states (candidate values) \tilde{C}_t should be added to the cell state. It also takes into account the current input x_t and the previous output h_{t-1} , and is passed through a sigmoid activation function. \tilde{C}_t represent the candidate values that are added to the cell state at time t . The forget gate activation vector f_t at time step t , which determines how much of the previous cell state should be retained. The cell state C_t at time t is updated based on the forget gate, input gate, and candidate values. The output gate o_t at time t determines how much of the cell state should be output. The output vector h_t at time t is calculated based on the cell state and the output gate values. h_{t-1} is the output vector at the previous time step $t - 1$. W_f, W_i, W_c, W_o are the weight matrices for the input vector x_t . U_f, U_i, U_c, U_o are the weight matrices for the output vector h_{t-1} . b_f, b_i, b_c, b_o are the bias vectors. The symbol \odot denotes a pointwise multiplication. The sigmoid function σ is used as the activation function for the gate vectors, and the hyperbolic tangent function \tanh is used for the candidate values and the output vector.

A.2.3 Evaluation Metrics

To evaluate the model performance, we used the following equations:

$$L_\delta(y, \hat{y}) = \begin{cases} \frac{1}{2}(y - \hat{y})^2, & \text{if } |y - \hat{y}| \leq \delta, \\ \delta(|y - \hat{y}| - \frac{1}{2}\delta), & \text{otherwise} \end{cases} \quad (\text{A.9a})$$

$$MSE = \frac{1}{N} \sum_{i=1}^N (y_i - \hat{y}_i)^2 \quad (\text{A.9b})$$

$$MAE = \frac{1}{N} \sum_{i=1}^N |y_i - \hat{y}_i| \quad (\text{A.9c})$$

$$RMSE = \sqrt{\frac{\sum_{i=1}^N (y_i - \hat{y}_i)^2}{N}} \quad (\text{A.9d})$$

$$MAPE = \frac{1}{N} \sum_{i=1}^N \left| \frac{y_i - \hat{y}_i}{\hat{y}_i} \right| \quad (\text{A.9e})$$

$$R = \frac{\sum_{i=1}^n (y_i - \bar{y})(\hat{y}_i - \bar{\hat{y}})}{\sqrt{\sum_{i=1}^n (y_i - \bar{y})^2} \sqrt{\sum_{i=1}^n (\hat{y}_i - \bar{\hat{y}})^2}} \quad (\text{A.9f})$$

where y is the true value, \hat{y} is the predicted value, and δ is a threshold in the Huber loss function that controls the trade-off between the mean squared error (MSE) and the mean absolute error (MAE). In this paper, it was set to 0.1, which was selected based on several experiments.

MSE is the mean squared error, which measures the difference between predicted and actual values by calculating the average of squared differences. It provides a measure of the average squared magnitude of the errors in your forecasts, which can be useful in penalizing larger errors more heavily than smaller errors.

MAPE is the mean absolute percentage error, which measures the difference between predicted and actual values by calculating the average of absolute differences. It provides a measure of the average magnitude of the errors, allowing to evaluate the overall accuracy of your forecasts.

RMSE is the root mean squared error, which measures the difference between predicted and actual values by taking the square root of the average of squared differences. It provides a measure of the accuracy of the forecasts in the same units as the original data, allowing to evaluate the magnitude of errors in the same scale as the data.

MAPE is the mean absolute percentage error, which measures the accuracy of a forecast by calculating the average of absolute percentage errors. It provides a measure of the accuracy of the forecasts in percentage terms, allowing to evaluate the magnitude of errors relative to the actual values. MSE, MAE, RMSE, and MAPE are often used in regression analysis to assess the accuracy of the model's predictions.

Finally, R is the Pearson correlation coefficient, which measures the strength and direction of the relationship between two continuous variables, and can provide an indication of the extent to which changes in one variable may be related to changes in the other.

A.2.4 Model Configuration

The configurations for the ML models shown in Figure ?? and their performance on the validation set and the test set for the SEP integral flux ≥ 10 MeV are presented in Table ???. The batch size was set to be 64 and the number of training epochs was set to be 100. The *EarlyStopping* callback function, with a *patience* of 10, is used to help prevent overfitting during the training process by stopping training when the monitored metric has stopped improving for a certain number of epochs. The *patience* parameter controls how many epochs the training will continue without improvement before it is stopped. This is useful because if the validation loss stops getting better, the model has probably overfitted the training data and is not generalizing effectively to new data. By stopping the training early, we can avoid wasting time and resources on further training that is unlikely to improve the model's performance.

We used the *ModelCheckpoint* callback function to save the best weights of the model during training so that they can be reused later. The *LearningRateScheduler* callback function allows to dynamically adjust the learning rate of the model during training using a function passed to it that will be called at the beginning of each epoch, and it should return the desired learning rate for that epoch. It can be useful when training deep neural networks, as it allows for a higher learning rate in the early stages of

training when the model is still far from convergence, and a lower learning rate as the model approaches convergence, which can help it to converge more accurately. The downside might be the longer training time.

Table A.2: Configuration of the ML model. (1) refers to the error value for 1-day forecasting. Same for (2) refers to 2-day forecasting, and (3) for 3-day forecasting. *In the 1D-CNN layer, 32 filters, a kernel size of 5, and strides of 1 were used.

Model Architecture	No. of Hidden Layers	No. of Hidden Neurons	Activation Function	Batch Size	Learning Rate	Epochs	Callbacks Functions	Validation Set			Testing Set				
								MAE	MSE	RMSE	MAPE	MAE	MSE		
Linear	—	—	—	64	0.001	100	EarlyStopping	0.312	0.141	0.376	87.883	0.143	0.045	0.211	60.689
Dense ML	2	32	ReLU	64	0.001	100	EarlyStopping	0.262 (1) 0.275 (2) 0.290 (3)	0.118 (1) 0.138 (2) 0.166 (3)	0.344 (1) 0.372 (2) 0.407 (3)	132.580 (1) 132.004 (2) 129.288 (3)	0.400 (1) 0.395 (2) 0.392 (3)	0.281 (1) 0.286 (2) 0.294 (3)	0.530 (1) 0.535 (2) 0.542 (3)	238.898 (1) 234.704 (2) 230.896 (3)
Simple RNN	2	32	Tanh	64	0.001	100	EarlyStopping ModelCheckpoint	0.143 (1) 0.171 (2) 0.264 (3)	0.035 (1) 0.063 (2) 0.118 (3)	0.187 (1) 0.251 (2) 0.343 (3)	70.990 (1) 68.699 (2) 72.500 (3)	0.178 (1) 0.171 (2) 0.200 (3)	0.052 (1) 0.071 (2) 0.084 (3)	0.228 (1) 0.266 (2) 0.289 (3)	69.624 (1) 78.075 (2) 67.416 (3)
Stateful RNN	3	32	Tanh	64	1.58e ⁻⁴	100	LearningRateScheduler EarlyStopping	0.203 (1) 0.349 (3)	0.060 (1) 0.173 (3)	0.244 (1) 0.416 (3)	56.390 (1) 82.819 (3)	0.155 (1) 0.223 (2)	0.039 (1) 0.079 (2)	0.197 (1) 0.282 (2)	59.988 (1) 64.159 (3)
Stateful LSTM	3	32	Tanh	64	1.58e ⁻⁴	100	EarlyStopping	0.095 (1) 0.151 (2) 0.174 (3)	0.021 (1) 0.048 (2) 0.076 (3)	0.146 (1) 0.220 (2) 0.275 (3)	40.335 (1) 48.937 (2) 55.662 (3)	0.098 (1) 0.134 (2) 0.166 (3)	0.020 (1) 0.042 (2) 0.071 (3)	0.141 (1) 0.205 (2) 0.267 (3)	41.781 (1) 57.860 (2) 68.025 (3)
Stateful Bi-LSTM	3	32	Tanh	64	1.58e ⁻⁴	100	EarlyStopping	0.149 (1) 0.190 (2) 0.249 (3)	0.043 (1) 0.074 (2) 0.120 (3)	0.207 (1) 0.272 (2) 0.347 (3)	58.151 (1) 60.154 (2) 67.988 (3)	0.170 (1) 0.211 (2) 0.229 (3)	0.049 (1) 0.090 (2) 0.108 (3)	0.221 (1) 0.300 (2) 0.329 (3)	71.059 (1) 92.727 (2) 87.049 (3)
1D-CNN LSTM	3	32 (5.1)*	ReLU	64	1.58e ⁻⁴	100	EarlyStopping	0.108 (1) 0.146 (2) 0.177 (3)	0.027 (1) 0.051 (2) 0.078 (3)	0.165 (1) 0.226 (2) 0.279 (3)	41.164 (1) 47.512 (2) 53.087 (3)	0.098 (1) 0.138 (2) 0.156 (3)	0.023 (1) 0.047 (2) 0.067 (3)	0.151 (1) 0.217 (2) 0.259 (3)	51.732 (1) 68.376 (2) 69.338 (3)

All the calculations and model runs were implemented under the framework of TensorFlow 2.3.0 (?) in Python 3.6.13. The models were executed on Ubuntu 20.04.1 LTS OS with 4 × GPUs (NVIDIA GeForce RTX 2080 Ti, 11019 MiB, 300 MHz). According to the Keras API guide (?), the requirements to use the cuDNN implementation are the activation function must be set to *tanh* and the recurrent activation must be set to *sigmoid*. We also set the seed number to 7 across all the model runs to maintain reproducibility.

Stateful RNNs can be difficult to work with when using callbacks in Keras because their hidden state must be manually managed across mini-batch updates. When training a stateful RNN in Keras, the hidden state is carried over from the previous epoch and can cause problems with certain callbacks, such as *EarlyStopping* or *ModelCheckpoint*. To work around this issue, one can use stateless RNNs or manually reset the hidden state at the end of each epoch, but this can be complex and prone to errors.

A.2.5 Skill Scores

Skill scores and ratios are commonly used in evaluating the performance of classification models, particularly in binary classification tasks. They provide insights into the model’s ability to correctly predict positive and negative instances. Here is a brief description of each skill score and ratio, along with their formulas:

- **True Positive (TP):** The number of data points or intervals correctly identified as positive by the model. It represents instances where both the model and the ground truth indicate the presence of an event.
- **True Negative (TN):** The number of intervals correctly identified as negative by the model. It represents instances where both the model and the ground truth indicate the absence of an event.
- **False Positive (FP):** The number of intervals incorrectly identified as positive by the model. It occurs when the model predicts an event, but the ground truth indicates its absence.
- **False Negative (FN):** The number of intervals incorrectly identified as negative by the model. It occurs when the model fails to detect an event that the ground truth indicates its presence.
- **Accuracy:** Represents the proportion of correct predictions out of total predictions.

$$Accuracy = \frac{TP + TN}{TP + TN + FP + FN} \quad (\text{A.10})$$

- **Precision:** Represents the proportion of positive predictions that are actually positive.

$$Precision = \frac{TP}{TP + FP} \quad (\text{A.11})$$

- **Probability of Detection (POD) or Recall:** Represents the model's ability to correctly identify positive instances.

$$POD = \frac{TP}{TP + FN} \quad (\text{A.12})$$

- **Probability of False Detection (POFD):** Measures the model's tendency to falsely predict positive instances when the ground truth indicates their absence.

$$POFD = \frac{FP}{FP + TN} \quad (\text{A.13})$$

- **False Alarm Rate (FAR):** Indicates the ratio of false positive predictions to the total number of positive instances.

$$FAR = \frac{FP}{FP + TP} \quad (\text{A.14})$$

- **Critical Success Index (CSI):** Measures the model's ability to correctly predict both positive and negative instances.

$$CSI = \frac{TP}{TP + FP + FN} \quad (\text{A.15})$$

- **True Skill Statistic (TSS):** Takes into account both the model's ability to detect positive instances and its ability to avoid false alarms.

$$TSS = POD - FAR \quad (\text{A.16})$$

- **Heidke Skill Score (HSS):** Evaluates the model's performance by comparing it with random chance. It takes into account the agreement between the model's predictions and the observed data, considering both true positive and true negative predictions.

$$HSS = \frac{TP + TN - C}{T - C} \quad (\text{A.17})$$

where

$$T = TP + TN + FP + FN$$

$$C = \frac{(TP + FP)(TP + FN) + (TN + FP)(TN + FN)}{T}$$

Bibliography

- 2017, Nature Astronomy, 1, 771
- Abe, O., Fakomiti, M., Igboama, W., et al. 2023, Advances in Space Research, 71, 2240
- Akansu, A. N. 1991, Optical Engineering, 30, 912
- Akasofu, S. I. 1981, Space Science Reviews, 28, 121
- Alharbi, F. R. & Csala, D. 2021, in 2021 International Conference on Electrical, Communication, and Computer Engineering (ICECCE), IEEE, 1–6
- Alielden, K. & Taroyan, Y. 2022, The Astrophysical Journal, 935, 66
- Altschuler, M. D. & Newkirk, G. 1969, Solar Physics, 9, 131
- Amari, T., Canou, A., & Aly, J.-J. 2014, Nature, 514, 465
- Aminalragia-Giamini, S., Raptis, S., Anastasiadis, A., et al. 2021, Journal of Space Weather and Space Climate, 11, 59
- Aran, A., Sanahuja, B., & Lario, D. 2006, Advances in Space Research, 37, 1240
- Aschwanden, M. J. 2002, Space Science Reviews, 101, 1
- Aschwanden, M. J. 2010, Solar Physics, 262, 235
- Badman, S. T., Carley, E., Cañizares, L. A., et al. 2022, The Astrophysical Journal, 938, 95
- Bale, S., Goetz, K., Harvey, P., et al. 2016, Space science reviews, 204, 49
- Bein, B. M., Berkebile-Stoiser, S., Veronig, A. M., et al. 2011, apj, 738, 191
- Benson, B., Pan, W. D., Prasad, A., Gary, G. A., & Hu, Q. 2020, Solar Physics, 295, 65
- Benz, A. & Thejappa, G. 1988
- Benz, A. O. 2017, Living reviews in solar physics, 14, 1
- Berghmans, D., Auchère, F., Long, D. M., et al. 2021, Astronomy & Astrophysics, 656, L4
- Besliu-Ionescu, D., Maris Muntean, G., & Dobrica, V. 2022, Solar Physics, 297, 65
- Biesecker, D., Myers, D., Thompson, B., Hammer, D., & Vourlidas, A. 2002, The Astrophysical Journal, 569, 1009
- Bonnin, X., Hoang, S., & Maksimovic, M. 2008, Astron. Astrophys., 489, 419
- Borovsky, J. E. & Denton, M. H. 2006, Journal of Geophysical Research: Space Physics, 111
- Boudjada, M. Y., Abou el Fadl, A., Galopeau, P. H., Al-Haddad, E., & Lammer, H. 2020, Advances in Radio Science, 18, 83
- Brewer, D. A., Barth, J. L., Label, K. A., Kauffman, W. J., & Giffin, G. 2002, Acta Astronautica, 51, 609
- Bruno, A. & Richardson, I. G. 2021, Solar Physics, 296, 36

- Camporeale, E. 2019, Space weather, 17, 1166
- Cane, H. V., Erickson, W., & Prestage, N. 2002, Journal of Geophysical Research: Space Physics, 107, SSH
- Cattell, C., Glesener, L., Leiran, B., et al. 2021, Astron. Astrophys., 650, A6
- Che, H. 2018, Journal of Physics: Conference Series, 1100, 012005
- Chen, N., Ip, W.-H., & Innes, D. 2013, The Astrophysical Journal, 769, 96
- Chen, P. 2016, Low-Frequency Waves in Space Plasmas, 379
- Chen, P. F., Wu, S. T., Shibata, K., & Fang, C. 2002, The Astrophysical Journal, 572, L99
- Chen, X., Kontar, E. P., Chrysaphi, N., et al. 2023, arXiv e-prints, arXiv:2306.09160
- Chhabra, S., Klimchuk, J. A., & Gary, D. E. 2021, The Astrophysical Journal, 922, 128
- Cohen, C. & Mewaldt, R. 2018, Space Weather, 16, 1616
- Curto, J. J., Blanca, M., & Martínez, E. 2008, Solar Physics, 250, 411
- Dabrowski, B., Flisek, P., Mikuła, K., et al. 2021, Remote Sensing, 13, 148
- Dalla, S., Swalwell, B., Battarbee, M., et al. 2017in (Cambridge University Press), 268–271
- Debrunner, H., Flückiger, E., Grädel, H., Lockwood, J., & McGuire, R. 1988, Journal of Geophysical Research: Space Physics, 93, 7206
- Del Zanna, G., Aulanier, G., Klein, K.-L., & Török, T. 2011, Astronomy & Astrophysics, 526, A137
- Delannée, C. & Aulanier, G. 1999, Solar Physics, 190, 107
- Desai, M. & Giacalone, J. 2016, Living Reviews in Solar Physics, 13, 3
- Dierckxsens, M., Tziotziou, K., Dalla, S., et al. 2015, Solar Physics, 290, 841
- Dulk, G., Steinberg, J.-L., Lecacheux, A., Hoang, S., & MacDowall, R. J. 1985, Astronomy and Astrophysics, 150, L28
- Dumbović, M., Veronig, A., Podladchikova, T., et al. 2021, Astronomy & Astrophysics, 652, A159
- Dungey, J. 1961, Journal of Geophysical Research, 66, 1043
- Eastwood, J., Nakamura, R., Turc, L., Mejnertsen, L., & Hesse, M. 2017, Space Science Reviews, 212, 1221
- Echer, E. & Gonzalez, W. 2022, Advances in Space Research, 70, 2830
- Echer, E., Tsurutani, B., & Gonzalez, W. 2013, Journal of Geophysical Research: Space Physics, 118, 385
- Engell, A., Falconer, D., Schuh, M., Loomis, J., & Bissett, D. 2017, Space Weather, 15, 1321
- Ergun, R., Larson, D., Lin, R., et al. 1998, The Astrophysical Journal, 503, 435
- Feynman, J., Armstrong, T., Dao-Gibner, L., & Silverman, S. 1990, Solar Physics, 126, 385
- Fletcher, L., Dennis, B. R., Hudson, H. S., et al. 2011, Space science reviews, 159, 19
- Florios, K., Kontogiannis, I., Park, S.-H., et al. 2018, Solar Physics, 293, 28
- Fox, N., Velli, M., Bale, S., et al. 2016, Space Science Reviews, 204, 7
- Fredkin, D. R., Rice, J. A., Colquhoun, D., & Gibb, A. 1995, Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences, 350, 353
- Gabriel, S., Evans, R., & Feynman, J. 1990, Solar physics, 128, 415

- Gieseler, J., Dresing, N., Palmroos, C., et al. 2023, Frontiers in Astronomy and Space Sciences, 9, 384
- Gonzalez, W., Joselyn, J.-A., Kamide, Y., et al. 1994, Journal of Geophysical Research: Space Physics, 99, 5771
- González, W. D., Echer, E., Clua-Gonzalez, A., & Tsurutani, B. T. 2007, Geophysical research letters, 34
- Goodfellow, I., Bengio, Y., & Courville, A. 2016, Deep learning (MIT press)
- Gopalswamy, N. 2022, Atmosphere, 13, 1781
- Gopalswamy, N., Akiyama, S., Yashiro, S., Michalek, G., & Lepping, R. 2008, Journal of Atmospheric and Solar-Terrestrial Physics, 70, 245
- Gopalswamy, N., Mäkelä, P., Yashiro, S., et al. 2017in , IOP Publishing, 012009
- Gopalswamy, N., Mäkelä, P., Yashiro, S., Akiyama, S., & Xie, H. 2022a, in 2022 3rd URSI Atlantic and Asia Pacific Radio Science Meeting (AT-AP-RASC), 1–4
- Gopalswamy, N. & Yashiro, S. 2011, apjl, 736, L17
- Gopalswamy, N., Yashiro, S., Akiyama, S., et al. 2022b, Journal of Geophysical Research: Space Physics, 127, e2022JA030404
- Graves, A. & Jaitly, N. 2014in (Beijing, China: PMLR), 1764–1772
- Graves, A. & Schmidhuber, J. 2005, Neural networks, 18, 602
- Greff, K., Srivastava, R. K., Koutník, J., Steunebrink, B. R., & Schmidhuber, J. 2016, IEEE transactions on neural networks and learning systems, 28, 2222
- Guo, F. & Giacalone, J. 2013, The Astrophysical Journal, 773, 158
- Gurnett, D. A. & Anderson, R. R. 1976, Science, 194, 1159
- Gurnett, D. A. & Anderson, R. R. 1977, Journal of Geophysical Research, 82, 632
- Hale, G. E., Ellerman, F., Nicholson, S. B., & Joy, A. H. 1919, Astrophys. J., 49, 153
- Harra, L., Brooks, D. H., Bale, S. D., et al. 2021, Astronomy & Astrophysics, 650, A7
- Harvey, J. W., Hill, F., Hubbard, R. P., et al. 1996, Science, 272, 1284
- Hochreiter, S. & Schmidhuber, J. 1997, Neural computation, 9, 1735
- Holman, G. D., Aschwanden, M. J., Aurass, H., et al. 2011, Space science reviews, 159, 107
- Holschneider, M., Kronland-Martinet, R., Morlet, J., & Tchamitchian, P. 1989, in Wavelets. Time-Frequency Methods and Phase Space, ed. J.-M. Combes, A. Grossmann, & P. Tchamitchian (Springer-Verlag), 286
- Hu, J., Li, G., Ao, X., Zank, G. P., & Verkhoglyadova, O. 2017, Journal of Geophysical Research: Space Physics, 122, 10
- Huang, X., Tan, H., Lin, G., & Tian, Y. 2018, 2018 International Conference on Artificial Intelligence and Big Data (ICAIBD), 185
- Ihianle, I. K., Nwajana, A. O., Ebenuwa, S. H., et al. 2020, IEEE Access, 8, 179028
- Ireland, J., Inglis, A. R., Shih, A. Y., et al. 2019, Solar Physics, 294, 158
- Jackson, B., Hick, P., Buffington, A., et al. 2010in , American Institute of Physics, 659–662
- Jang, S., Moon, Y.-J., Kim, R.-S., Lee, H., & Cho, K.-S. 2016, The Astrophysical Journal, 821, 95
- Kahler, S., Cliver, E., Cane, H., et al. 1987, in Proceedings of the 20th International Cosmic Ray Conference Moscow, Volume 3, p. 121

- Kahler, S., Kazachenko, M., Lynch, B., & Welsch, B. 2017in , IOP Publishing, 012011
- Kahler, S., Sheeley Jr, N., Howard, R., et al. 1984, Journal of Geophysical Research: Space Physics, 89, 9683
- Kahler, S. W., Cliver, E. W., & Ling, A. G. 2007, Journal of Atmospheric and Solar-Terrestrial Physics, 69, 43
- Kane, R. 2011, Indian Journal of Radio & Space Physics, 40, 7
- Kasapis, S., Zhao, L., Chen, Y., et al. 2022, Space Weather, 20, e2021SW002842
- Kay, C. & Gopalswamy, N. 2018, Journal of Geophysical Research: Space Physics, 123, 7220
- Ketkar, N. & Ketkar, N. 2017, Deep learning with python: a hands-on introduction, 97
- Kim, T., Park, E., Lee, H., et al. 2019, Nature Astronomy, 3, 397
- Kingma, D. P. & Ba, J. 2015, International Conference on Learning Representations (ICLR), San Diego
- Klassen, A., Karlický, M., & Mann, G. 2003a, Astronomy & Astrophysics, 410, 307
- Klassen, A., Pohjolainen, S., & Klein, K.-L. 2003b, Solar Physics, 218, 197
- Klein, K.-L. & Dalla, S. 2017, Space Science Reviews, 212, 1107
- Kolen, J. F. & Kremer, S. C. 2001, Gradient Flow in Recurrent Nets: The Difficulty of Learning LongTerm Dependencies (IEEE), 237–243
- Kong, D., Pan, G., Yan, X., Wang, J., & Li, Q. 2018, The Astrophysical Journal Letters, 863, L22
- Kontar, E. P., Chen, X., Chrysaphi, N., et al. 2019, The Astrophysical Journal, 884, 122
- Kontar, E. P., Emslie, A. G., Clarkson, D. L., et al. 2023, arXiv preprint arXiv:2308.05839
- Kontar, E. P., Yu, S., Kuznetsov, A., et al. 2017, Nature communications, 8, 1515
- Kóta, J., Manchester, W., Jokipii, J., De Zeeuw, D., & Gombosi, T. 2005in , American Institute of Physics, 201–206
- Kouloumvakos, A., Rodríguez-García, L., Gieseler, J., et al. 2022, Frontiers in Astronomy and Space Sciences, 9, 974137
- Kozarev, K., Nedal, M., Miteva, R., Dechev, M., & Zucca, P. 2022, Frontiers in Astronomy and Space Sciences, 9, 34
- Kozarev, K. A., Davey, A., Kendrick, A., Hammer, M., & Keith, C. 2017, Journal of Space Weather and Space Climate, 7, A32
- Kozarev, K. A., Dayeh, M. A., & Farahat, A. 2019, The Astrophysical Journal, 871, 65
- Kozarev, K. A., Evans, R. M., Schwadron, N. A., et al. 2013, The Astrophysical Journal, 778, 43
- Kozarev, K. A., Korreck, K. E., Lobzin, V. V., Weber, M. A., & Schwadron, N. A. 2011, The Astrophysical Journal, 733, L25
- Kozarev, K. A., Raymond, J. C., Lobzin, V. V., & Hammer, M. 2015, The Astrophysical Journal, 799, 167
- Kozarev, K. A. & Schwadron, N. A. 2016, apj, 831, 120
- Krucker, S., Hudson, H. S., Jeffrey, N. L. S., et al. 2011, The Astrophysical Journal, 739, 96
- Krupar, V., Szabo, A., Maksimovic, M., et al. 2020, Astrophys. J. Suppl., 246, 57
- Kwon, R.-Y., Zhang, J., & Olmedo, O. 2014, The Astrophysical Journal, 794, 148
- Laitinen, T. & Dalla, S. 2017, The Astrophysical Journal, 834, 127

- Lakhina, G. S. & Tsurutani, B. T. 2016, *Geoscience Letters*, 3, 1
- Lanzerotti, L. J. 2001, Washington DC American Geophysical Union Geophysical Monograph Series, 125, 11
- Laurenza, M., Cliver, E., Hewitt, J., et al. 2009, *Space Weather*, 7
- Lavasa, E., Giannopoulos, G., Papaioannou, A., et al. 2021, *Solar Physics*, 296, 107
- Le, G.-M. & Zhang, X.-F. 2017, *Research in Astronomy and Astrophysics*, 17, 123
- Lecacheux, A., Steinberg, J. L., Hoang, S., & Dulk, G. A. 1989, *Astron. Astrophys.*, 217, 237
- Lemen, J. R., Title, A. M., Akin, D. J., et al. 2012, *Solar Physics*, 275, 17
- Li, B. & Cairns, I. H. 2012, *The Astrophysical Journal*, 753, 124
- Li, G., Shalchi, A., Ao, X., Zank, G., & Verkhoglyadova, O. 2012, *Advances in Space Research*, 49, 1067
- Li, G., Zank, G., & Rice, W. 2003, *Journal of Geophysical Research: Space Physics*, 108
- Li, R. & Zhu, J. 2013, *Research in Astronomy and Astrophysics*, 13, 1118
- Lilensten, J., Coates, A. J., Dehant, V., et al. 2014, *The Astronomy and Astrophysics Review*, 22, 1
- Lin, R. 2005, *Advances in Space Research*, 35, 1857
- Lin, R. 2011, *Space science reviews*, 159, 421
- Liu, W. & Ofman, L. 2014, *Solar Physics*, 289, 3233
- Long, D. M., Bloomfield, D. S., Chen, P.-F., et al. 2017, *Solar physics*, 292, 1
- Long, D. M., Bloomfield, D. S., Gallagher, P. T., & Pérez-Suárez, D. 2014, *Solar Physics*, 289, 3279
- Long, D. M., DeLuca, E. E., & Gallagher, P. T. 2011, *apjl*, 741, L21
- Ma, S., Raymond, J. C., Golub, L., et al. 2011, *The Astrophysical Journal*, 738, 160
- MacDowall, R., Lara, A., Manoharan, P., et al. 2003, *Geophysical research letters*, 30
- Madjarska, M. S. 2019, *Living Reviews in Solar Physics*, 16, 2
- Magdalenić, J., Marqué, C., Zhukov, A. N., Vršnak, B., & Žic, T. 2010, *Astrophys. J.*, 718, 266
- Malandraki, O. E. & Crosby, N. B. 2018, Solar particle radiation storms forecasting and analysis: the HESPERIA HORIZON 2020 project and beyond
- Mann, G., Breitling, F., Vocks, C., et al. 2018, *A&A*, 611, A57
- Manu, V., Balan, N., Zhang, Q.-H., & Xing, Z.-Y. 2022, *Journal of Geophysical Research: Space Physics*, 127, e2022JA030747
- Melrose, D. 1980, *Space Science Reviews*, 26, 3
- Melrose, D. 2017, *Reviews of Modern Plasma Physics*, 1, 1
- Mierla, M., Inhester, B., Antunes, A., et al. 2010in , Copernicus GmbH, 203–215
- Mikić, Z., Linker, J. A., Schnack, D. D., Lionello, R., & Tarditi, A. 1999, *Physics of Plasmas*, 6, 2217
- Miteva, R., Klein, K.-L., Malandraki, O., & Dorrian, G. 2013, *Solar Physics*, 282, 579
- Miteva, R., Nedal, M., Samwel, S. W., & Temmer, M. 2023, *Universe*, 9, 179
- Moore, R. L., Sterling, A. C., Hudson, H. S., & Lemen, J. R. 2001, *The Astrophysical Journal*, 552, 833
- Morosan, D. E. & Gallagher, P. T. 2017, in *Planetary Radio Emissions VIII*, ed. G. Fischer, G. Mann, M. Panchenko, & P. Zarka, 357–368

- Morton, R., Tomczyk, S., & Pinto, R. 2015, Nature Communications, 6, 7813
- Muhr, N., Veronig, A. M., Kienreich, I. W., et al. 2014, Solar Physics, 289, 4563
- Nammous, M. K., Saeed, K., & Kobojeck, P. 2022, Journal of King Saud University - Computer and Information Sciences, 34, 764
- Nedal, M., Kozarev, K., Zhang, P., & Pietro, Z. 2023, A&A, 680, A106
- Nedal, M., Mahrous, A., & Youssef, M. 2019, Astrophysics and Space Science, 364, 161
- Newkirk, G. 1967, Annual Review of Astronomy and Astrophysics, 5, 213
- Newkirk, Gordon, J. 1961, *Astrophys. J.*, 133, 983
- Ng, C. K., Reames, D. V., & Tylka, A. J. 2012in , American Institute of Physics, 212–218
- Nieves-Chinchilla, T., Vourlidas, A., Raymond, J. C., et al. 2018, Solar Physics, 293, 1
- Nindos, A., Aurass, H., Klein, K. L., & Trottet, G. 2008, Sol. Phys., 253, 3
- Nitta, N. V., Schrijver, C. J., Liu, W., et al. 2013, The Astrophysical Journal, 776, 58
- Nymmk, R. 2007, Advances in Space Research, 40, 321
- Núñez, M. 2011, Space Weather, 9
- Odstrcil, D., Riley, P., & Zhao, X. 2004, Journal of Geophysical Research: Space Physics, 109
- Offringa, A. R., McKinley, B., Hurley-Walker, N., et al. 2014, Mon. Not. R. Astron. Soc., 444, 606
- Olah, C. 2015, Neural Networks, Types, and Functional Programming, [Blog post]
- Ontiveros, V. & Vourlidas, A. 2009, apj, 693, 267
- Oughton, E. J., Skelton, A., Horne, R. B., Thomson, A. W., & Gaunt, C. T. 2017, Space Weather, 15, 65
- Pala, Z. & Atici, R. 2019, Solar Physics, 294, 50
- Paouris, E., Vourlidas, A., Papaioannou, A., & Anastasiadis, A. 2021, Space Weather, 19, e2020SW002617
- Papaioannou, A., Anastasiadis, A., Kouloumvakos, A., et al. 2018, Solar Physics, 293, 1
- Papaioannou, A., Vainio, R., Raukunen, O., et al. 2022, Journal of Space Weather and Space Climate, 12, 24
- Park, J., Innes, D. E., Bucik, R., & Moon, Y.-J. 2013, The Astrophysical Journal, 779, 184
- Parker, E. N. 1960, Astrophysical Journal, 132, 821
- Patsourakos, S. & Vourlidas, A. 2012, Solar Physics, 281, 187
- Patsourakos, S., Vourlidas, A., & Stenborg, G. 2010, *Astrophys. J. Lett.*, 724, L188
- Pérez-Suárez, D., Higgins, P. A., Bloomfield, D. S., et al. 2011, Automated Solar Feature Detection for Space Weather Applications, 207–225
- Pesnell, W. D., Thompson, B. J., & Chamberlin, P. C. 2012, solphys, 275, 3
- Pick, M. 2006, in Solar and Heliospheric Origins of Space Weather Phenomena, ed. J.-P. Rozelot, Vol. 699, 119
- Podladchikova, O. & Berghmans, D. 2005, Solar Physics, 228, 265
- Pomoell, J. & Poedts, S. 2018, Journal of Space Weather and Space Climate, 8, A35
- Priest, E. & Forbes, T. 2007, Magnetic Reconnection

- Pulkkinen, T. 2007, Living Reviews in Solar Physics, 4, 1
- Pulupa, M., Bale, S., Bonnell, J., et al. 2017, Journal of Geophysical Research: Space Physics, 122, 2836
- Pulupa, M., Bale, S. D., Badman, S. T., et al. 2020, *Astrophys. J. Suppl.*, 246, 49
- Pysz, M. A., Foygel, K., Panje, C. M., et al. 2011, *Investigative radiology*, 46, 187
- Qiu, S., Zhang, Z., Yousof, H., et al. 2022, *Advances in Space Research*, 70, 2047
- Ramstad, R., Holmström, M., Futaana, Y., et al. 2018, *Geophysical Research Letters*, 45, 7306
- Reames, D. V. 1999, *Space Science Reviews*, 90, 413
- Reames, D. V. 2000in , American Institute of Physics, 289–300
- Reames, D. V. 2013, *Space Science Reviews*, 175, 53
- Reames, D. V. 2021
- Reid, H. A. 2020, *Frontiers in Astronomy and Space Sciences*, 7, 56
- Reid, H. A. & Kontar, E. P. 2018a, *Astronomy & Astrophysics*, 614, A69
- Reid, H. A. & Kontar, E. P. 2018b, *The Astrophysical Journal*, 867, 158
- Reid, H. A. & Vilmer, N. 2017, *Astronomy & Astrophysics*, 597, A77
- Reid, H. A. S. & Ratcliffe, H. 2014, *Research in Astronomy and Astrophysics*, 14, 773
- Reiner, M. J., Goetz, K., Fainberg, J., et al. 2009, *Solar Physics*, 259, 255
- Richardson, I., von Rosenvinge, T., & Cane, H. 2016, *Solar Physics*, 291, 2117
- Ripley, B. D. 1996, *Pattern Recognition and Neural Networks* (Cambridge University Press)
- Rodriguez, J., Onsager, T., & Mazur, J. 2010, *Geophysical Research Letters*, 37
- Rouillard, A. P., Plotnikov, I., Pinto, R. F., et al. 2016, *The Astrophysical Journal*, 833, 45
- Rouillard, A. P., Sheeley, N. R., Tylka, A., et al. 2012, *The Astrophysical Journal*, 752, 44
- Saint-Hilaire, P., Vilmer, N., & Kerdraon, A. 2012, *The Astrophysical Journal*, 762, 60
- Saiz, E., Cerrato, Y., Cid, C., et al. 2013, *Journal of Space Weather and Space Climate*, 3, A26
- Samwel, S. & Miteva, R. 2023, *Advances in Space Research*, 72, 3440
- Schatten, K. H., Wilcox, J. M., & Ness, N. F. 1969, *Solar Physics*, 6, 442
- Schrijver, C. J. 2015, *Space Weather*, 13, 524
- Schrijver, C. J., Kauristie, K., Aylward, A. D., et al. 2015, *Advances in Space Research*, 55, 2745
- Schrijver, C. J. & Siscoe, G. L. 2010a, *Heliophysics: Evolving solar activity and the climates of space and Earth* (Cambridge University Press)
- Schrijver, C. J. & Siscoe, G. L. 2010b, *Heliophysics: Evolving solar activity and the climates of space and Earth* (Cambridge University Press)
- Schuster, M. & Paliwal, K. K. 1997, *IEEE transactions on Signal Processing*, 45, 2673
- Schwenn, R. 2006, *Living reviews in solar physics*, 3, 1
- Selvakumaran, R., Veenadhari, B., Akiyama, S., et al. 2016, *Journal of Geophysical Research: Space Physics*, 121, 8188
- Shibata, K. & Magara, T. 2011, *Living Reviews in Solar Physics*, 8, 1

- Singh, P., Manure, A., Singh, P., & Manure, A. 2020, Learn TensorFlow 2.0: Implement Machine Learning and Deep Learning Models with Python, 1
- Sokolov, I. V., Roussev, I. I., Skender, M., Gombosi, T. I., & Usmanov, A. V. 2009, *The Astrophysical Journal*, 696, 261
- Stansby, D., Yeates, A., & Badman, S. T. 2020, *Journal of Open Source Software*, 5, 2732
- Starck, J.-L. & Murtagh, F. 2002, *Astronomical image and data analysis* (Springer Netherlands)
- Stepanyuk, O., Kozarev, K., & Nedal, M. 2022, *Journal of Space Weather and Space Climate*, 12, 20
- Sundermeyer, M., Alkhouri, T., Wuebker, J., & Ney, H. 2014, in *Proceedings of the 2014 conference on empirical methods in natural language processing (EMNLP)* (Association for Computational Linguistics), 14–25
- Suzuki, S. & Dulk, G. 1985, Cambridge: Cambridge University Press, 289
- Švestka, Z. 1995, *Advances in Space Research*, 16, 27
- Szenicer, A., Fouhey, D. F., Munoz-Jaramillo, A., et al. 2019, *Science Advances*, 5, eaaw6548
- Temmer, M. 2016, *Astronomische Nachrichten*, 337, 1010
- Temmer, M. 2021, *Living Reviews in Solar Physics*, 18, 4
- Thernisien, A. 2011, *The Astrophysical Journal Supplement Series*, 194, 33
- Thernisien, A., Vourlidas, A., & Howard, R. 2009, *Solar Physics*, 256, 111
- Thompson, B., Plunkett, S., Gurman, J., et al. 1998, *Geophysical Research Letters*, 25, 2465
- Thompson, B. & Young, C. 2016, *The Astrophysical Journal*, 825, 27
- Thompson, B. J. & Myers, D. C. 2009, *Astrophys. J. Suppl.*, 183, 225
- Thompson, W. T. 2006, *A&A*, 449, 791
- Trottet, G., Samwel, S., Klein, K.-L., Dudok de Wit, T., & Miteva, R. 2015, *Solar Physics*, 290, 819
- Truscott, P., Lei, F., Dyer, C., et al. 2000, in *2000 IEEE Radiation Effects Data Workshop. Workshop Record. Held in conjunction with IEEE Nuclear and Space Radiation Effects Conference (Cat. No. 00TH8527)*, IEEE, 147–152
- Tsurutani, B. T. 1985
- Tsurutani, B. T., Gonzalez, W. D., & Kamide, Y. 1997, *Surveys in geophysics*, 18, 363
- Vainio, R., Desorgher, L., Heynderickx, D., et al. 2009, *Space science reviews*, 147, 187
- Vainio, R. & Laitinen, T. 2008, *Journal of Atmospheric and Solar-Terrestrial Physics*, 70, 467
- van Diepen, G., Dijkema, T. J., & Offringa, A. 2018, *Astrophysics Source Code Library*, ascl
- van Haarlem, M. P., Wise, M. W., Gunst, A., et al. 2013, *Astronomy & astrophysics*, 556, A2
- Verbeeck, C., Delouille, V., Mampaey, B., & De Visscher, R. 2014, *Astronomy and Astrophysics*, 561, A29
- Verbeke, C., Mays, M. L., Kay, C., et al. 2022, *Advances in Space Research*
- Veronig, A., Muhr, N., Kienreich, I., Temmer, M., & Vršnak, B. 2010, *The Astrophysical Journal Letters*, 716, L57
- Vourlidas, A., Syntelis, P., & Tsinganos, K. 2012, *Solar Physics*, 280, 509
- Vourlidas, A., Wu, S. T., Wang, A. H., Subramanian, P., & Howard, R. A. 2003, *apj*, 598, 1392
- Vršnak, B. & Cliver, E. W. 2008, *Solar Physics*, 253, 215

- Vršnak, B., Žic, T., Vrbanec, D., et al. 2013, Solar physics, 285, 295
- Warmuth, A. 2015, Living Reviews in Solar Physics, 12, 1
- Webb, D. F. & Howard, T. A. 2012, Living Reviews in Solar Physics, 9, 1
- Whitman, K., Egeland, R., Richardson, I. G., et al. 2023, Advances in Space Research, 72, 5161, cOSPAR Space Weather Roadmap 2022: Scientific Research and Applications
- Wild, J. 1950a, Australian Journal of Chemistry, 3, 399
- Wild, J. 1950b, Australian Journal of Chemistry, 3, 541
- Wild, J. & McCready, L. 1950, Australian Journal of Chemistry, 3, 387
- Wild, J., Smerd, S., & Weiss, A. 1963, Annual Review of Astronomy and Astrophysics, 1, 291
- Wills-Davey, M., DeForest, C., & Stenflo, J. O. 2007, The Astrophysical Journal, 664, 556
- Wilson, J. W., Townsend, L. W., Chun, S. Y., et al. 1988, BRYNTRN: A baryon transport computer code, computation procedures and data base, Tech. rep.
- Wöllmer, M., Zhang, Z., Weninger, F., Schuller, B., & Rigoll, G. 2013, in 2013 IEEE International Conference on Acoustics, Speech and Signal Processing, IEEE, 6822–6826
- Wood, B., Howard, R., & Socker, D. 2010, The Astrophysical Journal, 715, 1524
- Wu, C.-C. & Lepping, R. P. 2016, Solar Physics, 291, 265
- Wu, C. S., Wang, C. B., Yoon, P. H., Zheng, H. N., & Wang, S. 2002, The Astrophysical Journal, 575, 1094
- Wuelser, J.-P., Lemen, J. R., Tarbell, T. D., et al. 2004, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 5171, Telescopes and Instrumentation for Solar Astrophysics, ed. S. Fineschi & M. A. Gummin, 111–122
- Xapsos, M. A., Stauffer, C. A., Jordan, T. M., Adams, J. H., & Dietrich, W. F. 2012, IEEE Transactions on Nuclear Science, 59, 1054
- Xie, H., Ofman, L., & Lawrence, G. 2004, Journal of Geophysical Research: Space Physics, 109
- Young, P. R., Tian, H., Peter, H., et al. 2018, Space Science Reviews, 214, 1
- Zhang, J., Cheng, X., & Ding, M.-d. 2012, Nature communications, 3, 747
- Zhang, J. & Dere, K. P. 2006, apj, 649, 1100
- Zhang, J., Richardson, I., Webb, D., et al. 2007, Journal of Geophysical Research: Space Physics, 112
- Zhang, P., Wang, C., Ye, L., & Wang, Y. 2019, Solar Physics, 294, 62
- Zhang, P., Wang, C. B., & Ye, L. 2018, Astronomy & Astrophysics, 618, A165
- Zhang, P., Zucca, P., Kozarev, K., et al. 2022a, The Astrophysical Journal, 932, 17
- Zhang, W., Zhao, X., Feng, X., et al. 2022b, Universe, 8, 30
- Zhu, H., Zhu, W., & He, M. 2022, Solar Physics, 297, 157