1. The probability of a leap year selected at random contain 53							
Sunday is:							
	(b) 1/7						
2. A bag contains	3 red and 2 bl	ue marbles. A	marble is drawn at				
random. The prob	ability of draw	i <mark>ng a black </mark> ba	ıll is :				
(a) 3/5	(b) 2/5	(c) 0/5	(d) 1/5				
3. The probability	that it will rair	n tomorrow is	0.85. What is the				
probability that it							
(a) 0.25	(b) 0.145	(c) 3/20	(d) none of these				
4. What is the pro	bability that a	number selec	eted from the numbers				
(1, 2 <u>, 3,</u> ,15)							
(a) 1/5	(b) 4/5	(c) 2/15	(d) 1/3				
5. What are the t	otal outcomes	when we thro	w three coins?				
(a) 4	` '		` '				
		number selec	ted at random from the				
numbers (1,2,3,	<u></u> 35) is :	•					
			5 (d) none of these				
7. The sum of the							
	(b) 1 (c) (
•	•	re given; choo	ose the correct answer				
for that which is n	ot possible.						
(a) 0.15	(b) 2/7	(c) 7/5	(d) none of these.				
9. If three coins a	are tossed simu	ult <mark>aneously,</mark> th	nan the probability of				
getting at least tw	o heads, is: 📮						
(a) 1/4	(b) 3/8	(c) ½	(d) 1/8				
10. A letter is ch	osen at randon	n from the let	ters of the word				
ASSASSINATIO	N. The prob	ability that the	e letter chosen has:				
(a) 6/13	(b) 7/13	(c) 1	(d) none of these.				
			_				
			etting an even number.				
(A) 2/3	(B) 1 ((C) 5/6	(D) 1/2				
12. Two coins are thrown at the same time. Find the probability of							
		same time. Fi	na the probability of				
getting both head		(D) 0					
(A) 3/4 (B) 1/4	(C) 1/2	(D) 0					
13. Two dice are thrown simultaneously. The probability of getting a							

1

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/9					
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.								
• • •	(B) 27/50	(C) 1/4	(D) 2°	9/100				
_	a blue ball is d	oalls and some ouble that of a (C) 15		-				
taken out a non-defect	16. A box of 600 bulbs contains 12 defective bulbs. One bulb is taken out at random from this box. Then the probability that it is non-defective bulb is:							
(A) 143/150 (B) 147/150 (C) 1/25 (D) 1/50 17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (A) 9/100 (B) 1/10 (C) 3/10 (D) 19/100								
18. What is (A) 1/7	=	y of getting 53 (C) 2/7	_					
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13								
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312 ,then the probability that it will point to an odd number is: (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12								
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4								

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:								
(A) 364/365	(B) 31/365	(C) 1/365	(D) 1/133225					
23. A number x is chosen at random from the numbers -2, -1, 0 , 1, 2. Then the probability that $x^2 < 2$ is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$								
a marble is dra red is 2/3, the	awn at random f	rom the jar, the pr white marbles in	-					
25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) $7/50$ (B) $4/25$ (C) $1/25$ (D) $2/25$								
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?								
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$					
27. Runs scored by batsman in 5 one day matches are 50, 70, 82, 93, and 20. The standard deviation is								
		c) 25.29	d) 25.69					
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.								
	b) 13, 18		d) 13, 16					
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is								
a) $^{1}/_{2}$	b) $^{1}/_{3}$		d) $^{1}/_{6}$					
		nd 3. The value of c) 27 d						
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?								

32.Out of the probability?		alues, which	one is not poss	ible in
a) $P(x) = 1$	b) ∑ x 5 d) P(x	P(x) = 3 x) = -0.5		
33.If E(x) = a) 2	2 and E(z) = 4 b) 6	c) 0	•	ufficient data
34.The cova	ariance of two	independen	t random variab	le is
a) 1	b) 0	c) - 1	d) Un	defined
, ,) = k ² – 8 then, b) 1			sufficient data
, ,	0.5 and x = 4, b) 0.5	• •	d) 2	
37.In a disc is always?	rete probabilit	y distributio	n, the sum of all	probabilities
a) 0	b) Infinite	c) 1	d) Und	defined
38.If the pr	obability of hi	tting the targ	get is 0.4, find m	ean and
	b) 0.6,	0.24	c) 0.4, 0.16	d) 0.6, 0.16
-	% and if 10 bo	-	ped from a plac pped, find mean .4, 0.16	
a) 2	e mean of toss b) 4 s the mean and	c) 8	d) 1 r standard norm	al distribution?

c) 5

d) 7

a) 3

b) 4

43.Mean of a random variable X is given by a) E(X) b) E(X2) c) E(X2) - (E(X))2 d) 44.Mean of a constant 'a' is a) 0 b) a c) a/2 d) 1 45.Variance of a constant 'a' is a) 0 b) a c) a/2 d) 1 46.Find the mean and variance of X?	•		and varia and varia		•				
a) E(X) b) E(X2) c) E(X2) - (E(X))2 d) 44.Mean of a constant 'a' is a) 0 b) a c) a/2 d) 1 45.Variance of a constant 'a' is a) 0 b) a c) a/2 d) 1 46.Find the mean and variance of X?								_ · d) (E(X))2	
a) 0 b) a c) a/2 d) 1 45. Variance of a constant 'a' is a) 0 b) a c) a/2 d) 1 46. Find the mean and variance of X?					_	•		d) (E(X))2	
a) 0 b) a c) a/2 d) 1 46.Find the mean and variance of X? x 0 1 2 3 4							d) 1		
x 0 1 2 3 4									
	46.F	ind the	mean and	variance	of X?				
(1) 1/0 2/0 2/0 2/0	г 	Х	0	1	2	3	4		
T(X) 1/9 2/9 3/9 2/9 1/9		f(x)	1/9	2/9	3/9	2/9	1/9		

c) 2, 2/3

d) 3, 2/3

47. Find the expectation of a random variable X?

b) 3, 4/3

	X	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

a) 2, 4/3

b) npq

c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2 d) \sqrt{npq}