


STRUCTURE /

| | |
|------|----------|
| NH | Ammonia |
| R-NH | 1° Amine |
| R-NH | 2° Amine |
| R-N | 3° Amine |



Pyramidal geometry

10. AMINES

PHYSICAL PROPERTIES /

PHYSICAL STATE: Lower aliphatic amines are gases, intermediate members are liquid (fishy odour), while higher members are solid.


SOLUBILITY: Lower aliphatic amines are soluble in water due to H-bonding, while higher amines ($>C$) are insoluble in water.

$$\text{Solubility} \propto \frac{1}{\text{Molecular weight}}$$

BOILING POINT: Primary and secondary amines form intermolecular H-bonding while tertiary does not.


$$1^\circ \text{ Amine} > 2^\circ \text{ Amine} > 3^\circ \text{ Amine}$$

CHEMICAL PROPERTIES /




PREPARATION /


Reduction of Nitro Compounds.




Ammonolysis



Reduction of Nitriles




Reduction of Amides




Hofmann Bromamide Degradation reaction

one carbon less amine is formed as compared to amide



Gabriel Phthalimide synthesis



Aromatic primary amines cannot be prepared by this method


BASIC NATURE /

Due to the presence of lone pair on nitrogen amines are basic.


Factors affecting basicity

- Inductive effect
- solvation effect
- Steric hindrance


In Gaseous Phase




In Aqueous Phase




Overall Basicity Order





TEST FOR AMINES / - HINSBERG'S TEST



ELECTROPHILIC SUBSTITUTION




DIAZONIUM SALTS



PHYSICAL PROPERTIES
 Colourless, soluble in water, decompose in dry state.
 $\text{C}_6\text{H}_5\text{N}_2^+\text{Cl}^-$ is readily soluble in water.

CHEMICAL REACTIONS



IMPORTANCE

In preparation of substituted aromatic compounds which cannot be prepared by direct substitution in benzene.