Introduction to quadratic equations

Tom Coleman

Summary

Solving quadratic equations of the form $ax^2 + bx + c$ is a core skill in mathematics. A guaranteed method to solve these is use of the quadratic formula. This guide explains the terminology and usage of the quadratic formula, as well as an explanation of why the quadratic formula works.

Before reading this guide, it is recommended that you read (Guide: Introduction to complex numbers).

What is a quadratic equation?

Equations in one variable come in all shapes and sizes.

One of the most important types of an equation is known as a **quadratic equation**. These appear almost everywhere in mathematics, from modelling projectile motion in mechanics to describing circles, ellipses, parabolas and hyperbolas in 2D space.

i Definition of quadratic equation, root

A quadratic equation is an equation that can be rearranged into the form

$$ax^2 + bx + c = 0$$

where x is a variable and a, b, c are real numbers with $a \neq 0$.

Values of x that satisfy the equation $ax^2 + bx + c = 0$ are known as **roots** of the equation.

Here, $a \neq 0$ as if it was, then the equation would no longer be a quadratic equation!

The general shape of a quadratic equation is known as a **parabola**. A figure of two parabolas is given in Figure 1; the left hand graph is if a>0, whereas the right hand graph is when a<0.

It is very important to be able to identify the variable in a quadratic equation, as well as the coefficients a,b,c. In the course of your mathematical study, it may be that the variable of a quadratic equation is not only letters like x,y,z, but squares or cubes like x^2 and y^3 , or even functions like e^x , $\cos(x)$, and $\sin(y)$.




Figure 1: A pair of parabolas. (left) A graph of a quadratic $ax^2 + bx + c$ where a > 0. (right) A graph of a quadratic $ax^2 + bx + c$ where a < 0.

i Example 1

You are given the quadratic equation $2x^2 + 4x - 8 = 0$. The variable of the quadratic equation is x, and the coefficients are a = 2, b = 4, c = -8.

Example 2

Here, the equation $y^4-10y^2+25=0$ may look like a quartic equation, but it is actually a quadratic equation. Using the laws of indices, you can rewrite the equation as $(y^2)^2-10y^2+25=0$. Therefore, the variable of the quadratic equation is y^2 , and the coefficients are a=1, b=-10, c=25.

i Example 3

You are given the equation $-e^{2x}+4e^x-5=0$. Using the laws of indices, you can rewrite the equation as $-(e^x)^2+4e^x-5=0$. The variable of the quadratic equation is e^x , and the coefficients are a=-1, b=4, c=-5. This is not the only solution to the coefficients; you can multiply the equation through by -1 to get $(e^x)^2-4e^x+5=0$, which gives a=1, b=-4 and c=5. Both solutions are equally valid.

i Example 4

You are given the equation $t+1=\frac{4}{t-3}$. This really is a quadratic equation! You can multiply both sides by x-3 to get

$$(t+1)(t+3) = 4$$

You can then expand the brackets to get

$$t^2 + t + 3t + 3 = 4$$

and so $t^2+4t+3=4$. Finally, you are able to subtract 4 from both sides to get $t^2+4t-1=0$. It follows that the variable of the quadratic equation is t, and the coefficients are a=1, b=4, c=-1.

Solving a quadratic equation

To solve the quadratic equation, you could use one of three methods:

- You could **factorise** the quadratic equation $ax^2 + bx + c = 0$ into linear equations (mx+n)(px+q), then work out the roots when each of these linear equations is zero. See (Guide: Factorisation) for more.
- You could **complete the square** in order to reduce the quadratic equation $ax^2 + bx + c = 0$ into the form $(x + b/2a)^2 = d$, and then solve from there (not forgetting the negative root). See (Guide: Completing the square) for more.
- You could **use the quadratic formula**; for a quadratic equation $ax^2 + bx + c = 0$, the two roots to the quadratic equation are given by

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Each method is equally valid, but some may involve more work than others. It is up to you to decide which method is best for each quadratic you encounter; but it is thoroughly recommended that if you are not sure which method is best, then the quadratic formula is the one to choose. See Guide: Using the quadratic formula for more.

The discriminant

What the roots of the quadratic formula look like are determined by the term b^2-4ac ; this term has a special name.

i The discriminant

The term $D=b^2-4ac$ is known as the **discriminant** of the quadratic equation $ax^2+bx+c=0$.

There are then three separate cases for solutions to quadratic equations.

• If $D=b^2-4ac$ is positive, then \sqrt{D} is a real number and the two roots of the quadratic equation $ax^2+bx+c=0$ are

$$x = \frac{-b + \sqrt{D}}{2a} \qquad \text{and} \qquad x = \frac{-b - \sqrt{D}}{2a}$$

These two roots are both real numbers and distinct from each other. You can observe this behaviour on a graph in Figure 2; the parabola crosses the x-axis in two places.




Figure 2: A pair of parabolas. (left) A graph of a quadratic $ax^2 + bx + c$ where a > 0 and D > 0. (right) A graph of a quadratic $ax^2 + bx + c$ where a < 0 and D > 0.

• If $D=b^2-4ac=0$ is zero, then $\sqrt{D}=0$. In this case, the two roots of the quadratic equation $ax^2+bx+c=0$ are

$$x = \frac{-b}{2a} \qquad \text{and} \qquad x = \frac{-b}{2a}$$

These two roots are given by the same real number. To be sure that you express both roots, you can write 'x=-b/2a twice'. You can observe this behaviour on a graph in Figure 3; the parabola touches the x-axis in exactly one place.

4




Figure 3: A pair of parabolas. (left) A graph of a quadratic $ax^2 + bx + c$ where a > 0 and D = 0. (right) A graph of a quadratic $ax^2 + bx + c$ where a < 0 and D = 0.

• If $D=b^2-4ac$ is negative, then \sqrt{D} is not a real number. In this case, the two roots of the quadratic equation are **complex numbers**. You can express the two roots of the quadratic equation by

$$x = \frac{-b + i\sqrt{-D}}{2a} \qquad \text{and} \qquad x = \frac{-b - i\sqrt{-D}}{2a}$$

where i is the imaginary unit (so $i^2=-1$; see (Guide: Introduction to complex numbers)). In a graph, the parabola does not cross the x-axis at all; this indicates that there are no real solutions to this quadratic equation. See Figure 4 for a picture.




Figure 4: A pair of parabolas. (left) A graph of a quadratic $ax^2 + bx + c$ where a > 0 and D > 0. (right) A graph of a quadratic $ax^2 + bx + c$ where a < 0 and D < 0.

Warning

You can use the discriminant to check how many roots a quadratic equation has in the variable given to you. However, this is at most a maximum number of solutions. Conditions on that variable may also reduce the number of valid solutions, particularly if you have real valued functions. For instance, since $e^x>0$ for all real number x, there are no solutions in x if you find $e^x=-1$.

Here's some examples of using the discriminant and known properties of functions to rule out solutions.

i Example 5

In Example 1, you identified the coefficients of $2x^2+4x-8=0$ as a=2,b=4,c=-8. Using these, you can work out the value of the discriminant $D=b^2-4ac$ as

$$D = (4)^2 - 4(2)(-8) = 16 + 64 = 80.$$

Since D=80, you can say that this quadratic equation has two distinct real roots in $x=r_1$ and $x=r_2$.

i Example 6

In Example 2, you identified the coefficients of $y^4-10y^2+25=0$ as a=1,b=-10,c=25, and the variable as y^2 . Using these, you can work out the value of the discriminant $D=b^2-4ac$ as

$$D = (-10)^2 - 4(1)(25) = 100 - 100 = 0.$$

Since D=0, you can say that this quadratic equation has at most one real root $u^2=r$.

Whether or not the equation itself has real solutions in y depends on whether r is positive or negative!

i Example 7

In Example 3, you identified the coefficients of $-e^{2x}+4e^x-5=0$ as a=-1,b=4,c=-5, and the variable as e^x . Using these, you can work out the value of the discriminant $D=b^2-4ac$ as

$$D = (4)^2 - 4(-1)(-5) = 16 - 20 = -4.$$

Since D=-4, you can say that this quadratic equation has complex roots. This equation therefore has no real solutions in x. This is because e^x is real for any real x; if e^x is complex, it follows that x cannot be real.

i Example 8

In Example 4, you rearranged the equation $t+1=\frac{4}{t-3}$ to $t^2+4t-1=0$, and therefore identified the coefficients as a=1,b=4,c=-1, and the variable as t. Using these, you can work out the value of the discriminant $D=b^2-4ac$ as

$$D = (4)^2 - 4(1)(-1) = 16 + 4 = 20.$$

Since D=20, you can say that this quadratic equation has two distinct real roots in $t. \ \ \,$

Quick check problems

- 1. What is the discriminant of the quadratic equation $x^2 x 1 = 0$?
- 2. You are given the quadratic equation $4h^2 h + 101 = 0$. Identify the variable, and the coefficients a,b,c.
- 3. You are given three statements below. Decide whether they are true or false.
- (a) The quadratic equation $m^2 + 4m + 4 = 0$ has two distinct real roots.
- (b) The quadratic equation $m^2 4m 4 = 0$ has exactly one real root.
- (c) The quadratic eqution $4m^2 + 4m + 4 = 0$ has no real roots.

For more questions on the subject, please click on this sentence.

Further reading

(Guide: Using the quadratic formula)