

# MCAC 201: Design and Analysis of Algorithms

Neelima Gupta

ngupta@cs.du.ac.in

January 5, 2025

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

# Asymptotic Notations 1: Big O


Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .



# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = 3n^2$  and  $g(n) = n^2 - 2n$ .

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = 3n^2$  and  $g(n) = n^2 - 2n$ .

$3n^2 \leq 3n^2 + n^2 - 8n$  for all  $n \geq 8$ ,

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = 3n^2$  and  $g(n) = n^2 - 2n$ .

$$\begin{aligned} 3n^2 &\leq 3n^2 + n^2 - 8n \text{ for all } n \geq 8, \\ &= 4(n^2 - 2n) \end{aligned}$$

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = 3n^2$  and  $g(n) = n^2 - 2n$ .

$$\begin{aligned} 3n^2 &\leq 3n^2 + n^2 - 8n \text{ for all } n \geq 8, \\ &= 4(n^2 - 2n) \end{aligned}$$

Thus,  $c = 4$  and  $n_0 = 8$ .



# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = n + 3\frac{n}{\log n}, g(n) = n$

$n + 3\frac{n}{\log n} \leq 4n$  for all  $n \geq 2$

Thus,  $c = 4, n_0 = 2$ .

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = n + 3\frac{n}{\log n}, g(n) = n$

$n + 3\frac{n}{\log n} \leq 4n$  for all  $n \geq 2$

Thus,  $c = 4, n_0 = 2$ .

- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = n + 3\frac{n}{\log n}, g(n) = n$

$n + 3\frac{n}{\log n} \leq 4n$  for all  $n \geq 2$

Thus,  $c = 4, n_0 = 2$ .

- **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

- ▶ **More Examples:**

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{7n^2 - 2n - 5}{5n^2 + 3n + 10} = \lim_{n \rightarrow \infty} \frac{7 - \frac{2}{n} - \frac{5}{n^2}}{5 + \frac{3}{n} + \frac{10}{n^2}} = \frac{7}{5}.$$

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

- ▶ **More Examples:**

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{7n^2 - 2n - 5}{5n^2 + 3n + 10} = \lim_{n \rightarrow \infty} \frac{7 - \frac{2}{n} - \frac{5}{n^2}}{5 + \frac{3}{n} + \frac{10}{n^2}} = \frac{7}{5}.$$

**Computing  $c$  and  $n_0$ :**  $7n^2 - 2n - 5 \leq \frac{7}{5}(5n^2 + 3n + 10)$  for all  $n \geq 1$ .

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

- ▶ **More Examples:**  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n+10}{n^2-100} = 0.$

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

- ▶ **More Examples:**  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n+10}{n^2-100} = 0$ .

**Computing  $c$  and  $n_0$ :**  $2(n^2 - 100) = n^2 + n^2 - 200 \geq n + 10$   
for  $n \geq \sqrt{210}$ .

$\geq n + 10$  for  $n \geq 1$ .  $c = 2, n_0 = \max\{1, \sqrt{210}\}$



# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

- ▶ **Notation:**  $f(n) \leq_{\infty} g(n)$  asymptotically.

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

- ▶ **Notation:**  $f(n) \leq_{\infty} g(n)$  asymptotically.  
By asymptotically we mean within constant multiplicative factor and for large  $n$

# Asymptotic Notations 1: Big O

Consider a set  $\mathcal{F}$  of non-negative, monotonically non-decreasing functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

- ▶ **Notation:**  $f(n) \leq_{\infty} g(n)$  asymptotically.

By asymptotically we mean within constant multiplicative factor and for large  $n$

We say that  $f(n)$  grows at most as fast as  $g(n)$  with  $n$  asymptotically.

## Asymptotic Notations 2: Omega


- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .

## Asymptotic Notations 2: Omega

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .

## Asymptotic Notations 2: Omega

- $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .



$$f(n) = \Omega(g(n))$$

## Asymptotic Notations 2: Omega

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = n^2 - 2n$  and  $g(n) = 3n^2$ .

$4(n^2 - 2n) = 3n^2 + n^2 - 8n$  for all  $n \geq 8$ ,

## Asymptotic Notations 2: Omega

- $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = n^2 - 2n$  and  $g(n) = 3n^2$ .

$$4(n^2 - 2n) = 3n^2 + n^2 - 8n \text{ for all } n \geq 8, \\ \geq 3n^2$$



## Asymptotic Notations 2: Omega

- $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$   
for all  $n \geq n_o$ .

Examples:

Let  $f(n) = n^2 - 2n$  and  $g(n) = 3n^2$ .

$$4(n^2 - 2n) = 3n^2 + n^2 - 8n \text{ for all } n \geq 8, \\ \geq 3n^2$$

Thus,  $c = 1/4$  and  $n_0 = 8$ .

## Asymptotic Notations 2: Omega

- $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$   
for all  $n \geq n_o$ .

Examples:

Let  $f(n) = n^2 - 2n$  and  $g(n) = 3n^2$ .

$$4(n^2 - 2n) = 3n^2 + n^2 - 8n \text{ for all } n \geq 8, \\ \geq 3n^2$$

Thus,  $c = 1/4$  and  $n_0 = 8$ .

## Asymptotic Notations 2: Omega

- $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = 3\frac{n^2}{\log n} + n, g(n) = n$

$3\frac{n^2}{\log n} + n \geq 4n$  for all  $n \geq 1$

Thus,  $c = 4, n_0 = 1$ .

- **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

## Asymptotic Notations 2: Omega

- $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = 3\frac{n^2}{\log n} + n, g(n) = n$

$3\frac{n^2}{\log n} + n \geq 4n$  for all  $n \geq 1$

Thus,  $c = 4, n_0 = 1$ .

- **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

## Asymptotic Notations 2: Omega

- $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = 3\frac{n^2}{\log n} + n, g(n) = n$

$3\frac{n^2}{\log n} + n \geq 4n$  for all  $n \geq 1$

Thus,  $c = 4, n_0 = 1$ .

- **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

## Asymptotic Notations 2: Omega

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

## Asymptotic Notations 2: Omega

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

- ▶ **More Examples:**

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{5n^2 + 3n + 10}{7n^2 - 2n - 5} = \lim_{n \rightarrow \infty} \frac{5 + \frac{3}{n} + \frac{10}{n^2}}{7 - \frac{2}{n} - \frac{5}{n^2}} = \frac{5}{7}.$$

## Asymptotic Notations 2: Omega

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_0 > 0 : f(n) \geq cg(n)$  for all  $n \geq n_0$ .
- ▶ **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

- ▶ **More Examples:**

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{5n^2 + 3n + 10}{7n^2 - 2n - 5} = \lim_{n \rightarrow \infty} \frac{5 + \frac{3}{n} + \frac{10}{n^2}}{7 - \frac{2}{n} - \frac{5}{n^2}} = \frac{5}{7}.$$

**Computing  $c$  and  $n_0$ :**  $(5n^2 + 3n + 10) \leq \frac{5}{7}(7n^2 - 2n - 5)$  for all  $n \geq 1$ .



## Asymptotic Notations 2: Omega

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

- ▶ **More Examples:**  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n^2 - 100}{n + 10} = \infty.$

## Asymptotic Notations 2: Omega

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

- ▶ **More Examples:**  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n^2 - 100}{n + 10} = \infty$ .

Computing  $c$  and  $n_0$ :

$$2(n^2 - 100) = (n^2 + 10) + n^2 - 210 \geq n^2 + 10 \text{ for } n \geq \sqrt{210}.$$
$$\geq n + 10 \text{ for } n \geq 1. \quad c = 1/2, n_0 = \max\{1, \sqrt{210}\}$$

## Asymptotic Notations 2: Omega

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

- ▶ **Notation:**  $f(n) \geq_{\infty} g(n)$  asymptotically.  
We say that  $f(n)$  grows at least as fast as  $g(n)$  with  $n$  asymptotically.

## Asymptotic Notations 3: theta


- ▶  $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_o > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_o$ .

## Asymptotic Notations 3: theta

- ▶  $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_o > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_o$ .

## Asymptotic Notations 3: theta

- $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_o > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_o$ .



$$f(n) = \theta(g(n))$$

## Asymptotic Notations 3: theta

- $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_o > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = n^2 - 2n$  and  $g(n) = 3n^2$ .

$\frac{1}{4}3n^2 \leq (n^2 - 2n) \leq 3n^2$  for all  $n \geq 8$ ,

## Asymptotic Notations 3: theta

- $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_o > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_o$ .

Examples:

Let  $f(n) = n^2 - 2n$  and  $g(n) = 3n^2$ .

$\frac{1}{4}3n^2 \leq (n^2 - 2n) \leq 3n^2$  for all  $n \geq 8$ ,

i.e.,      Thus,  $(1/4)g(n) \leq f(n) \leq g(n)$  for all  $n \geq 8$ .



## Asymptotic Notations 3: theta

- $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_0 > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_0$ .

### Examples:

Let  $f(n) = n^2 - 2n$  and  $g(n) = 3n^2$ .

$\frac{1}{4}3n^2 \leq (n^2 - 2n) \leq 3n^2$  for all  $n \geq 8$ ,

i.e., Thus,  $(1/4)g(n) \leq f(n) \leq g(n)$  for all  $n \geq 8$ .

Thus,  $c_1 = 1/4, c_2 = 1$  and  $n_0 = 8$ .

## Asymptotic Notations 3: theta

- $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_0 > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_0$ .

### Examples:

Let  $f(n) = n^2 - 2n$  and  $g(n) = 3n^2$ .

$\frac{1}{4}3n^2 \leq (n^2 - 2n) \leq 3n^2$  for all  $n \geq 8$ ,

i.e., Thus,  $(1/4)g(n) \leq f(n) \leq g(n)$  for all  $n \geq 8$ .

Thus,  $c_1 = 1/4, c_2 = 1$  and  $n_0 = 8$ .

## Asymptotic Notations 3: theta

- ▶  $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_0 > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_0$ .

Examples:

Let  $f(n) = \log_2 n, g(n) = \log_3 n$

$\log_2 n = \log_3 2 \log_3 n$  for all  $n$ .

Thus,  $c_1 = c_2 = \log_3 2, n_0 = 1$ .

- ▶ **Equivalent Definition:**  $f(n) = \theta(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c < \infty$$

$c$  is a (non-zero) positive finite constant.  $c$  is neither 0 nor  $\infty$ .

## Asymptotic Notations 3: theta

- $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_0 > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_0$ .

Examples:

Let  $f(n) = \log_2 n, g(n) = \log_3 n$

$\log_2 n = \log_3 2 \log_3 n$  for all  $n$ .

Thus,  $c_1 = c_2 = \log_3 2, n_0 = 1$ .

- **Equivalent Definition:**  $f(n) = \theta(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c < \infty$$

$c$  is a (non-zero) positive finite constant.  $c$  is neither 0 nor  $\infty$ .

- **Example:**

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{7n^2 - 2n - 5}{5n^2 + 3n + 10} = \lim_{n \rightarrow \infty} \frac{7 - \frac{2}{n} - \frac{5}{n^2}}{5 + \frac{3}{n} + \frac{10}{n^2}} = \frac{7}{5}.$$

## Asymptotic Notations 3: theta

- $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_0 > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_0$ .

Examples:

Let  $f(n) = \log_2 n, g(n) = \log_3 n$

$\log_2 n = \log_3 2 \log_3 n$  for all  $n$ .

Thus,  $c_1 = c_2 = \log_3 2, n_0 = 1$ .

- **Equivalent Definition:**  $f(n) = \theta(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c < \infty$$

$c$  is a (non-zero) positive finite constant.  $c$  is neither 0 nor  $\infty$ .

- **Example:**

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{7n^2 - 2n - 5}{5n^2 + 3n + 10} = \lim_{n \rightarrow \infty} \frac{7 - \frac{2}{n} - \frac{5}{n^2}}{5 + \frac{3}{n} + \frac{10}{n^2}} = \frac{7}{5}.$$

- **Notation:**  $f(n) =_{\infty} g(n)$  asymptotically.

## Asymptotic Notations 3: theta

- ▶  $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_o > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \theta(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c < \infty$$

$c$  is a (non-zero) positive finite constant.  $c$  is neither 0 nor  $\infty$ .

- ▶ **Example:**

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{7n^2 - 2n - 5}{5n^2 + 3n + 10} = \lim_{n \rightarrow \infty} \frac{7 - \frac{2}{n} - \frac{5}{n^2}}{5 + \frac{3}{n} + \frac{10}{n^2}} = \frac{7}{5}.$$

- ▶ **Notation:**  $f(n) =_{\infty} g(n)$  asymptotically. We say that  $f(n)$  grows at the same rate as  $g(n)$  with  $n$ , asymptotically.

## Asymptotic Notations 4: small o

- ▶  $f(n) = o(g(n))$  iff for every constant  $c > 0 \exists n_c > 0$  :  
 $f(n) < cg(n)$  for all  $n \geq n_c$ .

## Asymptotic Notations 4: small o

- ▶  $f(n) = o(g(n))$  iff for every constant  $c > 0 \exists n_c > 0$  :  
 $f(n) < cg(n)$  for all  $n \geq n_c$ .
- ▶ **Equivalent Definition:**  $f(n) = o(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$



## Asymptotic Notations 4: small o

- ▶  $f(n) = o(g(n))$  iff for every constant  $c > 0 \exists n_c > 0$  :  
 $f(n) < cg(n)$  for all  $n \geq n_c$ .
- ▶ **Equivalent Definition:**  $f(n) = o(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

- ▶ **Example:**  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n+10}{n^2-100} = 0$ .

## Asymptotic Notations 4: small o

- ▶  $f(n) = o(g(n))$  iff for every constant  $c > 0 \exists n_c > 0$  :  
 $f(n) < cg(n)$  for all  $n \geq n_c$ .
- ▶ **Equivalent Definition:**  $f(n) = o(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

- ▶ **Example:**  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{\log^2 n}{\sqrt{n}} = 0.$

## Asymptotic Notations 4: small o

- ▶  $f(n) = o(g(n))$  iff for every constant  $c > 0 \exists n_c > 0$  :  
 $f(n) < cg(n)$  for all  $n \geq n_c$ .
- ▶ **Equivalent Definition:**  $f(n) = o(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

- ▶ **Example:**  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n^{10}}{2^n} = 0$ .
- ▶ **Notation:**  $f(n) <_{\infty} g(n)$  asymptotically.  
We say that  $f(n)$  grows strictly slower than  $g(n)$  with  $n$ , asymptotically.

## Asymptotic Notations 4: small o

- ▶  $f(n) = o(g(n))$  iff for every constant  $c > 0 \exists n_c > 0 :$   
 $f(n) < cg(n)$  for all  $n \geq n_c$ .
- ▶ **Equivalent Definition:**  $f(n) = o(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

- ▶ **Example:**  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{2^n}{3^n} = 0$ .
- ▶ **Notation:**  $f(n) <_{\infty} g(n)$  asymptotically.  
We say that  $f(n)$  grows strictly slower than  $g(n)$  with  $n$ , asymptotically.

## Asymptotic Notations 5: small omega

- ▶  $f(n) = \omega(g(n))$  iff for every constant  $c > 0 \exists n_c > 0$  :  
 $f(n) > cg(n)$  for all  $n \geq n_c$

## Asymptotic Notations 5: small omega

- ▶  $f(n) = \omega(g(n))$  iff for every constant  $c > 0 \exists n_c > 0$  :  
 $f(n) > cg(n)$  for all  $n \geq n_c$
- ▶ **Equivalent Definition:**  $f(n) = \omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

## Asymptotic Notations 5: small omega

- ▶  $f(n) = \omega(g(n))$  iff for every constant  $c > 0 \exists n_c > 0$  :  
 $f(n) > cg(n)$  for all  $n \geq n_c$
- ▶ **Equivalent Definition:**  $f(n) = \omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

- ▶ **Example:**  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n^2 - 100}{n + 10} = \infty$ .

## Asymptotic Notations 5: small omega

- ▶  $f(n) = \omega(g(n))$  iff for every constant  $c > 0 \exists n_c > 0 :$   
 $f(n) > cg(n)$  for all  $n \geq n_c$
- ▶ **Equivalent Definition:**  $f(n) = \omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

- ▶ **Example:**  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n^2 - 100}{n + 10} = \infty.$
- ▶ **Notation:**  $f(n) >_{\infty} g(n)$  asymptotically.  
We say that  $f(n)$  grows strictly faster than  $g(n)$  with  $n$ , asymptotically.



## Some easy to prove Observations

1.  $f(n) \leq g(n)$  iff either  $f(n) < g(n)$  or  $f(n) = g(n)$  is true.

## Some easy to prove Observations

1.  $f(n) \leq g(n)$  iff either  $f(n) < g(n)$  or  $f(n) = g(n)$  is true.
2.  $f(n) \leq g(n)$  iff  $g(n) \geq f(n)$ .

## Some easy to prove Observations

1.  $f(n) \leq g(n)$  iff either  $f(n) < g(n)$  or  $f(n) = g(n)$  is true.
2.  $f(n) \leq g(n)$  iff  $g(n) \geq f(n)$ .
3.  $f(n) = g(n)$  iff  $f(n) \leq g(n)$  and  $f(n) \geq g(n)$ .

Let's compare some popular functions (use *L'Hôpital's* Rule wherever required)

1.  $f(n) = \log n$ ,  $g(n) = n$

Let's compare some popular functions (use *L'Hôpital's* Rule wherever required)

1.  $f(n) = \log n$ ,  $g(n) = n$

$$\lim_{n \rightarrow \infty} \frac{\log n}{n} = \lim_{n \rightarrow \infty} \frac{1}{n \cdot \ln 2} = 0$$

Let's compare some popular functions (use *L'Hôpital's* Rule wherever required)

1.  $f(n) = \log n$ ,  $g(n) = n$

$$\lim_{n \rightarrow \infty} \frac{\log n}{n} = \lim_{n \rightarrow \infty} \frac{1}{n \cdot \ln 2} = 0$$

2.  $f(n) = n^4$ ,  $g(n) = 2^n$

Let's compare some popular functions (use *L'Hôpital's Rule* wherever required)

1.  $f(n) = \log n$ ,  $g(n) = n$

$$\lim_{n \rightarrow \infty} \frac{\log n}{n} = \lim_{n \rightarrow \infty} \frac{1}{n \cdot \ln 2} = 0$$

2.  $f(n) = n^4$ ,  $g(n) = 2^n$

$$\lim_{n \rightarrow \infty} \frac{n^4}{2^n} = \lim_{n \rightarrow \infty} \frac{4n^3}{2^n} = \lim_{n \rightarrow \infty} \frac{12n^2}{2^n} = \lim_{n \rightarrow \infty} \frac{24n}{2^n} = \lim_{n \rightarrow \infty} \frac{24}{2^n} = 0$$

Let's compare some popular functions (use *L'Hôpital's Rule* wherever required)

1.  $f(n) = \log n$ ,  $g(n) = n$

$$\lim_{n \rightarrow \infty} \frac{\log n}{n} = \lim_{n \rightarrow \infty} \frac{1}{n \cdot \ln 2} = 0$$

2.  $f(n) = n^4$ ,  $g(n) = 2^n$

$$\lim_{n \rightarrow \infty} \frac{n^4}{2^n} = \lim_{n \rightarrow \infty} \frac{4n^3}{2^n} = \lim_{n \rightarrow \infty} \frac{12n^2}{2^n} = \lim_{n \rightarrow \infty} \frac{24n}{2^n} = \lim_{n \rightarrow \infty} \frac{24}{2^n} = 0$$

3.  $f(n) = \log^2 n$ ,  $g(n) = \sqrt{n}$



Let's compare some popular functions (use *L'Hôpital's Rule* wherever required)

1.  $f(n) = \log n$ ,  $g(n) = n$

$$\lim_{n \rightarrow \infty} \frac{\log n}{n} = \lim_{n \rightarrow \infty} \frac{1}{n \cdot \ln 2} = 0$$

2.  $f(n) = n^4$ ,  $g(n) = 2^n$

$$\lim_{n \rightarrow \infty} \frac{n^4}{2^n} = \lim_{n \rightarrow \infty} \frac{4n^3}{2^n} = \lim_{n \rightarrow \infty} \frac{12n^2}{2^n} = \lim_{n \rightarrow \infty} \frac{24n}{2^n} = \lim_{n \rightarrow \infty} \frac{24}{2^n} = 0$$

3.  $f(n) = \log^2 n$ ,  $g(n) = \sqrt{n}$

$$\lim_{n \rightarrow \infty} \frac{\log^2 n}{\sqrt{n}} = \lim_{n \rightarrow \infty} \frac{2 \log n}{n} \cdot 2\sqrt{n} = \lim_{n \rightarrow \infty} \frac{4 \log n}{\sqrt{n}} = \lim_{n \rightarrow \infty} \frac{4}{n} \cdot 2\sqrt{n} = 0$$

Let's compare some popular functions (use *L'Hôpital's Rule* wherever required)

1.  $f(n) = \log n$ ,  $g(n) = n$

$$\lim_{n \rightarrow \infty} \frac{\log n}{n} = \lim_{n \rightarrow \infty} \frac{1}{n \cdot \ln 2} = 0$$

2.  $f(n) = n^4$ ,  $g(n) = 2^n$

$$\lim_{n \rightarrow \infty} \frac{n^4}{2^n} = \lim_{n \rightarrow \infty} \frac{4n^3}{2^n} = \lim_{n \rightarrow \infty} \frac{12n^2}{2^n} = \lim_{n \rightarrow \infty} \frac{24n}{2^n} = \lim_{n \rightarrow \infty} \frac{24}{2^n} = 0$$

3.  $f(n) = \log^2 n$ ,  $g(n) = \sqrt{n}$

$$\lim_{n \rightarrow \infty} \frac{\log^2 n}{\sqrt{n}} = \lim_{n \rightarrow \infty} \frac{2 \log n}{n} \cdot 2\sqrt{n} = \lim_{n \rightarrow \infty} \frac{4 \log n}{\sqrt{n}} = \lim_{n \rightarrow \infty} \frac{4}{n} \cdot 2\sqrt{n} = 0$$

4.  $f(n) = 2^n$ ,  $g(n) = 3^n$

$$\lim_{n \rightarrow \infty} \frac{2^n}{3^n} = \lim_{n \rightarrow \infty} \left(\frac{2}{3}\right)^n = 0$$

Let's compare some popular functions (use *L'Hôpital's Rule* wherever required)

1.  $f(n) = \log n$ ,  $g(n) = n$

$$\lim_{n \rightarrow \infty} \frac{\log n}{n} = \lim_{n \rightarrow \infty} \frac{1}{n \cdot \ln 2} = 0$$

2.  $f(n) = n^4$ ,  $g(n) = 2^n$

$$\lim_{n \rightarrow \infty} \frac{n^4}{2^n} = \lim_{n \rightarrow \infty} \frac{4n^3}{2^n} = \lim_{n \rightarrow \infty} \frac{12n^2}{2^n} = \lim_{n \rightarrow \infty} \frac{24n}{2^n} = \lim_{n \rightarrow \infty} \frac{24}{2^n} = 0$$

3.  $f(n) = \log^2 n$ ,  $g(n) = \sqrt{n}$

$$\lim_{n \rightarrow \infty} \frac{\log^2 n}{\sqrt{n}} = \lim_{n \rightarrow \infty} \frac{2 \log n}{n} \cdot 2\sqrt{n} = \lim_{n \rightarrow \infty} \frac{4 \log n}{\sqrt{n}} = \lim_{n \rightarrow \infty} \frac{4}{n} \cdot 2\sqrt{n} = 0$$

4.  $f(n) = 2^n$ ,  $g(n) = 3^n$

$$\lim_{n \rightarrow \infty} \frac{2^n}{3^n} = \lim_{n \rightarrow \infty} \left(\frac{2}{3}\right)^n = 0$$

# Practice Questions

Use the equivalent definitions (limits) to prove the following:

1. Show that a polynomial of degree  $d$ , with positive leading coefficient is  $\Theta(n^d)$ .
2. For  $g(n) = f(n) + o(f(n))$ , show that  $g(n) = \Theta(f(n))$ .
3. Show that
  - a.  $\log n = o(n)$ .
  - b.  $\log^M n = o(n^\epsilon)$  where  $M$  and  $\epsilon$  are positive constants.
4.  $a^n = o(b^n)$  for all  $a < b$ .
5.  $\log_a n = \theta(\log_b n)$ .