SIH26143 Innovation Strategy

The strongest student-feasible opportunity in SIH26143 is **not** building another oil-spill detector. It is building an **uncertainty-aware oil-spill attribution workflow** that ranks suspect vessels, explains why they are ranked, and explicitly handles missing or spoofed AIS, rather than claiming definitive identification from a single SAR scene.

Research Landscape

Current research is strong in **oil-spill detection from SAR**, where deep learning now substantially outperforms older handcrafted methods, but false positives from look-alikes, boundary blur, speckle, and dataset bias remain central problems (Jafarzadeh et al., 2021; Shaban et al., 2021; Li et al., 2024; Li et al., 2023; Yekeen et al., 2020). Public benchmarking has improved the field, but the literature still warns that many models are trained on narrow custom datasets, making cross-scene comparison and operational generalization difficult (Krestenitis et al., 2019; Hasimoto-Beltrán et al., 2023).

Detection accuracy improves when models fuse more than raw SAR intensity. Multi-channel SAR inputs that include wind and texture reached very high pixel accuracy, and self-evolving or multiscale architectures improved F1 or segmentation quality in difficult scenes (Hasimoto-Beltrán et al., 2023; Li et al., 2023; Li et al., 2024). False alarms also fall when SAR is fused with quad-pol features, GNSS-R, optical data, or semantic context such as wind and likely pollution sources (Jamal et al., 2025; Wan et al., 2026; Benito-Ortiz et al., 2026; Dong et al., 2025).

Title: Existing Capability Layers

Layer	What Exists	Main Limitation
Spill detection	SAR DL segmentation and instance detection (Shaban et al., 2021; Yekeen et al., 2020)	Look-alikes and poor generalization
Spill characterization	Thickness, damping, emulsification proxies (Garcia-Pineda et al., 2019; Meng et al., 2024)	Operational coverage is limited
Drift modeling	GNOME, MEDSLIK-II, OpenDrift, Lagrangian hindcasts (Prasad et al., 2020; Pärt et al., 2023; Liubartseva et al., 2016)	Sensitive to forcing uncertainty
AIS analytics	Trajectory cleaning, prediction, anomaly detection (Liang et al., 2024; Wolsing et al., 2022; Nguyen & Fablet, 2021)	Missing/spoofed AIS
Fusion/attribution	SAR+AIS association and suspect ranking (Luo et al., 2024; Rodger & Guida, 2020; Chang et al., 2024)	Rarely uncertainty-aware or explainable

FIGURE 1 Current technical layers in oil-spill attribution systems

Spill characterization is advancing, but mostly as a **response** product rather than an attribution product. Time-series SAR can identify persistent thick oil and short-term drift patterns, while SAR, thermal, optical, hyperspectral, and polarimetric methods can estimate relative thickness, emulsification, or equivalent oil thickness under specific conditions (Espeseth et al., 2020; Garcia-Pineda et al., 2019; Jiao et al., 2021; Jiang et al., 2025; Meng et al., 2024). But these products often depend on favorable viewing conditions, special sensors, or calibration regimes that are not continuously available to a student system (Leifer et al., 2012; Scafutto et al., 2025; Wan et al., 2026).

Biggest Unsolved Problems

The core unsolved problem is that **attribution is much harder than detection**. Several studies can detect spills, simulate drift, or associate SAR ships with AIS, but only a small subset directly traces a spill back to a probable ship, and those methods are still case-driven rather than fully operationalized (Luo et al., 2024; Yang et al., 2019; Chang et al., 2024).

Backward origin reconstruction remains highly sensitive to ocean currents, wind history, and random walk parameterization. Studies repeatedly show that hindcast accuracy improves when local drift factors are calibrated, when higher-resolution ocean currents are used, and when Stokes drift or multi-model forcing is included (Yu et al., 2016; Prasad et al., 2020; Pärt et al., 2023; Liubartseva et al., 2016; Prasad et al., 2021). More recent work goes further by using neural correction of timely-but-inaccurate wind or sea-surface dynamic fields to approximate delayed reanalysis, specifically to improve timely backtracking and drift prediction (Li et al., 2024; Ren et al., 2025).

AIS is also not trustworthy enough to serve as a sole attribution backbone. Raw AIS has noise and missingness that require reconstruction, and anomaly literature shows that gaps can arise from congestion, poor reception, equipment failure, or intentional disabling (Liang et al., 2024; Ford et al., 2018). Separate work shows spoofing, stealth deviations, and switchoffs can be detected statistically, but these methods are not yet tightly integrated into oil-spill attribution pipelines (d'Afflisio et al., 2021; Ioannou et al., 2024; Ford et al., 2018).

Title: Highest-Value Gaps

Gap	Why It Persists	What It Means
Spill age estimation	Weak direct labels and confounded weathering signals (Espedal, 1999; De Souza Júnior et al., 2024)	Origin time is poorly constrained
Vessel attribution	Detection, drift, and AIS are built separately (Park et al., 2025; Liu et al., 2023)	Rankings are brittle
Explainability	Maritime XAI is emerging, not standard (Murugan et al., 2026; Zhang & Xu, 2025)	Evaluators will distrust black boxes
Uncertainty	Operational systems rarely show calibrated uncertainty (Liubartseva et al., 2016)	False confidence is dangerous

Gap	Why It Persists	What It Means
Human investigation	Most pipelines optimize automation, not analyst workflow (Sharma, 2026; ■ Ribeiro et al., 2025)	Weak demo and poor usability

FIGURE 2 Most important unsolved gaps for student innovation

Research Gaps to Target

The best innovation space is **system integration around uncertainty and evidence**, not inventing a wholly new remote sensor. Multi-source methods already show that combining SAR, optical, AIS, weather, and contextual data improves discrimination and surveillance quality (■ La et al., 2024; Liu et al., 2023; Şahin et al., 2026; ■ Chen & Qiu, 2026). What is still missing is a student-feasible pipeline that turns these heterogeneous inputs into a **forensic evidence graph** for one spill event.

A credible pipeline would treat each module as partial evidence: spill mask confidence, spill direction, spill age range, hindcast origin cloud, AIS gap score, spoofing score, vessel proximity score, route parallelism, and behavior anomaly score (■ Li et al., 2023; ■ Espeseth et al., 2020; Espedal, 1999; ■ Yu et al., 2017; ■ Ford et al., 2018; d'Afflisio et al., 2021; ■ Millati & Choi, 2026; ■ Han et al., 2021). That is different from current papers that usually optimize one piece at a time.

Explainability should be designed into attribution, not added after ranking. The literature on explainable oil-spill detection and maritime AI supports showing model rationales, confidence, alternatives, and rule traces, while dark-vessel and forensic AI work explicitly argues for "triage and analyst support" rather than autonomous accusation (■ Murugan et al., 2026; ■ Lee & Youn, 2025; Boato et al., 2025; Sharma, 2026).

A second strong gap is **time alignment**. SAR-AIS association degrades because ships move between acquisition and report times, but several studies show this can be improved with velocity compensation, historical traffic knowledge, classification-aided association, and asynchronous trajectory matching (Graziano et al., 2019; Mazzarella et al., 2015; ■ Rodger & Guida, 2020; Guo et al., 2023). For SIH, a practical novelty is to apply those ideas specifically to oil-spill attribution rather than general vessel monitoring.

Top 5 Innovation Opportunities

Title: Innovation Scorecard

Idea	Novelty /10	Feasibility /10	SIH Impact /10	Research Value /10	Risk /10
Uncertainty-aware vessel attribution engine	8	8	10	9	5
AIS-off / spoof-aware suspect ranking	8	7	9	8	6
Spill age + hindcast time-window estimator	7	7	8	8	6

Idea	Novelty /10	Feasibility /10	SIH Impact /10	Research Value /10	Risk /10
Evidence-graph explainability dashboard	7	9	9	7	4
Human-in-the-loop investigation workspace	6	9	8	7	3

FIGURE 3 Scored innovation opportunities for SIH26143

1. Uncertainty-aware vessel attribution engine. Existing solutions can find candidate polluters using SAR, AIS, and drift similarity, but they usually output a likely ship rather than a calibrated ranked explanation (Luo et al., 2024). Your innovation is to output **Top-N suspects with confidence bands and evidence breakdown**, using uncertainty from drift, trajectory prediction, and association modules (Capobianco et al., 2022; Lu et al., 2025; Singh et al., 2022). Demonstrate it on one historical event with ablations: without uncertainty, with uncertainty, and with analyst overrides.

2. AIS-off / spoof-aware suspect ranking. Most ship-tracing pipelines assume AIS is present enough to screen candidates, but the AIS anomaly literature shows intentional disabling and spoofing are common enough to matter operationally (d'Afflisio et al., 2021; Ioannou et al., 2024; Ford et al., 2018). Your innovation is to penalize or escalate candidates based on reconstructed gaps, switchoff probability, and spoof indicators rather than dropping them from the search space (Liang et al., 2024; Ioannou et al., 2024; d'Afflisio et al., 2021).

3. Spill age + hindcast time-window estimator. Age estimation exists through wind history, repeat-pass SAR, and weathering/backscatter relationships, but it is rarely integrated into vessel ranking (Espedal, 1999; De Souza Júnior et al., 2024; Karunathilake et al., 2022). A student-feasible contribution is to estimate a **time-of-release interval** from slick geometry, wind history, and drift consistency, then filter AIS trajectories only within that interval.

4. Evidence-graph explainability dashboard. Knowledge-graph and explainable maritime AI studies suggest that structured reasoning across heterogeneous evidence is underused and valuable (Liu et al., 2023; Lee & Youn, 2025). Your dashboard should show exactly why vessel A outranks vessel B: proximity, heading alignment, gap anomaly, hindcast overlap, and uncertainty.

5. Human-in-the-loop investigation workspace. Operational GIS systems exist, but most focus on response forecasting rather than forensic attribution workflow (Fetissov et al., 2021; Yang et al., 2025; Sorgente et al., 2020). Build a case board where an analyst can adjust weights, inspect alternate suspects, and export an investigation brief.

Title: Top 5 Realistic Innovations

Rank	Innovation	Why Realistic
1	Uncertainty-aware suspect ranking	Builds from available models and public data
2	AIS-off / spoof-aware attribution	Uses existing AIS anomaly methods
3	Spill age constrained hindcasting	Moderate modeling, strong demo value

Rank	Innovation	Why Realistic
4	Explainable evidence graph	High evaluator clarity
5	Human-in-the-loop GIS investigator	Productizes research into workflow

FIGURE 4 Top five realistic innovations for a student team

Killer Differentiators and Positioning

The top 3 killer differentiators should be:

- **Ranked suspect vessels with evidence and uncertainty**, not a binary accusation (Liubartseva et al., 2016; Zhang & Xu, 2025).
- **AIS-off and spoof-aware attribution**, so the system remains useful when the guilty vessel hides (Ford et al., 2018; Ioannou et al., 2024).
- **Interactive forensic replay**, showing spill detection, backward drift cone, candidate trajectories, and why the final ranking changed (Kulawiak et al., 2010; Fetissov et al., 2021; Oliveira et al., 2014).

The strongest positioning is **Oil Spill Attribution Platform**, with a subtitle like **Maritime Forensic Intelligence for Pollution Events**. “Oil Spill Detection System” undersells the novelty because many detectors already exist (Krestenitis et al., 2019; Yekeen et al., 2020). “Maritime Domain Awareness Platform” is too broad and dilutes the problem statement (Şahin et al., 2026). “Oil Spill Attribution Platform” is the clearest evaluator-facing frame because the hard problem in the literature is not just seeing the slick but linking it to a plausible vessel under uncertainty (Luo et al., 2024; Yu et al., 2016; Rodger & Guida, 2020).

SIH Evaluator Perspective

Title: SIH Evaluator Score

Dimension	Score /10
Innovation	8
Technical complexity	9
AI/ML depth	8
Real-world impact	9
Government relevance	10
Feasibility	7
Data availability	7

Dimension	Score /10
Scalability	8
Demo potential	9
Research potential	9
Startup potential	8
Differentiation	8

FIGURE 5 Likely SIH evaluator scoring profile

This project could win because it sits at the intersection of environment, maritime security, geospatial AI, and public-sector relevance, and because a strong demo can make the forensic reasoning visually obvious. It also has real research depth across SAR AI, drift modeling, AIS analytics, and explainability (Yekeen & Balogun, 2020; Yang et al., 2024; Şahin et al., 2026).

This project could lose if the team promises **definitive culprit identification**. The literature does not support certain attribution from one scene, especially with missing AIS, uncertain drift forcing, and look-alikes (Espedal, 1999; Liubartseva et al., 2016; Sharma, 2026). It could also lose if the demo is just a U-Net on Sentinel-1 plus a map of nearby vessels, because that is too close to existing work (Chang et al., 2024).

Evaluators will likely ask: How do you validate the true vessel? How do you handle AIS gaps versus intentional disabling? What public datasets do you use? How fast can the system run after a new SAR scene? How do you explain wrong rankings? What happens when multiple vessels traverse the same corridor? Those questions follow directly from the literature's known limits in association, uncertainty, and operational latency (Rodger & Guida, 2020; Liubartseva et al., 2016; Naz et al., 2020; Sharma, 2026).

Avoid claims like "identifies the guilty vessel with certainty," "works in real time everywhere," or "solves spoofed AIS completely." The evidence supports **probabilistic suspect ranking and analyst support**, not legal-grade automated accusation (Sharma, 2026; Ford et al., 2018).

A memorable demo would replay a known spill event end-to-end: detect slick, estimate age window, hindcast likely origin cone, reconstruct AIS gaps, rank three suspect vessels, and show why rank 1 beats rank 2. Then let the evaluator toggle off one evidence source and watch the ranking change.

The single most important shift in the evidence is from **oil-spill detection** toward **multi-source, uncertainty-aware forensic attribution**. The single biggest open question is whether a student-built system can produce rankings that are accurate enough on real cases to be trusted as an operational decision-support tool rather than only a research demo.

These search results were found and analyzed using Consensus, an AI-powered search engine for research. Try it at https://consensus.app. © 2026 Consensus NLP, Inc. Personal, non-commercial use only; redistribution requires copyright holders' consent.

References

Benito-Ortiz, M., Mata-Moya, D., Del-Rey-Maestre, N., Hernández-Fernández, C., & Jabonero-Ranera, P. (2026). Application of multi-resolution techniques, intelligent classification and semantic data fusion for the identification of oil spills in SAR imagery.. *Marine pollution bulletin, 232*, 120015. https://doi.org/10.1016/j.marpolbul.2026.120015

Boato, G., Montibeller, A., Delp, E., Verdoliva, L., & Miorandi, D. (2025). Don't Guess, Escalate: Towards Explainable Uncertainty-Calibrated AI Forensic Agents. *ArXiv, abs/2512.16614*. https://doi.org/10.48550/arxiv.2512.16614

Capobianco, S., Forti, N., Millefiori, L. M., Braca, P., & Willett, P. (2022). Recurrent Encoder–Decoder Networks for Vessel Trajectory Prediction With Uncertainty Estimation. *IEEE Transactions on Aerospace and Electronic Systems, 59*, 2554-2565. https://doi.org/10.1109/taes.2022.3216823

Chang, L., Chen, Y.-T., & Cheng, C.-M. (2024). Combining SAR and AIS to track oil discharge vessels using the improved U-Net. 1314313 - 1314313-8. https://doi.org/10.1117/12.3027230

Chen, H., & Qiu, R. (2026). Ship Trajectory Analysis Based on the Integration of Multi-Source Remote Sensing and AIS Data. *Journal of Physics: Conference Series, 3178*. https://doi.org/10.1088/1742-6596/3178/1/012070

d'Afflisio, E., Braca, P., & Willett, P. (2021). Malicious AIS Spoofing and Abnormal Stealth Deviations: A Comprehensive Statistical Framework for Maritime Anomaly Detection. *IEEE Transactions on Aerospace and Electronic Systems, 57*, 2093-2108. https://doi.org/10.1109/taes.2021.3083466

De Souza Júnior, J. M. N., Mendonça, L., Da Silva Costa, H., De Freitas, R. A. P., Casagrande, F., Da Silva Lindemann, D., Reis, R. A. D. N., Lentini, C., & De Cunha Lima, A. T. (2024). Dispersion analysis of the 2017 Persian Gulf oil spill based on remote sensing data and numerical modelling.. *Marine pollution bulletin, 205*, 116639. https://doi.org/10.1016/j.marpolbul.2024.116639

Dong, S., Feng, J., Gu, Z., Yin, K., & Long, Y. (2025). A Review of Artificial Intelligence and Remote Sensing for Marine Oil Spill Detection, Classification, and Thickness Estimation. *Remote Sensing*. https://doi.org/10.3390/rs17223681

Espedal, H. (1999). Satellite SAR oil spill detection using wind history information. *International Journal of Remote Sensing, 20*, 49-65. https://doi.org/10.1080/014311699213596

Espeseth, M. M., Jones, C. E., Holt, B., Brekke, C., & Skrunes, S. (2020). Oil-Spill-Response-Oriented Information Products Derived From a Rapid-Repeat Time Series of SAR Images. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, 13*, 3448-3461. https://doi.org/10.1109/jstars.2020.3003686

Fetissov, M., Aps, R., Goerlandt, F., Jänes, H., Kotta, J., Kujala, P., & Szava-Kovats, R. (2021). Next-Generation Smart Response Web (NG-SRW): An Operational Spatial Decision Support System for Maritime Oil Spill Emergency Response in the Gulf of Finland (Baltic Sea). *Sustainability*. https://doi.org/10.3390/su13126585

Ford, J. H., Peel, D., Kroodsma, D., Hardesty, B. D., Rosebrock, U., & Wilcox, C. (2018). Detecting suspicious activities at sea based on anomalies in Automatic Identification Systems transmissions. *PLoS ONE, 13*. https://doi.org/10.1371/journal.pone.0201640

Garcia-Pineda, O., Staples, G., Jones, C. E., Hu, C., Holt, B., Kourafalou, V. H., Graettinger, G., DiPinto, L., Ramirez, E., Streett, D., Cho, J., Swayze, G., Sun, S., Garcia, D., & Haces-Garcia, F. (2019). Classification of oil spill by thicknesses using multiple remote sensors. *Remote Sensing of Environment, 236*, 111421. https://doi.org/10.1016/j.rse.2019.111421

Graziano, M., Renga, A., & Moccia, A. (2019). Integration of Automatic Identification System (AIS) Data and Single-Channel Synthetic Aperture Radar (SAR) Images by SAR-Based Ship Velocity Estimation for Maritime Situational Awareness. *Remote. Sens., 11*, 2196. https://doi.org/10.3390/rs11192196

Guo, Y., Liu, R. W., Qu, J., Lu, Y., Zhu, F., & Lv, Y. (2023). Asynchronous Trajectory Matching-Based Multimodal Maritime Data Fusion for Vessel Traffic Surveillance in Inland Waterways. *IEEE Transactions on Intelligent Transportation Systems, 24*, 12779-12792. https://doi.org/10.1109/tits.2023.3285415

Han, X., Armenakis, C., & Jadidi, M. (2021). Modeling Vessel Behaviours by Clustering AIS Data Using Optimized DBSCAN. *Sustainability*. https://doi.org/10.3390/su13158162

Hasimoto-Beltrán, R., Canul-Ku, M., Díaz Méndez, G. M., Ocampo-Torres, F., & Esquivel-Trava, B. (2023). Ocean oil spill detection from SAR images based on multi-channel deep learning semantic segmentation.. *Marine pollution bulletin, 188*, 114651. https://doi.org/10.1016/j.marpolbul.2023.114651

Ioannou, G., Gaglione, D., Millefiori, L. M., Renga, A., Braca, P., & Willett, P. (2024). Dark-VADER: Detection of Anomalous AIS Message Delays for Maritime Situational Awareness. *2024 27th International Conference on Information Fusion (FUSION)*, 1-7. https://doi.org/10.23919/fusion59988.2024.10706391

Jafarzadeh, H., Mahdianpari, M., Homayouni, S., Mohammadimanesh, F., & Dabboor, M. (2021). Oil spill detection from Synthetic Aperture Radar Earth observations: a meta-analysis and comprehensive review. *GIScience & Remote Sensing, 58*, 1022 - 1051. https://doi.org/10.1080/15481603.2021.1952542

Jamal, S., Sun, G., Li, Y., Khan, A. Q., Zhang, Y., Verlanti, A., & Nunziata, F. (2025). CWCM-Net: A Novel Feature Fusion Method for Oil Spill Detection Using Single- and Quad-Polarization SAR Data. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, 18*, 19115-19128. https://doi.org/10.1109/jstars.2025.3591692

Jiang, Z., Zhang, J., Y., & Mao, X. (2025). Research on Remote Sensing Quantitative Inversion of Oil Spills and Emulsions Using Fusion of Optical and Thermal Characteristics. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, 18*, 8472-8489. https://doi.org/10.1109/jstars.2025.3547719

Jiao, J., Lu, Y., Hu, C., Shi, J., Sun, S., & Liu, Y. (2021). Quantifying ocean surface oil thickness using thermal remote sensing. *Remote Sensing of Environment, 261*, 112513. https://doi.org/10.1016/j.rse.2021.112513

Karunathilake, A., Ohashi, M., Kaneta, S., & Chiba, T. (2022). Monitoring the spatial dispersion of an oil slick by enhancing and noise-filtering SAR images using SENTINEL-1 satellite repeat-pass observations. *International Journal of Remote Sensing, 43*, 4187 - 4207. https://doi.org/10.1080/01431161.2022.2102949

Krestenitis, M., Orfanidis, G., Ioannidis, K., Avgerinakis, K., Vrochidis, S., & Kompatsiaris, Y. (2019). Oil Spill Identification from Satellite Images Using Deep Neural Networks. *Remote. Sens., 11*, 1762. https://doi.org/10.3390/rs11151762

Kulawiak, M., Prospathopoulos, A., Perivoliotis, L., Luba, M., Kioroglou, S., & Stepnowski, A. (2010). Interactive visualization of marine pollution monitoring and forecasting data via a Web-based GIS. *Comput. Geosci., 36*, 1069-1080. https://doi.org/10.1016/j.cageo.2010.02.008

La, T., Pelich, R., Li, Y., Matgen, P., & Chini, M. (2024). Monitoring of Spatio-Temporal Variations of Oil Slicks via the Collocation of Multi-Source Satellite Images. *Remote. Sens., 16*, 3110. https://doi.org/10.3390/rs16163110

Lee, C., & Youn, C.-H. (2025). In-Context Vision-Pattern-Language Model for Enhancing Vessel Activity Explanation. *IEEE Access, 13*, 94464-94477. https://doi.org/10.1109/access.2025.3572714

Leifer, I., Lehr, W., Simecek-Beatty, D., Bradley, E., Clark, R., Dennison, P., Hu, Y., Matheson, S., Jones, C. E., Holt, B., Reif, M., Roberts, D., Svejkovsky, J., Swayze, G., & Wozencraft, J. (2012). State of the art satellite and airborne marine oil spill remote sensing: Application to the BP Deepwater Horizon oil spill. *Remote Sensing of Environment, 124*, 185-209. https://doi.org/10.1016/j.rse.2012.03.024

Li, C., Yang, Y., Yang, X., Chu, D., & Cao, W. (2024). A Novel Multi-Scale Feature Map Fusion for Oil Spill Detection of SAR Remote Sensing. *Remote. Sens., 16*, 1684. https://doi.org/10.3390/rs16101684

Li, C., Kim, D.-J., Park, S., Kim, J., & Song, J. (2023). A self-evolving deep learning algorithm for automatic oil spill detection in Sentinel-1 SAR images. *Remote Sensing of Environment*. https://doi.org/10.1016/j.rse.2023.113872

Li, Y., Lyu, X., & Ren, P. (2024). Oil Spill Timely Backtracking Oriented by Wind Field Correction With Self-Attention Temporal Convolutional Networks. *IEEE Journal of Oceanic Engineering, 49*, 114-132. https://doi.org/10.1109/joe.2023.3314101

Liang, M., Su, J., Liu, R. W., & Lam, J. S. L. (2024). AISClean: AIS data-driven vessel trajectory reconstruction under uncertain conditions. *Ocean Engineering*. https://doi.org/10.1016/j.oceaneng.2024.117987

Liu, X., Zhang, Y., Zou, H., Wang, F., Cheng, X., Wu, W., Liu, X., & Li, Y. (2023). Multi-source knowledge graph reasoning for ocean oil spill detection from satellite SAR images. *Int. J. Appl. Earth Obs. Geoinformation, 116*, 103153. https://doi.org/10.1016/j.jag.2022.103153

Liubartseva, S., Coppini, G., Pinardi, N., Dominicis, M., Lecci, R., Turrisi, G., Creti', S., Martinelli, S., Agostini, P., Marra, P., & Palermo, F. (2016). Decision support system for emergency management of oil spill accidents in the Mediterranean Sea. *Natural Hazards and Earth System Sciences, 16*, 2009-2020. https://doi.org/10.5194/nhess-16-2009-2016

Lu, Y., Yang, K., Yang, D., Ding, H., Weng, J., & Liu, R. W. (2025). Graph Learning-Driven Multi-Vessel Association: Fusing Multimodal Data for Maritime Intelligence. *IEEE Transactions on Intelligent Transportation Systems, 27*, 5739-5754. https://doi.org/10.1109/tits.2026.3655007

Luo, D., Chen, P., Yang, J., Chen, X., Li, X., Zheng, G., Ren, L., & Zhao, Y. (2024). A new ship tracing technology from oil spills based on multi-source data.. *Marine pollution bulletin, 207*, 116808. https://doi.org/10.1016/j.marpolbul.2024.116808

Mazzarella, F., Vespe, M., & Santamaria, C. (2015). SAR Ship Detection and Self-Reporting Data Fusion Based on Traffic Knowledge. *IEEE Geoscience and Remote Sensing Letters, 12*, 1685-1689. https://doi.org/10.1109/lgrs.2015.2419371

Meng, T., Nunziata, F., Yang, X., Buono, A., Chen, K.-S., & Migliaccio, M. (2024). Scattering Model-Based Oil-Slick-Related Parameters Estimation From Radar Remote Sensing: Feasibility and Simulation Results. *IEEE Transactions on Geoscience and Remote Sensing, 62*, 1-12. https://doi.org/10.1109/tgrs.2024.3369023

Millati, P., & Choi, Y.-H. (2026). Abnormal Vessel Activity Detection Using a Two-Level Grid Representation of AIS Data. *IEEE Access, 14*, 22289-22303. https://doi.org/10.1109/access.2026.3662397

Murugan, J., Ramkumar, K., Kshirsagar, P., & Tak, T. K. (2026). Satellite-based oil spill detection using an explainable ViR-SC hybrid deep learning ensemble for improved accuracy and transparency. *Scientific Reports, 16*. https://doi.org/10.1038/s41598-026-37081-1

Naz, S., Iqbal, M., Mahmood, I., & Allam, M. (2020). Marine oil spill detection using Synthetic Aperture Radar over Indian Ocean.. *Marine pollution bulletin, 162*, 111921. https://doi.org/10.1016/j.marpolbul.2020.111921

Nguyen, D., & Fablet, R. (2021). A Transformer Network With Sparse Augmented Data Representation and Cross Entropy Loss for AIS-Based Vessel Trajectory Prediction. *IEEE Access, 12*, 21596-21609. https://doi.org/10.1109/access.2024.3349957

Oliveira, A., Jesus, G., Gomes, J., Rogeiro, J., Azevedo, A., Rodrigues, M., Fortunato, A., Dias, J., Tomas, L. M., Vaz, L., Oliveira, E. R., Alves, F., & Boer, S. (2014). An interactive WebGIS observatory platform for enhanced support of integrated coastal management. 507 - 512. https://doi.org/10.2112/si70-086.1

Park, H., Jung, J., Seo, M.-S., Choi, H., Cho, D., Park, S., & Choi, D.-G. (2025). AIS-LLM: A Unified Framework for Maritime Trajectory Prediction, Anomaly Detection, and Collision Risk Assessment With Explainable Forecasting. *IEEE Access, 14*, 106874-106891. https://doi.org/10.1109/access.2026.3705215

Pärt, S., Björkqvist, J., Alari, V., Maljutenko, I., & Uiboupin, R. (2023). An ocean-wave-trajectory forecasting system for the eastern Baltic Sea: Validation against drifting buoys and implementation for oil spill modeling.. *Marine pollution bulletin, 195*, 115497. https://doi.org/10.1016/j.marpolbul.2023.115497

Prasad, S. J., Francis, P. A., Nair, T. B. B., Shenoi, S., & Vijayalakshmi, T. (2020). Oil spill trajectory prediction with high-resolution ocean currents. *Journal of Operational Oceanography, 13*, 84 - 99. https://doi.org/10.1080/1755876x.2019.1606691

Prasad, S., Nair, T. B. B., & Vijayalakshmi, T. (2021). Oil spill trajectory prediction assessment using multi model ocean currents - case study of MT Dawn Oil spill 2017. *International Oil Spill Conference Proceedings*. https://doi.org/10.7901/2169-3358-2021.1.684276

Ren, P., Jia, Q., Xu, Q., Li, Y., Bi, F., Xu, J., & Gao, S. (2025). Oil Spill Drift Prediction Enhanced by Correcting Numerically Forecasted Sea Surface Dynamic Fields With Adversarial Temporal Convolutional Networks. *IEEE Transactions on Geoscience and Remote Sensing, 63*, 1-18. https://doi.org/10.1109/tgrs.2025.3528631

Ribeiro, C. V., Bedo, M. V. N., Mello, R., Paes, A., & De Oliveira, D. (2025). Detecting Anomalous Vessel Trajectories: A Collaborative Clustering-Based Approach. 84-97. https://doi.org/10.5753/sbbd.2025.247003

Rodger, M., & Guida, R. (2020). Classification-Aided SAR and AIS Data Fusion for Space-Based Maritime Surveillance. *Remote. Sens., 13*, 104. https://doi.org/10.3390/rs13010104

Şahin, B. B., Erdaş, Ç., & Sümer, E. (2026). Multi-sensor data fusion in maritime surveillance: a review on methods and applications. *PeerJ Computer Science*. https://doi.org/10.7717/peerj-cs.3765

Scafutto, R. D. M., Lassalle, G., Alves, M., De Paula Miranda, L., Costa, P. M., & De Souza Filho, C. R. (2025). Monitoring oil spill thickness and weathering using UAV-borne hyperspectral sensing.. *Marine pollution bulletin, 218*, 118134. https://doi.org/10.1016/j.marpolbul.2025.118134

Shaban, M., Salim, R., Khalifeh, H. A., Khelifi, A., Shalaby, A., El-Mashad, S. Y., Mahmoud, A. M., Ghazal, M., & El-Baz, A. (2021). A Deep-Learning Framework for the Detection of Oil Spills from SAR Data. *Sensors (Basel, Switzerland), 21*. https://doi.org/10.3390/s21072351

Sharma, A. (2026). DarkVesselNet: Multi-Modal Remote Sensing and Trajectory Reasoning for Dark Vessel Detection. *ArXiv, abs/2606.00445*. https://doi.org/10.48550/arxiv.2606.00445

Singh, S., Fowdur, J. S., Gawlikowski, J., & Medina, D. (2022). Leveraging Graph and Deep Learning Uncertainties to Detect Anomalous Maritime Trajectories. *IEEE Transactions on Intelligent Transportation Systems, 23*, 23488-23502. https://doi.org/10.1109/tits.2022.3190834

Sorgente, R., La Guardia, D., Ribotti, A., Arrigo, M., Signa, A., Pessini, F., Oliva, G., Pes, A., Perilli, A., & Di Maio, A. (2020). An Operational Supporting System for Oil Spill Emergencies Addressed to the the Italian Coast Guard. *Journal of Marine Science and Engineering*. https://doi.org/10.3390/jmse8121035

Wan, Y., Peng, L., Zhang, R., Zhang, R., & Wang, H. (2026). A Dual-Branch EfficientNetV2-S-Based Method for Marine Oil Spill Detection Using Multisource Satellite Data Fusion. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, 19*, 1549-1566. https://doi.org/10.1109/jstars.2025.3639503

Wolsing, K., Roepert, L., Bauer, J., & Wehrle, K. (2022). Anomaly Detection in Maritime AIS Tracks: A Review of Recent Approaches. *Journal of Marine Science and Engineering*. https://doi.org/10.3390/jmse10010112

Yang, D., Wu, L., Wang, S., Jia, H., & Li, K. X. (2019). How big data enriches maritime research – a critical review of Automatic Identification System (AIS) data applications. *Transport Reviews, 39*, 755 - 773. https://doi.org/10.1080/01441647.2019.1649315

Yang, Z., Chen, Z., & Lee, K. (2025). GOOSM: A GIS-based offshore oil spill management tool for enhanced response and preparedness.. *Marine pollution bulletin, 216*, 118009. https://doi.org/10.1016/j.marpolbul.2025.118009

Yang, Y., Liu, Y., Li, G., Zhang, Z., & Liu, Y. (2024). Harnessing the power of Machine learning for AIS Data-Driven maritime Research: A comprehensive review. *Transportation Research Part E: Logistics and Transportation Review*. https://doi.org/10.1016/j.tre.2024.103426

Yekeen, S., Balogun, A., & Yusof, K. (2020). A novel deep learning instance segmentation model for automated marine oil spill detection. *Isprs Journal of Photogrammetry and Remote Sensing, 167*, 190-200. https://doi.org/10.1016/j.isprsjprs.2020.07.011

Yekeen, S., & Balogun, A. (2020). Advances in Remote Sensing Technology, Machine Learning and Deep Learning for Marine Oil Spill Detection, Prediction and Vulnerability Assessment. *Remote. Sens., 12*, 3416. https://doi.org/10.3390/rs12203416

Yu, F., Li, J., Cui, S., Zhao, Y., Feng, Q.-J., & Chen, G. (2016). A hindcast method to simulate oil spill trajectories for the Bohai Sea, Northeast China. *Ocean Engineering, 124*, 363-370. https://doi.org/10.1016/j.oceaneng.2016.08.004

Yu, F., Li, J., Zhao, Y., Li, Q., & Chen, G. (2017). Calibration of backward-in-time model using drifting buoys in the East China Sea. *Oceanologia, 59*, 238-247. https://doi.org/10.1016/j.oceano.2017.01.003

Zhang, Z., & Xu, H. (2025). Explainable AI for Maritime Autonomous Surface Ships (MASS): Adaptive Interfaces and Trustworthy Human-AI Collaboration. *ArXiv, abs/2509.15959*. https://doi.org/10.48550/arxiv.2509.15959