

12.9.4.15

EE24BTECH11041 - Mohit

- 1) Find the equation of a curve passing through the point (0,0) and whose differential equation is $y' = e^x \sin x$

Solution:-

The given differential equation is:

$$y' = e^x \sin x \quad (1.1)$$

Integrating both sides:

$$y = \int e^x \sin x dx \quad (1.2)$$

Using integration by parts:

$$\int e^x \sin x dx = e^x \sin x - \int e^x \cos x dx \quad (1.3)$$

Let $I = \int e^x \sin x dx$. Then:

$$I = e^x \sin x - \int e^x \cos x dx \quad (1.4)$$

Now, solve $\int e^x \cos x dx$ using integration by parts again:

$$\int e^x \cos x dx = e^x \cos x - \int e^x (-\sin x) dx = e^x \cos x + \int e^x \sin x dx \quad (1.5)$$

Substituting back, we get:

$$I = e^x \sin x - (e^x \cos x + I) \quad (1.6)$$

$$I + I = e^x (\sin x - \cos x) \quad (1.7)$$

$$2I = e^x (\sin x - \cos x) \quad (1.8)$$

$$I = \frac{e^x (\sin x - \cos x)}{2} \quad (1.9)$$

Thus, the solution to the differential equation is:

$$y = \frac{e^x (\sin x - \cos x)}{2} + C \quad (1.10)$$

Using the initial condition $y(0) = 0$:

$$y(0) = \frac{e^0(\sin 0 - \cos 0)}{2} + C \quad (1.11)$$

$$0 = \frac{(0 - 1)}{2} + C \quad (1.12)$$

$$C = \frac{1}{2} \quad (1.13)$$

The final solution is:

$$y = \frac{e^x(\sin x - \cos x)}{2} + \frac{1}{2} \quad (1.14)$$

CODING LOGIC:-

The trapezoidal rule is as follows.

$$A = \int_a^b f(x) dx \approx h \left(\frac{1}{2}f(a) + f(x_1) + f(x_2) \cdots + f(x_{n-1}) + \frac{1}{2}f(b) \right) \quad (1.15)$$

$$h = \frac{b-a}{n} \quad (1.16)$$

$$A = j_n, \text{ where, } j_{i+1} = j_i + h \frac{f(x_{i+1}) + f(x_i)}{2} \quad (1.17)$$

$$\rightarrow j_{i+1} = j_i + h \left(\sqrt{x_{i+1}} + \sqrt{x_i} \right) \quad (1.18)$$

$$x_{i+1} = x_i + h \quad (1.19)$$

$$(1.20)$$

