

Integration: The process of finding an anti-derivative or integral of a function is called integration. It is the inverse process of differentiation. If $f(x)$ be a function of x related with another function $F(x)$ in such a way that

$$\frac{d}{dx}[F(x)] = f(x)$$

then

$$\int f(x)dx = F(x) + c$$

which is called an indefinite integral of $f(x)$ with respect to x where $f(x)$, $F(x)$ and c are called integrand, integral and constant of integration respectively.

And

$$\int_a^b f(x)dx = F(b) - F(a)$$

which is called the definite integral of $f(x)$ from a to b , and ' a ' is called the lower limit and ' b ' the upper limit of the definite integral.

Fundamental Properties:

1. $\int [f_1(x) \pm f_2(x) \pm \dots \dots \text{to } n \text{ terms}] dx = \int f_1(x)dx \pm \int f_2(x)dx \pm \dots \dots \text{to } n \text{ terms}.$
2. $\int cf(x)dx = c \int f(x)dx$

where c is a constant.

General rules of integration:

- | | |
|---|--|
| 1. $\int x^n dx = \frac{x^{n+1}}{n+1}$ where $(n \neq -1).$ | 2. $\int \frac{dx}{x^n} = -\frac{1}{(n-1)x^{n-1}}$ where $(n \neq 1).$ |
| 3. $\int \frac{dx}{\sqrt{x}} = 2\sqrt{x} + c.$ | 4. $\int dx = x + c.$ |
| 5. $\int \frac{dx}{x} = \ln x.$ | 6. $\int e^x dx = e^x.$ |
| 7. $\int e^{mx} dx = \frac{e^{mx}}{m}.$ | 8. $\int a^x dx = \frac{a^x}{\ln a}$ where $a > 0.$ |
| 9. $\int \sin mx dx = -\frac{\cos mx}{m}.$ | 10. $\int \sin x dx = -\cos x.$ |

$$11. \int \cos mx dx = \frac{\sin mx}{m}.$$

$$12. \int \cos x dx = \sin x.$$

$$13. \int \sec^2 x dx = \tan x.$$

$$14. \int \sec x \tan x dx = \sec x.$$

$$15. \int \cosec^2 x dx = -\cot x.$$

$$16. \int \cosec x \cot x dx = -\cosec x.$$

$$17. \int \tan x dx = \ln|\sec x|.$$

$$18. \int \cot x dx = \ln|\sin x|.$$

$$19. \int \frac{dx}{a^2 + x^2} = \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) \quad \text{where } a \neq 0.$$

$$20. \int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \ln \left| \frac{x-a}{x+a} \right| + c.$$

$$21. \int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \ln \left| \frac{a+x}{a-x} \right|.$$

$$22. \int \frac{dx}{\sqrt{x^2 + a^2}} = \ln \left(x + \sqrt{x^2 + a^2} \right).$$

$$23. \int \frac{dx}{\sqrt{x^2 - a^2}} = \ln \left(x + \sqrt{x^2 - a^2} \right).$$

$$24. \int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \left(\frac{x}{a} \right).$$

$$25. \int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \sec^{-1} \left(\frac{x}{a} \right)$$

$$26. \int \frac{dx}{x\sqrt{x^2 - 1}} = \sec^{-1} x$$

$$27. \int \cosec x dx = \ln|\cosec x - \cot x|.$$

$$28. \int \sec x dx = \ln|\sec x + \tan x|.$$

$$29. \int uv dx = u \int v dx - \int \left(\frac{du}{dx} \cdot \int v dx \right) dx.$$

$$30. \int \sqrt{x^2 + a^2} dx = \frac{x\sqrt{x^2 + a^2}}{2} + \frac{a^2}{2} \ln|x + \sqrt{x^2 + a^2}|.$$

$$31. \int \sqrt{x^2 - a^2} dx = \frac{x\sqrt{x^2 - a^2}}{2} - \frac{a^2}{2} \ln|x + \sqrt{x^2 - a^2}|.$$

$$32. \int \sqrt{a^2 - x^2} dx = \frac{x\sqrt{a^2 - x^2}}{2} + \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right).$$

$$33. \int \frac{f'(x)}{f(x)} dx = \ln f(x).$$

$$34. \int e^x [f(x) + f'(x)] dx = e^x f(x).$$

$$35. \int e^{ax} \sin bx dx = \frac{e^{ax} (a \sin bx - b \cos bx)}{a^2 + b^2}.$$

$$36. \int e^{ax} \cos bx dx = \frac{e^{ax} (a \cos bx + b \sin bx)}{a^2 + b^2}.$$

Indefinite Integration

Illustrative Examples:

Problem-01: $\int \sin^2 x dx$

$$\begin{aligned} \text{Sol}^n : \text{Let } I &= \int \sin^2 x dx \\ &= \frac{1}{2} \int 2 \sin^2 x dx \\ &= \frac{1}{2} \int (1 - \cos 2x) dx \\ &= \frac{1}{2} \left(x - \frac{\sin 2x}{2} \right) + c \end{aligned}$$

where c is an integrating constant.

Problem-02: $\int \tan^2 x dx$

$$\begin{aligned} \text{Sol}^n : \text{Let } I &= \int \tan^2 x dx \\ &= \int (\sec^2 x - 1) dx \\ &= (\tan x - x) + c. \end{aligned}$$

where c is an integrating constant.

Problem-03: $\int \frac{a \sin^2 x + b \cos^2 x}{\sin^2 x \cos^2 x} dx$

$$\begin{aligned} \text{Sol}^n : \text{Let } I &= \int \frac{a \sin^2 x + b \cos^2 x}{\sin^2 x \cos^2 x} dx \\ &= \int \left(\frac{a}{\cos^2 x} + \frac{b}{\sin^2 x} \right) dx \\ &= \int (a \sec^2 x + b \csc^2 x) dx \\ &= a \tan x - b \cot x + c \end{aligned}$$

where c is an integrating constant.

Problem-04: $\int \frac{\cos x - \cos 2x}{1 - \cos x} dx$

$$\begin{aligned} \text{Sol}^n : \text{Let } I &= \int \frac{\cos x - \cos 2x}{1 - \cos x} dx \\ &= \int \frac{\cos x - (\cos^2 x - \sin^2 x)}{1 - \cos x} dx \end{aligned}$$

Exercise-01: $\int \cos^2 x dx$.

$$\text{Ans: } \frac{1}{2} \left(x + \frac{\sin 2x}{2} \right) + c.$$

Exercise-02: $\int \cot^2 x dx$

$$\text{Ans: } -\cot x - x + c.$$

Exercise-03: $\int \frac{\sin^3 x + \cos^3 x}{\cos x + \sin x} dx$

$$\text{Ans: } x + \frac{1}{4} \cos 2x + c.$$

Exercise-04: $\int \frac{\sin x + \cos x}{\sqrt{1 + \sin 2x}} dx$

$$\text{Ans: } x + c.$$

$$\begin{aligned}
&= \int \frac{\cos x - \cos^2 x + \sin^2 x}{1 - \cos x} dx \\
&= \int \frac{\cos x(1 - \cos x) + (1 - \cos^2 x)}{1 - \cos x} dx \\
&= \int \left\{ \frac{\cos x(1 - \cos x)}{1 - \cos x} + \frac{(1 - \cos x)(1 + \cos x)}{1 - \cos x} \right\} dx \\
&= \int (\cos x + 1 + \cos x) dx \\
&= \int (1 + 2\cos x) dx \\
&= x + 2\sin x + c
\end{aligned}$$

where c is an integrating constant.

Problem-05: $\int \sqrt{1 - \sin 2x} dx$

$$\begin{aligned}
\text{Sol}^n : \text{Let } I &= \int \sqrt{1 - \sin 2x} dx \\
&= \int \sqrt{\cos^2 x + \sin^2 x - 2\sin x \cos x} dx \\
&= \int \sqrt{(\cos x - \sin x)^2} dx \\
&= \int (\cos x - \sin x) dx \\
&= \sin x + \cos x + c
\end{aligned}$$

where c is an integrating constant.

Problem-06: $\int \sqrt{1 + \cos x} dx$

$$\begin{aligned}
\text{Sol}^n : \text{Let } I &= \int \sqrt{1 + \cos x} dx \\
&= \int \sqrt{2\cos^2 \frac{x}{2}} dx \\
&= \sqrt{2} \int \cos \frac{x}{2} dx \\
&= \sqrt{2} \frac{\sin \frac{x}{2}}{\frac{1}{2}} + c \\
&= 2\sqrt{2} \sin \frac{x}{2} + c
\end{aligned}$$

where c is an integrating constant.

Exercise-05: $\int \sqrt{1 + \sin x} dx$.

$$\text{Ans: } 2 \left(\sin \frac{x}{2} - \cos \frac{x}{2} \right) + c .$$

Exercise-06: $\int \sqrt{1 - \cos 2x} dx$.

$$\text{Ans: } -\sqrt{2} \cos x + c .$$

Indefinite Integration

Problem-07: $\int \frac{dx}{1 + \sin x}$

Solⁿ : Let $I = \int \frac{dx}{1 + \sin x}$

$$\begin{aligned} &= \int \frac{(1 - \sin x)}{(1 + \sin x)(1 - \sin x)} dx \\ &= \int \frac{1 - \sin x}{1 - \sin^2 x} dx \\ &= \int \frac{1 - \sin x}{\cos^2 x} dx \\ &= \int \frac{1}{\cos^2 x} dx - \int \frac{\sin x}{\cos^2 x} dx \\ &= \int \sec^2 x dx - \int \sec x \tan x dx \\ &= \tan x - \sec x + c \end{aligned}$$

where c is an integrating constant.

Problem-08: $\int \cos^4 x dx$

Solⁿ : Let $I = \int \cos^4 x dx$

$$\begin{aligned} &= \frac{1}{4} \int (2 \cos^2 x)^2 dx \\ &= \frac{1}{4} \int (1 + \cos 2x)^2 dx \\ &= \frac{1}{4} \int (1 + 2 \cos 2x + \cos^2 2x) dx \\ &= \frac{1}{4} \int \left\{ 1 + 2 \cos 2x + \frac{1}{2} (2 \cos^2 2x) \right\} dx \\ &= \frac{1}{4} \int \left\{ 1 + 2 \cos 2x + \frac{1}{2} (1 + \cos 4x) \right\} dx \\ &= \frac{1}{4} \left[x + \sin 2x + \frac{x}{2} + \frac{\sin 4x}{8} \right] + c \\ &= \frac{1}{4} \left[\frac{3x}{2} + \sin 2x + \frac{\sin 4x}{8} \right] + c \\ &= \frac{3x}{8} + \frac{\sin 2x}{4} + \frac{\sin 4x}{32} + c \end{aligned}$$

where c is an integrating constant.

Exercise-07: $\int \frac{dx}{1 + \cos x}$.

Ans: $-\cot x + \cos ec x + c$.

Exercise-08: $1. \int \sin^4 x dx$.

Ans: $\frac{1}{32} \sin 4x - \frac{1}{4} \sin 2x + \frac{3}{8} x + c$.

Problem-09: $\int \frac{\sin x}{\sqrt{1+\cos x}} dx$

Solⁿ : Let $I = \int \frac{\sin x}{\sqrt{1+\cos x}} dx$

$$= \int \frac{2 \sin \frac{x}{2} \cos \frac{x}{2}}{\sqrt{2 \cos^2 \frac{x}{2}}} dx$$

$$= \int \frac{2 \sin \frac{x}{2} \cos \frac{x}{2}}{\sqrt{2} \cos \frac{x}{2}} dx$$

$$= \sqrt{2} \int \sin \frac{x}{2} dx$$

$$= -\sqrt{2} \frac{\cos \frac{x}{2}}{\frac{1}{2}} + c$$

$$= -2\sqrt{2} \cos \frac{x}{2} + c$$

where c is an integrating constant.

Problem-10: $\int \frac{dx}{\sin^2 x \cos^2 x}$

solⁿ : Let $I = \int \frac{dx}{\sin^2 x \cos^2 x}$

$$= \int \frac{\sin^2 x + \cos^2 x}{\sin^2 x \cos^2 x} dx$$

$$= \int \left(\frac{1}{\cos^2 x} + \frac{1}{\sin^2 x} \right) dx$$

$$= \int (\sec^2 x + \csc^2 x) dx$$

$$= \tan x - \cot x + c.$$

where c is an integrating constant.

Problem-11: $\int \frac{\sin^6 x + \cos^6 x}{\sin^2 x \cos^2 x} dx$

Solⁿ : Let $I = \int \frac{\sin^6 x + \cos^6 x}{\sin^2 x \cos^2 x} dx$

Exercise-09: $\int \frac{\sin 2x}{\sqrt{1-\cos 2x}} dx$

Ans: $\sqrt{2} \sin x + c.$

Exercise-09: $\int \frac{\sin^8 x - \cos^8 x}{1 - 2 \sin^2 x \cos^2 x} dx$

Ans: $-\frac{1}{2} \sin 2x + c.$

Indefinite Integration

$$\begin{aligned}&= \int \frac{(\sin^2 x)^3 + (\cos^2 x)^3}{\sin^2 x \cos^2 x} dx \\&= \int \frac{(\sin^2 x + \cos^2 x)^3 - 3 \sin^2 x \cos^2 x (\sin^2 x + \cos^2 x)}{\sin^2 x \cos^2 x} dx \\&= \int \frac{1 - 3 \sin^2 x \cos^2 x}{\sin^2 x \cos^2 x} dx \\&= \int \left(\frac{1}{\sin^2 x \cos^2 x} - 3 \right) dx \\&= \int \left(\frac{\sin^2 x + \cos^2 x}{\sin^2 x \cos^2 x} - 3 \right) dx \\&= \int \left(\frac{1}{\cos^2 x} + \frac{1}{\sin^2 x} - 3 \right) dx \\&= \int (\sec^2 x + \csc^2 x - 3) dx \\&= \tan x - \cot x - 3x + c.\end{aligned}$$

where c is an integrating constant.

Problem-12: $\int \frac{\cos^4 x - \sin^4 x}{\sqrt{1 + \cos 4x}} dx$

Solⁿ : Let $I = \int \frac{\cos^4 x - \sin^4 x}{\sqrt{1 + \cos 4x}} dx$

$$\begin{aligned}&= \int \frac{(\cos^2 x)^2 - (\sin^2 x)^2}{\sqrt{2 \cos^2 2x}} dx \\&= \frac{1}{\sqrt{2}} \int \frac{(\cos^2 x + \sin^2 x)(\cos^2 x - \sin^2 x)}{\cos 2x} dx \\&= \frac{1}{\sqrt{2}} \int \frac{\cos 2x}{\cos 2x} dx \\&= \frac{1}{\sqrt{2}} \int dx \\&= \frac{x}{\sqrt{2}} + c\end{aligned}$$

where c is an integrating constant.

Problem-13: $\int \frac{\sin 2x}{\sin 5x \sin 3x} dx$

solⁿ : Let $I = \int \frac{\sin 2x}{\sin 5x \sin 3x} dx$

$$= \int \frac{\sin(5x-3x)}{\sin 5x \sin 3x} dx$$

$$= \int \frac{\sin 5x \cos 3x - \cos 5x \sin 3x}{\sin 5x \sin 3x} dx$$

$$= \int \left(\frac{\cos 3x}{\sin 3x} - \frac{\cos 5x}{\sin 5x} \right) dx$$

$$= \int \frac{\cos 3x}{\sin 3x} dx - \int \frac{\cos 5x}{\sin 5x} dx$$

$$= \int (\cot 3x - \cot 5x) dx$$

$$= \frac{1}{3} \ln(\sin 3x) - \frac{1}{5} \ln(\sin 5x) + c.$$

where c is an integrating constant.

Problem-14: $\int \sin x \sin 2x \sin 3x dx$

solⁿ : Let $I = \int \sin x \sin 2x \sin 3x dx$

$$= \frac{1}{2} \int (2 \sin x \sin 2x) \sin 3x dx$$

$$= \frac{1}{2} \int \{ \cos(x-2x) - \cos(x+2x) \} \sin 3x dx$$

$$= \frac{1}{2} \int \{ \cos x - \cos 3x \} \sin 3x dx$$

$$= \frac{1}{4} \int \{ 2 \sin 3x \cos x - 2 \sin 3x \cos 3x \} dx$$

$$= \frac{1}{4} \int \{ 2 \sin 3x \cos x - \sin 6x \} dx$$

$$= \frac{1}{4} \int \{ \sin(3x+x) + \sin(3x-x) - \sin 6x \} dx$$

$$= \frac{1}{4} \int \{ \sin 4x + \sin 2x - \sin 6x \} dx$$

$$= \frac{1}{4} \left\{ -\frac{\cos 4x}{4} - \frac{\cos 2x}{2} + \frac{\cos 6x}{6} \right\} + c$$

$$= -\frac{\cos 4x}{16} - \frac{\cos 2x}{8} + \frac{\cos 6x}{24} + c.$$

Exercise-10: $\int \frac{\sin x}{\sin(x-a)} dx.$

Ans: $x \cos a + \sin a \ln [\sin(x-a)] + c.$

Exercise-11: $\int \sin 3x \sin 4x dx$

Ans: $\frac{1}{2} \sin x - \frac{1}{14} \sin 7x + c.$

Indefinite Integration

where c is an integrating constant.

Problem-15: $\int \frac{dx}{\sqrt{x} + \sqrt{x+1}}$

Solⁿ : Let $I = \int \frac{dx}{\sqrt{x} + \sqrt{x+1}}$

$$\begin{aligned} &= \int \frac{(\sqrt{x} - \sqrt{x+1})}{(\sqrt{x} + \sqrt{x+1})(\sqrt{x} - \sqrt{x+1})} dx \\ &= \int \frac{(\sqrt{x} - \sqrt{x+1})}{(\sqrt{x})^2 - (\sqrt{x+1})^2} dx \\ &= \int \frac{(\sqrt{x} - \sqrt{x+1})}{x - (x+1)} dx \\ &= \int \frac{(\sqrt{x} - \sqrt{x+1})}{x - x - 1} dx \\ &= - \int (\sqrt{x} - \sqrt{x+1}) dx \\ &= - \frac{x^{\frac{1}{2}+1}}{\frac{1}{2}+1} + \frac{(x+1)^{\frac{1}{2}+1}}{\frac{1}{2}+1} + c \\ &= \frac{2}{3}(x+1)^{\frac{3}{2}} - \frac{2}{3}x^{\frac{3}{2}} + c \end{aligned}$$

where c is an integrating constant.

Problem-15: $\int \frac{2 - \sin 2x}{1 - \cos 2x} dx$ **Exercise-13:** $\int \frac{dx}{\sqrt{x} - \sqrt{x-1}}$

Solⁿ : Let $I = \int \frac{2 - \sin 2x}{1 - \cos 2x} dx$ **Ans:** $\frac{2}{3}x^{\frac{3}{2}} + \frac{2}{3}(x-1)^{\frac{3}{2}} + c$

$$\begin{aligned} &= \int \frac{2 - 2 \sin x \cos x}{2 \sin^2 x} dx \\ &= \int \left(\frac{2}{2 \sin^2 x} - \frac{2 \sin x \cos x}{2 \sin^2 x} \right) dx \\ &= \int (\operatorname{cosec}^2 x - \cot x) dx \\ &= \int \operatorname{cosec}^2 x dx - \int \cot x dx \end{aligned}$$

$$= -\cot x - \ln(\sin x) + c$$

where c is an integrating constant.

Problem-17: $\int \frac{dx}{a \cos x + b \sin x}$

Solⁿ : Let $I = \int \frac{dx}{a \cos x + b \sin x}$

Let, $a = r \cos \theta$ and $b = r \sin \theta$

$$\therefore r = \sqrt{a^2 + b^2} \text{ and } \theta = \tan^{-1} \frac{b}{a}$$

$$= \frac{1}{r} \int \frac{dx}{\cos x \cos \theta + \sin x \sin \theta}$$

$$= \frac{1}{r} \int \frac{dx}{\cos(x - \theta)}$$

$$= \frac{1}{r} \int \sec(x - \theta) dx$$

$$= \frac{1}{r} \ln \{ \sec(x - \theta) + \tan(x - \theta) \} + c$$

$$= \frac{1}{\sqrt{a^2 + b^2}} \ln \left\{ \sec \left(x - \tan^{-1} \frac{b}{a} \right) + \tan \left(x - \tan^{-1} \frac{b}{a} \right) \right\} + c$$

where c is an integrating constant.

Method of substitution

Sometimes, the integration of given integral $\int f(x) dx$ is relatively difficult. In this case, we can replace x by $\phi(z)$ and dx by $\phi'(z) dz$ for integrating easily. This process is known as method of substitution.

Illustrative Examples:

Problem-01: $\int (a + bx)^n dx$

solⁿ : Let $I = \int (a + bx)^n dx$

Put $z = a + bx \therefore dz = b dx$

$$\Rightarrow \frac{1}{b} dz = dx$$

Now $I = \int z^n \frac{1}{b} dz$

$$= \frac{1}{b} \int z^n dz$$

Exercise-01: $\int \frac{2 \sin x}{5 + 3 \cos x} dx$

Ans: $-\frac{2}{3} \ln(5 + 3 \cos x) + c$

Indefinite Integration

$$\begin{aligned} &= \frac{1}{b} \frac{z^{n+1}}{n+1} + c \\ &= \frac{(a+bx)^{n+1}}{b(n+1)} + c \end{aligned}$$

where c is an integrating constant.

Problem-02: $\int \frac{dx}{x\sqrt{(x^2-a^2)}}$

solⁿ : Let $I = \int \frac{dx}{x\sqrt{(x^2-a^2)}}$

Put $x = a \sec \theta \therefore dx = a \sec \theta \tan \theta d\theta$

$$\begin{aligned} \text{Now } I &= \int \frac{a \sec \theta \tan \theta d\theta}{a \sec \theta \sqrt{(a^2 \sec^2 \theta - a^2)}} \\ &= \int \frac{a \sec \theta \tan \theta d\theta}{a \sec \theta a \sqrt{(\sec^2 \theta - 1)}} \\ &= \frac{1}{a} \int \frac{\sec \theta \tan \theta d\theta}{\sec \theta \sqrt{\tan^2 \theta}} \\ &= \frac{1}{a} \int \frac{\sec \theta \tan \theta d\theta}{\sec \theta \tan \theta} \\ &= \frac{1}{a} \int d\theta \\ &= \frac{1}{a} \theta + c \\ &= \frac{1}{a} \sec^{-1} \frac{x}{a} + c \end{aligned}$$

where c is an integrating constant.

Problem-03: $\int \frac{\sin^{-1} x}{\sqrt{1-x^2}} dx$

solⁿ : Let $I = \int \frac{\sin^{-1} x dx}{\sqrt{1-x^2}}$

Put $z = \sin^{-1} x \therefore dz = \frac{dx}{\sqrt{1-x^2}}$

Now $I = \int z dz$

Exercise-02: $\int \frac{dx}{x\sqrt{x^2-1}}$

Ans: $\sec^{-1} x + c$.

Exercise-03: $\int \frac{e^{m \tan^{-1} x}}{1+x^2} dx$

Ans: $\frac{e^{m \tan^{-1} x}}{m} + c$.

$$= \frac{z^2}{2} + c$$

$$= \frac{(\sin^{-1} x)^2}{2} + c$$

where c is an integrating constant.

Problem-04: $\int \frac{(1+x)e^x}{\cos^2(xe^x)} dx$

solⁿ : Let $I = \int \frac{(1+x)e^x}{\cos^2(xe^x)} dx$

Put $xe^x = z \therefore (1+x)e^x dx = dz$

Now $I = \int \frac{dz}{\cos^2 z}$

$$= \int \sec^2 z dz$$

$$= \tan z + c$$

$$= \tan(xe^x) + c$$

where c is an integrating constant.

Problem-05: $\int \frac{dx}{e^x + 1}$

solⁿ : Let $I = \int \frac{dx}{e^x + 1}$

$$= \int \frac{e^{-x}}{1 + e^{-x}} dx$$

Put $1 + e^{-x} = z \therefore -e^{-x} dx = dz$

Now $I = -\int \frac{dz}{z}$

$$= -\ln z + c$$

$$= -\ln(1 + e^{-x}) + c$$

where c is an integrating constant.

Problem-06: $\int \frac{\sqrt{x} + \ln x}{x} dx$

solⁿ : Let $I = \int \frac{\sqrt{x} + \ln x}{x} dx$

Exercise-04: $\int \frac{(x+1)(x+\ln x)^2}{x} dx$

Ans: $\frac{1}{3}(x+\ln x)^3 + c$.

Exercise-05: $\int \frac{\sin x}{(1 - \cos x)^2} dx$

Ans: $-\frac{1}{(1 - \cos x)} + c$.

Indefinite Integration

$$\begin{aligned} &= \int \left(\frac{1}{\sqrt{x}} + \frac{\ln x}{x} \right) dx \\ &= \int x^{-\frac{1}{2}} dx + \int \frac{\ln x}{x} dx \\ &= \frac{x^{\frac{-\frac{1}{2}+1}{- \frac{1}{2}+1}}}{-\frac{1}{2}+1} + \frac{1}{2} (\ln x)^2 + c \quad [\text{putting } \ln x = z \text{ in the 2nd part}] \\ &= 2\sqrt{x} + \frac{1}{2} (\ln x)^2 + c \end{aligned}$$

where c is an integrating constant.

Problem-07: $\int x\sqrt{1+x} dx$

solⁿ : Let $I = \int x\sqrt{1+x} dx$

Put $1+x = z \therefore dx = dz$

$$\begin{aligned} \text{Now } I &= \int (z-1) z^{\frac{1}{2}} dz \\ &= \int (z-1) z^{\frac{1}{2}} dz \\ &= \int \left(z^{\frac{3}{2}} - z^{\frac{1}{2}} \right) dz \\ &= \frac{2}{5} z^{\frac{5}{2}} - \frac{2}{3} z^{\frac{3}{2}} + c \\ &= \frac{2}{5} (1+x)^{\frac{5}{2}} - \frac{2}{3} (1+x)^{\frac{3}{2}} + c \end{aligned}$$

where c is an integrating constant.

Problem-08: $\int \frac{x}{\sqrt{x}+1} dx$

solⁿ : Let $I = \int \frac{x}{\sqrt{x}+1} dx$

Put $x = z^2 \therefore 2z dz = dx$

$$\begin{aligned} \text{Now } I &= 2 \int \frac{z^2 z}{z+1} dz \\ &= 2 \int \frac{z^3}{z+1} dz \end{aligned}$$

Exercise-06: $\int x \sqrt[3]{(1-x^2)^5} dx$

Ans: $-\frac{3}{16} (1-x^2)^{\frac{8}{3}} + c$.

Exercise-07: $\int \frac{dx}{\sqrt{x}-1}$

Ans: $2\sqrt{x} + 2\ln(\sqrt{x}-1) + c$.

$$\begin{aligned}
&= 2 \int \frac{z^3 + z^2 - z^2 - z + z + 1 - 1}{z + 1} dz \\
&= 2 \int \frac{z^2(z + 1) - z(z + 1) + (z + 1) - 1}{z + 1} dz \\
&= 2 \int \left[z^2 - z + 1 - \frac{1}{z + 1} \right] dz \\
&= 2 \left[\frac{z^3}{3} - \frac{z^2}{2} + z - \ln(z + 1) \right] + c \\
&= 2 \left[\frac{x^{3/2}}{3} - \frac{x}{2} + \sqrt{x} - \ln(\sqrt{x} + 1) \right] + c
\end{aligned}$$

where c is an integrating constant.

Problem-09: $\int \frac{x^3}{\sqrt{x-1}} dx$

solⁿ : Let $I = \int \frac{x^3}{\sqrt{x-1}} dx$

Put $x - 1 = z \therefore dx = dz$

Now $I = \int \frac{(z+1)^3}{\sqrt{z}} dz$

$$\begin{aligned}
&= \int \frac{(z^3 + 3z^2 + 3z + 1)}{\sqrt{z}} dz \\
&= \int \left(z^{\frac{5}{2}} + 3z^{\frac{3}{2}} + 3z^{\frac{1}{2}} + z^{-\frac{1}{2}} \right) dz \\
&= \frac{2}{7} z^{7/2} + \frac{6}{5} z^{5/2} + 2z^{3/2} + 2z^{1/2} + c \\
&= \frac{2}{7} (x-1)^{7/2} + \frac{6}{5} (x-1)^{5/2} + 2(x-1)^{3/2} + 2(x-1)^{1/2} + c.
\end{aligned}$$

where c is an integrating constant.

Problem-10: $\int \frac{\sqrt{\tan x}}{\sin x \cos x} dx$

solⁿ : Let $I = \int \frac{\sqrt{\tan x}}{\sin x \cos x} dx$

$$= \int \frac{\sqrt{\tan x}}{\tan x \cos^2 x} dx$$

Exercise-08: $\int \frac{x^2}{\sqrt{x+1}} dx$

Ans: $\frac{2}{5} (x+1)^{5/2} - \frac{4}{3} (x+1)^{3/2} + 2\sqrt{(x+1)} + c.$

Indefinite Integration

$$= \int \frac{\sec^2 x}{\sqrt{\tan x}} dx$$

Put $\tan x = z \quad \therefore \sec^2 x dx = dz$

Now $I = \int \frac{dz}{\sqrt{z}}$

$$= \frac{z^{-\frac{1}{2}+1}}{-\frac{1}{2}+1} + c$$

$$= \frac{z^{\frac{1}{2}}}{\frac{1}{2}} + c$$

$$= 2\sqrt{\tan x} + c$$

where c is an integrating constant.

Problem-11: $\int \frac{1}{\sqrt{x}} \cos \sqrt{x} dx$

solⁿ : Let $I = \int \frac{1}{\sqrt{x}} \cos \sqrt{x} dx$

Let, $\sqrt{x} = z$

$$\therefore \frac{1}{2\sqrt{x}} dx = dz$$

$$\Rightarrow \frac{1}{\sqrt{x}} dx = 2dz$$

Now $I = 2 \int \cos z dz$

$$= 2 \sin z + c$$

$$= 2 \sin \sqrt{x} + c$$

where c is an integrating constant.

Problem-12: $\int \frac{1 - \sin x}{x + \cos x} dx$

solⁿ : Let $I = \int \frac{1 - \sin x}{x + \cos x} dx$

Let, $x + \cos x = z$

$$\therefore (1 - \sin x) dx = dz$$

Exercise-09: $\int \frac{a \cos x - b \sin x}{a \sin x + b \cos x + d} dx$

Ans: $\ln(a \sin x + b \cos x + d) + c$

$$\begin{aligned}
\text{Now } I &= \int \frac{dz}{z} \\
&= \ln z + c \\
&= \ln(x + \cos x) + c
\end{aligned}$$

where c is an integrating constant.

Problem-13: $\int \frac{\sin 2x}{a \sin^2 x + b \cos^2 x} dx$

solⁿ : Let $I = \int \frac{\sin 2x}{a \sin^2 x + b \cos^2 x} dx$

Let, $a \sin^2 x + b \cos^2 x = z$

$$\therefore (2a \sin x \cos x - 2b \sin x \cos x) dx = dz$$

$$\Rightarrow (a - b) 2 \sin x \cos x dx = dz$$

$$\Rightarrow \sin 2x dx = \frac{1}{(a - b)} dz$$

$$\begin{aligned}
\text{Now } I &= \frac{1}{(a - b)} \int \frac{dz}{z} \\
&= \frac{1}{(a - b)} \ln z + c \\
&= \frac{1}{(a - b)} \ln(a \sin^2 x + b \cos^2 x) + c
\end{aligned}$$

where c is an integrating constant.

Problem-14: $\int \frac{\sin x \cos x}{\cos^4 x + \sin^4 x} dx$

solⁿ : Let $I = \int \frac{\sin x \cos x}{\cos^4 x + \sin^4 x} dx$

$$= \int \frac{\sin x \cos x}{\cos^4 x \left(1 + \frac{\sin^4 x}{\cos^4 x} \right)} dx$$

$$= \int \frac{\tan x \sec^2 x}{(1 + \tan^4 x)} dx$$

Put $\tan^2 x = z \therefore 2 \tan x \sec^2 x dx = dz$

Now $I = \frac{1}{2} \int \frac{dz}{1 + z^2}$

Exercise-10: $\int \frac{\sin 2x}{(a^2 \cos^2 x + b^2 \sin^2 x)^2} dx$

Ans: $\frac{1}{(a^2 - b^2)} \left\{ \frac{1}{a^2 \cos^2 x + b^2 \sin^2 x} \right\} + c$

Indefinite Integration

$$\begin{aligned} &= \frac{1}{2} \tan^{-1} z + c \\ &= \frac{1}{2} \tan^{-1} (\tan^2 x) + c \end{aligned}$$

where c is an integrating constant.

Problem-15: $\int \frac{x^2 + \sin^2 x}{1 + x^2} \sec^2 x dx$

solⁿ : Let $I = \int \frac{x^2 + \sin^2 x}{1 + x^2} \sec^2 x dx$

$$\begin{aligned} &= \int \frac{x^2 + 1 - \cos^2 x}{1 + x^2} \sec^2 x dx \\ &= \int \frac{(1 + x^2) \sec^2 x - 1}{1 + x^2} dx \\ &= \int \sec^2 x dx - \int \frac{dx}{1 + x^2} \\ &= \tan x - \tan^{-1} x + c \end{aligned}$$

where c is an integrating constant.

Problem-16: $\int \frac{dx}{(1 + x^2)^2}$

solⁿ : Let $I = \int \frac{dx}{(1 + x^2)^2}$

Put $x = \tan \theta \quad \therefore dx = \sec^2 \theta d\theta$

Now $I = \int \frac{\sec^2 \theta d\theta}{(1 + \tan^2 \theta)^2}$

$$\begin{aligned} &= \int \frac{\sec^2 \theta d\theta}{(\sec^2 \theta)^2} \\ &= \int \frac{\sec^2 \theta d\theta}{\sec^4 \theta} \\ &= \int \frac{d\theta}{\sec^2 \theta} \\ &= \int \cos^2 \theta d\theta \\ &= \frac{1}{2} \int 2 \cos^2 \theta d\theta \\ &= \frac{1}{2} \int (1 + \cos 2\theta) d\theta \end{aligned}$$

Exercise-11: $\int \frac{dx}{(1 + x^2)^{3/2}}$

Ans: $\frac{x}{\sqrt{1 + x^2}} + c$.

$$\begin{aligned}
&= \frac{1}{2} \left(\theta + \frac{\sin 2\theta}{2} \right) + c \\
&= \frac{1}{2} \left(\theta + \frac{1}{2} \frac{2 \tan \theta}{(1 + \tan^2 \theta)} \right) + c \\
&= \frac{1}{2} \left(\tan^{-1} x + \frac{x}{(1 + x^2)} \right) + c
\end{aligned}$$

where c is an integrating constant.

Problem-17: $\int \sqrt{\frac{x}{a-x}} dx$

solⁿ : Let $I = \int \sqrt{\frac{x}{a-x}} dx$

Put $x = a \sin^2 \theta \quad \therefore dx = 2a \sin \theta \cos \theta d\theta$

Now $I = \int \sqrt{\frac{a \sin^2 \theta}{a - a \sin^2 \theta}} 2a \sin \theta \cos \theta d\theta$

$$\begin{aligned}
&= \int \sqrt{\frac{\sin^2 \theta}{1 - \sin^2 \theta}} 2a \sin \theta \cos \theta d\theta \\
&= \int \sqrt{\frac{\sin^2 \theta}{\cos^2 \theta}} 2a \sin \theta \cos \theta d\theta \\
&= \int \frac{\sin \theta}{\cos \theta} 2a \sin \theta \cos \theta d\theta \\
&= a \int 2 \sin^2 \theta d\theta \\
&= a \int (1 - \cos 2\theta) d\theta \\
&= a \left(\theta - \frac{\sin 2\theta}{2} \right) + c \\
&= a(\theta - \sin \theta \cos \theta) + c \\
&= a \left(\theta - \sin \theta \sqrt{1 - \sin^2 \theta} \right) + c \\
&= a \left(\sin^{-1} \sqrt{\frac{x}{a}} - \sqrt{\frac{x}{a}} \sqrt{1 - \frac{x}{a}} \right) + c .
\end{aligned}$$

where c is an integrating constant.

Indefinite Integration

Problem-18: $\int \frac{\sqrt{x}}{\sqrt{a^3 - x^3}} dx$

solⁿ : Let $I = \int \frac{\sqrt{x}}{\sqrt{a^3 - x^3}} dx$

$$= \int \frac{\sqrt{x}}{\sqrt{\left(a^{\frac{3}{2}}\right)^2 - \left(x^{\frac{3}{2}}\right)^2}} dx$$

Put $x^{\frac{3}{2}} = z \therefore \frac{3}{2} \sqrt{x} dx = dz$

Now $I = \frac{2}{3} \int \frac{dz}{\sqrt{\left(a^{\frac{3}{2}}\right)^2 - z^2}}$

$$= \frac{2}{3} \sin^{-1} \left(\frac{z}{a^{\frac{3}{2}}} \right) + c$$
$$= \frac{2}{3} \sin^{-1} \left(\frac{x}{a} \right)^{\frac{3}{2}} + c$$

where c is an integrating constant.

Problem-19: $\int \frac{x^2 + 1}{x^4 + 1} dx$

solⁿ : Let $I = \int \frac{x^2 + 1}{x^4 + 1} dx$

$$= \int \frac{x^2 \left(1 + \frac{1}{x^2}\right)}{x^2 \left(x^2 + \frac{1}{x^2}\right)} dx$$
$$= \int \frac{1 + \frac{1}{x^2}}{x^2 + \frac{1}{x^2}} dx$$
$$= \int \frac{1 + \frac{1}{x^2}}{2 + \left(x - \frac{1}{x}\right)^2} dx$$

Exercise-12: $\int \frac{x^2 - 1}{x^4 + 1} dx$

Ans: $\frac{1}{2\sqrt{2}} \ln \left(\frac{x^2 + 1 - \sqrt{2}x}{x^2 + 1 + \sqrt{2}x} \right) + c$

$$= \int \frac{1 + \frac{1}{x^2}}{\left(\sqrt{2}\right)^2 + \left(x - \frac{1}{x}\right)^2} dx$$

Put $x - \frac{1}{x} = z \therefore \left(1 + \frac{1}{x^2}\right) dx = dz$

Now $I = \int \frac{dz}{\left(\sqrt{2}\right)^2 + z^2}$

$$= \frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{z}{\sqrt{2}} \right) + c$$

$$= \frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{x^2 - 1}{\sqrt{2}x} \right) + c$$

where c is an integrating constant.

Problem-20: $\int \sqrt{1 + \sec x} dx$

solⁿ : Let $I = \int \sqrt{1 + \sec x} dx$

$$= \int \sqrt{1 + \frac{1}{\cos x}} dx$$

$$= \int \sqrt{\frac{1 + \cos x}{\cos x}} dx$$

$$= \int \sqrt{\frac{2 \cos^2 \frac{x}{2}}{\cos^2 \frac{x}{2} - \sin^2 \frac{x}{2}}} dx$$

$$= \int \sqrt{\frac{2 \cos^2 \frac{x}{2}}{1 - \sin^2 \frac{x}{2} - \sin^2 \frac{x}{2}}} dx$$

$$= \int \frac{\sqrt{2} \cos \frac{x}{2}}{\sqrt{1 - 2 \sin^2 \frac{x}{2}}} dx$$

$$= \int \frac{\sqrt{2} \cos \frac{x}{2}}{\sqrt{1 - \left(\sqrt{2} \sin \frac{x}{2}\right)^2}} dx$$

Indefinite Integration

Put $\sqrt{2} \sin \frac{x}{2} = z \quad \therefore \sqrt{2} \cos \frac{x}{2} dx = 2dz$

Now
$$I = \int \frac{2dz}{\sqrt{1-z^2}}$$
$$= 2 \int \frac{dz}{\sqrt{1-z^2}}$$
$$= 2 \sin^{-1} z + c$$
$$= 2 \sin^{-1} \left(\sqrt{2} \sin \frac{x}{2} \right) + c.$$

where c is an integrating constant.

Problem-21: $\int \sqrt{\tan x} dx$

solⁿ : Let $I = \int \sqrt{\tan x} dx$

put $\tan x = z^2 \quad \therefore \sec^2 x dx = 2z dz$

$$\Rightarrow dx = \frac{2z dz}{\sec^2 x}$$
$$= \frac{2z dz}{1 + \tan^2 x}$$
$$= \frac{2z dz}{1 + z^4}$$

Now
$$I = \int \frac{2z^2 dz}{1 + z^4}$$
$$= \int \frac{(z^2 + 1) + (z^2 - 1)}{z^4 + 1} dz$$
$$= \int \frac{z^2 + 1}{z^4 + 1} dz + \int \frac{z^2 - 1}{z^4 + 1} dz$$
$$= \int \frac{1 + \frac{1}{z^2}}{z^2 + \frac{1}{z^2}} dz + \int \frac{1 - \frac{1}{z^2}}{z^2 + \frac{1}{z^2}} dz$$
$$= \int \frac{1 + \frac{1}{z^2}}{2 + \left(z - \frac{1}{z}\right)^2} dz + \int \frac{1 - \frac{1}{z^2}}{\left(z + \frac{1}{z}\right)^2 - 2} dz$$

Exercise-13: $\int \frac{dx}{x^4 + 1}$

Ans: $\frac{1}{2\sqrt{2}} \tan^{-1} \left(\frac{x^2 - 1}{\sqrt{2}x} \right) - \frac{1}{4\sqrt{2}} \ln \left(\frac{x^2 + 1 - \sqrt{2}x}{x^2 + 1 + \sqrt{2}x} \right) + c$

$$\begin{aligned}
&= \int \frac{1 + \frac{1}{z^2}}{\left(\sqrt{2}\right)^2 + \left(z - \frac{1}{z}\right)^2} dz + \int \frac{1 - \frac{1}{z^2}}{\left(z + \frac{1}{z}\right)^2 - \left(\sqrt{2}\right)^2} dz \\
&= I_1 + I_2 \quad \dots \dots \dots (1)
\end{aligned}$$

$$\text{where, } I_1 = \int \frac{1 + \frac{1}{z^2}}{\left(\sqrt{2}\right)^2 + \left(z - \frac{1}{z}\right)^2} dz$$

$$\text{Put } z - \frac{1}{z} = t \therefore \left(1 + \frac{1}{z^2}\right) dz = dt$$

$$\begin{aligned}
\text{Now } I_1 &= \int \frac{dt}{\left(\sqrt{2}\right)^2 + t^2} \\
&= \frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{t}{\sqrt{2}} \right) + c_1 \\
&= \frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{z^2 - 1}{\sqrt{2}z} \right) + c_1
\end{aligned}$$

$$\text{and } I_2 = \int \frac{1 - \frac{1}{z^2}}{\left(z + \frac{1}{z}\right)^2 - \left(\sqrt{2}\right)^2} dz$$

$$\text{Put } z + \frac{1}{z} = t \therefore \left(1 - \frac{1}{z^2}\right) dz = dt$$

$$\begin{aligned}
\text{Now } I_2 &= \int \frac{dt}{t^2 - \left(\sqrt{2}\right)^2} \\
&= \frac{1}{2\sqrt{2}} \ln \left(\frac{t - \sqrt{2}}{t + \sqrt{2}} \right) + c_2 \\
&= \frac{1}{2\sqrt{2}} \ln \left(\frac{z^2 + 1 - \sqrt{2}z}{z^2 + 1 + \sqrt{2}z} \right) + c_2
\end{aligned}$$

From (1) we have,

Indefinite Integration

$$I = \frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{z^2 - 1}{\sqrt{2}z} \right) + \frac{1}{2\sqrt{2}} \ln \left(\frac{z^2 + 1 - \sqrt{2}z}{z^2 + 1 + \sqrt{2}z} \right) + c \quad ; \text{ putting, } c = c_1 + c_2$$

$$= \frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{\tan x - 1}{\sqrt{2} \tan x} \right) + \frac{1}{2\sqrt{2}} \ln \left(\frac{\tan x + 1 - \sqrt{2} \tan x}{\tan x + 1 + \sqrt{2} \tan x} \right) + c$$

where c is an integrating constant.

Problem-22: $\int \frac{dx}{\sqrt[3]{x} \sqrt[3]{(1+x)^5}}$

Exercise-14: $\int \frac{dx}{x^{\frac{1}{2}} (1+x)^{\frac{5}{2}}}$

solⁿ : Let $I = \int \frac{dx}{\sqrt[3]{x} \sqrt[3]{(1+x)^5}}$

$$= \int \frac{dx}{x^{\frac{1}{3}} (1+x)^{\frac{5}{3}}}$$

Ans: $2\sqrt{\left(\frac{x}{1+x}\right)} - \frac{2}{3}\left(\frac{x}{1+x}\right)^{\frac{3}{2}} + c$

Put $1+x = zx \Rightarrow x = \frac{1}{z-1}$

or, $z = 1 + \frac{1}{x}$

$\therefore dz = -\frac{1}{x^2} dx \Rightarrow dx = -x^2 dz$

Now $I = -\int \frac{x^2 dz}{x^{\frac{1}{3}} z^{\frac{5}{3}} x^{\frac{5}{3}}}$

$$= -\int \frac{x^2 dz}{x^2 z^{\frac{5}{3}}}$$

$$= -\int z^{-\frac{5}{3}} dz$$

$$= -\frac{z^{-\frac{5}{3}+1}}{-\frac{5}{3}+1} + c$$

$$= -\frac{z^{-\frac{2}{3}}}{-\frac{2}{3}} + c$$

$$= \frac{3}{2} \left(\frac{1+x}{x} \right)^{-\frac{2}{3}} + c$$

$$= \frac{3}{2} \left(\frac{x}{1+x} \right)^{\frac{2}{3}} + c$$

where c is an integrating constant.

NOTE: Integrals of the type $\int \frac{dx}{x^m (a+bx)^n}$ where $m \neq 0, n \neq 0$ can be evaluated exactly in the same way.

Some Important Standard Integrals

Problem-01: $\int \frac{dx}{4x^2 + 4x + 5}$

solⁿ : Let $I = \int \frac{dx}{4x^2 + 4x + 5}$

$$= \frac{1}{4} \int \frac{dx}{x^2 + x + \frac{5}{4}}$$

$$= \frac{1}{4} \int \frac{dx}{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} + \frac{5}{4} - \frac{1}{4}}$$

$$= \frac{1}{4} \int \frac{dx}{\left(x + \frac{1}{2}\right)^2 + 1}$$

$$= \frac{1}{4} \tan^{-1} \left(x + \frac{1}{2} \right) + c$$

where c is an integrating constant.

Problem-02: $\int \frac{dx}{1+x-x^2}$

solⁿ : Let $I = \int \frac{dx}{1+x-x^2}$

$$= \int \frac{dx}{-x^2 + x + 1}$$

$$= \int \frac{dx}{-(x^2 - x - 1)}$$

$$= \int \frac{dx}{-\left(x^2 - 2x \cdot \frac{1}{2} + \frac{1}{4} - 1 - \frac{1}{4}\right)}$$

Exercise-01: $\int \frac{dx}{1+x+x^2}$

Ans: $\frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{2x+1}{\sqrt{3}} \right) + c$.

Exercise-02: $\int \frac{dx}{x^2 - x - 6}$

Ans: $\frac{1}{5} \ln \left| \frac{x-3}{x+2} \right| + c$.

Exercise-03: $\int \frac{dx}{x^2 + 7x - 18}$

Ans: $\frac{1}{11} \ln \left| \frac{x-2}{x+9} \right| + c$.

Indefinite Integration

$$\begin{aligned} &= \int \frac{dx}{\frac{5}{4} - \left(x - \frac{1}{2}\right)^2} \\ &= \int \frac{dx}{\left(\frac{\sqrt{5}}{2}\right)^2 - \left(x - \frac{1}{2}\right)^2} \\ &= \frac{1}{\sqrt{5}} \ln \left(\frac{\frac{\sqrt{5}}{2} + \left(x - \frac{1}{2}\right)}{\frac{\sqrt{5}}{2} - \left(x - \frac{1}{2}\right)} \right) + c \\ &= \frac{1}{\sqrt{5}} \ln \left(\frac{\sqrt{5} + 2x - 1}{\sqrt{5} - 2x + 1} \right) + c \end{aligned}$$

where c is an integrating constant.

Problem-03: $\int \frac{dx}{\sqrt{x^2 - 7x + 12}}$

solⁿ : Let $I = \int \frac{dx}{\sqrt{x^2 - 7x + 12}}$

$$\begin{aligned} &= \int \frac{dx}{\sqrt{x^2 - 2 \cdot \frac{7}{2}x + \left(\frac{7}{2}\right)^2 - \frac{49}{4} + 12}} \\ &= \int \frac{dx}{\sqrt{\left(x - \frac{7}{2}\right)^2 - \left(\frac{1}{2}\right)^2}} \\ &= \ln \left(\left(x - \frac{7}{2}\right) + \sqrt{\left(x - \frac{7}{2}\right)^2 - \left(\frac{1}{2}\right)^2} \right) + c \\ &= \ln \left(\left(\frac{2x - 7}{2}\right) + \sqrt{x^2 - 7x + 12} \right) + c \end{aligned}$$

where c is an integrating constant.

Problem-04: $\int \frac{dx}{\sqrt{x^2 - 4x + 3}}$

solⁿ : Let $I = \int \frac{dx}{\sqrt{x^2 - 4x + 3}}$

Exercise-04: $\int \frac{dx}{\sqrt{1 - x - x^2}}$

Ans: $\sin^{-1} \left(\frac{2x + 1}{\sqrt{5}} \right) + c$.

Exercise-05: $\int \frac{dx}{\sqrt{2ax - x^2}}$

Ans: $\sin^{-1} \left(\frac{x - a}{a} \right) + c$.

Exercise-06: $\int \frac{dx}{\sqrt{3x - x^2 - 2}}$

Ans: $\sin^{-1} (2x - 3) + c$.

$$\begin{aligned}
&= \int \frac{dx}{\sqrt{x^2 - 2x + 4 - 1}} \\
&= \int \frac{dx}{\sqrt{(x-2)^2 - (1)^2}} \\
&= \ln \left((x-2) + \sqrt{(x-2)^2 - 1} \right) + c \\
&= \ln \left((x-2) + \sqrt{x^2 - 4x + 3} \right) + c.
\end{aligned}$$

where c is an integrating constant.

Problem-05: $\int \sqrt{4-3x-2x^2} dx$

Exercise-07: $\int \sqrt{18x-65-x^2} dx$

solⁿ : Let $I = \int \sqrt{4-3x-2x^2} dx$

$$\begin{aligned}
&= \int \sqrt{4-2\left(x^2 + \frac{3}{2}x\right)} dx \\
&= \sqrt{2} \int \sqrt{2-\left(x^2 + 2x \cdot \frac{3}{4} + \frac{9}{16} - \frac{9}{16}\right)} dx \\
&= \sqrt{2} \int \sqrt{2+\frac{9}{16}-\left(x^2 + 2x \cdot \frac{3}{4} + \frac{9}{16}\right)} dx \\
&= \sqrt{2} \int \sqrt{\frac{41}{16}-\left(x+\frac{3}{4}\right)^2} dx \\
&= \sqrt{2} \int \sqrt{\left(\frac{\sqrt{41}}{4}\right)^2 - \left(x+\frac{3}{4}\right)^2} dx \\
&= \sqrt{2} \left\{ \frac{\left(x+\frac{3}{4}\right) \sqrt{\left(\frac{\sqrt{41}}{4}\right)^2 - \left(x+\frac{3}{4}\right)^2}}{2} + \frac{\left(\frac{\sqrt{41}}{4}\right)^2}{2} \sin^{-1} \left(\frac{\left(x+\frac{3}{4}\right)}{\frac{\sqrt{41}}{4}} \right) \right\} + c \\
&= \sqrt{2} \left\{ \frac{(4x+3) \sqrt{\frac{41}{16} - \left(x+\frac{3}{4}\right)^2}}{8} + \frac{41}{32} \sin^{-1} \left(\frac{4x+3}{\sqrt{41}} \right) \right\} + c \\
&= \sqrt{2} \left\{ \frac{(4x+3) \sqrt{2-\frac{3}{2}x-x^2}}{8} + \frac{41}{32} \sin^{-1} \left(\frac{4x+3}{\sqrt{41}} \right) \right\} + c
\end{aligned}$$

Indefinite Integration

$$= \frac{(4x+3)\sqrt{4-3x-2x^2}}{8} + \frac{41\sqrt{2}}{32} \sin^{-1}\left(\frac{4x+3}{\sqrt{41}}\right) + c$$

where c is an integrating constant.

Problem-06: $\int \sqrt{2ax-x^2} dx$

Exercise-08: $\int \sqrt{3x-x^2} dx$

sol^n : Let $I = \int \sqrt{2ax-x^2} dx$

$$\begin{aligned} &= \int \sqrt{a^2 - (x^2 - 2ax + a^2)} dx \\ &= \int \sqrt{a^2 - (x-a)^2} dx \\ &= \frac{(x-a)\sqrt{a^2 - (x-a)^2}}{2} + \frac{a^2}{2} \sin^{-1}\left(\frac{x-a}{a}\right) + c \\ &= \frac{(x-a)\sqrt{2ax-x^2}}{2} + \frac{a^2}{2} \sin^{-1}\left(\frac{x-a}{a}\right) + c \end{aligned}$$

where c is an integrating constant.

NOTE: Integrals of the type $\int \frac{dx}{ax^2+bx+c}$, $\int \frac{dx}{\sqrt{ax^2+bx+c}}$, $\int \sqrt{ax^2+bx+c} dx$ where $a \neq 0$ can be evaluated exactly in the same way.

Problem-07: $\int \frac{dx}{\sqrt{(x-\alpha)(x-\beta)}}$

Exercise-09: $\int \frac{dx}{\sqrt{(x-\alpha)(\beta-x)}}$

sol^n : Let $I = \int \frac{dx}{\sqrt{(x-\alpha)(x-\beta)}}$

Ans: $2 \sin^{-1}\left(\frac{\sqrt{x-\alpha}}{\sqrt{\beta-\alpha}}\right) + c$.

Put $x-\alpha = z^2 \therefore dx = 2zdz$ and $x = z^2 + \alpha$

Now $I = \int \frac{2zdz}{\sqrt{z^2(z^2 + \alpha - \beta)}}$

$$\begin{aligned} &= 2 \int \frac{dz}{\sqrt{z^2 + (\sqrt{\alpha - \beta})^2}} \\ &= 2 \ln \left| z + \sqrt{z^2 + (\sqrt{\alpha - \beta})^2} \right| + c \\ &= 2 \ln \left| \sqrt{x - \alpha} + \sqrt{x - \alpha + \alpha - \beta} \right| + c \\ &= 2 \ln \left| \sqrt{x - \alpha} + \sqrt{x - \beta} \right| + c. \end{aligned}$$

where c is an integrating constant.

Problem-08: $\int \frac{x+1}{x^2+4x+5} dx$

solⁿ : Let $I = \int \frac{x+1}{x^2+4x+5} dx$

Put $x+1 = l(2x+4) + m$; $\left[\text{Let, } px+q = l \times \text{diff. coeff of } (ax^2+bx+c) + m \right]$

$\therefore 1 = 2l, 1 = 4l + m$

$\Rightarrow l = \frac{1}{2}, m = -1$

Now $I = \int \frac{\frac{1}{2}(2x+4) - 1}{x^2+4x+5} dx$

$$= \frac{1}{2} \int \frac{(2x+4)}{x^2+4x+5} dx - \int \frac{1}{x^2+4x+5} dx$$

$$= \frac{1}{2} \int \frac{(2x+4)}{x^2+4x+5} dx - \int \frac{1}{(x+2)^2+1} dx$$

$$= \frac{1}{2} \ln(x^2+4x+5) - \tan^{-1}(x+2) + c.$$

where c is an integrating constant.

Problem-09: $\int \frac{x-2}{\sqrt{2x^2-8x+5}} dx$

solⁿ : Let $I = \int \frac{x-2}{\sqrt{2x^2-8x+5}} dx$

Put $x-2 = l(4x-8) + m$; $\left[\text{Let, } px+q = l \times \text{diff. coeff of } (ax^2+bx+c) + m \right]$

$\therefore 1 = 4l, -2 = -8l + m$

$\Rightarrow l = \frac{1}{4}, m = 0$

Now $I = \int \frac{\frac{1}{4}(4x-8)}{\sqrt{2x^2-8x+5}} dx$

$$= \frac{1}{4} \int \frac{dz}{\sqrt{z}} ; \text{ putting } 2x^2-8x+5 = z$$

$$= \frac{1}{4} \int z^{-\frac{1}{2}} dz$$

$$= \frac{1}{4} \frac{z^{-\frac{1}{2}+1}}{-\frac{1}{2}+1} + c$$

Exercise-10: $\int \frac{4x+15}{x^2+6x+10} dx$

Ans: $2 \ln(x^2+6x+10) + 3 \tan^{-1}(x+3) + c.$

Exercise-11: $\int \frac{x-1}{\sqrt{4+x^2-2x}} dx$

Ans: $\sqrt{x^2-2x+4} + c.$

Exercise-12: $\int \frac{2x+5}{\sqrt{x^2-2x+2}} dx$

Indefinite Integration

$$= \frac{1}{2} z^{\frac{1}{2}} + c$$

$$= \frac{1}{2} \sqrt{2x^2 - 8x + 5} + c$$

where c is an integrating constant.

NOTE: Integrals of the type $\int \frac{px+q}{ax^2+bx+c} dx$, $\int \frac{px+q}{\sqrt{ax^2+bx+c}} dx$ where $a \neq 0, p \neq 0$ can be evaluated exactly in the same way.

Problem-10: $\int \frac{dx}{(2x+1)\sqrt{(4x+3)}}$

solⁿ : Let $I = \int \frac{dx}{(2x+1)\sqrt{(4x+3)}}$

Put $4x+3 = z^2$ or $x = \frac{z^2-3}{4}$

$$\therefore dx = \frac{1}{2} z dz$$

Now $I = \int \frac{\frac{1}{2} z dz}{\left(\frac{z^2-3}{2} + 1\right) z}$ **Exercise-15:** $\int \frac{dx}{(1-x)\sqrt{(1+x)}}$

$$= \frac{1}{2} \int \frac{dz}{\left(\frac{z^2-3+2}{2}\right)} \text{ Ans: } \frac{1}{\sqrt{2}} \ln \left(\frac{\sqrt{2} + \sqrt{1+x}}{\sqrt{2} - \sqrt{1-x}} \right) + c.$$

$$= \int \frac{dz}{z^2-1}$$

$$= \frac{1}{2} \ln \left| \frac{z-1}{z+1} \right| + c$$

$$= \frac{1}{2} \ln \left| \frac{\sqrt{4x+3}-1}{\sqrt{4x+3}+1} \right| + c$$

where c is an integrating constant.

Problem-11: $\int \frac{xdx}{(1+x^2)\sqrt{(x^2-1)}}$

solⁿ : Let $I = \int \frac{xdx}{(1+x^2)\sqrt{(x^2-1)}}$

Put $x^2-1 = z^2$ or $x^2 = z^2+1$

Exercise-13: $\int \frac{dx}{(2+x)\sqrt{(1+x)}}$

Ans: $2 \tan^{-1}(\sqrt{1+x}) + c.$

Exercise-14: $\int \frac{dx}{(x-3)\sqrt{(x-2)}}$

Ans: $\ln \left(\frac{\sqrt{x-2}-1}{\sqrt{x-2}+1} \right) + c.$

Exercise-16: $\int \frac{xdx}{(x^2+2)\sqrt{(x^2+3)}}$

Ans: $\frac{1}{2} \ln \left(\frac{\sqrt{x^2+3}-1}{\sqrt{x^2+3}+1} \right) + c.$

$$\therefore xdx = zdz$$

$$\begin{aligned}\text{Now } I &= \int \frac{zdz}{(z^2+1+1)z} \\ &= \int \frac{dz}{z^2+2} \\ &= \int \frac{dz}{(\sqrt{2})^2+z^2} \\ &= \frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{z}{\sqrt{2}} \right) + c \\ &= \frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{x^2-1}}{\sqrt{2}} \right) + c\end{aligned}$$

where c is an integrating constant.

NOTE: Integrals of the type $\int \frac{dx}{(ax+b)\sqrt{(cx+d)}}$, $\int \frac{xdx}{(ax^2+b)\sqrt{(cx^2+d)}}$ where $a \neq 0, c \neq 0$ can be evaluated exactly in the same way.

Problem-12: $\int \frac{dx}{(1+x^2)\sqrt{(x^2+4)}}$

solⁿ : Let $I = \int \frac{dx}{(1+x^2)\sqrt{(x^2+4)}}$

Put $x = \frac{1}{z}$

$$\therefore dx = -\frac{1}{z^2} dz$$

$$\begin{aligned}\text{Now } I &= \int \frac{-\frac{1}{z^2} dz}{\left(\frac{1}{z^2}+1\right)\sqrt{\frac{1}{z^2}+4}} \\ &= -\int \frac{zdz}{(z^2+1)\sqrt{4z^2+1}}\end{aligned}$$

Again let $4z^2+1=t^2$ or, $z^2 = \frac{t^2-1}{4}$

$$\therefore zdz = \frac{1}{4} tdt$$

Exercise-17: $\int \frac{dx}{(x^2+1)\sqrt{(1-x^2)}}$

Ans: $-\frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{1-x^2}}{x\sqrt{2}} \right) + c$.

Indefinite Integration

$$\begin{aligned}
 \therefore I &= -\frac{1}{4} \int \frac{tdt}{t \left(\frac{t^2-1}{4} + 1 \right)} \\
 &= -\frac{1}{4} \int \frac{dt}{\left(\frac{t^2+3}{4} \right)} \\
 &= -\int \frac{dt}{3+t^2} \\
 &= -\int \frac{dt}{(\sqrt{3})^2 + t^2} \\
 &= -\frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{t}{\sqrt{3}} \right) + c \\
 &= -\frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{\sqrt{4z^2+1}}{\sqrt{3}} \right) + c \\
 &= -\frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{\sqrt{4+x^2}}{x\sqrt{3}} \right) + c
 \end{aligned}$$

where c is an integrating constant.

NOTE: Integrals of the type $\int \frac{dx}{(ax^2+b)\sqrt{cx^2+d}}$ where $a \neq 0, c \neq 0$ can be evaluated exactly in the same way.

Problem-13: $\int \frac{dx}{(1+x)\sqrt{(1+2x-x^2)}}$

solⁿ: Let $I = \int \frac{dx}{(1+x)\sqrt{(1+2x-x^2)}}$

Put $1+x = \frac{1}{z} \therefore dx = -\frac{1}{z^2} dz$

Now $I = \int \frac{-\frac{1}{z^2} dz}{\frac{1}{z} \sqrt{\left[1+2\left(\frac{1}{z}-1\right) - \left(\frac{1}{z}-1\right)^2 \right]}}$

$$= -\int \frac{dz}{z \sqrt{\left[1+\frac{2}{z}-2 - \left(\frac{1}{z^2}-\frac{2}{z}+1\right) \right]}}$$

Exercise-18: $\int \frac{dx}{(x-3)\sqrt{(x^2-6x+8)}}$

Ans: $\sec^{-1}(x-3) + c$.

Exercise-19: $\int \frac{dx}{(2x+3)\sqrt{(x^2+3x+2)}}$

Ans: $\sec^{-1}(2x+3) + c$.

Exercise-20: $\int \frac{dx}{(x-1)\sqrt{(x^2+2x+2)}}$

$$\begin{aligned}
&= -\int \frac{dz}{z\sqrt{\left(-\frac{1}{z^2} + \frac{4}{z} - 2\right)}} \\
&= -\int \frac{dz}{\sqrt{(-1 + 4z - 2z^2)}} \\
&= -\frac{1}{\sqrt{2}} \int \frac{dz}{\sqrt{\left(-\frac{1}{2} + 2z - z^2\right)}} \\
&= -\frac{1}{\sqrt{2}} \int \frac{dz}{\sqrt{-\frac{1}{2} - (z^2 - 2z)}} \\
&= -\frac{1}{\sqrt{2}} \int \frac{dz}{\sqrt{-\frac{1}{2} - (z^2 - 2z + 1) + 1}} \\
&= -\frac{1}{\sqrt{2}} \int \frac{dz}{\sqrt{\left[\frac{1}{2} - (z-1)^2\right]}} \\
&= -\frac{1}{\sqrt{2}} \int \frac{dz}{\sqrt{\left[\left(\frac{1}{\sqrt{2}}\right)^2 - (z-1)^2\right]}} \\
&= -\frac{1}{\sqrt{2}} \sin^{-1} \left(\frac{z-1}{\frac{1}{\sqrt{2}}} \right) + c \\
&= -\frac{1}{\sqrt{2}} \sin^{-1} \left(\frac{\frac{1}{(x+1)} - 1}{\frac{1}{\sqrt{2}}} \right) + c \\
&= -\frac{1}{\sqrt{2}} \sin^{-1} \left(-x\sqrt{2} / (x+1) \right) + c \\
&= \frac{1}{\sqrt{2}} \sin^{-1} \left(x\sqrt{2} / (x+1) \right) + c.
\end{aligned}$$

where c is an integrating constant.

NOTE: Integrals of the type $\int \frac{dx}{(px+q)\sqrt{ax^2+bx+c}}$ where $p \neq 0, a \neq 0$ can be evaluated exactly in the same way.

Integration by Parts

The formula for the integration of a product of two functions is referred to as integration by parts. *i.e.*,

$$\int (uv)dx = u \int vdx - \int \left\{ \frac{du}{dx} \int vdx \right\} dx .$$

While applying the above rule for integration by parts to the product of two functions, care should be taken to choose properly the first function, *i.e.*, the function not to be integrated.

Illustrative Examples:

Problem-01: $\int x e^x dx$

solⁿ : Let $I = \int x e^x dx$

$$= x \int e^x dx - \int \left\{ \frac{d}{dx}(x) \int e^x dx \right\} dx$$

$$= x e^x - \int 1 \cdot e^x dx$$

$$= x e^x - e^x + c$$

where c is an integrating constant.

Problem-02: $\int x^3 e^{-x} dx$

solⁿ : Let $I = \int x^3 e^{-x} dx$

$$= x^3 \int e^{-x} dx - \int \left\{ \frac{d}{dx}(x^3) \int e^{-x} dx \right\} dx$$

$$= -x^3 e^{-x} - \int \{ 3x^2 (-e^{-x}) \} dx$$

$$= -x^3 e^{-x} + 3 \int x^2 e^{-x} dx$$

$$= -x^3 e^{-x} + 3 \left[x^2 \int e^{-x} dx - \int \left\{ \frac{d}{dx}(x^2) \int e^{-x} dx \right\} dx \right]$$

$$= -x^3 e^{-x} + 3 \left[-x^2 e^{-x} - \int \{ 2x (-e^{-x}) \} dx \right]$$

$$= -x^3 e^{-x} - 3x^2 e^{-x} + 6 \int x e^{-x} dx$$

$$= -x^3 e^{-x} - 3x^2 e^{-x} + 6 \left[x \int e^{-x} dx - \int \left\{ \frac{dx}{dx} \int e^{-x} dx \right\} dx \right]$$

Exercise-01: $\int x^2 \cos x dx$

Ans: $x^2 \sin x + 2x \cos x - 2 \sin x + c$

Exercise-02: $\int x^n \ln x dx$

Ans: $\frac{x^{n+1}}{(n+1)^2} \{ (n+1) \ln x - 1 \} + c$

$$\begin{aligned}
&= -x^3 e^{-x} - 3x^2 e^{-x} + 6 \left[-x e^{-x} - \int 1 \cdot (-e^{-x}) dx \right] \\
&= -x^3 e^{-x} - 3x^2 e^{-x} - 6x e^{-x} + 6 \int e^{-x} dx \\
&= -x^3 e^{-x} - 3x^2 e^{-x} - 6x e^{-x} - 6e^{-x} + c
\end{aligned}$$

where c is an integrating constant.

Problem-03: $\int \tan^{-1} x dx$

$$sol^n : Let I = \int \tan^{-1} x dx$$

$$\begin{aligned}
&= \tan^{-1} x \int dx - \int \left\{ \frac{d}{dx} (\tan^{-1} x) \int dx \right\} dx \\
&= x \tan^{-1} x - \int \frac{1}{1+x^2} \cdot x dx \\
&= x \tan^{-1} x - \frac{1}{2} \int \frac{2x}{1+x^2} dx \\
&= x \tan^{-1} x - \frac{1}{2} \ln(1+x^2) + c
\end{aligned}$$

where c is an integrating constant.

Problem-04: $\int \frac{x e^x}{(1+x)^2} dx$

$$sol^n : Let I = \int \frac{x e^x}{(1+x)^2} dx$$

$$\begin{aligned}
&= \int \frac{(x+1-1) e^x}{(1+x)^2} dx \\
&= \int \frac{e^x}{1+x} dx - \int \frac{e^x}{(1+x)^2} dx \\
&= \frac{1}{1+x} \int e^x dx - \int \left\{ \frac{d}{dx} \left(\frac{1}{1+x} \right) \int e^x dx \right\} dx - \int \frac{e^x}{(1+x)^2} dx + c \\
&= \frac{e^x}{1+x} - \int \left\{ \frac{-1}{(1+x)^2} e^x \right\} dx - \int \frac{e^x}{(1+x)^2} dx + c \\
&= \frac{e^x}{1+x} + \int \frac{e^x}{(1+x)^2} dx - \int \frac{e^x}{(1+x)^2} dx + c \\
&= \frac{e^x}{1+x} + c
\end{aligned}$$

where c is an integrating constant.

Exercise-03: $\int \cos^{-1} x dx$

$$Ans: x \cos^{-1} x - \sqrt{1-x^2} + c$$

Exercise- 04: $\int e^x \frac{x^2+1}{(1+x)^2} dx$

$$Ans: e^x \left(\frac{x-1}{x+1} \right) + c$$

Indefinite Integration

Problem-05: $\int e^x \left(\frac{1-x}{1+x^2} \right)^2 dx$

solⁿ : Let $I = \int e^x \left(\frac{1-x}{1+x^2} \right)^2 dx$

$$= \int e^x \frac{1-2x+x^2}{(1+x^2)^2} dx$$

$$= \int e^x \left[\frac{(1+x^2)-2x}{(1+x^2)^2} \right] dx$$

$$= \int e^x \left[\frac{1}{1+x^2} - \frac{2x}{(1+x^2)^2} \right] dx$$

$$= \int \frac{e^x}{1+x^2} dx - \int e^x \frac{2x}{(1+x^2)^2} dx$$

$$= \frac{1}{1+x^2} \int e^x dx - \int \left\{ \frac{d}{dx} \left(\frac{1}{1+x^2} \right) \int e^x dx \right\} dx - \int e^x \frac{2x}{(1+x^2)^2} dx$$

$$= \frac{e^x}{1+x^2} + \int e^x \frac{2x}{(1+x^2)^2} dx - \int e^x \frac{2x}{(1+x^2)^2} dx$$

$$= \frac{e^x}{1+x^2} + c$$

where c is an integrating constant.

Problem-06: $\int \cos \sqrt{x} dx$

solⁿ : Let $I = \int \cos \sqrt{x} dx$

Put $\sqrt{x} = z \therefore \frac{1}{2\sqrt{x}} dx = dz \Rightarrow dx = 2z dz$ $+\frac{1}{8} \sin 2x + c$

Now $I = \int \cos z \cdot 2z dz$

$$= 2 \int z \cos z dz$$

$$= 2 \left[z \int \cos z dz - \int \left\{ \frac{dz}{dz} \int \cos z dz \right\} dz \right]$$

$$= 2 \left[z \sin z - \int \sin z dz \right]$$

Exercise- 05: $\int e^x \frac{x-1}{(1+x)^3} dx$

Ans: $e^x \frac{1}{(x+1)^2} + c$

Exercise-06: $\int x^2 \sin^2 x dx$

Ans: $\frac{x^3}{6} - \frac{1}{4} x^2 \sin 2x - \frac{1}{4} x \cos 2x$

$$\begin{aligned}
&= 2[z \sin z + \cos z] + c \\
&= 2[\sqrt{x} \sin \sqrt{x} + \cos \sqrt{x}] + c
\end{aligned}$$

where c is an integrating constant.

Problem-07: $\int x^2 \sin x \cos x dx$

$$\begin{aligned}
\text{sol}^n : \text{Let } I &= \int x^2 \sin x \cos x dx \\
&= \frac{1}{2} \int x^2 \cdot 2 \sin x \cos x dx \\
&= \frac{1}{2} \int x^2 \sin 2x dx \\
&= \frac{1}{2} \left[x^2 \int \sin 2x dx - \int \left\{ \frac{d}{dx}(x^2) \int \sin 2x dx \right\} dx \right] \\
&= \frac{1}{2} \left[-\frac{x^2}{2} \cos 2x + \int x \cos 2x dx \right] \\
&= -\frac{x^2}{4} \cos 2x + \frac{1}{2} \left[x \int \cos 2x dx - \int \left\{ \frac{d}{dx}(x) \int \cos 2x dx \right\} dx \right] \\
&= -\frac{x^2}{4} \cos 2x + \frac{1}{2} \left[\frac{x}{2} \sin 2x - \frac{1}{2} \int \sin 2x dx \right] \\
&= -\frac{x^2}{4} \cos 2x + \frac{1}{2} \left[\frac{x}{2} \sin 2x + \frac{1}{4} \cos 2x \right] + c \\
&= -\frac{x^2}{4} \cos 2x + \frac{x}{4} \sin 2x + \frac{1}{8} \cos 2x + c
\end{aligned}$$

where c is an integrating constant.

Problem-08: $\int \frac{\ln(\ln x)}{x} dx$

$$\text{sol}^n : \text{Let } I = \int \frac{\ln(\ln x)}{x} dx$$

$$\text{Put } \ln x = z \quad \therefore \frac{1}{x} dx = dz$$

$$\text{Now } I = \int \ln z dz$$

$$\begin{aligned}
&= \ln z \int dz - \int \left\{ \frac{d}{dz}(\ln z) \int dz \right\} dz \\
&= z \ln z - \int \frac{1}{z} \cdot z dz
\end{aligned}$$

Indefinite Integration

$$\begin{aligned} &= z \ln z - \int dz \\ &= z \ln z - z + c \\ &= \ln x \ln(\ln x) - \ln x + c \end{aligned}$$

where c is an integrating constant.

Problem-09: $\int \frac{x}{1 + \cos x} dx$

Solⁿ : Let $I = \int \frac{x}{1 + \cos x} dx$

$$\begin{aligned} &= \int \frac{x}{2 \cos^2 \frac{x}{2}} dx \\ &= \frac{1}{2} \int x \sec^2 \frac{x}{2} dx \\ &= \frac{1}{2} \left[x \int \sec^2 \frac{x}{2} dx - \int \left\{ \frac{d}{dx}(x) \int \sec^2 \frac{x}{2} dx \right\} dx \right] \\ &= \frac{1}{2} \left[x \frac{\tan \frac{x}{2}}{1/2} - \int \frac{\tan \frac{x}{2}}{1/2} dx \right] + c \\ &= x \tan \frac{x}{2} - \int \tan \frac{x}{2} dx + c \\ &= x \tan \frac{x}{2} - \frac{\ln \left| \sec \frac{x}{2} \right|}{1/2} + c \\ &= x \tan \frac{x}{2} - 2 \ln \left| \sec \frac{x}{2} \right| + c \end{aligned}$$

where c is an integrating constant.

Problem-10: $\int e^{ax} \cos bxdx$

Solⁿ : Let $I = \int e^{ax} \cos bxdx$

$$\begin{aligned} &= e^{ax} \int \cos bxdx - \int \left\{ \frac{d}{dx}(e^{ax}) \int \cos bxdx \right\} dx \\ &= \frac{e^{ax} \sin bx}{b} - \int \left\{ ae^{ax} \frac{\sin bx}{b} \right\} dx \end{aligned}$$

Exercise-07: $\int \frac{x + \sin x}{1 + \cos x} dx$

Ans: $x \tan \frac{x}{2} + c$

Exercise-08: $\int e^{ax} \sin(bx + d) dx$

Ans: $\frac{e^{ax} [a \sin(bx + d) - b \cos(bx + d)]}{a^2 + b^2} + c$

$$\begin{aligned}
&= \frac{e^{ax} \sin bx}{b} - \frac{a}{b} \int e^{ax} \sin bx dx \\
&= \frac{e^{ax} \sin bx}{b} - \frac{a}{b} \left[e^{ax} \int \sin bx dx - \int \left\{ \frac{d}{dx} (e^{ax}) \int \sin bx dx \right\} dx \right] \\
&= \frac{e^{ax} \sin bx}{b} - \frac{a}{b} \left[\frac{-e^{ax} \cos bx}{b} + \frac{a}{b} \int e^{ax} \cos bx dx \right] \\
&= \frac{e^{ax} \sin bx}{b} + \frac{a}{b^2} e^{ax} \cos bx - \frac{a^2}{b^2} I \\
\therefore I + \frac{a^2}{b^2} I &= \frac{e^{ax} \sin bx}{b} + \frac{a}{b^2} e^{ax} \cos bx \\
\Rightarrow \frac{I(a^2 + b^2)}{b^2} &= \frac{be^{ax} \sin bx + ae^{ax} \cos bx}{b^2} \\
\Rightarrow I &= \frac{e^{ax} (a \cos bx + b \sin bx)}{a^2 + b^2} \\
\therefore I &= \frac{e^{ax} (a \cos bx + b \sin bx)}{a^2 + b^2} + c
\end{aligned}$$

where c is an integrating constant.

Integration of Trigonometric Functions

Problem-01: $\int \frac{dx}{5 + 4 \cos x}$

Solⁿ : Let $I = \int \frac{dx}{5 + 4 \frac{1 - \tan^2 x/2}{1 + \tan^2 x/2}}$

$$= \int \frac{dx}{\frac{5 + 5 \tan^2 x/2 + 4 - 4 \tan^2 x/2}{1 + \tan^2 x/2}}$$

$$= \int \frac{1 + \tan^2 x/2}{9 + \tan^2 x/2} dx$$

$$= \int \frac{\sec^2 x/2}{3^2 + \tan^2 x/2} dx$$

Exercise-01: $\int \frac{dx}{2 + \cos x}$

Ans: $\frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{1}{\sqrt{3}} \tan \frac{x}{2} \right) + c$

Indefinite Integration

Put $\tan \frac{x}{2} = z \therefore \sec^2 \frac{x}{2} dx = 2dz$

Now $I = 2 \int \frac{dz}{3^2 + z^2}$

$$= \frac{2}{3} \tan^{-1} \frac{z}{3} + c$$
$$= \frac{2}{3} \tan^{-1} \left(\frac{1}{3} \tan \frac{x}{2} \right) + c$$

where c is an integrating constant.

Problem-02: $\int \frac{dx}{2 + 3 \cos 2x}$

Solⁿ : Let $I = \int \frac{dx}{2 + 3 \frac{1 - \tan^2 x}{1 + \tan^2 x}}$

$$= \int \frac{dx}{\frac{2 + 2 \tan^2 x + 3 - 3 \tan^2 x}{1 + \tan^2 x}}$$
$$= \int \frac{1 + \tan^2 x}{5 - \tan^2 x} dx$$
$$= \int \frac{\sec^2 x}{5 - \tan^2 x} dx$$

Put $\tan x = z \therefore \sec^2 x dx = dz$

Now $I = \int \frac{dz}{5 - z^2}$

$$= \int \frac{dz}{(\sqrt{5})^2 - z^2}$$
$$= \frac{1}{2\sqrt{5}} \ln \left(\frac{\sqrt{5} + z}{\sqrt{5} - z} \right) + c$$
$$= \frac{1}{2\sqrt{5}} \ln \left(\frac{\sqrt{5} + \tan x}{\sqrt{5} - \tan x} \right) + c$$

where c is an integrating constant.

Exercise-02: $\int \frac{dx}{3 + 5 \cos x}$

Ans: $\frac{1}{4} \ln \left| \frac{2 + \tan \frac{x}{2}}{2 - \tan \frac{x}{2}} \right| + c$

Problem-03: $\int \frac{dx}{4+5\sin x}$

Solⁿ : Let $I = \int \frac{dx}{4+5\sin x}$

$$= \int \frac{dx}{4+5 \cdot \frac{2 \tan \frac{x}{2}}{1+\tan^2 \frac{x}{2}}}$$

$$= \int \frac{dx}{\frac{4+4 \tan^2 \frac{x}{2}+10 \tan \frac{x}{2}}{1+\tan^2 \frac{x}{2}}}$$

$$= \int \frac{1+\tan^2 \frac{x}{2}}{4 \tan^2 \frac{x}{2}+10 \tan \frac{x}{2}+4} dx$$

$$= \int \frac{\sec^2 \frac{x}{2}}{4 \tan^2 \frac{x}{2}+10 \tan \frac{x}{2}+4} dx$$

Put $\tan \frac{x}{2} = z \quad \therefore \sec^2 \frac{x}{2} dx = 2dz$

Now $I = \int \frac{2dz}{4z^2+10z+4}$

$$= \frac{1}{2} \int \frac{dz}{\left(z^2 + \frac{5}{2}z + 1\right)}$$

$$= \frac{1}{2} \int \frac{dz}{z^2 + 2 \cdot \frac{5}{4}z + \left(\frac{5}{4}\right)^2 - \frac{9}{16}}$$

$$= \frac{1}{2} \int \frac{dz}{\left(z + \frac{5}{4}\right)^2 - \left(\frac{3}{4}\right)^2}$$

$$= \frac{1}{2} \left[\frac{1}{2 \cdot \frac{3}{4}} \ln \left| \frac{\left(z + \frac{5}{4}\right) - \frac{3}{4}}{\left(z + \frac{5}{4}\right) + \frac{3}{4}} \right| \right] + c$$

Exercise-03: $\int \frac{dx}{3+2\cos x}$

Ans: $\frac{2}{\sqrt{5}} \tan^{-1} \left\{ \frac{3 \tan \left(\frac{x}{2}\right) + 2}{\sqrt{5}} \right\} + c$

Indefinite Integration

$$= \frac{1}{3} \ln \left| \frac{\tan \frac{x}{2} + \frac{1}{2}}{\tan \frac{x}{2} + 2} \right| + c$$

where c is an integrating constant.

Problem-04: $\int \frac{11 \cos x - 16 \sin x}{2 \cos x + 5 \sin x} dx$

Solⁿ : Let $I = \int \frac{11 \cos x - 16 \sin x}{2 \cos x + 5 \sin x} dx$

Put $11 \cos x - 16 \sin x = l(2 \cos x + 5 \sin x) + m(-2 \sin x + 5 \cos x) + n$

Comparing coefficient of $\cos x$, $\sin x$ and constant terms, we get

$$2l + 5m = 11; 5l - 2m = -16; n = 0$$

Solving,

$$l = -2; m = 3; n = 0$$

Now
$$\begin{aligned} I &= \int \frac{-2(2 \cos x + 5 \sin x) + 3(-2 \sin x + 5 \cos x)}{2 \cos x + 5 \sin x} dx \\ &= -2 \int \frac{2 \cos x + 5 \sin x}{2 \cos x + 5 \sin x} dx + 3 \int \frac{-2 \sin x + 5 \cos x}{2 \cos x + 5 \sin x} dx \\ &= -2 \int dx + 3 \int \frac{-2 \sin x + 5 \cos x}{2 \cos x + 5 \sin x} dx \\ &= -2x + 3 \ln(2 \cos x + 5 \sin x) + c \end{aligned}$$

where c is an integrating constant.

Exercise-04: $\int \frac{2 \sin x + 3 \cos x}{7 \sin x - 2 \cos x} dx$

Ans: $\frac{8x}{53} + \frac{25}{53} \ln|7 \sin x - 2 \cos x| + c$

Exercise-05: $\int \frac{2 \sin x + 3 \cos x}{3 \sin x + 4 \cos x} dx$

Ans: $\frac{18x}{25} + \frac{1}{25} \ln|3 \sin x + 4 \cos x| + c$