

Doctor Information

Dr. Jeba Tasnia

Patient Section

Manual Entry Prescription

Rx

Medicine Details	Dosage
Napa Paracetamol Indication: Paracetamol is indicated for fever, common cold and influenza, headache, toothache, earache, bodyache, myalgia, neuralgia, dysmenorrhoea, sprains, colic pain, back pain, post-operative pain, postpartum pain, inflammatory pain and post vaccination pain in children. It is also indicated for rheumat... Pharmacology: Paracetamol has analgesic and antipyretic properties with weak anti-inflammatory activity. Paracetamol (Acetaminophen) is thought to act primarily in the CNS, increasing the pain threshold by inhibiting both isoforms of cyclooxygenase, COX-1, COX-2, and COX-3 enzymes involved in prostaglandin (PG... Dosage: Tablet: Adult: 1-2 tablets every 4 to 6 hours up to a maximum of 4 gm (8 tablets) daily. Children (6-12 years): ½ to 1 tablet 3 to 4 times daily. For long term treatment it is wise not to exceed the dose beyond 2.6 gm/day. Extended Release Tablet: Adults & Children over 12 years: Two tablets, swa... Side Effects: Side effects of paracetamol are usually mild, though haematological reactions including thrombocytopenia, leucopenia, pancytopenia, neutropenia, and agranulocytosis have been reported. Pancreatitis, skin rashes, and other allergic reactions occur occasionally.. Contraindications: It is contraindicated in known hypersensitivity to Paracetamol.. Pregnancy & Lactation: Pregnancy category B according to USFDA. This drug should be used during pregnancy only if clearly needed. Storage: N/A.	1+0+1
Omeprol Omeprazole Indication: Omeprazole is indicated for the treatment of- Gastric and duodenal ulcer NSAID-associated duodenal and gastric ulcer As prophylaxis in patients with a history of NSAID-associated duodenal and gastric ulcer Gastro-esophageal reflux disease Long-term management of acid reflux disease ... Read more ... Pharmacology: Omeprazole, a substituted benzimidazole, is an inhibitor of gastric acid secretion. It inhibits gastric acid secretion by blocking hydrogen-potassium-adenosine triphosphatase (H+/K+ ATPase) enzyme system in the gastric parietal cell. After oral administration, the onset of the antisecretory effec... Dosage: Oral- Benign gastric and duodenal ulcer: 20 mg once daily for 4 weeks in duodenal ulceration, 8 weeks in gastric ulceration; in severe or recurrent cases, dose to be increased to 40 mg daily; maintenance dose for recurrent duodenal ulcer, 20 mg once daily; in prevention of relapse in duodenal ulc... Side Effects: Omeprazole is generally well tolerated. Nausea, abdominal colic, paresthesia, dizziness and headache have been stated to be generally mild and transient and not requiring a reduction in dosage.. Contraindications: Omeprazole is contraindicated in patients with known hypersensitivity to any of the components of the formulation.. Pregnancy & Lactation: US FDA pregnancy category of Omeprazole is C. However, results from three prospective epidemiological studies indicate no adverse effects of Omeprazole on pregnancy or on the health of the fetus/newborn child. There is no information available on the passage of Omeprazole into breast milk or its ... Storage: N/A.	1+1+1

