

Doctor Information

Dr. Dr. Tahira Jahan

Patient Section

Manual Prescription Entry

Rx

Medicine Details	Dosage
Ketosteril <i>Amino Acid + Calcium</i> Indication: Prevention and therapy of damages due to faulty or deficient protein metabolism in chronic renal insufficiency in connection with limited protein in food of 40 g/day (for adults) ie, generally in patients with a glomerular filtration rate (GFR) 25 mL/minutes. Side Effects: Hypercalcemia may develop. In this case it is recommended to decrease Vitamin D intake. If hypercalcaemia persists, reduce the dosage of this tablet as well as other source of calcium. The serum calcium level should be monitored regularly. Ensure sufficient calorie intake. No experience has been gained so far with the administration in paediatric patients.	1-0-1
Robentyl <i>Dicycloverine Hydrochloride</i> Indication: Dicycloverine is indicated in: Functional bowel/irritable bowel syndrome Urinary incontinence secondary to unstable detrusor muscle Infantile colic GIT spasm Colicky abdominal pain Diverticulitis Abdominal colic Side Effects: Insomnia, mydriasis, cycloplegia, increased ocular tension, urinary hesitancy, palpitations, dyspnea.	1-1-1
Elimate Plus <i>Permethrin + Crothamiton</i> Indication: This lotion is indicated for the treatment of scabies and pruritus. Side Effects: Occasional reports of burning or stinging sensation which passes quickly. This is usually mild and occurs more frequently in patients with severe scabies. Transient signs and symptoms of skin irritation including rash or itching, erythema, edema and eczema. These are generally considered to be part of the natural history of scabies.	1-0-0