

Doctor Information

Dr. Tasnuva Shreya

Patient Section

Manual Entry Prescription

Rx

Medicine Details	Dosage
Gabapen Gabapentin Indication: Gabapentin is indicated for- Neuropathic Pain. Adjunctive therapy in partial seizure and secondary generalized seizure.. Pharmacology: Gabapentin is an anti-convulsant. It is a structural analog of gamma-amino-butyric-acid (GABA). All pharmacological actions following administration of Gabapentin are due to the activity of parent compound. Gabapentin binds with the alpha-2-delta subunit of voltage gated L-type Calcium channel, a... Dosage: Neuropathic Pain: The treatment may be initiated as a single 300 mg dose on Day-1, than 300 mg twice on Day-2 and 300 mg thrice on Day-3. The dose can be subsequently be titrated up as needed for pain relief to a daily dose of 1800 mg (divided TID). Epilepsy: Patients over 12 years of age: the ef... Side Effects: Fatigue, Dizziness, ataxia, weight gain, peripheral edema, dry mouth and somnolence may occur.. Contraindications: Gabapentin is contraindicated in patients who have known hypersensitivity to the drug.. Pregnancy & Lactation: Pregnancy category C. it should be used during pregnancy only if potential benefits justifies the potential risk to the fetus. Gabapentin may be secreted through the breast milk. So it should be used during lactation only if potential benefits justifies the potential risk to the baby.. Storage: N/A.	1-0-1
Advel Ibuprofen Indication: Ibuprofen is indicated in the following indications- Rheumatoid arthritis Osteoarthritis Gouty arthritis Juvenile polyarthritis Ankylosing spondylitis Synovitis Low back pain Dysmenorrhoea Fever Migraine Soft tissue injuries Pain & Inflammation in dental and musculoskeletal origin.. Pharmacology: Ibuprofen has a high level of anti-inflammatory, anti-pyretic, and analgesic activity. The analgesic effects of Ibuprofen are due to both a peripheral and a central effect. Ibuprofen is a potent inhibitor of the enzyme cyclooxygenase, which thus results in a marked reduction in prostaglandin synt... Dosage: Adults: The dose is initially, 400 mg 3 times daily. A dose of 2400 mg daily should not be exceeded. Children: 3-6 months (body-weight over 5 kg): ½ tsp (2.5 ml) 3 times daily; max. 30 mg/kg daily in 3-4 divided doses. 6 months-1 year: ½ tsp (2.5 ml) 3-4 times daily; max. Side Effects: Upset stomach, vomiting, heartburn, nausea may occur.. Contraindications: Ibuprofen is contraindicated in patients who have shown the previous hypersensitivity to Ibuprofen, and in patients with severe or active peptic ulceration.. Pregnancy & Lactation: Adverse effects of Ibuprofen on the developing fetus cannot be fully excluded. Ibuprofen should not be used during pregnancy and for nursing mothers unless the potential benefits to the mothers outweigh the potential risks.. Storage: N/A.	1-0-0
Lamictal Lamotrigine Indication: Lamotrigine is indicated for: Epilepsy-combination therapy in patients aged 2 years and older: partial-onset seizures primary generalized tonic-clonic seizures generalized seizures of Lennox-Gastaut syndrome Epilepsy-monotherapy in patients aged 16 years and older. Adults with Bipolar Disorder..	0-0-1

Medicine Details	Dosage
Pharmacology: The exact mechanism of action of lamotrigine is not fully elucidated, as it may exert cellular activities that contribute to its efficacy in a range of conditions. Although chemically unrelated, lamotrigine actions resemble those of phenytoin and carbamazepine, inhibiting voltage-sensitive sodium... Dosage: Epilepsy- Table-1: Escalation Regimen for Lamotrigine in Patients Older than 12 Years with Epilepsy Weeks 1 & 2: In patients not taking Carbamazepine, Phenytoin, Phenobarbital, Primidone or Valproate: 25 mg every day In patients taking Valproate: 25 mg every other day In patients taking Carbamazepine... Side Effects: Common side effects of Lamotrigine include-dizziness, tremor, headache, rash, blurred vision, fever, lack of coordination, abdominal pain, infections, sleepiness, back pain, vomiting, diarrhea, tiredness, insomnia, dry mouth, stuffy nose, sore throat.. Contraindications: N/A. Pregnancy & Lactation: Pregnancy Category C. Lamotrigine is excreted in human milk. Because of the potential for serious adverse reactions in nursing infants from Lamotrigine, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.. Storage: N/A.	

* Auto-generated prescription (summary view). Full details stored in system.