

Doctor Information

Dr. Jeba Tasnia

Patient Section

Manual Entry Prescription

Rx

Medicine Details	Dosage
Seclo Omeprazole Indication: Omeprazole is indicated for the treatment of- Gastric and duodenal ulcer NSAID-associated duodenal and gastric ulcer As prophylaxis in patients with a history of NSAID-associated duodenal and gastric ulcer Gastro-esophageal reflux disease Long-term management of acid reflux disease ... Read more ... Pharmacology: Omeprazole, a substituted benzimidazole, is an inhibitor of gastric acid secretion. It inhibits gastric acid secretion by blocking hydrogen-potassium-adenosine triphosphatase (H+/K+ ATPase) enzyme system in the gastric parietal cell. After oral administration, the onset of the antisecretory effect... Dosage: Oral- Benign gastric and duodenal ulcer: 20 mg once daily for 4 weeks in duodenal ulceration, 8 weeks in gastric ulceration; in severe or recurrent cases, dose to be increased to 40 mg daily; maintenance dose for recurrent duodenal ulcer, 20 mg once daily; in prevention of relapse in duodenal ulcer... Side Effects: Omeprazole is generally well tolerated. Nausea, abdominal colic, paresthesia, dizziness and headache have been stated to be generally mild and transient and not requiring a reduction in dosage.. Contraindications: Omeprazole is contraindicated in patients with known hypersensitivity to any of the components of the formulation.. Pregnancy & Lactation: US FDA pregnancy category of Omeprazole is C. However, results from three prospective epidemiological studies indicate no adverse effects of Omeprazole on pregnancy or on the health of the fetus/newborn child. There is no information available on the passage of Omeprazole into breast milk or its ... Storage: N/A.	1-0-1
Ribovit Riboflavin Indication: To prevent and to treat deficiencies of vitamin B2 such as cheilosis, angular stomatitis, glossitis, keratitis, seborrhoeic dermatitis.. Pharmacology: Riboflavin or vitamin B2 is an easily absorbed, water-soluble micronutrient with a key role in maintaining human health. Like the other B vitamins, it supports energy production by aiding in the metabolising of fats, carbohydrates, and proteins. Vitamin B2 is also required for red blood cell formation... Dosage: For treating low levels of riboflavin (riboflavin deficiency) in adults: 5-30 mg of riboflavin daily in divided doses. For preventing migraine headaches: 400 mg of riboflavin (Vitamin B2) per day. It may take up to three months to get best results. For preventing cataracts: approximately 2.6 mg o... Side Effects: Riboflavin is likely safe for most people. In some people, Riboflavin can cause the urine to turn a yellow- orange color. When taken in high doses, Riboflavin might cause diarrhea, an increase in urine, and other side effects.. Contraindications: Hypersensitivity to Riboflavin.. Pregnancy & Lactation: Riboflavin is likely safe for pregnant or breastfeeding women when taken in the amounts recommended.. Storage: N/A.	1-1-1