1. The probability of a leap year selected at random contain 53								
Sunday is:								
	6 (b) 1/7							
2. A bag contai	ins 3 red and 2 blu	e marbles. A ı	marble is drawn at					
random. The pr	obability of drawir	ng a black ball	is:					
(a) 3/5 (b) 2/5 (d) 1/5								
3. The probabil	lity that it will rain	tomorrow is 0).85. What is the					
probability that	it will not rain tom	orrow						
(a) 0.25	(b) 0.145	3/20	(d) none of these					
4. What is the p	probability that a n	umber select	ed from the numbers					
(1, 2, 3,,1	(5) is a multiple of	4?						
1/5	(b) 4/5	(c) 2/15	(d) 1/3					
5. What are the	e total outcomes v	when we throw	three coins?					
(a) 4	(b) 5	8 ((d) 7					
• •	* *		ed at random from the					
numbers (1,2,3,	-							
		(c) 13/35	(d) none of these					
	the probability of a		* *					
(a) 2	(c) 0	(d) non	e of these.					
			se the correct answer					
for that which is	• .							
		7/5	(d) none of these.					
* *	* *	* *	an the probability of					
getting at least		•	,					
	(b) 3/8	1/2	(d) 1/8					
` ,	chosen at random		• •					
♦ ASSASSINAT	ΓΙΟΝ�. The proba	bility that the	letter chosen has:					
(6/13	(b) 7/13	(c) 1	(d) none of these.					
	(10)	(5)	(3) 112112 21 3122					
11. A dice is thr	rown. Find the prol	bability of get	ting an even number.					
(A) 2/3	(B) 1 (C							
(* ') =/ '	(-) . (-)	,, ,, ,,	., _					
12. Two coins are thrown at the same time. Find the probability of								
getting both he		(D) 0						
(A) 3/4 (1)	/4 (C) 1/2	(D) 0						
13. Two dice are thrown simultaneously. The probability of getting a								

1

sum of 9 is:

(A) 1/10	(B) 3/10	(1/9	(D) 4/9)
	ds are number rime number.	red from 1 to 10	00. Find the p	robability of
(A) 3/4		1/4	(D) 2	29/100
of drawing blue balls i	a blue ball is d	ouble that of a		the probability the number of
(A) 5	, ,	•	` ,	
	t random from	ntains 12 defections this box. Then		
(A) 143/150	0 (147	7/150 (C)	1/25	(D) 1/50
mixed thore the probabi	oughly. One ca ility that the nu	mbers 2 to 101 ard is drawn from mber on card is (C) 3/10	m this box ra s a perfect so	ndomly, then quare.
18. What is (A) 1/7	-	ey of getting 53	-	
probability	of getting a kir	a well shuffled ng of red suit. C) 7/52 (D)		ards. Find the
equally like	ly to come to r ,then the prob	nsists of spinni est pointing to ability that it wi	one of the nu	ımber odd number is:
21. A game its outcome result i.e. the probability	e consists of to e each time. A hree heads or t	ssing a one rup ryan wins if all t three tails and l lose the game	ee coin 3 tin he tosses givoses otherwi	nes and noting ve the same

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:								
•		(C) 1/365) 1/133225					
2. Then the pro	x is chosen at rapped to the company of the company	< 2 is?	umbers -2, -1, 0 , 1,					
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 8 (D) 7								
Then the proba	25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) 7/50 (B) 4/25 (C) 1/25 2/25							
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?								
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	$\frac{4}{21}$					
			nes are 50, 70, 82,					
	e standard devia b) 25.49	c) 25.29	d) 25.69					
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.								
	13, 18		d) 13, 16					
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is								
$\frac{1}{2}$ 30. X is a varia	b) $^{1}/_{3}$	c) ¹ / ₄ d 3. The value of (c) 27	$\begin{array}{c} & \text{d) } ^1/_6 \\ \textbf{E(X^2) is } \underline{\qquad} . \\ \textbf{9} \end{array}$					
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?								

a) 3	b) 4	c) 5	7					
32.Out of the following values, which one is not possible in probability? a) $P(x) = 1$ b) $\sum x P(x) = 3$ c) $P(x) = 0.5$ d) $P(x) = -0.5$								
33.If E(x) = a) 2	2 and E(z) = 4, b) 6	then E(z - x) =? c) 0	Insufficient data					
34.The cova	ariance of two i	ndependent rand	om variable is					
1	b) 0	c) - 1	d) Undefined					
35.If Σ P(x) a) 0	= k ² – 8 then, 1	the value of k is?	d) Insufficient data	ì				
• •	0.5 and x = 4, t b) 0.5	hen E(x) = ?	d) 2					
37.In a discrete probability distribution, the sum of all probabilities is always?								
a) 0 b) Infinite c) 1 Undefined 38.If the probability of hitting the target is 0.4, find mean and variance.								
a) 0.4, 0.24	b) 0.6, 0	.24 —0	.4, 0.16 d) 0.6, 0.16)				
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6								
40. Find the mean of tossing 8 coins. a) 2								

a) Mean is (c) Mean is (,						
42.Variance of a random variable X is given by E (X) b) E(X2) c) E(X2) - (E(X))2									
43.Mean of a random variable X is given by a) $E(X)$ b) $E(X2)$ $E(X2) - (E(X))2$ d) $E(X)$									
44.Mean of	44.Mean of a constant 'a' is 0 b) a c) a/2 d) 1								
45.Variance of a constant 'a' is . a) 0									
46.Find the mean and variance of X?									
Х	0	1	2	3	4				
f(x)	1/9	2/9	3/9	2/9	1/9				
2, 4/3 47.Find the	2, 4/3 b) 3, 4/3 c) 2, 2/3 d) 3, 2/3 47.Find the expectation of a random variable X?								

	Х	0	1	2	3		
	f(x)	1/6	2/6	2/6	1/6		
a) ().5		1.5			c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by



c) np2q

d) npq2

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**

a)
$$P(X = x) = nCx px qx$$

$$P(X = x) = nCx px q(n-x)$$

c)
$$P(X = x) = xCn qx p(n-x)$$

d)
$$P(x = x) = xCn pn qx$$

50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2

