

SECTION - A

Q.1] (A) 60° ✓

Q.2] (c) a segment ✓

Q.3] (c) $\frac{132}{7} \text{ cm}^2$ ✓

Q.4] (B) -2, 2 ✓

Q.5] (c) 2.3cm ✓

Q.6] (D) $\frac{1}{6}$ ✓

Q.7] (B) 2 ✓

(A) One point only ✓

(B) 5 units ✓

) (D) ✓

) (C) $x(x+1) + 8 = (x+2)(x-2)$ ✓

) (D) more than 3 ✓

) (C) 424.5 ✓

) (A) 8 ✓

) (D) 20° ✓

) (B) $\frac{5}{3}$ ✓

3.17] (A) (B) (-3, 0)



3.18] (A) $\frac{23}{3}$



3.19] (D) Assertion (A) is false, but Reason (R) is true



3.20] (A) Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of Assertion (A).



SECTION - B

Q.2]

$$\text{Given, } 5 \sin^2 60^\circ + 3 \cos^2 30^\circ - \sec^2 45^\circ$$

Substituting with all the appropriate well values.]

$$* \sin 60^\circ = \sqrt{3}/2 ; \cos 30^\circ = \sqrt{3}/2 \text{ and } \sec 45^\circ = \sqrt{2}$$

$$\rightarrow 5 \left(\frac{\sqrt{3}}{2}\right)^2 + 3 \left(\frac{\sqrt{3}}{2}\right)^2 - (\sqrt{2})^2$$

$$\rightarrow 5 \left(\frac{3}{4}\right) + 3 \left(\frac{3}{4}\right) - 2$$

$$\rightarrow \frac{15}{4} + \cancel{\frac{9}{4}} - 2$$

$$\rightarrow \frac{24}{4} - 2$$

$$\rightarrow 6 - 2$$

$$\rightarrow \underline{\underline{4}}$$

∴ After evaluating " $5 \sin^2 60^\circ + 3 \cos^2 30^\circ - \sec^2 45^\circ$ ", we obtain ' 4 '.

(a) Given polynomial, $8x^2 + 14x + 3$

[It is written in the form $ax^2 + bx + c = 0$]

$$\text{So, } a = 8; b = 14 \text{ and } c = 3$$

Now, we know that

$$\rightarrow \alpha + \beta = -\frac{b}{a} = -\frac{14}{8} = -\frac{14}{8} = -\frac{7}{4} \quad \text{--- eq(1)}$$

$$\rightarrow \alpha \cdot \beta = \frac{c}{a} = \frac{3}{8} \quad \text{--- eq(2)}$$

We need to find the value of $\frac{1}{\alpha} + \frac{1}{\beta}$

$\rightarrow \frac{1}{\alpha} + \frac{1}{\beta}$ can be rewritten as given below through cross multiplication

$$\rightarrow \frac{\beta + \alpha}{\alpha \beta} \quad \text{--- eq(3)}$$

P.T.O

[Poor quality of paper]

P.T.O

Now, by substituting the value of ' $\alpha + \beta$ ' and ' $\alpha \times \beta$ ' from eq① and ② respectively in eq③

We obtain,

$$\rightarrow \frac{-7}{4}$$

$$\frac{3}{8}$$


$$\rightarrow \frac{-7 \times 8}{4 \times 3}$$

$$\rightarrow \frac{-7 \times 2}{3}$$

$$\rightarrow \frac{-14}{3}$$

$\therefore 1/\alpha + 1/\beta$ is equal to $-14/3$.

Q.23]



Given, $\triangle OPG$ and $\triangle OSR$ with $PQ \parallel RS$

To prove, $OP \times OR = OG \times OS$

Proof, As $PQ \parallel RS$

So, $\angle OPG = \angle OSR$ [Alternate interior angles]

$\angle OGP = \angle ORS$ [Alternate interior angles]

And, $\angle POG = \angle ROS$ [Vertically opposite angles]

Now, in $\triangle OPQ$ and $\triangle OSR$

$$\angle OPG = \angle OSR$$

$$\angle OGP = \angle ORS$$

$$\angle POQ = \angle ROS$$

[Reason written above before]

By, AAA similarity criterion $\triangle OPQ \sim \triangle OSR$

$$\text{So, } \frac{OP}{OS} = \frac{PQ}{SR} = \frac{OQ}{OR} \quad [\text{By CPCT}]$$

$$\rightarrow \frac{OP}{OS} = \frac{OQ}{\cancel{OR}}$$

$$\rightarrow OP \times OR = OQ \times OS$$

Hence Proved

Q.2u]

Given, HCF (306, 1314) = 18

To find, Lcm of (306, 1314)

As we know,

$$\text{product of two no.} = \text{HCF} \times \text{Lcm} \quad [\text{Respective no. must be same}]$$

$$\rightarrow 306 \times 1314 = 18 \times \text{Lcm}$$


$$\rightarrow \text{Lcm} = \frac{306 \times 1314}{18}$$

$$\rightarrow \text{Lcm} = 17 \times 1314$$

$$\rightarrow \text{Lcm} = \underline{\underline{22338}}$$

\therefore Lcm of (306, 1314) is equal to 22338.

Q.26]



Given, a circle with centre O and diameter AB.

Two tangents XY and PQ at ends A and B respectively.

To prove, XY || PQ

Proof, As XY and PQ are tangents to the circle.

So,

$OA \perp XY$ [Radius from the centre of a circle is
 $OB \perp PQ$ [perpendicular to tangent at point of contact]

∴ So, $\angle OAX = \angle OAY = 90^\circ$ [Perpendicular] — eq ①
 $\angle OBP = \angle OBQ = 90^\circ$ — eq ② [Perpendicular]

By comparing eq ① and ②, we get

$$\rightarrow \angle OAX = \angle OBQ — \text{pair 1}$$

$$\rightarrow \angle OBP = \angle OAY — \text{pair 2}$$

As pair 1 and pair 2 of angles are equal
 indicating alternate ~~interior~~ angles are equal, thus
 $\underline{XY \parallel PQ}$.

Hence Proved

SECTION-C

Q.2c)

Given, $\sqrt{3}$ is an irrational number

To prove, $2+5\sqrt{3}$ is an irrational number.

Proof, Let $2+5\sqrt{3}$ be a rational number

(Then, we can write $2+5\sqrt{3}$ in form of fraction)

$$\rightarrow 2+5\sqrt{3} = \frac{a}{b} \quad \left[\begin{array}{l} \text{Here } a \text{ and } b \text{ are co-prime integers} \\ \text{and } b \neq 0. \end{array} \right]$$

• Subtracting 2 from both the sides [L.H.S and R.H.S].

$$\rightarrow 5\sqrt{3} = \frac{a-2}{b}$$

$$\rightarrow \sqrt{3} = \frac{a-2b}{b}$$

• Dividing both sides by '5'. We get

$$\rightarrow \sqrt{3} = \frac{a-2b}{5b}$$

* As a and b are integers, thus $(a-2b)/5b$ is also a rational no. [integer] but as given, $\sqrt{3}$ is an irrational no. (in question)

∴ This contradicts with the fact.

\rightarrow Irrational \neq Rational

∴ Our Supposition was wrong, $2+5\sqrt{3}$ is an irrational no.

Hence Proved

Q.27] (a) Given equation,

$$2x^2 + 2x + 9 = 0$$

[It is written in the form $ax^2 + bx + c = 0$]

So, $a = 2$; $b = 2$ and $c = 9$

Now using quadratic formula [to find real roots if exist]

$$\frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \rightarrow \text{eq. ①}$$

$$\rightarrow b^2 - 4ac$$

[Substituting all the appropriate values] we get,

$$\rightarrow 4 - (4)(2)(9)$$

$$\rightarrow 4 - 72$$

$$\rightarrow -68 < 0 \text{ (zero)}$$

As the value of discriminant is negative [less than 0].

\therefore The Real roots doesn't exist for the given equation.

Q.28]

To prove,

$$\frac{1 + \sec \theta}{\sec \theta} = \frac{\sin^2 \theta}{1 - \cos \theta}$$

$$\text{Proof, L.H.S.} = \frac{1 + \sec \theta}{\sec \theta}$$

$$= 1 + \frac{1}{\cos \theta}$$

$$\frac{1}{\cos \theta}$$

$$= \frac{\cos\theta + 1}{\cos\theta}$$

$$= \frac{1}{\cos\theta}$$

$$= \frac{(\cos\theta + 1)(\cos\theta)}{(\cos\theta)}$$

$$= \underline{\cos\theta + 1}$$

$$\text{Now, R.H.S.} = \frac{\sin^2\theta}{1 - \cos\theta}$$

$$= \frac{1 - \cos^2\theta}{1 - \cos\theta}$$

$$= \frac{(-\cos\theta)(1 + \cos\theta)}{(1 - \cos\theta)} \quad \left[\begin{array}{l} \text{Using identity: } a^2 - b^2 \\ = (a+b)(a-b) \end{array} \right]$$

$$= 1 + \cos\theta$$

$$\therefore \text{L.H.S.} = \text{R.H.S.} = 1 + \cos\theta$$

Hence Proved

Q.29]

Given, Deck of 52 playing set cards.

From that deck black queens and red kings are removed.

[No. of black queens = 2 ; No. of red kings = 2]

$$\begin{aligned}\text{Total no. of cards left now} &= 52 - (2+2) \\ &= 52 - 4 \\ &= 48\end{aligned}$$

i] Probability of an ace :-

Total no. of ace in a deck = 4

Total no. of cards left = 48

$$P(\text{Selected card is an ace}) = \frac{\text{Total no. of favourable outcome}}{\text{Total no. of outcome}}$$

$$\rightarrow \frac{\text{Total no. of ace cards}}{\text{Total no. of cards}} = \frac{4}{48} = \frac{1}{12} = 0.08\bar{3}$$

∴ Probability for the selected card to be an ace is
 '1/12' OR '0.08\bar{3}'.

ii) Probability of a 'jack of red colour'.

$$P(\text{Selected card is a 'jack of red colour'}) = \frac{\text{Total no. of red jack cards}}{\text{Total no. of cards}}$$

$$\text{Total no. of jack of red colour} = 2$$

$$\text{So, } \rightarrow \frac{2}{48} = \frac{1}{24} = 0.041\bar{6}$$

∴ The probability of a selected card to be a 'jack of red colour' is ' $\frac{1}{24}$ ' OR ' $0.041\bar{6}$ '.

(iii) a king of spade-


Total no. of king of spade = 1


Total no. of cards = 48

$$P(\text{a king of spade}) = \frac{\text{Total no. of king of spade}}{\text{Total no. of cards}}$$


$$= \frac{1}{48} = 0.0208\bar{3}$$

∴ Probability of a king of spade card to be selected is ' $0.0208\bar{3}$ ' OR ' $\frac{1}{48}$ '.





Q.30]

(90° at centre) ~~so~~

Given, a circle with centre O and radius 10cm, Subtends ^
So, $\theta = 90^\circ$

To find, Area of minor sector and area of major sector

$$\text{Area of minor sector} = \frac{\theta}{360^\circ} \times \pi r^2$$

$$= \frac{90^\circ}{360^\circ} \times 3.14 \times (10)^2$$

$$= \frac{1}{4} \times 3.14 \times 100$$

$$= \frac{1}{4} \times 314$$

$$= \left(\frac{314}{4} \right) \text{ cm}^2$$

$$= \underline{\underline{78.5}} \text{ cm}^2$$

\therefore The area of minor sector is '78.5 cm²'.

Now, area of major sector $= \frac{\theta}{360} \times \pi r^2$

$$= \frac{(360^\circ - \theta)}{360^\circ} \times \pi r^2$$

$$= \frac{(360^\circ - 90^\circ)}{360^\circ} \times 3.14 \times \cancel{(10)^2}$$


$$= \frac{270^\circ}{360^\circ} \times 3.14 \times 100$$

$$= \frac{3}{4} \times 314$$

$$= 3 \times 78.5 \longrightarrow \underline{\underline{235.5}} \text{ cm}^2$$

∴ The area of major sector is '235.5cm²'.

Q.31] (b)



Given, a circle with centre O and $\angle ROP = 40^\circ$ with PR as a tangent.

To find, $\angle POQ = ?$

As, PR is a tangent to circle.

So, $OP \perp PR$ [The radius of circle is perpendicular to the tangent at the point of contact]

So, $\angle OPR = 90^\circ$

Now, $\angle OPR = \angle OPQ + \angle QPR = 90^\circ$

$$\rightarrow \angle OPQ + \angle QPR = 90^\circ$$

$$\rightarrow \angle OPQ + 40^\circ = 90^\circ \quad [\angle QPR = 40^\circ \text{ is given}]$$

$$\rightarrow \angle OPQ = 50^\circ$$

Also, OP and OQ are radius of a circle, so
 $OP = OQ$

then in $\triangle OPQ$

$\angle OPQ = \angle OQP$ [Angles opposite to the equal side
 are equal]

$$\text{So, } \angle OPQ = \angle OQP = 50^\circ - \text{eq } \textcircled{1}$$

Now, using angle sum property of triangle:

$$\rightarrow \angle OPQ + \angle OQP + \angle POQ = 180^\circ$$


$$\rightarrow 50^\circ + 50^\circ + \angle POQ = 180^\circ \quad [\text{from eq. } \textcircled{1}]$$

$$\rightarrow \angle POG = 100^\circ - 80^\circ$$

\therefore The measure of $\angle POG$ is 80° .

SECTION-D

Q.32]



Given, $\triangle ABC$ and $\triangle PQR$ with median BD and GM respectively.

Also, $\triangle ABC \sim \triangle PQR$

To prove, $\frac{AB}{PQ} = \frac{BD}{GM}$

Proof, As BD and QM are the medians, they will divide the AC and PR respectively in equal manner. So, $AP = PC$ AND $PR = QR$, $PM = MR$

As, $\triangle ABC \sim \triangle PQR$

$$\text{So, } \frac{AB}{PQ} = \frac{BC}{QR} = \frac{AC}{PR}$$

And, $\angle A = \angle P$; $\angle B = \angle Q$ and $\angle C = \angle R$

$$\rightarrow \angle ABC = \angle PQR; \angle BCA = \angle QRP \text{ and } \angle BAC = \angle PQR$$

$$\text{Now, } \frac{AB}{PQ} = \frac{AC}{PR}$$

$$\rightarrow \frac{AB}{PQ} = \frac{\frac{1}{2}AC}{\frac{1}{2}PR}$$

$$\rightarrow \frac{AB}{PQ} = \frac{AD}{PM} \quad [\text{As } BD \text{ and } QM \text{ are medians}]$$

Now in $\triangle ABD$ and $\triangle PQM$

$$\rightarrow \frac{AB}{PQ} = \frac{AD}{PM}$$

$$\rightarrow \angle BAC = \angle QPR \rightarrow \angle BAD = \angle QPM$$

So, by SAS similarity criterion, $\triangle ABD \sim \triangle PQM$

Now, $\frac{AB}{PQ} = \frac{BD}{QM}$ [By CPCT]

Hence Proved.

Q.33]

Given, two cubes with each volume 125 cm^3 and they are later joined end to end.

To find, Volume and surface area of newly formed cuboid.

Now, Volume of cube = $a^3 = 125 \text{ cm}^3$ [given]

$$\text{So, } a^3 = 125 \text{ cm}^3$$

$$\rightarrow a = \sqrt[3]{125}$$

$$\rightarrow a = (125)^{1/3}$$

$$\rightarrow a = [(\pm 5)^3]^{1/3}$$

$$\rightarrow a = 5 \text{ cm}$$

\therefore The dimension of each of the ^{side of} cube is 5cm.

Now, the surface area of resulting cuboid:-

Dimensions of the cuboid:-

Length of cuboid :- $5\text{cm} + 5\text{cm} = 10\text{cm}$ [as two cubes are joined]

Breadth of cuboid :- Remains same = 5cm

Height of cuboid :- Remains same = 5cm

Now, using

$$\text{T.S.A of cuboid} = 2(lb + bh + lh)$$

[Substituting all appropriate values]

$$\rightarrow 2[(10 \times 5) + (5 \times 5) + (10 \times 5)] \text{ cm}^2$$

$$\rightarrow 2[50 + 25 + 50] \text{ cm}^2$$

$$\rightarrow 2[125] \text{ cm}^2$$

$$\rightarrow \underline{\underline{250 \text{ cm}^2}}$$

∴ The surface area of resulting cuboid is 250 cm².

Now, Volume of cuboid = $l \times b \times h$

[Substituting with all appropriate values]

$$\rightarrow (10 \times 5 \times 5) \text{ cm}^3$$

$$\rightarrow \underline{\underline{250 \text{ cm}^3}}$$

∴ The volume of resulting cuboid is also 250 cm³.

Q.34]

Given, $S_7 = 91$ and $S_{17} = 561$

To find, sum of first n terms $\rightarrow S_n$
 n^{th} term $\rightarrow a_n$

$$S_7, \quad S_7 = 91$$

Now using formula

$$S_n = \frac{n}{2} [2a + (n-1)d]$$

$$\rightarrow S_7 = 91 = \frac{7}{2} [2a + (7-1)d]$$

$$= \frac{7}{2} [2a + 6d]$$

$$\rightarrow 91 = 7 [a + 3d]$$

$$\rightarrow 13 = a + 3d \quad \text{--- eq(1)}$$

$$\text{Again, } S_{17} = 561 = \frac{17}{2} [2a + (17-1)d]$$

$$= \frac{17}{2} [2a + 16d]$$

$$\rightarrow 561 = 17 [a + 8d]$$

$$\rightarrow 33 = a + 8d \quad \text{--- eq (2)}$$

Now, by subtracting eq (1) from eq (2),

$$\rightarrow a + 8d = 33$$

$$- a + 3d = 13$$

$$5d = 20$$

$$\boxed{d = 4} \quad \text{--- eq (3)}$$

Substituting the value of 'd' or common diff. in eq (1)

$$\rightarrow a + 8d = 33$$

$$\rightarrow a + 3(a) = 13$$

$$\rightarrow \boxed{a = 1} \quad \text{--- eq (4)} \quad a \rightarrow \text{first term of A.P.}$$

Now, sum of first n terms:-

$$\rightarrow S_n = \frac{n}{2} [2a + (n-1)d]$$

[Substituting the values of a and d from eq ③ and ④ respectively]

$$\rightarrow S_n = \frac{n}{2} [2(1) + (n-1)(4)]$$

$$= \frac{n}{2} [2 + (n-1)(4)]$$

$$= \frac{n}{2} \times 2 [1 + (n-1)(2)]$$

$$= n [1 + 2n - 2]$$

$$= n [2n - 1]$$

$$= \underline{\underline{2n^2 - n}}$$

of an AP


\therefore The sum of 'n' terms ^ is ' $2n^2 - n$ '.

Now, n^{th} term:

$$\begin{aligned} a_n &= a + (n-1)d \\ &= 1 + (n-1)4 \\ &= 1 + 4n - 4 \\ &= \underline{\underline{4n - 3}} \end{aligned}$$

\therefore n^{th} term of an A.P. is ' $4n - 3$ '.

Q. 357



In the given figure,

$DB \rightarrow$ pedestal

$AD \rightarrow$ Statue

$C \rightarrow$ point on the ground from where it is observed

$AB \rightarrow 1.6m + DB$

Given, height of a statue = $1.6m = AD$

angle of elevation to top of statue :- 60°

angle of elevation to top of pedestal :- 45°

Need to find, height of pedestal = $DB = ?$

$$\text{Now, } DB = h \quad [\text{let}] - \text{eq } ①$$

$$CB = x \quad [\text{let}] - \text{eq } ②$$

Now, in $\triangle DBC$

$$\tan 45^\circ = \frac{DB}{BC}$$

$$\rightarrow l = \frac{h}{x} \quad [\text{from eq } ① \text{ and } ②]$$

$$\rightarrow \underline{xc = h} \quad - \text{eq } ③$$

✓

Now, in $\triangle ABC$

$$\tan 60^\circ = \frac{AB}{BC}$$

$$\frac{\sqrt{3}}{1} = \frac{1.6m + h}{h} \quad [\text{from eq } ③ \quad BC = x = h]$$

$$\rightarrow \sqrt{3}h = 1.6m + h$$

$$\rightarrow \sqrt{3}h - h = 1.6m$$

$$\rightarrow h(\sqrt{3}-1) = 1.6m$$

$$\rightarrow h = 1.6m / (\sqrt{3}-1)$$

$$\rightarrow h = \frac{1.6}{(\sqrt{3}-1)}$$

$\rightarrow h = [$ Now rationalizing the denominator]

So, rationalizing factor = $\sqrt{3}+1$

$$\rightarrow h = \frac{(1.6)(\sqrt{3}+1)}{\cancel{(\sqrt{3}-1)}(\sqrt{3}+1)}$$

$$\rightarrow h = \frac{(1.6)(\sqrt{3}+1)}{3-1}$$

$$= \frac{1.6(\sqrt{3}+1)}{2}$$

$$h = 0.8(\sqrt{3}+1) \text{ m}$$

$$h = 0.8(1.732+1) \text{ m} \quad [\text{Given, } \sqrt{3}=1.732]$$

$$= (0.8)(2.732) \text{ m}$$

$$= 2.1856 \text{ m}$$

∴ Height of the pedestal [DB: h₀] is 2.1856 m.

SECTION-B

Q.36] (i) Given, price of notebook :- ₹x
price of pen :- ₹y

Equation → $3x + 2y = 80$ — eq①
 ~~$4x + 3y = 110$~~ — eq②

(ii) Multiplying eq① by 3 and eq② by 2, we get

$$\begin{array}{rcl} \rightarrow & 9x + 6y = 240 & \left[\text{Subtracting eq① from ②, using}\right. \\ & - 8x + 6y = 220 & \left. \text{elimination method}\right] \\ & \hline & x = 20 \end{array}$$

∴ Price of one notebook is "₹20".

(iii) Substituting value of x in eq① → $3(20) + 2y = 80$

$$\rightarrow 2y = 20 \rightarrow y = ₹10$$

∴ The value of pen is ₹10.

$$\text{Now, value of 6 notebooks + 3 pens} = 6(20) + 3(10) = \underline{\underline{₹150}}$$

Q.37] (i) Model class of the data :- 10-15 [as it has the highest frequency]

So, upper limit of model class = 15

(ii) Total frequency = $n = 80$
 And, $n/2 = 80/2 = 40$

Class interval	frequency	Cumulative frequency
0-5	13	13
5-10	16	29
10-15	22	51
15-20	18	69
20-25	11	80

(iii) (a) mode of NAV of mutual funds.

model class :- 10-15

l , lower limit :- 10

h , class size is ~~5~~ 5

f_0 :- 22

f_1 :- 16

f_2 :- 18

Now using,

$$\begin{aligned}
 \text{med mode} &= l + \left(\frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h \\
 &= 10 + \left(\frac{22 - 16}{44 - 16 - 18} \right) \times 5 \\
 &= 10 + \left(\frac{6}{10} \right) \times 5 \\
 &= 13
 \end{aligned}$$

∴ mode NAV of mutual funds is 13.

Q.38) (i) Position of pole C :- (5, 4)

(ii) Distance of pole B $(6, 6)$ from corner of park O $(0, 0)$.
 Using distance formula -

$$\rightarrow \sqrt{(0-6)^2 + (0-6)^2}$$

$$\rightarrow \sqrt{72}$$

$$\rightarrow 6\sqrt{2} \text{ units [Distance]}$$

(iii) (b) Distance between poles A (2, 7) and C (5, 4).

Using distance formula $\rightarrow \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

$$\rightarrow \sqrt{(5-2)^2 + (4-7)^2}$$

$$\rightarrow \sqrt{9+9}$$

$$\rightarrow \sqrt{18}$$

$$\rightarrow 3\sqrt{2} \text{ units}$$

\therefore distance between poles A and C is $3\sqrt{2}$ units.

RIGHT WORK

$$\frac{1 + \cos\theta}{1 - \cos\theta} \rightarrow \frac{1 + \cos\theta}{\cos\theta}$$

$$\frac{1 + \cos\theta + 1}{\cos\theta}$$

~~$\frac{11}{25} \times 5$~~

$10 + \frac{5}{2}$

12.5

$$\begin{array}{r} + 0.8 \\ \hline 2.1, 856 \end{array}$$

$$\begin{array}{l} 1 + \sec \theta \cos \theta \\ = 1 + \frac{1}{\cos \theta} \times \cos \theta \\ = 1 + \cos \theta \end{array}$$

$$3m = 7 + 1c$$

$$m = \frac{23}{3}$$

$$S \sqrt{3}/4$$

$$\begin{array}{r} \times 17 \\ \hline 233 \\ + 18 \\ \hline 412 \end{array}$$

$$25 - 1c \rightarrow 9$$

$$(3)^2 - 5(3) + 4$$

$$9 - 15 + 4$$

$$\rightarrow -2$$

$$x(x+1) + 8 = (x+2)(x-2)$$

$$\rightarrow x^2 + x + 8 = x^2 - 4$$

$$a_{23} - a_{19} = 32 \rightarrow a + 22d - (a + 18d)$$

$$a + 22d - a - 18d \rightarrow 4d = 32 \rightarrow d = 8$$

$$x = \frac{-1}{a} + \frac{3}{2} = 1 - \cos^2 \theta$$

$$(1 + \cos \theta) (1 - \cos \theta)$$


$$-6/2 -3,0$$

$$1 + \frac{1}{\cos \theta} \rightarrow \frac{\cos \theta + 1}{\cos^2 \theta}$$

$$\frac{1}{\cos \theta}$$

$$\frac{5}{7} \neq \frac{2}{5} = \frac{6}{8}$$

$$1 + \cos \theta$$



Best

$$\begin{array}{r} 78.5 \\ \times 3 \\ \hline 235.5 \end{array}$$

$$\begin{array}{r} 2.732 \\ + 0.8 \\ \hline 21856 \end{array}$$


$$\frac{1}{7} \times 22 \times 36^{\circ}$$

$$\sqrt{3}/4 \sqrt{1}/4 \sqrt{2}/4 \sqrt{3}/4 \sqrt{1}/4$$

$$\begin{array}{ccccccc} & & & 0 & 30^{\circ} & 45^{\circ} & 60^{\circ} & 90^{\circ} \\ & & & \times & \frac{5}{27} & \times & \frac{8}{216} & \times \\ & & & & \sqrt{3}/4 & & 1/1/\sqrt{2} & \end{array}$$

$$1080 = 2^x \times 3^y \times 5^z$$

$$216 = 2^x \times 3^y$$



$$(2 \times 3)^3$$

$$27 \times 3$$

$$8 \times 3$$

$$\sqrt{3}/4$$

$$\frac{5}{27} \times \frac{8}{216}$$

$$\sqrt{3}/4$$

$$\sqrt{1}/4$$

$$\sqrt{2}/4$$

$$\sqrt{3}/4$$

$$\sqrt{1}/4$$

$$\frac{5 \pm 3}{2}, 1, 4$$

$$(2, -1), (-1, 5) \rightarrow \sqrt{(-1-2)^2 + (-5+1)^2}$$

$$\rightarrow \sqrt{9+16} = \sqrt{25} = 5$$

$$\times 306$$

$$171884$$

$$394200$$

$$402084$$

$$\begin{array}{r} 8x^2 + 12x + 2x + 3 \\ 4x(2x+3) + 1 \\ (4x+1)(2x+3) \end{array}$$

$$x = \frac{-1}{a} + \frac{3}{2} = 1 - \cos^2 \theta$$

$$1 + \cos \theta (1 - \cos \theta)$$

$$1 + \cos \theta$$

$$1 + \cos \theta$$