

Abstract

In this pieces of notes, we will go through the concepts related to functions and the meaning of graphs of the functions.

Contents

1	Function and its graph	5
1.1	What is a Function?	5
1.1.1	Function as an input-output pair	6
1.1.2	Defining a function with variables	8
1.1.3	Domain, Co-domain and Range	9
1.2	Graph of a function	13
1.2.1	Blobs-and-arrows diagram for discrete functions	13
1.2.2	The pairing table and xy-coordination	15
1.3	Fundamental properties of a graph	17
1.3.1	points lying on the graph	17
1.3.2	Special intersections: x-intercept and x-intercepts	17
1.3.3	general intersection	17
1.4	Solving equations using graphs of functions	17
1.4.1	Homogeneous equations	17
1.4.2	Non-homogeneous equations	17
1.4.3	Simultaneous equations	17
1.5	Transformation of functions	17
1.5.1	Translation	17
1.5.2	Dilation	17
1.6	Challenging questions	17
2	Linear functions	18
2.1	Fundamental concepts of points	18
2.1.1	Distance between two points	18
2.1.2	Mid-point and Division points	18
2.2	Different forms of a linear function	18
2.2.1	Slope of a line	18
2.2.2	Two-point form	18
2.2.3	Point-slope form	18

2.2.4	Slope-intercept form	18
2.2.5	General form	18
2.2.6	Special: Intercept-form	18
2.3	Parallel lines and Perpendicular lines	18
2.3.1	Point of intersection	18
2.3.2	Parallel lines	18
2.3.3	Perpendicular lines	18
2.3.4	Number of intersections	18
2.4	Angle of elevation and depression	18
2.5	Additional content: Point-line distance	18
2.6	Linear inequalities	19
2.6.1	One variable inequality	19
2.6.2	Two variable inequality	19
2.6.3	Linear programming	19
2.7	Challenging questions	19
3	Quadratic functions	20
3.1	Solving Quadratic Equations	20
3.1.1	The square-root method	20
3.1.2	General form and Completing the square method	20
3.1.3	The quadratic formula	20
3.2	Solvability of Quadratic Equations	20
3.2.1	Discriminant: the factor affecting solvability	20
3.2.2	Application of solvability	20
3.3	Relation between coefficients and roots	20
3.3.1	What does it mean by a root?	20
3.3.2	The Vieta formula	20
3.3.3	Forming quadratic equations with given roots	20
3.4	Vertex form of a quadratic function	20
3.4.1	Obtaining vertex form using Transformation	20
3.4.2	Relation between vertex form and general form	20
3.4.3	Opening direction	20
3.4.4	The axis of symmetry	20
3.4.5	The extremum of quadratic functions	20
3.5	Quadratic inequalities	20

3.5.1	Boolean algebra	20
3.5.2	Solving quadratic inequalities	20
3.6	Challenging questions	20
4	Polynomial functions	21
4.1	Arithmetic rules for polynomials	21
4.1.1	Addition, Subtraction and Multiplication	21
4.1.2	Division	21
4.2	Divisibility of polynomials	21
4.2.1	Division algorithm	21
4.2.2	Remainder Theorem	21
4.2.3	Factor Theorem and General Vieta formula	21
4.3	G.C.D. and L.C.M.	21
4.3.1	Greatest common divisor	21
4.3.2	Least common multiple	21
4.4	Positional notation	21
4.4.1	Binary, Octal, Decimal and Hexadecimal	21
4.4.2	Conversion between different bases	21
4.5	Challenging questions	21
5	Exponential and Logarithmic functions	22
5.1	Exponential functions	22
5.1.1	Law of Exponential algebra	22
5.1.2	Seeing surds as fractional exponents	22
5.1.3	Solving equations using exponentiation method	22
5.2	Logarithmic functions	22
5.2.1	What is Logarithm?	22
5.2.2	Law of Logarithmic algebra	22
5.2.3	Solving equations using Logarithmic method	22
5.3	Challenging questions	22
6	Sequence as a function of natural numbers	23
6.1	What is a sequence?	23
6.2	Arithmetic sequence	23
6.2.1	General form	23

6.2.2	Summation of arithmetic sequence	23
6.3	Geometric sequence	23
6.3.1	General form	23
6.3.2	Summation on Geometric sequence	23
6.3.3	Infinite sum of Geometric sequence	23
6.4	Additional content: Arithmetic-Geometric sequence	23
6.5	Challenging questions	23

1 Function and its graph

‘Functions describes the world!’, one Professor in Mathematics of Massachusetts Institute of Technology (a.k.a. MIT) said that. His speech was greatly influential, as I have never heard such conclusive thinking about functions. In fact, in the past few years, whenever I was studying in schools, my thought about functions is always only about projecting elements from one set to another set, but what he said had a big impact to my knowledge about functions.

What function talks about, is a subjection of one elements to one another. It can be thought of as a pointing action started from element A to element B , which not so far away, if we could think of subjecting a lot of, a bunch of, or a list of, whatever, objects from one collection to another collection, and they can all be matched under this pointing action, then it is a so-called function.

For example, in an Indian factory, the production of food undergoes many different process. Those can all be called functions. Let say a raw material comes to the factory first, it then be put to a machine to chop into many small pieces. It is the chopping function inside the factory. Next, the chopped material will be put into a pool of yellowish-brownish liquid and be stirred by dirty hands. It is the Mixing function in the factory. After that, the liquid will be drained on the dirty floor and be stepped on by Indian workers so that they can be smell freshed. It is the flavouring function in the factory. Finally, it will be sold to stores, which is the selling function.

Another example is what our body does. We eat and drink, going down the digestive system, and we sit on a toilet. Although we stupid human knows nothing about how the digestive system works, we could still name the conversion from food to poops a digestion, which means the digestive function representing the process in our body.

So we know that function as an english word represents the naming of a process of conversion, it is the time to explore how Math functions works.

1.1 What is a Function?

A function is defined as follow:

Definition 1.1 (Function). *Given an input x and an output y , a **function** is a relation between x and y so that we can write $y = f(x)$ to represent the relationship.*

Essential Practice 1.1.0.1. *Write down functions for the following input-output variables:*

1. u as input and v as output;
2. b as input and a as output;

3. n as input and 1 as output (which we call it a constant function);
4. x^2 as input and y as output;
5. xy as input and z as output;
6. 2^x as input and k as output;
7. \sqrt{p} as input and q as output;

Remark. It is notable that we may write functions as $y = g(x)$, $y = h(x)$, $y = d(x)$, ... as we want. The ‘naming’ of a function is always definitive and up to user’s construction.

We can also apply functions after functions. To do so, we have to talk about the following:

Definition 1.2 (Composite functions). Let f and g be functions such that f takes x as input and y as output, and g takes y as input and z as output. Then we can say there is a function h that takes x as input and z as output. In other words, h is a **composite function** such that $z = h(x) = g(f(x))$.

Essential Practice 1.1.0.2. Let f and g be functions such that $b = f(a)$, $c = g(b)$. Write a function for a as input and c as output.

1.1.1 Function as an input-output pair

We may now consider how functions carry things to things using arrow notations. We may use a stroked arrow \mapsto to emphasis the carrying process. Let’s take a look at the following examples.

Example. Given a function that undergoes the process of adding one to the given element. We can say

$$\dots, 1 \mapsto 2, 2 \mapsto 3, 3 \mapsto 4, \dots$$

In other words, we may write

$$x \mapsto x + 1$$

to generalize the process of adding one to the given element, which is how we usually write to describe any functions.

The example shows that we can generalize the process using algebraic notation, in which it is usually in the form of

$$x \mapsto f(x)$$

with the function f . It is equivalent to say that

$$\dots, 1 \mapsto f(1), 2 \mapsto f(2), \dots$$

but what's important is how explicit the mapping process is done. More examples could be viewed to familiarize with it.

Example. *To describe a function undergoes the process ‘multiplying the element by two and add one to it’, we may examine that*

$$\dots, 0 \mapsto 1, 1 \mapsto 3, 2 \mapsto 5, \dots$$

so that it is equivalent to write

$$x \mapsto 2x + 1$$

as a generalization of the function. It is equivalent to write $f(x) := 2x + 1$ to emphasize that we shall call the function f as a naming for the given process, as long as we can write

$$x \mapsto f(x), f(x) := 2x + 1$$

Essential Practice 1.1.1.1. *Examine the following functions from 0 to 3, and generalize the function using algebraic notation. You may either choose writing $x \mapsto \square$ directly or $x \mapsto f(x), f(x) := \square$.*

1. *Multiply the element by 3 and then add 2 to it.*
2. *Divide the element by 10 and then Subtract 7 from it.*
3. *Multiply the element by a and then add b to it.*
4. *Squaring the element.*
5. *Multiply 3 to the square of the element, an then add 5 to it.*
6. *Add 1 to the element first, then multiply the square of the result by 6, and then add 1 to it.*
7. *Subtract 8 from the element first, then divide the square of the result by 2, and then add 4 to it.*
8. *Subtract h from the element first, then multiply the square of the result by a , and then add k to it.*

It is notable that a function can only give one output for each input, which makes the next page a fruitful discussion.

1.1.2 Defining a function with variables

So far, we have learned how writing a generalization of a function is, and have we used algebraic notation to shorten the examination, one suggest we can always write algebraic notation for a function definition. In addition, we also find that a function cannot have more than one output, as we see functions as a pointing process from one element to another element. So we have the following definition:

Definition 1.3 (Function). A **function** f of x defines the pointing process $x \mapsto f(x)$ is a one-to-one pointing process, which can have only one output.

It is important to note that $f(x)$ is a function of x if and only if one x produce one $f(x)$. The x is called a *dummy variable*, which is a variable that can be changed all the time. For example, writing $f(y)$ or $f(z)$ still makes sense to say a function f , but not of x .

From now on, we can determine whether a given relation is a function or not.

Example. Given the relation $y = mx + c$. Since one x can produce only one y , y is a function of x ; on the other hand, we also see one y produces only one x , so x is also a function of y .

Example. Given the relation $y = x^2$. Since one x can produce only one y , y is a function of x ; however, one y may produce more than one x , say if $y = 4$ then x can be 2 or -2 , so x is not a function of y .

Essential Practice 1.1.2.1. Determine whether the following given relation between x and y is a function of one another or not. Provide counterexample if it is not a function.

1. $y = -x$;
2. $y = 4x + 3$;
3. $y = \frac{1}{x}$;
4. $y = \frac{x}{6}$;
5. $y^2 = x$;
6. $y^2 = x^2$;
7. $y^3 = 4x^2 - 3$.

1.1.3 Domain, Co-domain and Range

For explicit definition of a function, we need the following concepts to help with: *domain*, *co-domain* and *range*.

A **domain** is where the input comes from, which is usually half-customized and half-restricted. For example, $f(x) = \frac{1}{x}$ can have input of negative real numbers, positive real numbers, any complex numbers except 0. This means that 0 is naturally restricted by the operation of $\frac{1}{x}$, but other than 0, we can choose freely our input from all complex numbers. Thus, the largest domain of $f(x) = \frac{1}{x}$ is all complex numbers except 0. However, it is not saying that the domain of $f(x) = \frac{1}{x}$ must be all complex numbers except 0, we can still put restrictions on our own, what means by customize, like all real numbers except 0 or all positive real numbers except 0 as its domain, is still a possible choice. Hence, we shall usually talk about the *greatest possible domain of a function* if we need to find the natural restrictions, and the *domain of a function* if we are going to define our source of input.

Essential Practice 1.1.3.1. Find the greatest possible domain of the following functions if (i) the output is restricted to complex numbers and (ii) the output is restricted to be real numbers:

1. $f(x) := x$;
2. $f(x) := ax + b$;
3. $f(x) := \frac{a}{x}$;
4. $f(x) := x^2$;
5. $f(x) := a(x - h)^2 + k$;
6. $f(x) := \sqrt{x}$;
7. $f(x) := \sqrt[3]{x}$;
8. $f(x) := \frac{1}{\sqrt{x}}$;

A **co-domain** is where the output can go to. It is more likely a limitation of the output of the function so that we know where our target is. Similarly, we shall usually talk about the *greatest possible co-domain of the function* if we need to find the natural restrictions, and the *co-domain of the function* if we are going to define our target output.

Essential Practice 1.1.3.2. Find the greatest possible co-domain of the following functions if the input is unrestricted:

1. $f(x) := x;$
2. $f(x) := ax + b;$
3. $f(x) := \frac{a}{x};$
4. $f(x) := x^2;$
5. $f(x) := a(x - h)^2 + k;$
6. $f(x) := \sqrt{x};$
7. $f(x) := \sqrt[3]{x};$
8. $f(x) := \frac{1}{\sqrt{x}};$

With domain and co-domain, we can now define a function in a more explicit manner. Writing a function with where the inputs are in and where the outputs to go, we have a nice notation - an arrow \rightarrow emphasizing the direction from domain to co-domain. In general, we will write

$$f : D \rightarrow R$$

to specify the function f is a function goes from domain D to co-domain R . The formal way to define a function is like below:

$$f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto f(x)$$

which reads ‘a function f sending a real number x to a real number $f(x)$ ’.

Example. For a function sending a natural number n to a natural number 2^n , we may write its definition as

$$f : \mathbb{N} \rightarrow \mathbb{N}, n \mapsto 2^n$$

Example. For a function sending an integer n to a rational number 2^n , we may write its definition as

$$f : \mathbb{Z} \rightarrow \mathbb{Q}, n \mapsto 2^n$$

We shall see both example shows the same function process $f(n) := 2^n$ but different domain and codomain. Therefore, we acknowledge that although both functions are having the same process, they are indeed representing different things, which yields they are in fact different functions.

From this point of view, we can further examine the so-called **range** of a function, which is the exact target region of the function under the codomain. We shall define some set notation to present its meaning.

Definition 1.4 (Union and Intersection). *Let A and B be sets. The **union** of A and B is defined as*

$$A \cup B := \{x : x \in A \text{ or } x \in B\}$$

*while the **intersection** of A and B is defined as*

$$A \cap B := \{x : x \in A \text{ and } x \in B\}$$

In fact, we say union is the collection of the objects which are either in set A or set B , or simply say it is the joined set of two sets. We can take a look at the following examples:

Example. *Let $A = \{1, 2, 3, 4\}$, $B = \{3, 4, 5\}$, then*

$$A \cup B = \{1, 2, 3, 4, 5\}$$

Example. *Let $A = \{1, 3, 5, 7, 9, \dots\}$ be the set of all positive odd numbers, $B = \{2, 4, 6, 8, 10, \dots\}$ be the set of positive even numbers, then*

$$A \cup B = \{1, 2, 3, 4, 5, 6, \dots\}$$

which is the set of all positive numbers. Sometimes, we may write the set by specifying the property of the elements as following:

$$A \cup B = \{x : x \text{ is a positive integer}\}$$

For intersection, it is generally speaking the collection of repeated elements in both sets, or we can say the sharing elements. We can take a look at the following examples:

Example. *Let $A = \{1, 2, 3, 4\}$, $B = \{3, 4, 5\}$, then*

$$A \cap B = \{3, 4\}$$

Example. *Let $A = \{1, 3, 5, 7, 9, \dots\}$ be the set of all positive odd numbers, $B = \{2, 4, 6, 8, 10, \dots\}$ be the set of positive even numbers, then*

$$A \cap B = \emptyset$$

which is the set with no element, an empty set.

We may represent the two definitions by shading regions in a Venn-diagram. Suppose we are calling a set by enclosing the region by a circle, then we have the following representation.




Figure 1: Union(Left) and Intersection(Right) of two sets

Essential Practice 1.1.3.3. Let $A = \{1, 3, 4, 6, 7\}$, $B = \{3, 4, 5, 7, 8\}$, then find $A \cup B$ and $A \cap B$.

Essential Practice 1.1.3.4. Prove the following identities:

1. $A \cap (B \cup C) \equiv (A \cap B) \cup (A \cap C)$;
2. $A \cup (B \cap C) \equiv (A \cup B) \cap (A \cup C)$.

Definition 1.5 (Range). The **range** of a function $f : D \rightarrow R$, denoted by $\mathbf{Ran}(f)$, is the set $f(D) \cap R$, where $f(D) := \{f(x) : x \in D\}$.

Usually, a teacher in high school aims to discuss the range of a function rather than the co-domain of a function, as what he shall teach is the size of the output, but not where the output shall be in. However, the difference between the co-domain of a function and the range of a function is we can further define the concept of a well-defined function with the concept of range.

Example. For a function f defined as

$$f : \mathbb{N} \rightarrow \mathbb{N}, n \mapsto 2^n$$

The range of f , denoted by $\mathbf{Ran}(f)$, is the set of all possible outcomes of $f(n)$ with natural numbers n . That is, the set

$$\{2^n : n \in \mathbb{N}\} = \{2, 4, 8, 16, 32, \dots\}$$

Example. For a function f defined as

$$f : \mathbb{Z} \rightarrow \mathbb{Q}, n \mapsto 2^n$$

$\mathbf{Ran}(f)$ is the set of all possible outcomes of $f(n)$ with integers n . That is, the set

$$\{2^n : n \in \mathbb{Z}\} = \{\dots, \frac{1}{4}, \frac{1}{2}, 1, 2, 4, \dots\}$$

We must observe that the two ranges of the same function process f are different when their co-domains are different. This shows the importance of discussion of co-domain when we are defining ranges of functions.

Essential Practice 1.1.3.5. *Find the range of the following functions:*

1. $f : \mathbb{N} \rightarrow \mathbb{N}, n \mapsto n;$
2. $f : \mathbb{N} \rightarrow \mathbb{N}, n \mapsto 3^n;$
3. $f : \mathbb{N} \rightarrow \mathbb{N}, n \mapsto 3^n - n^3;$
4. $f : \mathbb{Z} \rightarrow \mathbb{Z}, n \mapsto n;$
5. $f : \mathbb{Z} \rightarrow \mathbb{Z}, n \mapsto n^2;$
6. $f : \mathbb{Z} \rightarrow \mathbb{Q}, n \mapsto 5^n;$
7. $f : \mathbb{Q} \rightarrow \mathbb{Q}, n \mapsto n;$
8. $f : \mathbb{Z} \rightarrow \mathbb{Q}, n \mapsto n/2;$
9. $f : \mathbb{N} \rightarrow \mathbb{Q}, n \mapsto n/10;$
10. $f : \mathbb{Z} \rightarrow \mathbb{R}, n \mapsto \pi n^2;$

Essential Practice 1.1.3.6. *Are the functions in previous practice all well-defined? Which of them are not?*

1.2 Graph of a function

Graphing has been an essential skill in understanding mathematical objects. We usually say it is a mathematical modeling technique. Through graphing, we can see the relationship between the input and output clearly, whether they are related, increasing or decreasing. It is also easier to draw conclusion to estimations by valid graphs. This section aims to build the concept of representing functions by graphs.

1.2.1 Blobs-and-arrows diagram for discrete functions

We shall first take a step backward to a simpler function - a *discrete function* with finite inputs. This will help the construction very much.

Definition 1.6 (Discrete functions). A **discrete function** is a function with direct indication of the function process for each element in domain. That is, for each element $x \in D$, the output $f(x)$ is assigned to co-domain directly.

A discrete function is usually with a discrete domain, and random assignment.

Example. Let $D := \{1, 2, 3, 4, 5\}$ be the domain of a function f . Suppose f is defined by

$$f(x) := \begin{cases} 1 & , x = 1 \\ 4 & , x = 2 \\ 3 & , x = 3 \\ 2 & , x = 4 \\ 5 & , x = 5 \end{cases}$$


In this case, f is a discrete function.

To represent the above function using a graph, it is recommended to use a *blobs-and-arrows* diagram.

Definition 1.7 (Blobs-and-arrows diagram). A **blobs-and-arrows** diagram is a diagram representing a function by denoting the domain and co-domain by two circles, and for each element in the circle of domain, a pointer arrow over-set with f pointing to one of the element in the co-domain, to emphasize the relation between the two elements are input and output pair.

It is undesired to read through such complicated definition of the diagram. Let's see the following example.


Example. Recall the function in previous example, we could draw the blobs-and-arrows diagram as shown.



In the figure above, the name of the function is hung over the main diagram, while the left ellipse denote the set of elements of domain D and the right ellipse denote the range of elements of codomain R . The letter R could also be interpreted as the range of f .

For a discrete function with finitely many inputs, where we will take them as a sequence, it can also be used to represent it efficiently. We just need to have some modification.

Axiom. Let $f : D \rightarrow R$ be a function. Let $a_1, a_2, a_3, \dots, a_n$ be a sequence of numbers in D and $b_1, b_2, b_3, \dots, b_n$ be a sequence of numbers in R . Suppose $b_1 = f(a_1), b_2 = f(a_2), \dots, b_n = f(a_n)$ defines the one-to-one correspondence from D to R . Then the following blobs-and-arrows diagram describes f formally:



Essential Practice 1.2.1.1. Draw the blobs-and-arrows diagram for the following discrete functions:

1. $f(1) = 3, f(2) = 6, f(3) = 7, f(4) = 2, f(5) = 3.$

$$2. f(x) := \begin{cases} 1 & , x = 1, 2, 3, 4, 5 \\ 2 & , x = 6, 7 \\ 3 & , x = 8, 9, 10 \\ 4 & , x = 11, 12, 13, 14, 15 \\ 5 & , x = 16, 17, 18, 19 \\ 6 & , x = 20 \end{cases}$$

1.2.2 The pairing table and xy-coordination

To extend our concept of function from discrete version to continuous version, that is, infinitely many numbers can be plugged into the function for calculation, we may need to ‘fill up’ the holes of the domain.

Let say we have 1 and 2 in the domain D , then all the real numbers within 1 and 2 also in D . This is the process of **continuation** or **extension**, and the whole set of real numbers within 1 and 2 is called the **interval** between 1 and 2. We can denote the interval by $[1, 2]$ if 1 and 2 are included in the meaning, or $(1, 2)$ if 1 and 2 are excluded from the meaning. To better distinct the difference between $[1, 2]$ and $(1, 2)$, we call the previous one the **closed interval** between 1 and 2, while the latter one the **open interval** between 1 and 2.

From this point of view, a real number line is used to represent the concept of any intervals. Let's define the real number line and discuss intervals using a number line.

Definition 1.8 (Real number line). *A **real number line** is a straight line with ordered property so that one of its direction is increasing, and the other direction is decreasing.*

We usually draw a horizontal real number line for convenience.

For the sake of simplicity and connected-understanding of open and closed intervals, we may put the desired parenthesis in the real number line to denote the meaning. Following that a closed interval is in a block-typed while open interval is in a round-typed, we can directly place them on the real number line.

Example. *The open interval $(0, 1)$.*

Example. *The open interval $(-1.5, 5.5)$.*

Example. *The closed interval $[0.5, 1]$.*

Example. *The closed interval $[-100, -67]$.*

Essential Practice 1.2.2.1. *Draw the mentioned interval on the provided real number line with suitable parenthesis.*

1. $[0, 1]$.
2. $[\alpha, \beta]$.
3. $(0, 1)$.
4. (α, β) .

It is of course we could draw it vertically with the same meaning, by indicating upward as positive direction and downward as negative direction. Consider a function $y = f(x)$ defines the relation between x and y coordinate, we can combine two real number lines in an orthogonal way, like a cross, to present the coordination in a sensible way. This is called the **continuous graph of a function**, or we simply call it the *graph of a function* if there's no ambiguity of what we are discussing.

In order to draw the graph of a function, we need to figure out the valid pairs of x and y first, so that the points could be addressed correctly for the function. We will make use of a **pairing table** so that each value of x in the domain is paired with a unique value of y in the co-domain. We will show its use with some examples.

Example. *Let G be the graph of $y = x + 1$. Then we have*

1.3 Fundamental properties of a graph

1.3.1 points lying on the graph

1.3.2 Special intersections: x-intercept and x-intercepts

1.3.3 general intersection

1.4 Solving equations using graphs of functions

1.4.1 Homogeneous equations

1.4.2 Non-homogeneous equations

1.4.3 Simultaneous equations

1.5 Transformation of functions

1.5.1 Translation

1.5.2 Dilation

1.6 Challenging questions

2 Linear functions

2.1 Fundamental concepts of points

2.1.1 Distance between two points

2.1.2 Mid-point and Division points

2.2 Different forms of a linear function

2.2.1 Slope of a line

2.2.2 Two-point form

2.2.3 Point-slope form

2.2.4 Slope-intercept form

2.2.5 General form

2.2.6 Special: Intercept-form

2.3 Parallel lines and Perpendicular lines

2.3.1 Point of intersection

2.3.2 Parallel lines

2.3.3 Perpendicular lines

2.3.4 Number of intersections

2.4 Angle of elevation and depression

2.5 Additional content: Point-line distance

By definition, the distance between two points on a \mathbb{R}^2 plane is

Definition 2.1 (Distance between two points). *Let $A(x_1, y_1)$ and $B(x_2, y_2)$ be two points on \mathbb{R}^2 -plane. The distance between A and B is computed by the formula*

$$\text{dist}(A, B) := \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$

Now let $L : ax + by + c = 0$ be a straight line and $P(x_0, y_0)$ be a point on \mathbb{R}^2 -plane. Note that with the following axiom

Axiom. *The distance between a point P and a line L is defined by the shortest distance between P and a point on L*

$$\text{dist}(P, L) := \inf\{\text{dist}(P, Q) : Q \in L\}$$

we can choose the perpendicular displacement of P from L to define the distance. Using the line Γ perpendicular to L passing through P , we can find such Q by following computation:

$$\begin{cases} L : ax + by + c = 0 \\ \Gamma : bx - ay + (ay_0 - bx_0) = 0 \end{cases} \implies Q\left(\frac{b^2x_0 - aby_0 - ac}{a^2 + b^2}, \frac{a^2y_0 - abx_0 - bc}{a^2 + b^2}\right)$$

Therefore, the distance between P and Q is

$$\begin{aligned} \text{dist}(P, Q) &= \frac{1}{a^2 + b^2} \sqrt{[(a^2 + b^2)x_0 - (b^2x_0 - aby_0 - ac)]^2 + [(a^2 + b^2)y_0 - (a^2y_0 - abx_0 - bc)]^2} \\ &= \frac{1}{a^2 + b^2} \sqrt{(a^2x_0 + aby_0 + ac)^2 + (b^2y_0 + abx_0 + bc)^2} \\ &= \frac{\sqrt{a^2 + b^2} \sqrt{(ax_0 + by_0 + c)^2}}{a^2 + b^2} \\ &= \frac{\sqrt{(ax_0 + by_0 + c)^2}}{\sqrt{a^2 + b^2}} \\ &= \left| \frac{ax_0 + by_0 + c}{\sqrt{a^2 + b^2}} \right| \end{aligned}$$

2.6 Linear inequalities

2.6.1 One variable inequality

2.6.2 Two variable inequality

2.6.3 Linear programming

2.7 Challenging questions

3 Quadratic functions

3.1 Solving Quadratic Equations

3.1.1 The square-root method

3.1.2 General form and Completing the square method

3.1.3 The quadratic formula

3.2 Solvability of Quadratic Equations

3.2.1 Discriminant: the factor affecting solvability

3.2.2 Application of solvability

3.3 Relation between coefficients and roots

3.3.1 What does it mean by a root?

3.3.2 The Vieta formula

3.3.3 Forming quadratic equations with given roots

3.4 Vertex form of a quadratic function

3.4.1 Obtaining vertex form using Transformation

3.4.2 Relation between vertex form and general form

3.4.3 Opening direction

3.4.4 The axis of symmetry

3.4.5 The extremum of quadratic functions

3.5 Quadratic inequalities

3.5.1 Boolean algebra

3.5.2 Solving quadratic inequalities

3.6 Challenging questions

4 Polynomial functions

4.1 Arithmetic rules for polynomials

4.1.1 Addition, Subtraction and Multiplication

4.1.2 Division

4.2 Divisibility of polynomials

4.2.1 Division algorithm

4.2.2 Remainder Theorem

4.2.3 Factor Theorem and General Vieta formula

4.3 G.C.D. and L.C.M.

4.3.1 Greatest common divisor

4.3.2 Least common multiple

4.4 Positional notation

4.4.1 Binary, Octal, Decimal and Hexadecimal

4.4.2 Conversion between different bases

4.5 Challenging questions

5 Exponential and Logarithmic functions

5.1 Exponential functions

5.1.1 Law of Exponential algebra

5.1.2 Seeing surds as fractional exponents

5.1.3 Solving equations using exponentiation method

5.2 Logarithmic functions

5.2.1 What is Logarithm?

5.2.2 Law of Logarithmic algebra

5.2.3 Solving equations using Logarithmic method

5.3 Challenging questions

6 Sequence as a function of natural numbers

6.1 What is a sequence?

6.2 Arithmetic sequence

6.2.1 General form

6.2.2 Summation of arithmetic sequence

6.3 Geometric sequence

6.3.1 General form

6.3.2 Summation on Geometric sequence

6.3.3 Infinite sum of Geometric sequence

6.4 Additional content: Arithmetic-Geometric sequence

6.5 Challenging questions